

IMA Journal of Numerical Analysis (2018) **00**, 1–24
doi:10.1093/imanum/dry033

Runge–Kutta convolution coercivity and its use for time-dependent boundary integral equations

LEHEL BANJAI*

*The Maxwell Institute for Mathematics in the Sciences, School of Mathematical & Computer Sciences,
Heriot-Watt University, EH14 4AS Edinburgh, UK*

*Corresponding author: l.banjai@hw.ac.uk

AND

CHRISTIAN LUBICH

*Mathematisches Institut, Universität Tübingen, Auf der Morgenstelle, D-72076 Tübingen, Germany
lubich@na.uni-tuebingen.de*

[Received on 27 February 2017; revised on 3 May 2018]

A coercivity property of temporal convolution operators is an essential tool in the analysis of time-dependent boundary integral equations and their space and time discretizations. It is known that this coercivity property is inherited by convolution quadrature time discretization based on A-stable multistep methods, which are of order at most 2. Here we study the question as to which Runge–Kutta-based convolution quadrature methods inherit the convolution coercivity property. It is shown that this holds without any restriction for the third-order Radau IIA method, and on permitting a shift in the Laplace domain variable, this holds for all algebraically stable Runge–Kutta methods and hence for methods of arbitrary order. As an illustration the discrete convolution coercivity is used to analyse the stability and convergence properties of the time discretization of a nonlinear boundary integral equation that originates from a nonlinear scattering problem for the linear wave equation. Numerical experiments illustrate the error behaviour of the Runge–Kutta convolution quadrature time discretization.

Keywords: Runge–Kutta convolution quadrature; coercivity; stability; boundary integral equation; wave equation.

1. Introduction

This paper is concerned with a discrete coercivity property that ensures the stability of time discretizations of boundary integral equations for wave equations, also in situations such as the following:

- nonlinear boundary integral equations; and
- boundary integral equations coupled with a wave equation in an interior domain, with an explicit time discretization in the domain.

For convolution quadrature based on A-stable multistep methods (which have approximation order at most 2) it is known from the study by [Banjai *et al.* \(2015\)](#) that the coercivity property is preserved under time discretization uniformly in the temporal step size. Here we study the preservation of convolution coercivity under time discretization by Runge–Kutta convolution quadrature. Up to a shift in the Laplace variable and a corresponding reformulation of the boundary integral equation for an exponentially scaled solution function, we show that the convolution coercivity property is preserved by all convolution

quadratures based on algebraically stable Runge–Kutta methods, which include in particular Radau IIA methods of arbitrary order. Without any such shift and exponential scaling the convolution coercivity is shown to be preserved by the two-stage Radau IIA method of order 3.

We illustrate the use of the discrete convolution coercivity by the stability and convergence analysis of the Runge–Kutta convolution quadrature time discretization of a nonlinear boundary integral equation for a nonlinear scattering problem for the acoustic wave equation. This problem has been studied with different numerical methods in the study by [Banjai & Rieder \(2018\)](#).

The discrete convolution coercivity is not needed for the corresponding linear scattering problem, because there the convolution quadrature time discretization of the linear boundary integral equation can be interpreted as a convolution quadrature discretization of the convolution operator that maps the data to the solution. Therefore, known bounds of the Laplace transform of the solution operator and known error bounds of convolution quadrature yield stability and error bounds ([Lubich, 1994](#); [Sayas, 2016](#)). The same argument can also be used for the coupling of a linear wave equation in an interior domain with the boundary integral equation that describes transparent boundary conditions, provided that the convolution quadrature for the boundary integral equation is based on the same (implicit) time discretization method as for the wave equation in the interior domain. This precludes explicit time-stepping in the interior. For the coupling of convolution quadrature on the boundary with an explicit time discretization in the interior, the discrete convolution coercivity as considered in the present paper is needed; see [Banjai *et al.* \(2015\)](#), [Kovács & Lubich \(2017\)](#) and [Eberle \(2018\)](#) for the coupling of implicit BDF2 convolution quadrature on the boundary with explicit leapfrog time-stepping in the domain for acoustic, elastic and electro-magnetic wave equations, respectively.

The paper is organised as follows:

In [Section 2](#) we recall the continuous-time convolution coercivity, which is related to a coercivity property of the Laplace transform of the (distributional) convolution kernel that holds uniformly for all values of the Laplace-domain frequency variable in a (possibly shifted) right half-plane.

In [Section 3](#) we study the preservation of the convolution coercivity under time discretization by Runge–Kutta convolution quadrature. This preservation depends on the numerical range of the Runge–Kutta differentiation symbol, which is shown to lie in the right half-plane for algebraically stable Runge–Kutta methods. With a matrix-function inequality that is obtained as an extension of a theorem of von Neumann, we then prove our main result, [Theorem 2](#), which yields the discrete convolution coercivity.

[Section 4](#) recapitulates error bounds of Runge–Kutta convolution quadrature shown in the study by [Banjai *et al.* \(2011\)](#).

In [Section 5](#) we apply our results to the time discretization of the wave equation with a nonlinear impedance boundary condition. We study only semidiscretization in time, but note that this could be extended to full discretization with the techniques of [Banjai & Rieder \(2018\)](#). The error behaviour is illustrated by numerical experiments in [Section 6](#). In the numerical experiments it is observed that the convolution quadrature based on the three-stage Radau IIA method performs well even without the shift and exponential scaling, which is more favourable than our theoretical results.

2. Coercivity of temporal convolutions

The following coercivity result is given in the study by [Banjai *et al.* \(2015\)](#), where it is used as a basic result in studying boundary integral operators for the acoustic wave equation; see also [Kovács & Lubich \(2017\)](#) for Maxwell's equation and [Eberle \(2018\)](#) for elastic wave equations. The result can be viewed as a time-continuous operator-valued extension of a theorem of Herglotz from 1911, which states that

an analytic function has positive real part on the unit disk if and only if convolution with its coefficient sequence is a positive semidefinite operation.

Let V be a complex Hilbert space and V' its dual, and let $\langle \cdot, \cdot \rangle$ denote the anti-duality between V and V' . Let $L(s) : V \rightarrow V'$ and $R(s) : V \rightarrow V$ be analytic families of bounded linear operators for $\operatorname{Re} s > \sigma$, continuous for $\operatorname{Re} s \geq \sigma$. We assume the uniform bounds, with some real exponent μ ,

$$\|L(s)\|_{V' \leftarrow V} \leq M(1 + |s|)^\mu, \quad \|R(s)\|_{V \leftarrow V} \leq M(1 + |s|)^\mu, \quad \operatorname{Re} s > \sigma. \quad (2.1)$$

This polynomial bound guarantees that L is the Laplace transform of a distribution ℓ . If we write $L(s) = s^k L_k(s)$ with an integer $k > \mu + 1$, then the Laplace inversion formula

$$\ell_k(t) = \frac{1}{2\pi i} \int_{\sigma' + i\mathbb{R}} e^{st} L_k(s) \, ds, \quad t \geq 0 \quad (\sigma' > \sigma)$$

defines a continuous and exponentially bounded function ℓ_k , which has ℓ as its k th distributional derivative. We write the convolution with ℓ as

$$u(t) = L(\partial_t)f(t) = (\ell * f)(t) = \left(\frac{d}{dt}\right)^k \int_0^t \ell_k(t - \tau)f(\tau) \, d\tau, \quad t > 0,$$

for functions f on $[0, T]$ whose extension to $t < 0$ by 0 is k times continuously differentiable. Similarly, we consider the convolution $R(\partial_t)f$.

THEOREM 2.1 (Banjai *et al.*, 2015, Lemma 2.2). Let $\alpha \geq 0$. In the above situation, the following statements are equivalent:

1. $\operatorname{Re} \langle v, L(s)v \rangle \geq \alpha \|R(s)v\|^2$ for all $v \in V$, $\operatorname{Re} s > \sigma$.
2. $\int_0^\infty e^{-2\sigma t} \operatorname{Re} \langle f(t), L(\partial_t)f(t) \rangle \, dt \geq \alpha \int_0^\infty e^{-2\sigma t} \|R(\partial_t)f(t)\|^2 \, dt$ for all $f \in C^k([0, \infty), V)$ with finite support and $f^{(j)}(0) = 0$ for $0 \leq j < k$, and for all $t \geq 0$.

Property 1 is known to be satisfied for the Laplace transforms of various boundary integral operators for wave equations (Bamberger & Ha-Duong, 1986; Laliena & Sayas, 2009; Banjai *et al.*, 2015; Sayas, 2016; Kovács & Lubich, 2017), and it is a fundamental property in the study of boundary integral equations for wave equations.

We are interested in time discretizations of the convolution operators $L(\partial_t)$ and $R(\partial_t)$ that preserve this coercivity property. It was shown in the study by Banjai *et al.* (2015) that this is achieved by convolution quadrature based on A-stable multistep methods such as the first- and second-order backward differentiation formulae. In Theorem 3.1 below we will show that the coercivity property is also preserved by convolution quadrature based on certain Runge–Kutta methods such as the third-order, two-stage Radau IIA method. For the particular case $\sigma = 0$, it will be shown to be preserved for all algebraically stable Runge–Kutta methods.

3. Preserving coercivity by Runge–Kutta convolution quadrature

3.1 Runge–Kutta differentiation symbol and convolution quadrature

An m -stage Runge–Kutta discretization of the initial value problem $y' = f(t, y)$, $y(0) = y_0$, is given by

$$Y_{ni} = y_n + \tau \sum_{j=1}^m a_{ij} f(t_n + c_j h, Y_{nj}), \quad i = 1, \dots, m,$$

$$y_{n+1} = y_n + \tau \sum_{j=1}^m b_j f(t_n + c_j h, Y_{nj}),$$

where $\tau > 0$ is the time step, $t_n = n\tau$, and the internal stages Y_{ni} and grid values y_n are approximations to $y(t_n + c_i\tau)$ and $y(t_n)$, respectively. In the following we use the notation

$$\mathcal{A} = (a_{ij})_{i,j=1}^m, \quad b = (b_1, \dots, b_m)^T, \quad \mathbb{1} = (1, 1, \dots, 1)^T.$$

We always assume that the Runge–Kutta matrix \mathcal{A} is invertible.

As has been shown in the studies by Lubich & Ostermann (1993), Schädle *et al.* (2006), Banjai & Lubich (2011) and Banjai *et al.* (2011), and in applications to wave propagation problems further in the studies by Wang & Weile (2011), Banjai *et al.* (2012), Banjai & Kachanovska (2014) and Banjai & Rieder (2018), Runge–Kutta methods can be used to construct convolution quadrature methods that enjoy favourable properties. Here one uses the *Runge–Kutta differentiation symbol*

$$\Delta(\zeta) = \left(\mathcal{A} + \frac{\zeta}{1 - \zeta} \mathbb{1} b^T \right)^{-1}, \quad \zeta \in \mathbb{C} \text{ with } |\zeta| < 1. \quad (3.1)$$

This is well defined for $|\zeta| < 1$ if $R(\infty) = 1 - b^T \mathcal{A}^{-1} \mathbb{1}$ satisfies $|R(\infty)| \leq 1$. In fact, the Sherman–Morrison formula then yields

$$\Delta(\zeta) = \mathcal{A}^{-1} - \frac{\zeta}{1 - R(\infty)\zeta} \mathcal{A}^{-1} \mathbb{1} b^T \mathcal{A}^{-1}.$$

To formulate the Runge–Kutta convolution quadrature for $L(\partial_t)g$, we formally replace in $L(s)$ the differentiation symbol s by $\Delta(\zeta)/\tau$ and expand the operator-valued matrix function

$$L\left(\frac{\Delta(\zeta)}{\tau}\right) = \sum_{n=0}^{\infty} \mathbf{W}_n(L) \zeta^n,$$

where in the case of $L(s) : V \rightarrow V'$ we have the convolution quadrature matrices $\mathbf{W}_n(L) : V^m \rightarrow (V')^m$. For the discrete convolution with these matrices we use the notation

$$(L(\partial_t^\tau)f)_n = \sum_{j=0}^n \mathbf{W}_{n-j}(L) f_j$$

for any sequence $\mathbf{f} = (f_n)$ in V^m . For vectors of function values of a function $f : [0, T] \rightarrow V$ given as $\mathbf{f}_n = (f(t_n + c_i\tau))_{i=1}^m$, the i th component of the vector $(L(\partial_i^\tau)\mathbf{f})_n$ is considered as an approximation to $(L(\partial_i)f)(t_n + c_i\tau)$.

In particular if $c_m = 1$, as is the case with Radau IIA methods, then the continuous convolution at t_{n+1} is approximated by the last component of the discrete block convolution:

$$(L(\partial_i)f)(t_{n+1}) \approx e_m^T (L(\partial_i^\tau)\mathbf{f})_n,$$

where $e_m = (0, \dots, 0, 1)^T$ is the m th unit vector. We recall the composition rule

$$L_2(\partial_i^\tau) L_1(\partial_i^\tau)\mathbf{f} = (L_2L_1)(\partial_i^\tau)\mathbf{f}.$$

For $\lambda \in \mathbb{C}$ the convolution quadrature $(\partial_i^\tau - \lambda)^{-1}\mathbf{f}$ (which is to be interpreted as $L(\partial_i^\tau)\mathbf{f}$ for the multiplication operator $L(s) = (s - \lambda)^{-1}$) contains the internal stages of the Runge–Kutta approximation to the linear differential equation $y' = \lambda y + f$ with initial value $y(0) = 0$.

Results on the order of convergence of this approximation are given in the studies by Lubich & Ostermann (1993), Banjai & Lubich (2011) and Banjai *et al.* (2011). The result of the study by Banjai *et al.* (2011), which is relevant for operators $L(s)$ arising in wave propagation, will be restated and extended to the internal stages in Section 4.

3.2 Numerical range of the Runge–Kutta differentiation symbol

We now consider methods that are *algebraically stable*:

- All weights b_i are positive.
- The symmetric matrix with entries $b_i a_{ij} + b_j a_{ji} - b_i b_j$ is positive semidefinite.

Gauss methods and Radau IIA methods are widely used classes of methods that satisfy this condition. We refer the reader to Hairer & Wanner (1996) for background literature on Runge–Kutta methods and their stability notions.

We consider the weighted inner product on \mathbb{C}^m ,

$$(u, v) = \sum_{i=1}^m b_i \bar{u}_i v_i, \quad u, v \in \mathbb{C}^m. \tag{3.2}$$

We have the following characterisation.

LEMMA 3.1 For an algebraically stable Runge–Kutta method and for the b -weighted inner product (3.2),

$$\operatorname{Re}(w, \Delta(\zeta)w) \geq 0 \quad \text{for all } w \in \mathbb{C}^m, |\zeta| < 1.$$

Conversely, if the differentiation symbol of a Runge–Kutta method with positive weights b_i satisfies this inequality, then the method is algebraically stable.

Proof. With a different notation, this is shown in [Lubich & Ostermann \(1987, p. 232, \(6.19\)\)](#). For the convenience of the reader we include the short proof. Since for $v = \Delta(\zeta)w$ we have $(w, \Delta(\zeta)w) = (\Delta(\zeta)^{-1}v, v)$, it suffices to show that

$$\operatorname{Re}(v, \Delta(\zeta)^{-1}v) \geq 0 \quad \text{for all } v \in \mathbb{C}^m, |\zeta| < 1.$$

We rewrite

$$\Delta(\zeta)^{-1} = \mathcal{A} + \frac{\zeta}{1-\zeta} \mathbb{1}b^T = \mathcal{C} + \frac{1}{2} \frac{1+\zeta}{1-\zeta} \mathbb{1}b^T, \quad \text{with } \mathcal{C} = \mathcal{A} - \frac{1}{2} \mathbb{1}b^T,$$

and observe that (cf. [Hairer & Lubich, 1984](#))

$$\begin{aligned} (\mathbb{1}b^T v, v) &= \left| \sum_{i=1}^m b_i v_i \right|^2 \\ 2 \operatorname{Re}(\mathcal{C}v, v) &= \sum_{i,j=1}^m (b_i a_{ij} + b_j a_{ji} - b_i b_j) \bar{v}_i v_j. \end{aligned}$$

Since $\operatorname{Re}(1+\zeta)/(1-\zeta) > 0$ for $|\zeta| < 1$ the result follows. \square

In this paper we will need a stronger positivity property, for which we show the following order barrier and a positive result for the two-stage Radau IIA method, which is of order 3 and has the coefficients

$$\mathcal{A} = \begin{pmatrix} 5/12 & -1/12 \\ 3/4 & 1/4 \end{pmatrix}, \quad b^T = (3/4, 1/4).$$

LEMMA 3.2 (a) For the two-stage Radau IIA method and for the b -weighted inner product (3.2) and corresponding norm $|\cdot|$ we have

$$\operatorname{Re}(w, \Delta(\zeta)w) \geq \frac{1}{2}(1 - |\zeta|^2)|w|^2 \quad \text{for all } w \in \mathbb{C}^m, |\zeta| \leq 1. \quad (3.3)$$

(b) For none of the Gauss methods with two or more stages and none of the Radau IIA methods with three or more stages, there exists $c > 0$ such that for all sufficiently small $\Delta > 0$,

$$\operatorname{Re}(w, \Delta(\zeta)w) \geq c\Delta |w|^2 \quad \text{for all } w \in \mathbb{C}^m, |\zeta| \leq e^{-\Delta}. \quad (3.4)$$

Clearly, (3.3) implies (3.4) with c arbitrarily close to 1 for small Δ . We further note that the implicit Euler method and the implicit midpoint rule (which are the one-stage Radau IIA and Gauss methods, respectively) also satisfy (3.4).

Proof.

(a) For the two-stage Radau IIA method we find

$$\Delta(\zeta) = \frac{1}{2} \begin{pmatrix} 3 & 1-4\zeta \\ -9 & 5+4\zeta \end{pmatrix}.$$

Denoting the diagonal matrix of the weights by $\mathcal{B} = \text{diag}(3/4, 1/4)$, we note

$$\text{Re}(w, \Delta(\zeta)w) = \bar{w}^T \mathcal{B}^{1/2} \cdot \mathcal{B}^{-1/2} \frac{1}{2} (\mathcal{B} \Delta(\zeta) + \Delta(\bar{\zeta})^T \mathcal{B}) \mathcal{B}^{-1/2} \cdot \mathcal{B}^{1/2} w.$$

We obtain the hermitian matrix

$$\mathcal{B}^{-1/2} \frac{1}{2} (\mathcal{B} \Delta(\zeta) + \Delta(\bar{\zeta})^T \mathcal{B}) \mathcal{B}^{-1/2} = \frac{1}{2} \begin{pmatrix} 3 & -\sqrt{3}(1+2\zeta) \\ -\sqrt{3}(1+2\bar{\zeta}) & 5+4\text{Re}\zeta \end{pmatrix},$$

which has the trace $4 + 2\text{Re}\zeta$ and the determinant $3(1 - |\zeta|^2)$. It follows that both eigenvalues are positive and bounded by 6, and hence the smaller eigenvalue is bounded from below by $3(1 - |\zeta|^2)/6 = (1 - |\zeta|^2)/2$. This yields the inequality (3.3).

- (b) The proof uses the W-transformation of Hairer & Wanner, see Hairer & Wanner (1996, p. 77). For each of the m -stage Gauss and Radau IIA methods, there exists an invertible real $m \times m$ matrix W with first column $\mathbb{1}$ such that, with the diagonal matrix \mathcal{B} of the weights b_i ,

$$W^T \mathcal{B} W = I_m$$

or in other words, $W^T \mathcal{B}^{1/2}$ is an orthogonal matrix (with respect to the Euclidean inner product), and

$$\mathcal{A} = XW^{-1},$$

where $X = \frac{1}{2}e_1 e_1^T - \beta_m e_m e_m^T$ is a skew-symmetric matrix with $\beta_m = 0$ for the Gauss method and $\beta_m > 0$ for the Radau IIA method. We write

$$\begin{aligned} \text{Re}(w, \Delta(\zeta)w) &= \text{Re} \bar{w}^T \mathcal{B} \Delta(\zeta) w \\ &= \text{Re} \bar{w}^T \mathcal{B}^{1/2} \cdot \mathcal{B}^{1/2} W \cdot W^T \mathcal{B}^{1/2} \cdot \mathcal{B}^{1/2} \Delta(\zeta) \mathcal{B}^{-1/2} \cdot \mathcal{B}^{1/2} W \cdot W^T \mathcal{B}^{1/2} \cdot \mathcal{B}^{1/2} w \\ &= \text{Re} \bar{w}^T \mathcal{B}^{1/2} \cdot \mathcal{B}^{1/2} W \cdot W^T \mathcal{B} \Delta(\zeta) W \cdot W^T \mathcal{B}^{1/2} \cdot \mathcal{B}^{1/2} w, \end{aligned}$$

where we note

$$W^T \mathcal{B} \Delta(\zeta) W = W^{-1} \Delta(\zeta) W = (W^{-1} \Delta(\zeta)^{-1} W)^{-1}.$$

Now, by the definition of $\Delta(\zeta)$ and the above-mentioned property of $W^{-1} \mathcal{A} W = X$ together with $W^T b = e_1$, the matrix $W^{-1} \Delta(\zeta)^{-1} W$ is the sum of a skew-hermitian matrix plus a rank 1 or rank 2 matrix for Gauss or Radau IIA methods, respectively, and by the Sherman–Morrison–Woodbury formula so is its inverse:

$$W^T \mathcal{B} \Delta(\zeta) W = Y + Z(\zeta),$$

where Y is skew-hermitian and $Z(\zeta)$ is of rank 1 or 2 for Gauss or Radau IIA methods, respectively. If $w \neq 0$ is in the null space of $Z(\zeta)W^T \mathcal{B}$, which is of codimension 1 or 2 for Gauss or Radau, respectively, then we obtain from the above formulas that

$$\text{Re}(w, \Delta(\zeta)w) = 0$$

in contradiction to (3.4). □

As we will show in Theorem 3.1 below, Runge–Kutta convolution quadrature with (3.4) preserves the coercivity property of Theorem 2.1 for arbitrary abscissa $\sigma \geq 0$, while general algebraically stable methods preserve it in the case $\sigma = 0$. Before we state and prove this theorem in Section 3.4, we need an auxiliary result of independent interest.

3.3 A matrix-function inequality related to a theorem by von Neumann

We consider again a complex Hilbert space V and its dual V' , with the anti-duality denoted by $\langle \cdot, \cdot \rangle$. On \mathbb{C}^m we consider an inner product (\cdot, \cdot) and associated norm $|\cdot|$. An inner product on V^m and the anti-duality between V^m and $(V')^m$ are induced in the usual way: for Kronecker products $a \otimes u$ and $b \otimes v$ with $a, b \in \mathbb{C}^m$ and $u, v \in V$ one defines $(a \otimes u, b \otimes v) = (a, b)(u, v)$ and extends this to a sesquilinear form on $V^m \times V^m$, and in the same way one proceeds for the anti-duality $\langle \cdot, \cdot \rangle$ on $V^m \times (V')^m$.

LEMMA 3.3 On the Hilbert space V , let $L(s) : V \rightarrow V'$ and $R(s) : V \rightarrow V$ be analytic families of bounded linear operators for $\operatorname{Re} s > \sigma$, continuous for $\operatorname{Re} s \geq \sigma$, such that (2.1) is satisfied and for some $\alpha \geq 0$,

$$\operatorname{Re} \langle v, L(s)v \rangle \geq \alpha \|R(s)v\|^2, \quad \text{for all } v \in V, \operatorname{Re} s \geq \sigma.$$

Let the matrix $S \in \mathbb{C}^{m \times m}$ be such that

$$\operatorname{Re} (w, Sw) \geq \sigma |w|^2, \quad \text{for all } w \in \mathbb{C}^m.$$

Then,

$$\operatorname{Re} \langle v, L(S)v \rangle \geq \alpha \|R(S)v\|^2, \quad \text{for all } v \in V^m.$$

This result can be viewed as an extension of a theorem of von Neumann (1951) (see also Hairer & Wanner (1996, p. 179)), which corresponds to the particular case where $L(s)$ is the identity operator on V (when V is identified with V' with the anti-duality given by the inner product on V).

Proof. The proof adapts Crouzeix's proof of von Neumann's theorem as given in Hairer & Wanner (1996, p. 179 f.). Without loss of generality we assume here $\sigma = 0$.

First we note that for a *diagonal* matrix S the result holds trivially, and so it does for a *normal* matrix S , which is diagonalised by a similarity transformation with a unitary matrix.

For a nonnormal matrix S we consider the matrix-valued complex function

$$S(z) = \frac{z}{2}(S + S^*) + \frac{1}{2}(S - S^*)$$

and we observe that $S = S(1)$ and

$$\operatorname{Re} (w, S(z)w) = (\operatorname{Re} z) \operatorname{Re} (w, Sw).$$

Together with the condition on S this shows that the numerical range of $S(z)$ is in the right complex half-plane for $\operatorname{Re} z \geq 0$, and hence all eigenvalues of $S(z)$ have non-negative real part. Therefore, the operator functions $L(S(z))$ and $R(S(z))$ are well defined for $\operatorname{Re} z \geq 0$.

If $\operatorname{Re} z = 0$, then the matrix $S(z)$ is normal, and hence the desired inequality is valid for $S(z)$ with $\operatorname{Re} z = 0$. The function

$$\varphi(z) = \alpha \|R(S(z))v\|^2 - \operatorname{Re} \langle v, L(S(z))v \rangle$$

is subharmonic, since the last term is harmonic as the real part of an analytic function and the first term is the inner product of an analytic function with itself, which is subharmonic (as is readily seen by computing the Laplacian and noting that the real and imaginary parts of the analytic function are harmonic). Hence, by the maximum principle (or its Phragmén–Lindelöf-type extension to polynomially bounded subharmonic functions on the half-plane),

$$\varphi(1) \leq \sup_{\operatorname{Re} z=0} \varphi(z) \leq 0,$$

which is the desired inequality. □

3.4 Preserving the convolution coercivity under discretization

We are now in the position to formulate the main result of this paper and give its proof, which relies on the auxiliary results of the previous subsections.

THEOREM 3.1 Let the m -stage Runge–Kutta method satisfy (3.4) for some inner product (\cdot, \cdot) , as in particular is the case for the two-stage Radau IIA method. In the situation of Theorem 2.1, Condition 1 of that theorem implies, for sufficiently small step size $\tau > 0$ and with $\tilde{\sigma} = \sigma/c$,

$$\tau \sum_{n=0}^{\infty} e^{-2\tilde{\sigma}n\tau} \operatorname{Re} \langle f_n, (L(\partial_i^\tau) f)_n \rangle \geq \alpha \tau \sum_{n=0}^{\infty} e^{-2\tilde{\sigma}n\tau} \|(R(\partial_i^\tau) f)_n\|^2,$$

for every sequence $f = (f_n)_{n \geq 0}$ in V^m with finitely many nonzero entries. Moreover, in the case $\sigma = 0$ this inequality holds for every algebraically stable Runge–Kutta method, with $\tilde{\sigma} = 0$ and with respect to the b -weighted inner product (2) on \mathbb{C}^m .

Proof. The proof uses Parseval’s formula and combines Lemma 3.3 with Lemmas 3.1 and 3.2. By (3.4), with $\tilde{\sigma} = \sigma/c$ and $\rho = e^{-\tilde{\sigma}\tau}$ we have with respect to the inner product weighted by the b_i that

$$\operatorname{Re} \left(w, \frac{\Delta(\rho e^{i\theta})}{\tau} w \right) \geq \sigma |w|^2 \quad \text{for all } w \in \mathbb{C}^m, \theta \in \mathbb{R}. \tag{3.5}$$

We abbreviate

$$\widehat{L}(\theta) = L \left(\frac{\Delta(\rho e^{i\theta})}{\tau} \right)$$

and similarly $\widehat{R}(\theta)$. We denote the Fourier series

$$\widehat{f}(\theta) = \sum_{n=0}^{\infty} \rho^n e^{in\theta} f_n.$$

By Parseval’s formula and the definition of the convolution quadrature weights $W_n(L)$,

$$\sum_{n=0}^{\infty} \operatorname{Re} \left\langle \rho^n f_n, \sum_{j=0}^n \rho^{n-j} W_{n-j}(L) \rho^j f_j \right\rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \langle \widehat{f}(\theta), \widehat{L}(\theta) \widehat{f}(\theta) \rangle d\theta.$$

Here (3.5) used in Lemma 3.3 yields

$$\operatorname{Re} \langle \widehat{f}(\theta), \widehat{L}(\theta) \widehat{f}(\theta) \rangle \geq \alpha \|\widehat{R}(\theta) \widehat{f}(\theta)\|^2.$$

Moreover, again by Parseval's formula,

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \|\widehat{R}(\theta) \widehat{f}(\theta)\|^2 d\theta = \sum_{n=0}^{\infty} \rho^{2n} \left\| \sum_{j=0}^n W_{n-j}(R) f_j \right\|^2,$$

which yields the result. \square

4. Error bounds of Runge–Kutta convolution quadrature

In this section we restate the result of the study by Banjai *et al.* (2011) and extend it to cover the approximation properties of the internal stages, which will be needed in the next section. To avoid restating the list of properties required for the underlying Runge–Kutta method we state the results just for the Radau IIA methods, which appear to be the most important class of Runge–Kutta methods to be used for convolution quadrature.

Let $K(s)$, for $\operatorname{Re} s > \sigma > 0$, be an analytic family of operators between Hilbert spaces V and W (or just Banach spaces are sufficient here), such that for some real exponent μ and $\nu \geq 0$ the operator norm is bounded as follows:

$$\|K(s)\| \leq M(\sigma) \frac{|s|^\mu}{(\operatorname{Re} s)^\nu} \quad \text{for all } \operatorname{Re} s > \sigma. \quad (4.1)$$

THEOREM 4.1 (Banjai *et al.*, 2011, Theorem 3). Let K satisfy (4.1) and consider the Runge–Kutta convolution quadrature based on the Radau IIA method with m stages. Let $r > \max(2m - 1 + \mu, 2m - 1, m + 1)$ and $f \in C^r([0, T], V)$ satisfy $f(0) = f'(0) = \dots = f^{(r-1)}(0) = 0$. Then, there exists $\bar{\tau} > 0$ such that for $0 < \tau \leq \bar{\tau}$ and $t_n = n\tau \in [0, T]$,

$$\begin{aligned} & \left\| e_m^T (K(\partial_t^\tau) f)_n - (K(\partial_t) f)(t_{n+1}) \right\| \\ & \leq C h^{\min(2m-1, m+1-\mu+\nu)} \left(\|f^{(r)}(0)\| + \int_0^t \|f^{(r+1)}(\tau)\| d\tau \right). \end{aligned}$$

The constant C is independent of τ and f , but does depend on $\bar{\tau}$, T , and the constants in (4.1).

The proof in the study by Banjai *et al.* (2011) is readily extended to yield the following error bound for the internal stages. Note that here the full order $2m - 1$ is replaced by the stage order plus one, $m + 1$. We give the result for $m \geq 2$ stages, so that $m + 1 \leq 2m - 1$. (For $m = 1$, the implicit Euler method, one can use the previous result.)

THEOREM 4.2 Let K satisfy (4.1) and consider the Runge–Kutta convolution quadrature based on the Radau IIA method with $m \geq 2$ stages. Let $r > \max(m + 1 + \mu, m + 1)$ and $f \in C^r([0, T], V)$ satisfy

$f(0) = f'(0) = \dots = f^{(r-1)}(0) = 0$. Then, there exists $\bar{\tau} > 0$ such that for $0 < \tau \leq \bar{\tau}$ and $t_n = n\tau \in [0, T]$,

$$\begin{aligned} & \| (K(\partial_t^\tau) f)_n - (K(\partial_t) f(t_n + c_i \tau))_{i=1}^m \| \\ & \leq C h^{\min(m+1, m+1-\mu+v)} \left(\|f^{(r)}(0)\| + \int_0^t \|f^{(r+1)}(\tau)\| d\tau \right). \end{aligned}$$

The constant C is independent of τ and f , but does depend on $\bar{\tau}$, T , and the constants in (4.1).

5. Application to the time discretization of the wave equation with a nonlinear impedance boundary condition

5.1 A nonlinear scattering problem

We consider the wave equation on an exterior smooth domain $\Omega^+ \subset \mathbb{R}^3$. Following the study by [Banjai & Rieder \(2018\)](#) we search for a function $u(\cdot, t) \in H^1(\Omega^+)$ satisfying the weak form of the wave equation

$$\ddot{u} = \Delta u \quad \text{in } \Omega^+ \tag{5.1}$$

with zero initial conditions and with the nonlinear boundary condition

$$\partial_\nu^+ u = g(\dot{u} + \dot{u}^{\text{inc}}) - \partial_\nu^+ u^{\text{inc}} \quad \text{on } \Gamma, \tag{5.2}$$

where ∂_ν^+ is the outer normal derivative on the boundary Γ of Ω^+ , where $g : \mathbb{R} \rightarrow \mathbb{R}$ is a given monotonically increasing function and where $u^{\text{inc}}(x, t)$ is a given solution of the wave equation. The interpretation is that the total wave $u^{\text{tot}} = u + u^{\text{inc}}$ is composed of the incident wave u^{inc} and the unknown scattered wave u . The nonlinear boundary condition is a simplification of nonlinear acoustic boundary conditions as investigated in the study by [Graber \(2012\)](#) and of boundary conditions in electromagnetism obtained by asymptotic approximations of thin layers of nonlinear materials, see [Haddar & Joly \(2002\)](#). Though simpler, the nonlinear boundary condition (5.2) is sufficiently complex that the analysis requires the convolution quadrature coercivity preservation property.

One approach to solve this exterior problem is to determine the Dirichlet and boundary data from boundary integral equations on Γ and then to compute the solution at points of interest $x \in \Omega^+$ from the Kirchhoff representation formula. Here we are interested in the stability and convergence properties of the numerical approximation when the time discretization in the boundary integral equation and in the representation formula is done by Runge–Kutta convolution quadrature. Since our interest in this paper is the aspect of time discretization, we will not address the space discretization by boundary elements, though with some effort this could also be included; cf. [Banjai & Rieder \(2018\)](#).

5.2 Boundary integral equation and representation formula

Using the standard notation of the boundary integral operators for the Helmholtz equation $s^2 u - \Delta u = 0$ ($\text{Re } s > 0$) as used, for example, in the works by [Laliena & Sayas \(2009\)](#), [Banjai et al. \(2015\)](#), [Sayas \(2016\)](#) and [Banjai & Rieder \(2018\)](#), we denote by

$$S(s) : H^{-1/2}(\Gamma) \rightarrow H^1(\mathbb{R}^3 \setminus \Gamma) \quad \text{and} \quad D(s) : H^{1/2}(\Gamma) \rightarrow H^1(\mathbb{R}^3 \setminus \Gamma)$$

the single-layer and double-layer potential operators, respectively, and by $V(s)$, $K(s)$, $K^T(s)$, $W(s)$ the corresponding boundary integral operators that form the Calderón operator

$$B(s) = \begin{pmatrix} sV(s) & K(s) \\ -K^T(s) & s^{-1}W(s) \end{pmatrix} : H^{-1/2}(\Gamma) \times H^{1/2}(\Gamma) \rightarrow H^{1/2}(\Gamma) \times H^{-1/2}(\Gamma) \quad (5.3)$$

and the related operator

$$B_{\text{imp}}(s) = B(s) + \begin{pmatrix} 0 & -\frac{1}{2}I \\ \frac{1}{2}I & 0 \end{pmatrix}, \quad (5.4)$$

where the suffix imp stands for impedance. With these operators the solution u is determined by first solving, for

$$\varphi = -\partial_v^+ u \quad \text{and} \quad \psi = \gamma^+ u$$

(where γ^+ is the trace operator on Ω^+), the time-dependent boundary integral equation (see [Banjai & Rieder, 2018](#))

$$B_{\text{imp}}(\partial_t) \begin{pmatrix} \varphi \\ \psi \end{pmatrix} + \begin{pmatrix} 0 \\ g(\psi + \dot{u}^{\text{inc}}) \end{pmatrix} = \begin{pmatrix} 0 \\ \partial_v^+ u^{\text{inc}} \end{pmatrix}. \quad (5.5)$$

The solution u is then obtained from the representation formula

$$u = S(\partial_t)\varphi + D(\partial_t)\partial_t^{-1}\psi. \quad (5.6)$$

We will address the question as to what are the approximation properties when the temporal convolutions in (5.5) and (5.6) are discretised by Runge–Kutta convolution quadrature. Since this will not turn out fully satisfactory, we will further consider time-differentiated versions of (5.5).

The following coercivity property was proved in the study by [Banjai et al. \(2015\)](#).

LEMMA 5.1 ([Banjai et al., 2015](#), Lemma 3.1). Let $\langle \cdot, \cdot \rangle_\Gamma$ denote the anti-duality pairing between $H^{-1/2}(\Gamma) \times H^{1/2}(\Gamma)$ and $H^{1/2}(\Gamma) \times H^{-1/2}(\Gamma)$. There exists $\beta > 0$ so that the Calderón operator (5.3) satisfies

$$\operatorname{Re} \left\langle \begin{pmatrix} \varphi \\ \psi \end{pmatrix}, B(s) \begin{pmatrix} \varphi \\ \psi \end{pmatrix} \right\rangle_\Gamma \geq \beta c_\sigma \left(\|s^{-1}\varphi\|_{H^{-1/2}(\Gamma)}^2 + \|s^{-1}\psi\|_{H^{1/2}(\Gamma)}^2 \right)$$

for $\operatorname{Re} s \geq \sigma > 0$ and for all $\varphi \in H^{-1/2}(\Gamma)$ and $\psi \in H^{1/2}(\Gamma)$, with $c_\sigma = \min(1, \sigma^2) \sigma$.

Since $B_{\text{imp}}(s)$ differs from $B(s)$ by a skew-hermitian matrix the same estimate then also holds for $B_{\text{imp}}(s)$. Note that Lemma 5.1 yields property 1 of Theorem 2.1 for the Calderón operator $B(s)$ and for the multiplication operator $R(s) = s^{-1}$.

5.3 Time discretization by Runge–Kutta convolution quadrature

Using the notation of Section 3.1, the boundary integral equation (5.5) is discretised in time with a step size $\tau > 0$ over a time interval $(0, T)$ with $T = N\tau$ by

$$B_{\text{imp}}(\partial_t^\tau) \begin{pmatrix} \varphi^\tau \\ \psi^\tau \end{pmatrix} + \begin{pmatrix} 0 \\ g(\Psi^\tau + \dot{u}^{\text{inc}}) \end{pmatrix} = \begin{pmatrix} 0 \\ \partial_v^+ u^{\text{inc}} \end{pmatrix}, \quad (5.7)$$

where $(\varphi^\tau, \psi^\tau) = (\varphi_n, \psi_n)_{n=0}^{N-1}$ with $(\varphi_n, \psi_n) = (\varphi_{n,i}, \psi_{n,i})_{i=1}^m$ and $(\varphi_{n,i}, \psi_{n,i}) \approx (\varphi(t_n + c_i\tau), \Psi(t_n + c_i\tau))$ is the numerical approximation that is to be computed, and $\dot{\mathbf{u}}^{\text{inc}} = (\dot{\mathbf{u}}_n^{\text{inc}})_{n=0}^{N-1}$ with $\dot{\mathbf{u}}_n^{\text{inc}} = (\dot{u}_n^{\text{inc}}(t_n + c_i\tau))_{i=1}^m$. The function g acts componentwise. At the n th time step, a nonlinear system of equations of the following form needs to be solved:

$$B_{\text{imp}}\left(\frac{\Delta(0)}{\tau}\right) \begin{pmatrix} \varphi_n \\ \psi_n \end{pmatrix} + \begin{pmatrix} 0 \\ g(\psi_n + \dot{\mathbf{u}}_n^{\text{inc}}) \end{pmatrix} = \dots,$$

where the dots represent known terms. This has a unique solution because the eigenvalues of $\Delta(0) = \mathcal{A}^{-1}$ have positive real part, and Lemma 5.1 and the monotonicity of g then yield the unique existence of the solution by the Browder–Minty theorem; cf. Banjai & Rieder (2018) for the analogous situation for multistep-based convolution quadrature.

As an alternative to (5.7), we further consider the time discretization of the time-differentiated boundary integral equation:

$$B_{\text{imp}}(\partial_t^\tau) \begin{pmatrix} \dot{\varphi}^\tau \\ \dot{\psi}^\tau \end{pmatrix} + \begin{pmatrix} 0 \\ g'(\psi^\tau + \dot{\mathbf{u}}^{\text{inc}})(\dot{\psi}^\tau + \ddot{\mathbf{u}}^{\text{inc}}) \end{pmatrix} = \begin{pmatrix} 0 \\ \partial_v^+ \dot{\mathbf{u}}^{\text{inc}} \end{pmatrix}, \tag{5.8}$$

which is now solved for the approximations $(\dot{\varphi}^\tau, \dot{\psi}^\tau)$ (where the dot is just suggestive notation) to $(\dot{\varphi}, \dot{\psi})$ (where the dot means again time derivative). Here we define $\Psi^\tau = (\partial_t^\tau)^{-1} \dot{\psi}^\tau$ and the same for φ^τ . Furthermore, $\ddot{\mathbf{u}}^{\text{inc}}$ contains the values $\ddot{u}^{\text{inc}}(t_n + c_i\tau)$.

We can go even further and consider the time discretization of the twice-differentiated boundary integral equation:

$$B_{\text{imp}}(\partial_t^\tau) \begin{pmatrix} \ddot{\varphi}^\tau \\ \ddot{\psi}^\tau \end{pmatrix} + \begin{pmatrix} 0 \\ g'(\psi^\tau + \dot{\mathbf{u}}^{\text{inc}})(\ddot{\psi}^\tau + \ddot{\mathbf{u}}^{\text{inc}}) + g''(\psi^\tau + \dot{\mathbf{u}}^{\text{inc}}) \cdot (\dot{\psi}^\tau + \dot{\mathbf{u}}^{\text{inc}})^2 \end{pmatrix} = \begin{pmatrix} 0 \\ \partial_v^+ \ddot{\mathbf{u}}^{\text{inc}} \end{pmatrix}, \tag{5.9}$$

where again the dots on the approximation $(\ddot{\varphi}^\tau, \ddot{\psi}^\tau)$ are suggestive notation, and we set $\ddot{\Psi}^\tau = (\partial_t^\tau)^{-1} \ddot{\psi}^\tau$ and $\Psi^\tau = (\partial_t^\tau)^{-2} \ddot{\psi}^\tau$, and the same for φ^τ .

Finally, at any point $x \in \Omega^+$ of interest we compute the approximation to the solution value $u(x, t_n + c_i\tau)$ by using the same Runge–Kutta convolution quadrature for discretizing the representation formula (5.6):

$$\mathbf{u}^\tau = S(\partial_t^\tau) \varphi^\tau + D(\partial_t^\tau) (\partial_t^\tau)^{-1} \Psi^\tau. \tag{5.10}$$

5.4 Error bounds for the linear case

We consider first the case of a linear impedance function

$$g(\xi) = \alpha\xi \quad \text{with} \quad \alpha \geq 0.$$

Let $\mathbf{u}^\tau = (\mathbf{u}_n)_{n=0}^{N-1}$ with $\mathbf{u}_n = (u_{n,i})_{i=1}^m$ be the solution approximation obtained by the discretised representation formula (5.10) with either of the discretised boundary integral equations (5.7), (5.8) or (5.9). The discretization is done by Runge–Kutta convolution quadrature based on the Radau IIA method with m stages.

Here we obtain the following optimal-order pointwise error bounds for x bounded away from Γ .

PROPOSITION 5.1 Suppose that in a neighbourhood of the boundary Γ , the incident wave u_{inc} together with its extension by 0 to $t < 0$ is sufficiently regular. For $x \in \Omega^+$ with $\text{dist}(x, \Gamma) \geq \Delta > 0$, the following optimal-order error bound is satisfied in the linear situation described above: for $0 \leq t_n = n\tau \leq T$,

$$|u_n(x) - u(x, t_n)| \leq C(\Delta, T) \tau^{2m-1}.$$

Proof. We denote

$$B_\alpha(s) = B_{\text{imp}}(s) + \begin{pmatrix} 0 & 0 \\ 0 & \alpha I \end{pmatrix}.$$

By Lemma 5.1 $B_\alpha(s)$ is invertible for $\alpha \geq 0$ with the bound, for $\text{Re } s \geq \sigma > 0$,

$$\|B_\alpha(s)^{-1}\| \leq C(\sigma) \frac{|s|^2}{\text{Re } s}. \quad (5.11)$$

The exact solution $u(x, t)$ is given by the representation formula (5.6) with

$$\begin{pmatrix} \varphi \\ \Psi \end{pmatrix} = B_\alpha^{-1}(\partial_t) \begin{pmatrix} 0 \\ \partial_v^+ u^{\text{inc}} - \alpha \dot{u}^{\text{inc}} \end{pmatrix}.$$

For $x \in \Omega^+$ we define the operators $S_x(s) : H^{-1/2}(\Gamma) \rightarrow \mathbb{C}$ and $D_x(s) : H^{1/2}(\Gamma) \rightarrow \mathbb{C}$ by

$$S_x(s)\varphi = (S(s)\varphi)(x) \quad \text{and} \quad D_x(s)\Psi = (D(s)\Psi)(x).$$

These operators are bounded for $\text{Re } s \geq \sigma > 0$ and $\text{dist}(x, \Gamma) \geq \Delta > 0$ by

$$\|S_x(s)\|_{\mathbb{C} \leftarrow H^{-1/2}(\Gamma)} \leq C(\sigma, \Delta) |s| e^{-\Delta \text{Re } s} \quad (5.12)$$

$$\|D_x(s)\|_{\mathbb{C} \leftarrow H^{1/2}(\Gamma)} \leq C(\sigma, \Delta) |s| e^{-\Delta \text{Re } s}. \quad (5.13)$$

The first bound is proved in the study by [Banjai et al. \(2011, Lemma 6\)](#) and the second bound is proved similarly.

We thus have

$$u(x, t) = (M_x(\partial_t)f)(t)$$

with

$$M_x(s) = (S_x(s), D_x(s)s^{-1})B_\alpha(s)^{-1} \quad \text{and} \quad f = \begin{pmatrix} 0 \\ \partial_v^+ u^{\text{inc}} - \alpha \dot{u}^{\text{inc}} \end{pmatrix}.$$

With the above operator bounds we obtain for $\text{Re } s \geq \sigma > 0$ and $\text{dist}(x, \Gamma) \geq \Delta > 0$

$$\|M_x(s)\|_{\mathbb{C} \leftarrow H^{1/2}(\Gamma) \times H^{-1/2}(\Gamma)} \leq C(\sigma, \Delta) \frac{|s|^3}{\text{Re } s} e^{-\Delta \text{Re } s}. \quad (5.14)$$

By the composition rule the numerical solution obtained by (5.7) and (5.10) is given as

$$\mathbf{u}^\tau(x) = M_x (\partial_t^\tau) \mathbf{f},$$

where \mathbf{f} contains the values of f at the points $t_n + c_i\tau$. If we take instead (5.8) or (5.9), then we have

$$\mathbf{u}^\tau(x) = M_x (\partial_t^\tau) (\partial_t^\tau)^{-1} \dot{\mathbf{f}} \quad \text{or} \quad \mathbf{u}^\tau(x) = M_x (\partial_t^\tau) (\partial_t^\tau)^{-2} \ddot{\mathbf{f}},$$

respectively. In view of (5.14) Theorem 4.1 then yields the result. □

The situation is different if we consider the $H^1(\Omega^+)$ norm of the error.

PROPOSITION 5.2 Suppose that in a neighbourhood of the boundary Γ , the incident wave u_{inc} together with its extension by 0 to $t < 0$ is sufficiently regular. Then the following error bounds are satisfied in the linear situation described above: for $0 \leq t_n = n\tau \leq T$,

$$\|u_n - u(\cdot, t_n)\|_{H^1(\Omega^+)} \leq C(T) \tau^k$$

with

$k = m + 1/2$	if (5.7) is used,
$k = \min(2m - 1, m + 3/2)$	if (5.8) is used,
$k = \min(2m - 1, m + 5/2)$	if (5.9) is used.

Proof. Consider the Laplace transformed wave equation (5.1)

$$\begin{aligned} -\Delta \hat{u} + s^2 \hat{u} &= 0 \quad \text{in } \Omega^+, \\ \partial_\nu^+ \hat{u} - \alpha s \hat{u} &= \hat{f} \quad \text{on } \Gamma, \end{aligned} \tag{5.15}$$

where \hat{u} is the Laplace transform of u and \hat{f} the Laplace transform of

$$f = \partial_\nu^+ u^{inc} - \alpha u^{inc}.$$

We will require the estimate, see [Bamberger & Ha-Duong \(1986, Equation \(2.9\)\)](#),

$$\|\partial_\nu^+ \hat{u}\|_{H^{-1/2}(\Gamma)} \leq C(\sigma) |s|^{1/2} \|\hat{u}\|_{|s|, \Omega^+}, \tag{5.16}$$

with $\text{Re } s \geq \sigma > 0$ and the scaled H^1 norm

$$\|\hat{u}\|_{|s|, \Omega^+}^2 = \|\nabla \hat{u}\|_{L^2(\Omega^+)}^2 + |s|^2 \|\hat{u}\|_{L^2(\Omega^+)}^2.$$

Testing (5.15) with $s\hat{u}$ integrating by parts and taking the real part gives

$$\operatorname{Re} s \|\hat{u}\|_{|s|, \Omega^+}^2 = -\operatorname{Re} \langle \partial_v^+ \hat{u}, s\gamma^+ \hat{u} \rangle_\Gamma \leq C(\sigma) |s|^{1/2} \|\hat{u}\|_{|s|, \Omega^+} \|\hat{\Psi}\|_{H^{1/2}(\Gamma)},$$

where $\hat{\Psi} = s\gamma^+ \hat{u}$ is the Laplace transform of Ψ . Making use of $\|\hat{u}\|_{H^1(\Omega^+)} \leq C(\sigma) \|\hat{u}\|_{|s|, \Omega^+}$ and the bound (5.11) gives

$$\|\hat{u}\|_{H^1(\Omega^+)} \leq C(\sigma) \frac{|s|^{5/2}}{(\operatorname{Re} s)^2} \|\hat{f}\|_{H^{-1/2}(\Gamma)}.$$

The stated result then follows from Theorem 4.1. □

5.5 Convergence for the nonlinear problem

There are several aspects that make the error analysis of the nonlinear problem more intricate:

- The numerical solution can no longer be interpreted as a mere convolution quadrature for an appropriate operator $K(s)$ acting on the data (i.e., the incident wave).
- We need to impose regularity assumptions on the solution rather than the data.
- Convolution coercivity now plays an important role in ensuring the stability of the time discretization.

We assume strict monotonicity of the nonlinear function $g : \mathbb{R} \rightarrow \mathbb{R}$: there exists $\beta > 0$ such that

$$(\xi - \eta)(g(\xi) - g(\eta)) \geq \beta |\xi - \eta|^2 \quad \text{for all } \xi, \eta \in \mathbb{R}. \tag{5.17}$$

Furthermore, we assume that the pointwise application of g maps $H^{1/2}(\Gamma)$ to $H^{-1/2}(\Gamma)$. As is shown in the study by [Banjai & Rieder \(2018\)](#) by Sobolev embeddings, this is satisfied if $g(\xi)$ grows at most cubically as $|\xi| \rightarrow \infty$. Similar assumptions on the nonlinearity are also made in the study by [Graber \(2012\)](#), which investigates nonlinear acoustic boundary conditions.

In the following we write for a step size $\tau > 0$ and a sequence $\mathbf{e} = (\mathbf{e}_n)_{n=0}^{N-1}$ with $\mathbf{e}_n = (e_{n,i})_{i=1}^m$ and $e_{n,i}$ in a Hilbert space V

$$\|\mathbf{e}\|_{\ell_2^{\tau}(0:N;V^m)} = \tau \sum_{n=0}^{N-1} \sum_{i=1}^m \|e_{n,i}\|_V^2.$$

We denote the numerical solution by $\mathbf{u}^\tau = (u_{n,i})$ and the corresponding values of the exact solution by $\mathbf{u} = (u(t_n + c_i\tau))$, where in both cases $n = 0, \dots, N-1$ and $i = 1, \dots, m$.

We have the following error bound for the nonlinear problem. Here the restriction to the two-stage Radau IIA method stems from Lemma 3.2.

PROPOSITION 5.3 Let the nonlinear function g be continuous, strictly monotone and have at most cubic growth. Suppose that the solution u to the problems (5.1) and (5.2) is sufficiently regular.

Consider the time discretization (5.7) and (5.10) by the two-stage Radau IIA convolution quadrature method. Then, there is $\bar{\tau} > 0$ such that for step sizes $0 < \tau \leq \bar{\tau}$, the error in the boundary values satisfies the bound

$$\|\gamma^+ \mathbf{u}^\tau - \gamma^+ \mathbf{u}\|_{\ell_2^{\tau}(0:N;H^{1/2}(\Gamma)^2)} \leq C\tau^3, \tag{5.18}$$

and the error in the exterior domain is bounded by

$$\|\mathbf{u}^\tau - \mathbf{u}\|_{\ell_2^3(0:N;H^1(\Omega^+)^2)} \leq C\tau^{3/2}. \tag{5.19}$$

The constants C are independent of τ and N with $0 < N\tau \leq T$, but depend on T .

Proof. We eliminate φ in the system of boundary integral equations (5.5) to arrive at a boundary integral equation for Ψ ,

$$L(\partial_t)\Psi + g(\Psi + \dot{u}^{\text{inc}}) = \partial_\nu^+ u^{\text{inc}}, \tag{5.20}$$

where

$$L(s) = s^{-1} \left(W(s) - \left(\frac{1}{2}I - K^T(s) \right) V(s)^{-1} \left(\frac{1}{2}I - K(s) \right) \right) = -s^{-1} \text{DtN}^+(s)$$

with the exterior Dirichlet-to-Neumann operator $\text{DtN}^+(s)$. It follows from Propositions 17 and 18 (and their proofs) in the study by [Laliena & Sayas \(2009\)](#) that, for $\text{Re } s \geq \sigma > 0$, there exist $C(\sigma)$ and $\alpha(\sigma) > 0$ such that

$$\|L(s)\|_{H^{-1/2}(\Gamma) \leftarrow H^{1/2}(\Gamma)} \leq C(\sigma) \frac{|s|}{\text{Re } s}, \tag{5.21}$$

$$\text{Re} \langle \Psi, L(s)\psi \rangle \geq \alpha(\sigma) \frac{\text{Re } s}{|s|^2} \|\Psi\|_{H^{1/2}(\Gamma)}^2 \quad \text{for all } \Psi \in H^{1/2}(\Gamma), \tag{5.22}$$

where $\langle \cdot, \cdot \rangle$ denotes the anti-duality pairing between $H^{-1/2}(\Gamma)$ and $H^{1/2}(\Gamma)$.

Thanks to the composition rule we can do the same for the numerical discretization (5.7) and reduce the numerical system to an equation for Ψ^τ , which is just the convolution quadrature time discretization of (5.20),

$$L(\partial_t^\tau)\Psi^\tau + g(\Psi^\tau + \dot{u}^{\text{inc}}) = \partial_\nu^+ u^{\text{inc}}. \tag{5.23}$$

For the error $\boldsymbol{\varepsilon} = \Psi^\tau - \Psi$ with $\Psi = \left((\Psi(t_n + c_i\tau))_{i=1}^m \right)_{n=0}^{N-1}$ we then have the error equation

$$L(\partial_t^\tau)\boldsymbol{\varepsilon} + g(\Psi^\tau + \dot{u}^{\text{inc}}) - g(\Psi + \dot{u}^{\text{inc}}) = \mathbf{d} \tag{5.24}$$

with the defect

$$\mathbf{d} = L(\partial_t^\tau)\Psi - \left((L(\partial_t)\Psi(t_n + c_i\tau))_{i=1}^m \right)_{n=0}^{N-1},$$

which is the convolution quadrature error for $L(\partial_t)\Psi$. By Theorem 4.2 and our assumption of a sufficiently regular $\Psi = \gamma^+ \dot{u}$ this is bounded by

$$\|\mathbf{d}_n\|_{H^{-1/2}(\Gamma)} \leq C\tau^3 \quad \text{for } 0 \leq n\tau \leq T.$$

Since we can apply the same argument also to spatial derivatives of Ψ (in the assumed case of a smooth boundary Γ) we even have

$$\|\mathbf{d}_n\|_{H^{1/2}(\Gamma)} \leq C\tau^3.$$

We test (5.24) with $\boldsymbol{\varepsilon}$, multiply with $e^{-2\tilde{\sigma}t}$ with $\tilde{\sigma} = 1/T$ and integrate from 0 to T . With (5.22) and the Runge–Kutta convolution coercivity as given by Theorem 3.1 and with the strict monotonicity (5.24) we conclude that

$$\begin{aligned} & \alpha\tau \sum_{n=0}^N e^{-2\tilde{\sigma}n\tau} \left\| \left((\partial_t^\tau)^{-1} \boldsymbol{\varepsilon} \right)_n \right\|_{H^{1/2}(\Gamma)}^2 + \beta\tau \sum_{n=0}^N e^{-2\tilde{\sigma}n\tau} \|\boldsymbol{\varepsilon}_n\|_{L_2(\Gamma)}^2 \\ & \leq \tau \sum_{n=0}^N e^{-2\tilde{\sigma}n\tau} \langle \boldsymbol{\varepsilon}_n, \mathbf{d}_n \rangle \end{aligned} \tag{5.25}$$

and estimate further

$$\langle \boldsymbol{\varepsilon}_n, \mathbf{d}_n \rangle \leq \|\boldsymbol{\varepsilon}_n\|_{L_2(\Gamma)} \|\mathbf{d}_n\|_{L_2(\Gamma)} \leq \frac{\beta}{2} \|\boldsymbol{\varepsilon}_n\|_{L_2(\Gamma)} + \frac{1}{2\beta} \|\mathbf{d}_n\|_{L_2(\Gamma)}^2.$$

We thus find the stability estimate

$$\left\| (\partial_t^\tau)^{-1} \boldsymbol{\varepsilon} \right\|_{\ell_2^2(0:N;H^{1/2}(\Gamma)^2)} + \|\boldsymbol{\varepsilon}\|_{\ell_2^2(0:N;L_2(\Gamma)^2)} \leq C \|\mathbf{d}\|_{\ell_2^2(0:N;L_2(\Gamma)^2)}.$$

Since $(\partial_t^\tau)^{-1} \boldsymbol{\varepsilon} = \gamma^+ \mathbf{u}^\tau - \gamma^+ \mathbf{u}$, this proves (5.18).

Let us denote by $M(s) = S(s)V^{-1}(s) : H^{1/2}(\Gamma) \rightarrow H^1(\Omega^+)$ the operator that maps Dirichlet data in $H^{1/2}(\Gamma)$ to the corresponding solution $\hat{u} \in H^1(\Omega^+)$ of the Helmholtz equation $s^2\hat{u} - \Delta\hat{u} = 0$. By Sayas (2016, Equation (3.10)) this is bounded for $\text{Re } s \geq \sigma > 0$ by

$$\|M(s)\|_{H^1(\Omega^+) \leftarrow H^{1/2}(\Gamma)} \leq C(\sigma) \frac{|s|^{3/2}}{\text{Re } s}.$$

We then have

$$\begin{aligned} \mathbf{u}^\tau - \mathbf{u} &= M(\partial_t^\tau) \gamma^+ \mathbf{u}^\tau - \left((M(\partial_t) \gamma^+ u(t_n + c_i\tau))_{i=1}^2 \right)_{n=0}^{N-1} \\ &= M(\partial_t^\tau) (\gamma^+ \mathbf{u}^\tau - \gamma^+ \mathbf{u}) + \left(M(\partial_t^\tau) \gamma^+ \mathbf{u} - \left((M(\partial_t) \gamma^+ u(t_n + c_i\tau))_{i=1}^2 \right)_{n=0}^{N-1} \right). \end{aligned}$$

By Theorem 4.2 and the bound for M the last term is bounded by $O(\tau^{5/2})$ in the $H^1(\Omega^+)$ norm. The first term is only $O(\tau^{3/2})$ since we lose a factor $\tau^{3/2}$ from the $O(\tau^3)$ error bound for $\gamma^+ u$ because of the $O(|s|^{3/2})$ bound of $M(s)$; this follows from Lemma 5.2 in the study by Banjai & Lubich (2011) and Parseval’s identity. \square

In a similar way we obtain the following results for the alternative discretizations (5.8) and (5.9):

(i) In addition to Proposition 5.3 assume that g has bounded second derivatives. With the discretization (5.8) instead of (5.7) the error bound in the $H^1(\Omega^+)$ norm improves to $O(\tau^{5/2})$, and the ℓ_2^τ error in a point x bounded away from the boundary Γ is at most $O(\tau)$.

(ii) In addition to Proposition 5.3 assume that g has bounded second and third derivatives. With the discretization (5.9) instead of (5.7) the error bound in the $H^1(\Omega^+)$ norm improves to $O(\tau^3)$, and the ℓ_2^ξ error in a point x bounded away from the boundary Γ is at most $O(\tau^2)$.

The proofs of these error bounds are very similar to that of Proposition 5.3, using in addition a discrete Gronwall inequality at the end of the estimation of ϵ , and an $O(|s|^3)$ bound for the norm of the operator from $H^{1/2}(\Gamma) \rightarrow \mathbb{C}$ that maps Dirichlet data to the solution of the Helmholtz equation $s^2\hat{u} - \Delta\hat{u} = 0$ at a point $x \in \Omega^+$ bounded away from Γ , for s in a right half-plane. Since our main concern here is to illustrate the use of the convolution coercivity we omit the details of these extensions.

REMARK 5.1 If we set $\tilde{L}(s) = L(s + \sigma)$ and $\tilde{\psi}(t) = e^{-\sigma t}\Psi(t)$ for some $\sigma > 0$ then the boundary integral equation (5.20) is equivalent to

$$(\tilde{L}(\partial_t)\tilde{\Psi})(t) + e^{-\sigma t}g(e^{\sigma t}\tilde{\Psi}(t) + \hat{u}^{\text{inc}}(t)) = e^{-\sigma t}\partial_v^+ u^{\text{inc}}(t). \tag{5.26}$$

By (5.22) we then have the coercivity estimate for $\tilde{L}(s)$ for all $\text{Re } s \geq 0$ (and not just for $\text{Re } s \geq \sigma$):

$$\text{Re} \langle \Psi, \tilde{L}(s)\Psi \rangle \geq \alpha(\sigma) \frac{\text{Re } s + \sigma}{|s + \sigma|^2} \|\Psi\|_{H^{1/2}(\Gamma)}^2 \quad \text{for all } \Psi \in H^{1/2}(\Gamma).$$

By Theorem 3.1 the coercivity estimate for the convolution quadrature approximation of $\tilde{L}(\partial_t)\tilde{\Psi}$ is then obtained for every algebraically stable Runge–Kutta method (and not just the two-stage Radau IIA method). Hence, by discretising the shifted boundary integral equation (5.26) on an interval $[0, T]$ with shift $\sigma = 1/T$, we obtain Runge–Kutta-based convolution quadrature time discretizations of arbitrarily high order of convergence (assuming sufficient regularity of the exact solution). We remark that similar shifts are familiar in the convergence analysis of space-time Galerkin methods for time-dependent boundary integral equations (Bamberger & Ha-Duong, 1986). As in that case numerical experiments indicate that implementing the shift may not be necessary in practical computations, although this is not backed by theory.

6. Numerical experiments

6.1 Scattering by the unit sphere

In these experiments we let Ω^+ be the exterior of the unit sphere and the trace of the incident wave u^{inc} on the sphere be space independent. As constant functions are eigenfunctions of all the integral operators on the sphere (Nédélec, 2001) the solution will also be constant in space. The eigenvalue for the combined operator $L(s)$ in (5.20) is given by

$$L(s)\hat{\Psi} = -s^{-1}\text{DtN}^+(s)\hat{\Psi} = \left(1 + \frac{1}{s}\right)\hat{\Psi}, \tag{6.1}$$

for any $\hat{\Psi}$ constant in space. This operator will reflect well the behaviour of scattering by a convex obstacle, but not that of a general scatterer. For this reason we concentrate on the corresponding interior problem with

$$L^-(s)\hat{\Psi} = s^{-1}\text{DtN}^-(s)\hat{\Psi} = \left(-\frac{1}{s} + \frac{1 + e^{-2s}}{1 - e^{-2s}}\right)\hat{\Psi}, \tag{6.2}$$




FIG. 1. Convergence of the error for the two-stage Radau IIA method with the two different nonlinear impedance conditions.

again for $\hat{\Psi}$ constant. Treating both these operators as scalar, complex valued functions of s we see that both have a better behaviour than the general operators, see (5.21) and (5.22). Namely,

$$|L(s)| \leq C(\sigma), \quad \operatorname{Re} L(s) \geq 1$$

and

$$|L^-(s)| \leq C(\sigma), \quad \operatorname{Re} L^-(s) \geq \alpha(\sigma).$$

As the operator $L(s)$ is too simple, in the numerical experiments we only consider the scalar, nonlinear equation

$$L^-(\partial_t)\Psi + g(\Psi + \dot{u}^{\text{inc}}) = 0. \quad (6.3)$$

Even though these operators are of such a simple form, due to the nonlinearity the exact solution is not available. Nevertheless, a highly accurate solution is not expensive to evaluate and can be used to compute the error in the ℓ_2^τ norm at time steps t_n . We have performed the numerical experiments with the following choices of g and u^{inc}

$$g_1(\xi) = \frac{1}{4}\xi + \xi|\xi|, \quad g_2(\xi) = \frac{1}{4}\xi + \xi^3, \quad u^{\text{inc}}(t) = 2e^{-10(t-5/2)^2}$$

and with final time $T = 6$. Note that g_1 is once continuously differentiable whereas g_2 is infinitely differentiable. The data u^{inc} is not causal, but it is vanishingly small for $t < 0$ and we have found that this discrepancy has no significant effect on the results.

In Fig. 1 we show the convergence of the two-stage Radau IIA convolution quadrature in the ℓ_2^τ norm at t_n . As expected, for the smooth nonlinear boundary condition we obtain full order of convergence. For the nonsmooth boundary condition the solution is less smooth and the order of convergence is reduced. To investigate this further the solution and its first derivative are shown in Fig. 2. Note that the two solutions have a similar shape, but a closer look at the derivative in Fig. 3 reveals that one is smooth and the other only once continuously differentiable.




FIG. 2. We show the solution and its first derivative. On the left is the solution with the once continuously differentiable impedance g_1 and on the right with the smooth impedance g_2 .




FIG. 3. A closer look at the derivatives of the solutions reveals the differing smoothness.




FIG. 4. A closer look at the derivatives of the solutions reveals the differing smoothness.




FIG. 5. Scattering of a plane wave by an L-shaped domain with a nonlinear impedance condition.

For the exterior as $\operatorname{Re} L(s) \geq 1$ the theory also applies to higher order Radau IIA methods. This is, however, not the case with $L^-(s)$. We nevertheless perform experiments with the three-stage Radau IIA method and obtain good results as shown in Fig. 4. For the smooth case we again obtain full convergence order, which in this case is of order 5. For the less smooth case we obtain the same reduced convergence order as with the two-stage Radau IIA method, but a better error constant.

6.2 A fully nonscalar example

We end the paper with a two-dimensional example that requires the full BEM discretization in space. The domain is an L-shape and the incident wave is a plane wave. Piecewise linear boundary element space is used to approximate the Dirichlet trace Ψ and piecewise constant boundary element space to approximate the Neumann trace φ and the time-discretization is performed using the two-stage Radau IIA method. We have set the incident wave to

$$u^{\text{inc}}(x, t) = -2e^{-100(x_1+t-3)^2}, \quad x = (x_1, x_2), \quad t \geq 0,$$

and the boundary function g to the less smooth case g_1 . The images of the solution are shown in Fig. 5.

To estimate the error we fix the dimension of both boundary element spaces to 1.2×10^3 degrees




FIG. 6. Convergence of the error (6.4) with decreasing τ_j .

of freedom and compute the solutions until time $T = 2$ with decreasing time steps $\tau_j = T/N_j$, $N_j = 5, 10, 20, 40, 80$. The error is then estimated by computing

$$\text{error}_j = \|\gamma^+ u^{\tau_{j+1}} - \gamma^+ u^{\tau_j}\|_{\ell_2^x(0:N;H^{1/2}(\Gamma))} = \left(\tau_j \sum_{n=0}^{N_j} \|\gamma^+ u^{\tau_{j+1}}(t_n) - \gamma^+ u^{\tau_j}(t_n)\|_{H^{1/2}(\Gamma)}^2 \right)^{1/2}. \quad (6.4)$$

The $H^{1/2}$ norm is computed by using the equivalence of norms $\|\phi\|_{H^{1/2}(\Gamma)}^2 \sim \text{Re} \langle \phi, W(1)\phi \rangle_\Gamma$. This equivalence follows from the continuity properties of the hypersingular operator and the coercivity of $W(s)$ on $H^{1/2}(\Gamma)$ for $\text{Re } s > 0$; see [Sayas \(2016\)](#). The convergence of this error estimate is shown in Fig. 6. The convergence order is not as clear as in the scalar case, but as expected, indicates a lower than optimal order.

Acknowledgements

We thank Ernst Hairer for helpful discussions.

Funding

Deutsche Forschungsgemeinschaft, Sonderforschungsbereich 1173.

REFERENCES

- BAMBERGER, A. & HA-DUONG, T. (1986) Formulation variationnelle espace-temps pour le calcul par potentiel retardé d’une onde acoustique. *Math. Meth. Appl. Sci.*, **8**, 405–435.
- BANJAI, L. & KACHANOVSKA, M. (2014) Fast convolution quadrature for the wave equation in three dimensions. *J. Comput. Phys.*, **279**, 103–126.
- BANJAI, L. & LUBICH, C. (2011) An error analysis of Runge-Kutta convolution quadrature. *BIT*, **51**, 483–496.

- BANJAI, L. & RIEDER, A. (2018) Convolution quadrature for the wave equation with a nonlinear impedance boundary condition. *Math. Comp.*, **87**, 1783–1819.
- BANJAI, L., LUBICH, C. & MELENK, J. M. (2011) Runge-Kutta convolution quadrature for operators arising in wave propagation. *Numer. Math.*, **119**, 1–20.
- BANJAI, L., LUBICH, C. & SAYAS, F.-J. (2015) Stable numerical coupling of exterior and interior problems for the wave equation. *Numer. Math.*, **129**, 611–646.
- BANJAI, L., MESSNER, M. & SCHANZ, M. (2012) Runge-Kutta convolution quadrature for the boundary element method. *Comput. Methods Appl. Mech. Engrg.*, **245/246**, 90–101.
- EBERLE, S. (2018) The elastic wave equation and the stable numerical coupling of its interior and exterior problems. *ZAMM*, doi:10.1002/zamm.201600236.
- GRABER, P. J. (2012) Uniform boundary stabilization of a wave equation with nonlinear acoustic boundary conditions and nonlinear boundary damping. *J. Evol. Equ.*, **12**, 141–164.
- HADDAR, H. & JOLY, P. (2002) Stability of thin layer approximation of electromagnetic waves scattering by linear and nonlinear coatings. *J. Comput. Appl. Math.*, **143**, 201–236.
- HAIRER, E. & LUBICH, C. (1984) On the stability of Volterra-Runge-Kutta methods. *SIAM J. Numer. Anal.*, **21**, 123–135.
- HAIRER, E. & WANNER, G. (1996) *Solving Ordinary Differential Equations II. Stiff and Differential-Algebraic Problems*. Springer Series in Computational Mathematics, vol. 14, 2nd edn. Berlin: Springer, pp. xvi+614.
- KOVÁCS, B. & LUBICH, C. (2017) Stable and convergent fully discrete interior-exterior coupling of Maxwell's equations. *Numer. Math.*, **137**, 91–117.
- LALIENA, A. R. & SAYAS, F.-J. (2009) Theoretical aspects of the application of convolution quadrature to scattering of acoustic waves. *Numer. Math.*, **112**, 637–678.
- LUBICH, C. (1994) On the multistep time discretization of linear initial-boundary value problems and their boundary integral equations. *Numer. Math.*, **67**, 365–389.
- LUBICH, C. & OSTERMANN, A. (1987) Multigrid dynamic iteration for parabolic equations. *BIT*, **27**, 216–234.
- LUBICH, C. & OSTERMANN, A. (1993) Runge-Kutta methods for parabolic equations and convolution quadrature. *Math. Comp.*, **60**, 105–131.
- NÉDÉLEC, J.-C. (2001) *Acoustic and Electromagnetic Equations. Integral Representations for Harmonic Problems*. Applied Mathematical Sciences, vol. 144. New York: Springer, pp. x+316.
- VON NEUMANN, J. (1951) Eine Spektraltheorie für allgemeine Operatoren eines unitären Raumes. *Math. Nachr.*, **4**, 258–281.
- SAYAS, F.-J. (2016) *Retarded Potentials and Time Domain Boundary Integral Equations*. Springer Series in Computational Mathematics, vol. 50. Heidelberg: Springer.
- SCHÄDLE, A., LÓPEZ-FERNÁNDEZ, M. & LUBICH, C. (2006) Fast and oblivious convolution quadrature. *SIAM J. Sci. Comput.*, **28**, 421–438 (electronic).
- WANG, X. & WEILE, D. S. (2011) Implicit Runge-Kutta methods for the discretization of time domain integral equations. *IEEE Trans. Antennas and Propagation*, **59**, 4651–4663.