

Accepted manuscript of article in Journal of Strength and Conditioning Research

Authors' Response

Muniz-Pumares, Daniel¹; Karsten, Bettina²; Triska, Christoph^{3,4}; Glaister, Mark⁵

¹ School of Life and Medical Sciences, University of Hertfordshire, Hatfield, UK.

² Department of Exercise and Sport Science, LUNEX International University of Health, Exercise and Sports, Differdingen, Luxemburg.

³ Centre for Sport Science and University Sports, University of Vienna, Vienna, Austria

⁴ Austrian Institute of Sports Medicine, Vienna, Austria

⁵ Faculty of Sport, Health and Applied Science, St Mary's University, Twickenham, UK.

Corresponding author:

Daniel Muniz-Pumares

School Life and Medical Sciences

College Lane

University of Hertfordshire

Hatfield

AL10 9EU

United Kingdom

Telephone: 0170 728 3495 (Ext. 77666)

Email: d.muniz@herts.ac.uk

We would like to thank Pettitt and colleagues (10) for their comments regarding our review on the different methods for calculating critical power (CP) and W' . The main criticism from Pettitt et al. (10) reflects on our conclusion that the 3-min all-out test (3MT) should be interpreted with 'a degree of caution' (7).

We agree that W' exhibits a large variability irrespective of the method used to determine this parameter, although this may be reduced in certain testing conditions (12). We also appreciate that several studies have reported a good agreement between 3MT and the 'Gold Standard' approach for both CP and W' . Crucially, however, others have reported that the 3MT may overestimate CP (4). This seems to occur primarily as fitness levels increase (1), in turn affecting the predictions of time-trial performance (9).

The 3MT may be able to differentiate the upper boundary of steady-state exercise (see for example (5)). However, other studies have shown that during exercise at 3MT-determined CP, physiological responses may not reach a steady state (3). Furthermore, exercise at CP can typically be sustained in excess of 20 min, but when CP is determined from the 3MT, task failure has been shown to occur after ~12.5 min (2).

Pettitt et al. (10) questioned whether the attainment of maximum oxygen uptake ($\dot{V}O_{2max}$) should be implemented as a test criterion for the 3MT as the test benefits from being a time-efficient test (as opposed to the more time-consuming conventional approach). We acknowledge that verifying $\dot{V}O_{2max}$ may compromise the strength of this aspect of the 3MT. However, if the option is available, we encourage a $\dot{V}O_{2max}$ attainment verification. Pettitt et al. (10) highlighted that a verification bout following the ramp test may be required to determine a 'true' $\dot{V}O_{2max}$ (11), so the attainment of $\dot{V}O_{2max}$ during the 3MT can be confirmed. However, as $\dot{V}O_2$ should theoretically reach $\dot{V}O_{2max}$ during the 3MT (5,14; cf. 13), it is plausible to use the 3MT as the verification bout.

In summary, we do not advocate that the 3MT should not be used. Indeed, the test offers a time-saving alternative to estimate CP and W' where other protocols may be cumbersome (e.g. 6,8). However, reflecting on the strengths but also on the limitations of the 3MT, we would like to uphold our conclusion that the 3MT should be interpreted with a degree of caution, particularly in elite athletes.

References

1. Bartram, J, Thewlis, D, Martin, DT, and Norton, KI. Predicting critical power in elite cyclists: Questioning validity of the 3-min all-out test. *Int J Sports Physiol Perform* 12: 783–787, 2017.

2. Bergstrom, HC, Housh, TJ, Zuniga, JM, Traylor, DA, Lewis, RW, Camic, CL, et al. Responses during exhaustive exercise at critical power determined from the 3-min all-out test. *J Sports Sci* 31: 537–45, 2013.
3. Bergstrom, HC, Housh, TJ, Zuniga, JM, Traylor, DA, Lewis, RW, Camic, C, et al. Metabolic and neuromuscular responses at critical power from the 3-min all-out test. *Appl Physiol Nutr Metab* 38: 7–13, 2013.
4. Bergstrom, HC, Housh, TJ, Zuniga, JM, Traylor, DA, Lewis, RW, Camic, CL, et al. Differences among estimates of critical power and anaerobic work capacity derived from five mathematical models and the three-minute all-out test. *J Strength Cond Res* 28: 592–600, 2014.
5. Burnley, M, Doust, JH, and Vanhatalo, A. A 3-min all-out test to determine peak oxygen uptake and the maximal steady state. *Med Sci Sport Exerc* 38: 1995–2003, 2006.
6. Muniz-Pumares, D, Karsten, B, Triska, C, and Glaister, M. Methodological approaches and related challenges associated with the determination of critical power and W' . *J Strength Cond Res* 33: 584–596, 2019.
7. Muniz-Pumares, D, Pedlar, C, Godfrey, R, and Glaister, M. A comparison of methods to estimate anaerobic capacity: Accumulated oxygen deficit and W' during constant and all-out work-rate profiles. *J Sports Sci* 35: 2357–2364, 2017.
8. Clark, I, Vanhatalo, A, Thompson, C, Wylie, LJ, Bailey, SJ, Kirby, B, et al. Changes in the power-duration relationship following prolonged exercise: estimation using conventional and all-out protocols and relationship to muscle glycogen. *J Appl Physiol*, 2019. Published ahead of print.
9. Nicolò, A, Bazzucchi, I, and Sacchetti, M. Parameters of the 3-minute all-out test: Overestimation of competitive-cyclist time-trial performance in the severe-intensity domain. *Int J Sports Physiol Perform* 12: 655–661, 2017.
10. Pettitt, RW, Jamnick, NA, Kramer, M, and Dicks, ND. A Different Perspective of the 3-min all-out Exercise Test: Manuscript Clarification. *J Strength Cond Res*, 2019.
11. Poole, DC and Jones, AM. Measurement of the maximum oxygen uptake $\dot{V}O_{2max}$: $\dot{V}O_{2peak}$ is no longer acceptable. *J Appl Physiol* 122: 997–1002, 2017.
12. Triska, C, Karsten, B, Heidegger, B, Koller-Zeisler, B, Prinz, B, Nimmerichter, A, et al. Reliability of the parameters of the power-duration relationship using maximal effort time-trials under laboratory conditions. *PLoS One* 12:e0189776, 2017.
13. Vanhatalo, A, Doust, JH, and Burnley, M. Determination of critical power using a 3-min all-out cycling test. *Med Sci Sports Exerc* 39: 548–55, 2007
14. Vanhatalo, A, Poole, DC, DiMenna, FJ, Bailey, SJ, and Jones, AM. Muscle fiber recruitment and the slow component of O_2 uptake: constant work rate vs. all-out sprint exercise. *Am J Physiol Regul Integr Comp Physiol* 300: R700-7, 2011.

