Determination of the Key Parameters influencing Dislodgeable foliar Pesticide Residues 'DFR'

<sup>1</sup>Mohamed Badawy, <sup>1</sup>Kathleen Lewis, <sup>2</sup>Neil Morgan, <sup>2</sup>Robert Mason <sup>1</sup>School of Life and Medical sciences, University of Hertfordshire, UK <sup>2</sup>Syngenta Ltd.,UK

Email: mb18ado@herts.ac.uk

syngenta

What does dislodgeable foliar residue DFR' mean?

Dislodgeable foliar residue (DFR) is the amount of pesticide residues which are deposited on plant leaves and may dislodge after pesticide application by people during the performance of various tasks and can subsequently deposited on human skin and clothes that may cause potential risk.



What is The current challenge of DFR to the industry?

The default values of DFR and other parameters in the risk assessment become more precautionary in the regulatory landscape, the requirements for specific DFR studies are increasing. These studies are costly, time consuming and limited by the crops seasons due to the unacceptability of extrapolation between DFR studies among crops.

Dissipation data resulting from DFR studies are used to evaluate the risk associated with post application exposure to pesticides and to establish chemical transfer coefficient (TC) to determine restricted – entry intervals of cultivated crops.









Factors that may affects DFR

### Application rate

Addressing the relation between the application rate and deposition rate, which may increase the potential accumulations of residues on leaves

#### Rain fall

Pesticides residues on plant leaves may be influenced by different precipitation intensities

## Formulation type

The role of different formulation types and coformulants with different range of active ingredient particle sizes may have an impact on dislodge-ability

### Temperature

The effect of temperature on foliar persistence of pesticides as a factor

### Leaf texture and shapes

Different leaf shapes
(ovate, oblong, linear,
cordate, Reniform etc.)
and textures from smooth
to hairy may have an
impact on different
dislodge-ability

Developing a Lab method for testing the effect of formulation types on DFR

**EC** Formulation

EC +Adjuvant
EC+ co-formulants


WP Formulation

WP +Adjuvant
WP+ co-formulants

### Micro pipetting Application



### Track sprayer Application







# Determination of the Key Parameters influencing Dislodgeable foliar Pesticide Residues `DFR' Mohamed Badawy



What does dislodgeable foliar residue DFR mean?





What is The current challenge of DFR to the industry?



harvesting orchard fruit



pruning orchard fruit



thinning orchard fruit



harvesting grapes




harvesting ornamentals



harvesting fruiting veg

### Factors that may affects DFR




Application rate



Rain fall



Formulation type & coformulants



Temperature



Leaf texture and shapes