

Accepted manuscript

Article accepted for publication in the British Journal of Sports Medicine, 9 August 2019.

International Scientific Tendinopathy Symposium Consensus, Groningen (ICON 2019). Part 1. Agreement and Recommendations re: Clinical Terminology

Authors:

ICON 2019: International Scientific Tendinopathy Symposium Consensus: Clinical Terminology

Authors:

Alexander Scott¹, Kipling Squier¹, Håkan Alfredson², Roald Bahr^{3,4}, Jill Cook⁵, Brooke K Coombes⁶, Robert-Jan de Vos⁷, Siu N (Amy) Fu⁸, Alison Grimaldi⁹, Jeremy Lewis^{10,11}, Nicola Maffuli¹², Stig Peter Magnusson¹³, Peter Malliaras¹⁴, Seán Mc Auliffe⁴, Edwin HG Oei¹⁵, Craig Purdam⁵, Jonathan D Rees^{16,17}, Ebonie Rio⁷, Cathy Speed¹⁸, Karin Gravare Silbernagel¹⁹, Inge van den Akker-Scheek^{20,21}, Bill Vicenzino²², Adam Weir^{23,24}, Jennifer Moriatis Wolf²⁵, Johannes Zwerver²⁰

¹ Department of Physical Therapy, Faculty of Medicine, University of British Columbia, 2177 Wesbrook Mall Vancouver, Canada

² Sports Medicine Unit, University of Umeå, Sweden, University of Umeå, Sweden Umeå, Sweden

³ Oslo Sports Trauma Research Center, Norwegian School of Sport Sciences Oslo, Norway

⁴ Aspetar Orthopedic and Sports Medicine Hospital, Doha, Qatar

⁵ La Trobe University Sport and Exercise Medicine Research Centre, Kingsbury Drive, Bundoora Melbourne, Australia

⁶ Griffith University, 170 Kessels Road, Nathan, QLD 4111 Brisbane, Australia

⁷ Department of Orthopaedics and Sports Medicine, Erasmus MC University Medical Centre, Rotterdam, The Netherlands.

⁸ Department of Rehabilitation Sciences, The Hong Kong Polytechnic University, Yuk Choy Road, Hung Hom, Kowloon Hong Kong, Hong Kong (SAR)

⁹ Physiotec, Tarragindi Brisbane, Australia

¹⁰ University of Hertfordshire, School of Health and Midwifery Hertfordshire, UK

¹¹ Central London Community Healthcare NHS Trust, London London, UK

¹² Department of Musculoskeletal Disorders, Faculty of Medicine and Surgery, University of Salerno, 84081 Baronissi Salerno, Italy

¹³ Department of Physical Therapy, Bispebjerg Hospital Copenhagen, Denmark

¹⁴ Department of Physiotherapy, Monash University Melbourne, Australia

¹⁵ Erasmus MC, University Medical Center; Department of Radiology & Nuclear Medicine, Dr. Molewaterplein 40 Rotterdam, The Netherlands.

¹⁶ Fortius Clinic, 17 Fitzhardinge Street London, UK

¹⁷ Centre for Sports & Exercise Medicine, Queen Mary University of London, London UK

¹⁸ Professor of Sports Medicine & Human Performance, Cardiff metropolitan university, Cardiff, UK Cardiff, UK

¹⁹ Department of Physical Therapy, University of Delaware, 540 South College Avenue Newark, Delaware, USA

²⁰ University of Groningen, University Medical Center Groningen, Department of Sport and Exercise Medicine, Hanzeplein 1, PO Box 30.001, 9700RB Groningen, The Netherlands.

²¹ University of Groningen, University Medical Center Groningen, Department of Orthopaedics, Hanzeplein 1, PO Box 30.001, 9700RB Groningen, The Netherlands.

²² The University of Queensland: School of Health & Rehabilitation Sciences, St Lucia Campus: Therapies Annexe (84-A) Brisbane, Australia

²³ Aspetar Sports Groin Pain Centre, Aspetar Orthopaedic and Sports Hospital Doha, Qatar

²⁴ Sport medicine and exercise clinic Haarlem (SBK), Haarlem, The Netherlands.

²⁵ Department of Orthopaedic Surgery, University of Chicago, USA

Background

Persistent tendon pain that impairs function has inconsistent medical terms that can influence choice of treatment (1). When a person is told they have **tendinopathy** by clinician A or **tendinitis** by clinician B, they might feel confused or be alarmed at receiving what they might perceive as two different diagnoses. This may lead to loss of confidence in their health professional and likely adds to uncertainty if they were to search for information about their condition. Clear and uniform terminology also assists inter-professional communication. Inconsistency in terminology for painful tendon disorders is a problem at numerous anatomical sites.

Historically, the term “tendinitis” was first used to describe tendon pain, thickening, and impaired function (Supplementary File, Figure S1). The term “tendinosis” has also been used in a small number of publications, some of which were very influential (2,3). Subsequently, “tendinopathy” emerged as the most common term for persistent tendon pain (4,5). To our knowledge, experts (clinicians and researchers) or patients have never engaged in a formal process to discuss the terminology we use. We believe that health professionals have not yet agreed on the appropriate terminology for painful tendon conditions.

The authors of this paper, a group of international clinical and research experts from different disciplines, aimed to achieve a consensus in terminology for persistent tendon disorders. We ran a Delphi and consensus process that culminated in a face-to-face meeting at the 5th International Scientific Tendinopathy Symposium (ISTS) in Groningen, the Netherlands, on September 26th, 2018. [placeholder for sentence about the other two papers and accompanying editorial if they get accepted] Here, we present the resulting consensus statements on terminology for persistent tendon pain.

Methods

Our two-stage Delphi design, consensus process consisted of an online survey followed by a face-to-face meeting. One of us (AS) drafted 11 statements about terminology relevant to typical clinical presentations and clinical findings. Following several teleconferences by the project committee (AS, BV, HZ, ER, AW, SM), these statements were further refined. The role of the committee members was to develop and execute the Delphi process. The final document we sent out for Stage 1 of the consensus process (see below) featured 10 statements (Supplementary File, Appendix 2).

Stage 1

We e-mailed invitations to 30 tendinopathy experts. These experts were unanimously selected by the core author group, according to 4 goals:

1. Encourage participation by both men and women or male and female experts;
2. Choose from among top ranked experts (Web of Science and Expertscape searches) who have published original research on tendinopathy;

3. Achieve representation from multiple disciplines (sports physicians, rheumatologists, physiotherapists, epidemiologists, clinicians, researchers, surgeons, radiologists);

4. Have broad geographic representation.

The invitation letter explained the process and the rules for participation, including an option to participate in both the online survey and the in-person meeting, or the online survey alone (Supplementary File, Appendix 3). Of 30 invitees, twenty-eight people with a clinical and/or research interest responded and agreed to participate. Of the two who did not agree to participate, one did not respond within the timeframe and one withdrew after agreeing to participate due to personal reasons, unrelated to the survey. The experts were sent a link to an anonymized survey (Survey Monkey, San Mateo, CA). For each statement the experts could choose from agree, disagree, or don't know/ not sure, as well as an optional space for comments. One author (BV) compiled the survey results and shared them with the core author group for analysis. Statements that achieved consensus (70% or higher agreed or disagreed) were not discussed further (Supplementary File, Appendices 4 and 5).

Stage 2

Five of the project committee members (AS, BV, HZ, ER, SM) and 10 of the experts who also completed the survey attended the in-person meeting in Groningen (15 participants). All participants funded their own travel and accommodation and received no payments for attending. We distributed a summary report of the survey results to attendees before the in-person meeting. This report included response rates for each question, and free-text entries (Appendix 5).

At the meeting, AS facilitated an open discussion using Power Point to display the 5 statements about which on-line participants (n=28) had not achieved consensus after the online survey. After discussion, the group excluded two statements from further analysis due to the possibility of perceived lack of diverse disciplinary representation at the in-person group (#5, imaging, and #10, rotator cuff). For the other statements, a vote was cast by secret ballot. At this step, the voting options were yes/agreed or no/disagreed. The votes were counted by two individuals in the room and the results announced and recorded (Supplementary File, Appendix 6).

A draft of the manuscript was circulated to all 28 authors and all comments/edits were accepted. Three experts declined to be listed as a co-author on the manuscript, and communicated this to one of us by email (BV). Each of them stated they did not disagree with the statements, but did not feel they had contributed enough to warrant authorship.

Results

The ICON 2019: International Scientific Tendinopathy Symposium Consensus on Clinical Terminology is shown in Table 2.

Table 2. ICON 2019: International Scientific Tendinopathy Symposium Consensus: Clinical Terminology

Tendinopathy is the preferred term for persistent tendon pain and loss of function related to mechanical loading.

Tendon tear (partial or complete) refers to a macroscopic* discontinuity of a load-bearing tendon.

Imaging is not always necessary for a diagnosis of tendinopathy.

Patellar tendinopathy is the preferred term for persistent patellar tendon pain and loss of function related to mechanical loading.

Achilles tendinopathy is the preferred term for persistent Achilles tendon pain and loss of function related to mechanical loading.

Peroneal (fibularis) tendinopathy is the preferred term for persistent peroneal (fibularis) tendon pain and loss of function related to mechanical loading.

Persistent tendon pain and loss of function related to mechanical loading of the medial or lateral elbow tendons should be referred to as medial or lateral elbow tendinopathy.

* large enough to be visible without magnification

Discussion

Tendinopathy is the preferred term for persistent tendon pain and loss of function related to mechanical loading.

This definition of tendinopathy emphasizes pain and function. Using the term tendinopathy can help frame treatment and improve communication between professionals and patients. With this statement we hope that the term tendinopathy will continue to supersede other terms. The term does not imply the presence of a particular pathological or biochemical process which cannot routinely be assessed clinically.

Tendinosis - a label without a disease?

We did not agree on the statement that tendinosis refers to a loss of microstructure (as opposed to macroscopic tearing). Tissue changes seen on imaging or histology and labelled as tendinosis may represent a snap-shot of a number of morphological processes with uncertain clinical relevance. In some situations, a loss of collagen alignment could be physiological (e.g. as a result of aging, adaptation or repair) rather than pathological. We cannot recommend using the term tendinosis unless future work can establish a consensus on its definition.

Tendon tear (partial or complete) refers to a macroscopic discontinuity of load-bearing tendon

Tears represent a clinical entity for some tendons which is distinct from tendinopathy, and which may require different management to that of tendinopathy. The ICON group defined tears as a macroscopic (as opposed to microscopic) discontinuity of load-bearing tendon. Future research should examine whether tears, partial tears and tendinopathy can be reproducibly distinguished. If they could be, it may have clinical utility (i.e. guide treatment).

We need to discover if diagnosing partial tears leads to improved patient outcomes, and in which situations it adds no value. We believe there is not a validated means of identifying partial tears on imaging for all tendons.

Does a tendinopathy at each region in the body warrant its own name?

Tendon pain occurs in several anatomical sites such as (but not limited only to) the Achilles, patellar, peroneal (fibularis), and elbow. We did not address nomenclature at other tendons, such as tibialis posterior or gluteal tendons.

The term 'rotator cuff tendinopathy' was on our list of options but the group did not achieve consensus on that term as being ideal for shoulder pain and loss of function because a proportion of participants felt that the diagnosis could not be made clinically because of the variety of nociceptive structures adjacent/near to the rotator cuff tendons. With respect to this, the Groningen statement is consistent with previous Delphi studies and clinical guidelines which include tendinopathy as part of a spectrum of subacromial or rotator cuff related shoulder pain and injury (6-8). Those guidelines recommend the terms subacromial pain (or impingement) syndrome for patients with painful shoulder tendons and loss of function; the term rotator cuff related shoulder pain has also been proposed (8).

Future work

Imaging can provide additional information which can assist with diagnosis. If terminology is needed to describe morphological tendon changes as seen with imaging or histology, we recommend that clinicians use objective measures (if their validity and reliability can be demonstrated) such as increased or decreased signal intensity (for MR images), collagen disorientation or hypoechogenicity (in the case of ultrasound images), etc. An ISTS imaging subgroup was formed to explore/consider issues related to the role of imaging for diagnosis in for tendinopathy and relevant terminology.

Conclusion

We recommend that clinicians and patients use the term tendinopathy to refer to persistent tendon pain related to mechanical loading.

Contributions

AS provided study proposal, compiling of contributing authors' comments, preparation of the first draft, consensus methods and analysis. BV contributed to consensus methods, writing, data collection and analysis. HZ, ER, AW and SM contributed to consensus methods and writing. All others contributed with Delphi participation and writing.

Acknowledgements

Hamed Mousavi, Olivier Dams, Lucas Rabello

Competing Interests

Authors had no competing interests and received no funding.

References

- 1 Nickel B, Barratt A, Copp T, et al. Words do matter: a systematic review on how different terminology for the same condition influences management preferences. *BMJ Open* 2017;7:014129.
- 2 Kraushaar BS, Nirschl RP. Tendinosis of the elbow (tennis elbow). Clinical features and findings of histological, immunohistochemical, and electron microscopy studies. *J Bone Joint Surg Am* 1999;81:259-278.
- 3 Kannus P, Jozsa L. Histopathological changes preceding spontaneous rupture of a tendon. A controlled study of 891 patients. *J Bone Joint Surg Am* 1991;73:1507-1525.
- 4 Khan KM, Cook JL, Kannus P, et al. Time to abandon the "tendinitis" myth. *BMJ* 2002;324:626-627.
- 5 Maffulli N, Khan KM, Puddu G. Overuse tendon conditions: time to change a confusing terminology. *Arthroscopy* 1998;14:840-843.
- 6 Diercks R, Bron C, Dorrestijn O, et al. Guideline for diagnosis and treatment of subacromial pain syndrome: a multidisciplinary review by the Dutch Orthopaedic Association. *Acta Orthop* 2014;85:314-322.
- 7 Huisstede BM, Miedema HS, Verhagen AP, et al. Multidisciplinary consensus on the terminology and classification of complaints of the arm, neck and/or shoulder. *Occup Environ Med* 2007;64:313-319.
- 8 Lewis J. Rotator cuff related shoulder pain: Assessment, management and uncertainties. *Man Ther* 2016;23:57-68.