


Assessing risks and mitigating impacts of harmful algal blooms on mariculture and marine fisheries

Andrew Ross Brown¹ , Martin Lilley¹, Jamie Shutler², Chris Lowe², Yuri Artioli³, Ricardo Torres³, Elisa Berdalet⁴ and Charles R. Tyler¹

1 College of Life and Environmental Sciences, University of Exeter, Exeter, UK

2 College of Life and Environmental Sciences, University of Exeter, Penryn, UK

3 Plymouth Marine Laboratory, Plymouth, UK

4 Institute of Marine Sciences (ICM-CSIC), Barcelona, Spain

Correspondence

Andrew Ross Brown, College of Life and Environmental Sciences, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter, Devon, EX4 4QD, UK. Email: ross.brown@exeter.ac.uk

Received 27 August 2019; accepted 18 November 2019.

Abstract

Aquaculture is the fastest growing food sector globally and protein provisioning from aquaculture now exceeds that from wild capture fisheries. There is clear potential for the further expansion of marine aquaculture (mariculture), but there are associated risks. Some naturally occurring algae can proliferate under certain environmental conditions, causing deoxygenation of seawater, or releasing toxic compounds (phycotoxins), which can harm wild and cultured finfish and shellfish, and also human consumers. The impacts of these so-called harmful algal blooms (HABs) amount to approximately 8 \$billion/yr globally, due to mass mortalities in finfish, harvesting bans preventing the sale of shellfish that have accumulated unsafe levels of HAB phycotoxins and unavoids human health costs. Here, we provide a critical review and analysis of HAB impacts on mariculture (and wild capture fisheries) and recommend research to identify ways to minimise their impacts to the industry. We examine causal factors for HAB development in inshore versus offshore locations and consider how mariculture itself, in its various forms, may exacerbate or mitigate HAB risk. From a management perspective, there is considerable scope for strategic siting of offshore mariculture and holistic Environmental Approaches for Aquaculture, such as offsetting nutrient outputs from finfish farming, via the co-location of extractive shellfish and macroalgae. Such pre-emptive, ecosystem-based approaches are preferable to reactive physical, chemical or microbiological control measures aiming to remove or neutralise HABs and their phycotoxins. To facilitate mariculture expansion and long-term sustainability, it is also essential to evaluate HAB risk in conjunction with climate change.

Key words: food production, food quality, HABs, mariculture, phycotoxins, risk mitigation.

Introduction

Managing global food security is one of the greatest challenges of the twenty-first century. Currently, around 820 million people (1 in 9 people) suffer from malnutrition (FAO, IFAD, UNICEF, WFP and WHO, 2018) and this is projected to rise as the human population grows from 7.6 to a projected 11.2 billion by 2100 (UN, 2017). While agricultural productivity and yields from wild capture fisheries have plateaued or are in decline, aquaculture has grown substantially over the last forty years, particularly in Asia, a

region which now supplies ~90% of the global aquaculture market (FAO, 2018). Future food production in all sectors, however, may be limited by increasing climate variability, including extremes in rainfall intensity and temperature. These changes in climate in combination with increasing human population numbers, pollution events, impaired nutrient cycling, outbreaks of disease and pestilence are likely to result in future shortfalls in food production (FAO, 2018; FAO, IFAD, UNICEF, WFP and WHO, 2018). For aquaculture production, one of the most critical threats is the occurrence of harmful algal blooms (HABs).

Increasing frequency of HABs is associated with climate change, nutrient enrichment and habitat disturbance and is leading to growing impacts, including the poisoning or asphyxiation of finfish, shellfish and poisoning of human consumers (Hallegraeff 1993; GESAMP, 2001; Smayda 2004; Anderson 2012; Berdalet *et al.* 2016). Harmful algal blooms can also cause a variety of other impacts affecting water quality, water flow and amenity value. Therefore, estimating the economic costs of HABs is complex and requires consideration of many different issues (see reviews by Berdalet *et al.* 2016; Adams *et al.* 2018). Among the biggest economic impacts of HABs are precautionary closures of fisheries and aquaculture farms to prevent human poisoning (see Section 'Global distribution and characterisation of HABs affecting human health through seafood consumption' on human poisoning). Annual costs of precautionary closures (US\$ at first point of sale) are estimated at \$3–4 billion: >\$0.03 billion in the UK (ASIMUTH, 2014); \$0.9–1.2 billion in the EU (Hoagland & Scatista 2006; S-3 EuroHAB, 2019); \$0.1–1.0 billion in Korea, Japan and China (Kim 2006; Trainer & Yoshida 2014); and >\$0.10 billion in the USA (Hoagland *et al.* 2002). Furthermore, the worldwide economic impacts of marine phyco-toxins on human health are estimated to be approximately \$4 billion a year (GESAMP, 2001; references in Berdalet *et al.* 2016). These estimates are very much 'best approximations' rather than detailed economic assessments (as conceded by some of the authors, e.g., Hoagland & Scatista 2006; Adams *et al.* 2018). According to conservative epidemiological assessments, around 2000 cases of HAB-related food poisonings occur each year globally, following human consumption of contaminated finfish or shellfish, and around 15% of these cases prove fatal (FAO, 2012; CTA, 2013). The proportion of farmed versus wild-caught finfish and shellfish that contain phycotoxins and subsequently poison human consumers is not currently known.

Food fish production from aquaculture (80 million tonnes, US\$232 billion per year) now exceeds capture fisheries (Table 1, adapted from FAO, 2018). Growth projections see this production from aquaculture rising by 37%, from 70 million tonnes to 109 million tonnes, by 2030 (FAO, 2018), with a significant contribution coming from the global expansion of mariculture (Kapetsky *et al.* 2013). Food fish production from mariculture currently amounts to

28.7 million tonnes, of which more than half comes from bivalve shellfish. Bivalves are among the most sustainable mariculture products, since they derive their food entirely from naturally occurring food sources, predominantly marine planktonic microalgae. The growth of these algae is fuelled by natural (and anthropogenic) nutrient supplies from land runoff and coastal upwelling (Huston & Wolverton 2009). Farming of aquatic plants and algae, dominated by seaweeds (macroalgae), has also increased recently to >30 million tonnes (FAO, 2018), worth an estimated US \$11.7 billion. The largest share of seaweed production is for human food products (polysaccharide carbohydrates and micronutrients), and the remainder is for animal feeds, fertilisers and biopolymers (Nayar & Bott 2014).

Around 200 marine species are currently farmed, with the greatest variety in tropical seas (FAO, 2015; Froehlich *et al.* 2016). Species can be divided into two broad categories: (i) fed species, including finfish and some crustaceans and (ii) 'extractive' species, including (a) unfed filter-feeding bivalves, algal grazers, detritivores and (b) autotrophic plants, mainly macroalgae. Each of these categories has different environmental susceptibilities, interactions and installation planning issues (Gentry *et al.* 2016), particularly at inshore sites (≤ 1 km from the coast). At inshore sites, mariculture is directly influenced by anthropogenic activities (agricultural and urban runoff, municipal and industrial effluent inputs, ships and mariculture itself), which potentially increase HAB risk (Anderson *et al.* 2008; Anderson 2012). Recent calculations have suggested that current seafood consumption could be met by extending mariculture offshore, into less than 1% of Exclusive Economic Zones belonging to coastal states (Gentry *et al.* 2017). Some HABs, however, originate in open oceanic waters (Davidson *et al.* 2009, 2016; Trainer *et al.* 2012; Shutler *et al.* 2015; Gobler *et al.* 2017), indicating that some algal species may present similar or even greater risks as mariculture moves offshore.

Mariculture represents the nexus of environment–food–health systems, with food productivity and quality depending on clean coastal waters and healthy intact marine ecosystems (FAO, IFAD, UNICEF, WFP and WHO, 2018). To ensure long-term sustainable growth of the industry, a collection of interconnecting issues covering biosecurity, economic and environmental aspects (including climate

Table 1 Global food fish production from aquaculture in 2016

Aquaculture production	Finfish	Molluscs	Crustacea	Other	Total for aquaculture	Total as % of total food fish
By weight (million tonnes)	54.1	17.1	7.9	1.0	80*	53
By value (billion US\$)	138.5	29.2	57.1	6.8	232	64

*Mariculture currently provides 36% (28.7 million tonnes) of food fish production from all forms of aquaculture (including freshwater and recirculating systems) and is dominated by molluscs (17.1 million tonnes; FAO, 2018).

change and HABs) need to be addressed (De Silva & Soto 2009; Lovatelli *et al.* 2013). Here, we critically review national and international HAB monitoring data records and published literature, to evaluate the occurrences, causes and impacts of HABs on shellfish and finfish mariculture in inshore and offshore waters. We identify environmental factors contributing to HAB risk and establish whether mariculture practices themselves can influence (increase or reduce) risks of HAB occurrence and impact. Methods for predicting and mitigating HAB risk are then reviewed. The risks of HABs to wild capture fisheries, as well as mariculture, are considered in this review also, since mariculture has the potential to attract and promote aggregations of wild finfish and shellfish. Building improved understanding of HAB risk for these related industries is of paramount importance to ensure future marine food security and safety.

Impacts of HABs on marine fisheries and mariculture

Nature of HABs and their impacts

Harmful algal blooms are proliferations of certain microalgae, macroalgae or blue/green algae (cyanobacteria), which under favourable environmental conditions reach certain levels that can have negative impacts on humans or the aquatic environment (Hallegraeff 1993; Anderson 2012; Bresnan *et al.* 2013; GlobalHAB, 2017). Some HAB species or strains synthesise phycotoxins that are ingested by marine plankton grazers and potentially bioaccumulate in higher food chain organisms, including humans. Epiphytic HAB species including *Prorocentrum lima*, *Ostreopsis* spp. and *Gambierdiscus* spp. have the potential to contaminate seaweeds, but human poisonings are generally caused by the consumption of seaweed grazing herbivorous shellfish, finfish or their predators, rather than from direct consumption of seaweeds. Globally, around 300 HAB species have been identified, of which more than a third, mainly in the dinoflagellate group, are known to produce toxins that are harmful to aquatic organisms and/or to humans consuming them (<http://www.marinespecies.org/hab/index.php>; Anderson 2012). Toxin production can vary between different genetic strains for some HAB species (e.g. Touzet *et al.* 2010; Cochlan *et al.* 2012) and/or different environmental conditions (Fehling *et al.* 2004; Wells *et al.* 2005). Poisoning syndromes in humans, responsible HAB genera, phycotoxin groups, and shellfish, finfish and macro-algal vectors of these phycotoxins are summarised in Section 'Global distribution and characterisation of HABs affecting human health through seafood consumption' (Table 2). Other metabolites may also be generated from these toxins, many of which have not been characterised in terms of chemical structure, potency or public health

significance (Wiese *et al.* 2010; Anderson 2012). Other HAB species cause harm to fish through gill clogging or via the production of fish toxins (ichthyotoxins). Also, when the blooms decay, the degradation of the accumulated algal biomass by bacteria results in oxygen depletion, affecting aquatic ecosystems as a whole (Smayda 2004; Svendsen *et al.* 2018).

Global distribution and characterisation of HABs affecting human health through seafood consumption

Information concerning the global occurrence and impact of HAB events is recorded in the Harmful Algae Event Database (HAEDAT, <http://haedat.iode.org>). Bivalve molluscs, which filter and feed directly on microalgae, including HAB species, are the principal vectors for shellfish poisoning in humans. Crustaceans that prey upon intoxicated bivalves, including crabs and lobsters (Shumway 1995; James *et al.* 2010), and also carnivorous finfish (Friedman *et al.* 2017) can also bioaccumulate and in turn act as important vectors for phycotoxins. Table 2 summarises the principal poisoning syndromes that result from humans ingesting intoxicated shellfish or finfish and the respective geographical areas of highest incidence.

The phycotoxins associated with each poisoning syndrome (column 1 of Table 2) are neurotoxins, and they are heat-stable (and thus unaffected by cooking), underlining their risk to human health. Global maps of reported shellfish poisonings are illustrated in Manfrin *et al.* (2012), and selected references on poisoning syndromes can be found in Berdalet *et al.* (2016). Microalgae can produce a broader spectrum of toxic compounds than illustrated in Table 2 and include yessotoxins (YTXs) and pectenotoxins (PTXs) that mainly cause diarrhoea (Reguera *et al.* 2014). An increasing number of toxic compounds derived from algae are being detected as monitoring and analytical tools become more advanced, including brevetoxins (Turner *et al.* 2015) and cyclic imines (Davidson *et al.* 2015).

Occurrences and impacts of HABs on marine organisms in fisheries and mariculture

Evidence on the occurrence and impacts of HAB on marine fisheries and mariculture is being gathered by ongoing regional programmes (e.g. Maguire *et al.* 2016), national programmes (e.g. UK FSA, <https://www.food.gov.uk/business-guidance/biotoxin-and-phytoplankton-monitoring>) and global (GlobalHAB, 2017) programmes (see section 'In situ monitoring'). However, despite the increasing coordination and integration of HAB monitoring programmes and research, not all incidents are captured and records may not always tally between local and global databases (e.g. HAEDAT). Some HABs are difficult to detect, notably

Table 2 Most common foodborne poisoning syndromes in humans caused by HABs and details concerning their occurrence and impacts

Poisoning syndrome	Symptoms	Causal phycotoxins	Mechanism of toxicity	Responsible HAB species	Principal vectors	Impacts (examples)	Global hot spots ^d
Amnesic shellfish poisoning (ASP)	Memory loss, brain damage ^a	Domoic acid (DO)	Agonism of the neurotransmitter glutamate	<i>Pseudo-nitzschia</i> spp.	Scallops, e.g., <i>Pecten maximus</i> Crabs, e.g., <i>Metacarcinus magister</i>	Scallop harvesting bans (months) ^b Collapse of Californian Dungeness crab fishery 2015–2016 ^c	Pacific, Atlantic coasts of North and Central America, Atlantic Europe
Paralytic shellfish poisoning (PSP)	Confused speech, tingling burning sensations, nausea, diarrhoea ^e	Saxitoxins (STXs)	Inhibition of voltage-dependent sodium channels ^e	<i>Alexandrium catenella</i> , <i>A. minutum</i> , <i>Gymnodinium catenatum</i> , <i>Pyrodinium bahamense</i> var. <i>compressum</i>	Mussels, clams, oysters, crabs, lobsters	Some 2000 PSP cases are reported per year globally (for all principal vectors), with occasional fatal consequences in humans ^e	North and South America and Canada, Africa, Europe (North Sea Mediterranean), and Australasia
Diarrhetic shellfish poisoning (DSP)	Diarrhoea, nausea, vomiting and abdominal cramps ^f	Okadaic acid (OA), <i>Dinophysis</i> toxins ^f (DTXs)	Inhibition of protein phosphatases in intestine and neurons ^f	<i>Dinophysis</i> spp. <i>Prorocentrum</i> spp.	Mussels, clams, oysters Edible crabs (<i>Cancer pagurus</i>)	Harvesting bans for bivalves in Europe (weeks-months) ^g Closure of edible crab fishery in Norway (weeks-months) ^h	Reported globally and particularly in NW Europe
Azaspiracid poisoning (AZP)	Diarrhoea, nausea, vomiting and abdominal cramps ⁱ	Azaspiracids (AZAs)	Modulation of gamma amino butyric acid (GABA) ⁱ	<i>Amphidomataceae</i> : <i>Amphidoma</i> , <i>Azadinium</i>	Mussels, king scallops and edible crabs ^j	Harvesting bans (months) for shellfisheries (principal vectors) and mariculture in Atlantic Europe ^j	Norway coast, UK and Atlantic coast of France and Spain
Neurotoxic shellfish poisoning (NSP)	Loss of motor control, nausea muscular ache, including abdominal ^k	Brevetoxins (BTXs)	Inhibition of voltage-dependent sodium channels ^k	<i>Karenia</i> spp.	Clams, oysters and mussels ^l	Seafood poisoning. The formation of toxic aerosols by wave action also produces respiratory irritation and asthma-like symptoms	East and West coasts of North America, Florida and the Gulf of Mexico
Ciguatera fish poisoning (CFP)	Gastrointestinal, neurologic and cardiac distress ^m	Ciguatoxin (CTX), maitotoxin (MTX)	Agonism of voltage-gated sodium channels	<i>Gambierdiscus</i> spp.	Herbivorous fish (grazing HABs on macrophytes macroalgae) and their predators	CFP is one of the most common poisoning syndromes resulting from the consumption of contaminated finfish ^m	Caribbean, Florida, East Africa, Madagascar, Northern Australia, Pacific Islands

References: (a) Lundholm *et al.* (1994); (b) Campbell *et al.* (2003); (c) California Ocean Science Trust (2016); (d) Manfrin *et al.* (2012); (e) Anderson (2012); (f) Munday (2013); (g) Reguera *et al.* (2014); (h) Castberg *et al.* (2004); (i) Furey *et al.* (2010); (j) Twiner *et al.* (2008); (k) Kirkpatrick *et al.* (2004); (l) Watkins *et al.* (2008); (m) Friedman *et al.* (2017).

for species which bloom below the sea surface and evade *in situ* monitoring and satellite imaging (Shutler *et al.* 2015). It is also often difficult to attribute cause(s) to observed impacts on complex marine systems, particularly when they involve cryptic species and nonspecific mechanisms, such as the depletion of dissolved oxygen and suffocation of (shell)fish by HABs such as *Karenia mikimotoi* (Davidson *et al.* 2009; Shutler *et al.* 2015). Since the 1960s, the number of hypoxic or anoxic 'dead zones' in coastal waters has doubled every decade (Diaz & Rosenberg 2008). This has occurred in conjunction with increasing eutrophication caused by nutrient enrichment and excessive algal growth. In some cases, notable asphyxiation impacts on finfish and shellfish have been attributed to high biomass blooming HAB species such as *Phaeocystis* spp., *Karenia* spp. and *Aureococcus anophagefferens* (Peperzak & Poelman 2008; Davidson *et al.* 2009; Gobler *et al.* 2011).

Evidence of acute toxicity from HABs on finfish and shellfish in wild fisheries and mariculture

Harmful algal bloom species from different taxonomic groups with few commonalities (dinoflagellates, dictyophytes, haptophytes, prymnesiophytes, raphidophytes) have been implicated in major finfish kills in marine fisheries and mariculture. In some cases, the toxicity can be transmitted up the food chain to seabirds and marine mammals. Widely cultured finfish species affected by HABs include Atlantic salmon (*Salmo salar*), Rainbow trout (*Oncorhynchus mykiss*) and Yellowtail amberjack/kingfish (*Seriola quinqueradiata*) (reviewed by Landsberg 2002; Clément *et al.* 2016). Nevertheless, the mechanisms of toxicity for 'fish killing HABs' are not well understood. An example illustrating the complexity associated with HAB toxicity in finfish is presented for *Heterosigma akashiwo*. Here, effects may be due to the production of reactive oxygen species, brevetoxin-like compound(s), excessive mucus production that impedes oxygen exchange, gill tissue damage by mucocysts and/or haemolytic activity. Uncertainties arise when there are differences in the toxicity of wild HAB populations versus laboratory cultures; for example, reduced toxicity has been shown to result from the long-term culturing of *H. akashiwo* (Cochlan *et al.* 2012). There may also be variability in mucocyst production by different strains of microalgae (in the case of *Pseudochattonella farcimen*; Andersen *et al.* 2015).

Marine fisheries (and other wildlife). Some of the largest and most regular finfish (and other wildlife) kills occur annually along Florida's Gulf coast. Here, epidemiological assessments have attributed these to brevetoxin poisonings from blooms of the dinoflagellate *Karenia brevis* (Landsberg *et al.* 2009; Flaherty & Landsberg 2011). A recent bloom of *K. brevis* lasted over a year, beginning in

November 2017, extending for a distance of 150–200 miles along Florida's Gulf coast and killed hundreds of tonnes of marine life, including thousands of small fish, numerous large fish (including groupers and a 21-ft whale shark) and marine mammals, including dolphins (Pickett 2018). The 2017–2018 bloom is one of the longest and most severe outbreaks recorded over the last 70 years and illustrates the scale of impacts possible from a single HAB outbreak (Krimsky *et al.* 2018). Elsewhere, for example in the UK (1978, 1980) and Ireland (1976, 1978, 1979 and 2005), major finfish and shellfish kills have been attributed to *Karenia mikimotoi* (a.k.a. *Gyrodinium (or Gymnodinium) aureolum*; e.g. Silke *et al.* 2005; Mitchell & Rodgers 2007). These blooms have caused widespread death of wild and cultured fish, through either acute toxicity attributed to phycotoxins with neurotoxic, haemolytic or cytotoxic effects, or via oxygen depletion caused by decaying blooms (e.g. Boalch 1979; Jenkinson & Connors 1980; Jones *et al.* 1982).

Saxitoxin produced by *Alexandrium* spp. may also be lethal to larvae and juveniles of commercially important finfish and shellfish species, such as Atlantic mackerel (*Scomber scombrus*) and American lobster (*Homarus americanus*; Robineau *et al.* 1991). Biomagnification of saxitoxin in the marine food chain has also been linked to significant fish kills, and both seabird and marine mammal deaths (Pitcher & Calder 2000; Sephton *et al.* 2007).

Mariculture. Harmful algal blooms often lead to finfish kills in caged environments, where the fish cannot escape phycotoxins or oxygen depletion from the decaying algal biomass. Risks from HABs are particularly high for finfish confined in sheltered inshore embayments, where the HABs may be concentrated by onshore winds and currents. As an example of this, between 1972 and 1982 in the Seto Inland Sea, Japan, at least 21.8 million cultured yellowtail amberjack (*Seriola quinqueradiata*) were killed by the raphidophyte *Chatonella antiqua* (Okaichi 1989). In 1972, the economic loss for the summer outbreak amounted to US \$70 million. Since then, annual losses have been lower, but recurring severe impacts have continued (Fukuyo *et al.* 2002). Recurring threats have been reported also from another toxic raphidophyte, *H. akashiwo*, causing finfish kills in Iceland, Spain, British Columbia and Chile (Landsberg 2002). The losses caused by outbreaks of *H. akashiwo* to wild and net-penned finfish off Puget Sound, Washington, have been estimated to cost in the region of US\$2–6 million per episode. The outbreaks of *H. akashiwo* are believed to have been increasing generally in scope and magnitude in various global regions over the past two decades (Landsberg 2002).

Originating offshore around the UK (Davidson *et al.* 2009; Shutler *et al.* 2015), high biomass blooms

(>1000 cells/mL) of *Karenia mikimotoi* have been increasingly frequent and have been associated with significant finfish kills, including for caged fish in inshore waters (Jenkinson & Connors 1980; Silke *et al.* 2005; Davidson *et al.* 2009). Farmed shellfish including mussels, oysters and clams (*Tapes semidecussata*) in the UK and Ireland, and hatchery raised juvenile bivalve spat have also periodically suffered significant mortalities, along with crustaceans and other benthic invertebrates, in conjunction with *K. mikimotoi* blooms (Raine *et al.* 2001; Silke *et al.* 2005).

Evidence of chronic toxicity from HABs in wild fisheries and mariculture

Symptoms of chronic toxicity in finfish are wide ranging for different HABs. These symptoms include liver pathologies caused by ciguatoxins released from *Gambierdiscus* spp. and microcystins produced by *Microcystis* spp., gill pathologies caused by cytotoxins from, for example *Prymnesium* spp. and *Heterosigma* spp., narcosis (loss of balance and swimming ability) caused by neurotoxins from *Karenia* spp. and paralysing saxitoxin from *Alexandrium* spp., and excess gill mucus production, for example, caused by *Chaetoceros* spp. (review by Burkholder 1998; Svendsen *et al.* 2018).

Chronic sublethal effects of HAB toxins in bivalve molluscs include reduction in feeding rates in scallops and oysters (e.g. caused by exposure to *Prorocentrum minimum*), reduction in growth and byssus production in blue mussels (*Mytilus edulis*), growth reduction in Eastern oysters (*Crassostrea virginica*), for example caused by *Gymnodinium aurelium/Karenia mikimotoi* (Burkholder 1998) and by *Alexandrium tamarense* (Li *et al.* 2002), reproductive impairment in blue mussels and Bay scallops (*Argopecten irradians*), for example caused by *Chrysochromulina polylepis*, reduction in the recruitment of juvenile Bay scallops, for example caused by *Karenia brevis* (reviewed by Burkholder 1998; Basti *et al.* 2018). Thus, in addition to toxin accumulation rendering shellfish unsafe for harvesting for human consumption, toxin presence can have a longer term effect, impacting on shellfish abundance and time taken to grow to marketable size. Slower pumping and filtering rates are also likely to increase the time taken to evacuate toxic material from shellfish tissues. Most shellfish species can eliminate phycotoxins within a few weeks, but retention of some toxins (e.g. saxitoxins) in some species, such as sea scallops (*Placopecten magellanicus*) and Atlantic surfclams (*Spisula solidissima*), can last up to 5 years (Shumway *et al.* 1990; Landsberg 2002). Harmful algal blooms also have the potential to impact adversely on the supply of larval 'seed' or 'spat' for aquaculture. Examples of this include *Karenia brevis* impacting on larval recruitment in Bay scallops (Burkholder 1998), Pacific oysters (*Crassostrea gigas*) and Northern quahog (*Mercenaria*

mercenaria); (Rolton *et al.* 2018). For these shellfisheries, the estimated annual economic losses due to *K. brevis* along Florida's Gulf coast alone are estimated to be up to US\$6 million (NOAA 2004; Adams 2017). *Karenia brevisulcata* has also been shown to be toxic to larvae of Greenshell mussel (*Perna canaliculus*), Pacific oyster and New Zealand abalone (*Haliotis iris*); (Shi *et al.* 2012).

Consumption of intoxicated finfish and shellfish can also lead to chronic toxicity in organisms higher in marine food chains. For example, domoic acid derived from *Pseudo-nitzschia* sp. can cause neuropathic injury in both finfish and shellfish eating mammals and birds (Lefebvre *et al.* 2007; Ramsdell & Zabka 2008; Soliño *et al.* 2019).

Environmental factors contributing to HAB risk

Environmental factors promoting HABs

HABs are natural phenomena within the seasonal cycles of planktonic micro-organisms in aquatic ecosystems (Glibert *et al.* 2005; Shumway *et al.* 2018). In recent decades, harmful events appear to be increasing in frequency, duration and impact globally. Verifying them is a research priority (GlobalHAB, 2017; e.g. Wells *et al.* 2015, 2019). Apparent increased frequencies of HABs may be due to a combination of factors (see Figure 1) including: (i) warming sea surface temperatures, and associated water column stratification and range extensions of tropical organisms, including toxic species; (ii) increased frequency and intensity of storm events and flooding and associated increasing nutrient inputs, upwelling intensities and wider HAB dispersal; (iii) increasing anthropogenic pressures on the marine environment, notably land- and sea-based nutrient enrichment and disturbance of coastal habitats; and (iv) increased awareness and improvements in HAB monitoring systems (Hallegraeff 1993; Raine *et al.* 2008; Anderson 2012; Bresnan *et al.* 2013; Wells *et al.* 2015; Gobler *et al.* 2017; Anderson *et al.* 2019).

Evaluating HAB risk in any 'system' is highly challenging, since environmental drivers include a range of physical, chemical and biological factors, which can combine to influence (i) the initiation/development of a HAB; (ii) its impact/toxicity and (iii) the termination of a HAB (Roelke & Buyukates 2001; Anderson *et al.* 2012a). These factors operate from micro- (mm) to meso- (10–100 km) to macro (>100 km) spatial scales and over a range of temporal scales (from seconds to minutes and from days to months; Dickey 2001). For example, an abundant supply of dissolved nutrients, calm sea state, warming, increasing stratification and increased sunlight over a period of weeks may allow the algae to grow in high concentrations, and then, dramatic and significantly increased turbulent sea state (causing increased vertical mixing) over several hours can result in bloom termination (e.g. Shutler *et al.* 2015). The challenge of




Figure 1 Environmental factors promoting HABs. Complex interactions among environmental factors (solar radiation, wind, waves, tides, rainfall, nutrients), ecological and trophic interactions and biological processes (e.g. cyst formation) can facilitate the proliferation of phytoplankton in general and harmful algal species as well. Excess and unbalanced nutrient supply and habitat alteration can increase the risk of HAB occurrence. HABs negatively impact mariculture production and product quality. (However, some mariculture practices can mitigate the occurrence and impact of HABs, for example, through the use of integrated multitrophic aquaculture approaches; see Figure 2).

understanding HAB occurrence and toxicity is further complicated by ecological interactions between HAB species and other members of plankton communities, which vary both spatially and temporally in species composition, genetic diversity and physiological status (Anderson *et al.* 2012a; Davidson 2014). Despite these complexities, some of the key factors driving HAB dynamics are well characterised and are outlined in sections ‘Environmental factors contributing to HAB initiation and toxicity’, ‘Environmental factors contributing to HAB termination’ and ‘Regulation of HABs by filter feeding shellfish’ below.

Environmental factors contributing to HAB initiation and toxicity

The pre-requisites for any HAB event are the presence of algal cells, spores or cysts; suitable conditions of light and nutrients for their growth and reproduction; and physical conditions that facilitate their accumulation in favourable growing conditions. Cells can accumulate either by horizontal transport (advection) in water bodies by wind and/or tide, or by resuspension from sediments by wave action,

or upwelling of bottom water (e.g. Farrell *et al.* 2012; Pitcher *et al.* 2017). The source of propagules that initiate blooms may be local, or distant, though the origin of propagules for any particular harmful bloom is typically difficult to determine. There is evidence that HABs in some areas originate in the ocean, rather than in coastal embayments (Hinder *et al.* 2011; Whyte *et al.* 2014; Berdalet *et al.* 2017; Pitcher *et al.* 2017). The majority of HABs, including dinoflagellates and diatoms, are holoplanktonic, relying on vegetative cells to survive inhospitable conditions and to seed blooms. In some cases, when growth conditions are suboptimal, highly toxic HABs such as *Alexandrium* spp. reproduce sexually and form resting cysts. These cysts settle on sediments (Smayda & Trainer 2010) and then undergo resuspension during storms or coastal upwelling, enabling (re)colonisation of existing and new areas (e.g. Anderson *et al.* 1994; Pitcher *et al.* 2017).

Nutrient availability is another key requirement for HAB initiation and maintenance. Most HAB species are primarily photoautotrophs, and their requirements for autotrophic growth include inorganic nitrogen (N), phosphorus (P) and silicate (Si, in the case of diatoms).

High-biomass HABs in estuaries and coastal zones have been linked to elevated inorganic nutrient inputs (eutrophication; Rabalais *et al.* 2010; Paerl *et al.* 2014) and organic nutrients (e.g. urea from fertilisers, following heavy precipitation and land runoff, Heisler *et al.* 2008). However, the effects of nutrient inputs may be confounded by many other factors, including natural occurrence of HABs, transport of HAB species via mariculture and other marine activities, variable meteorological forcing, and longer-term climate change (Callaway *et al.* 2012; Gowen *et al.* 2012). There is increasing evidence that many HAB species can use dissolved and particulate organic forms of N and P (through prey ingestion), in addition to autotrophy; this combination of trophic modes is termed mixotrophy (Burkholder 1998; Anderson *et al.* 2002; Lin *et al.* 2018). Mixotrophic HAB species are therefore able to proliferate both under high organic N concentrations and by engulfing prey under nutrient limited conditions. Examples of mixotrophic HAB species include low biomass (100–1000 cells/L) blooming dinoflagellates, such as *Alexandrium* spp. (Anderson *et al.* 2012b; Lee *et al.* 2016) and *Dinophysis* spp. (Jacobson & Andersen 1994), and also high biomass (>10 000 cells/L) blooming species such as *Pseudo-nitzschia* spp. (Loureiro *et al.* 2008) and *A. anophagefferens* (Gobler *et al.* 2011). Furthermore, changes in nutrient ratios (far from the classic stoichiometric Redfield N:P ratio of 16:1) may be important in stimulating the growth of some HABs and influencing their toxin content (Anderson *et al.* 2002; Kudela *et al.* 2010; Glibert *et al.* 2014a) and responses may be highly species-specific (Wells *et al.* 2015).

Reduced turbulent mixing and increased thermal stratification are key factors promoting HABs, especially those comprised of dinoflagellates. Water column stratification and nutrient enrichment caused by river plumes, jets, upwelling areas and tidal fronts are also particularly conducive for HAB development (Pitcher *et al.* 2017). Phytoplankton and other planktonic organisms tend to collect passively in boundary layers in stratified water bodies – motile dinoflagellate HAB species have the added advantage of being able to visit both nutrient-rich deeper water and irradiance-saturated shallower water either side of these boundary layers (e.g. Smayda 1997). HABs are also more likely to occur in sheltered zones of lagoons, estuaries and coasts, as a result of increased water residence times, warmer temperatures and increased penetration of photosynthetically active radiation (PAR; e.g. Smayda 1989). Although strong turbulent mixing may be disadvantageous to bloom development by causing the breakup of chains of individuals and by inhibiting cell division (Estrada & Berdalet 1997), low level turbulence can enhance nutrient availability by facilitating increased transfer of molecules in or out of plankton cells, especially in passively floating

diatoms (Peters *et al.* 2006). Other biological processes, including inter-cell quorum sensing and encounter rates with competitors and grazers (Gowen *et al.* 2012), are also modulated by fine scale turbulence and this can also favour HABs (e.g. Berdalet *et al.* 2017).

Environmental factors contributing to HAB termination

Advection and dispersion of HABs, increasing turbulent shear forces breaking up cells, and/or nutrient limitation are all understood to contribute to the termination of HABs (Gentien *et al.* 2007; Lenés *et al.* 2013), and consequently, HAB prediction models are often driven by these physical processes and biogeochemical fluxes. However, models that only include these processes often ‘over-predict’ HAB duration, indicating that inter-species biotic interactions play important roles in terminating harmful blooms (Roelke & Buyukates 2001; Lenés *et al.* 2013; Davidson *et al.* 2016).

Plankton grazers or predators play an important role in regulating the abundance of marine planktonic microalgae, including HAB species. In nutrient limited (oligotrophic) offshore marine environments, mesozooplankton (e.g. copepods 0.2–20 mm) consume 10–40% of marine phytoplankton, while micro-zooplankton (20–200 µm) consume around 60–70% (Calbet 2008). In temperate nutrient-rich (eutrophic) upwelling and estuarine ecosystems, micro-sized heterotrophic and mixotrophic dinoflagellates (including HAB species) can dominate phytoplankton grazing (Calbet 2008). More detailed, mechanistic understanding concerning how and to what extent grazers regulate or terminate HABs is lacking. Plankton community interactions can vary markedly in temperate waters displaying a seasonal succession of different blooming species and also in (sub)tropical waters with relative constant standing stocks of microplankton. In both cases, food web dynamics can alternate between resource (bottom-up) and predatory (top-down) control (Calbet 2008) and outcomes for HABs are highly situation-specific (Turner & Tester 1997).

Marine parasitic microbes (micro- and nano-sized protists 10–100 µm, pico-sized bacteria 0.2–10 µm and femto-sized viruses ≤ 0.1 µm) target all of the main phytoplankton groups (Gachon *et al.* 2010). They have been shown to play a significant role in terminating some major algal blooms (Wilson *et al.* 2002) and have also been linked to the decline of HABs (Chambouvet *et al.* 2008; Roth *et al.* 2008; Jones *et al.* 2011). In turn, this has prompted research into the microbial control and bioremediation of HABs (Brussaard 2004; Sun *et al.* 2018; see section ‘Spatial and temporal planning to minimise HAB risk’). Larger micro-sized parasites such as the dinoflagellate *Amoebophyra* spp. may also be responsible for the termination

(Rosetta & McManus 2003; Montagnes *et al.* 2008) or regulation (Nishitani *et al.* 1985) of dinoflagellate HABs such as *Alexandrium* spp.

Adaptive responses in HAB species to avoid or combat grazers and parasites include sensing and moving away from grazers (Wolrhaf 2013); adapting/optimising colony size (chain length) versus swimming speed (Selander *et al.* 2012); synthesising and releasing phycotoxins and/or other allelochemicals (Stüken *et al.* 2011; Anderson 2012); undergoing or prolonging encystment (Rengefors *et al.* 1998; Toth *et al.* 2004); and undergoing auto-lysis (i.e. programmed cell death) (Franklin *et al.* 2006; Lenés *et al.* 2013). Combinations of mechanisms underlying predator–prey and host–parasite interactions can vary greatly since algal prey/host and predator/parasite niches are highly species-specific (Amin *et al.* 2015; Ramanan *et al.* 2016).

Regulation of HABs by filter feeding shellfish

Filter-feeding shellfish can exert considerable (top-down) grazing pressure, limiting phytoplankton (and zooplankton) biomass, particularly in shallow, well mixed estuaries and coastal waters, where bottom-living bivalves can come into contact with and filter the majority of the water column (Newell 2004; Lucas *et al.* 2016). Bivalves such as mussels, suspended on ropes hanging vertically in the water column can also be effective at filtering plankton at deeper water sites (Stadmark & Conley 2011; Hedberg *et al.* 2018). Physical factors such as water column exchange, turbulent mixing, temperature and stratification, and the influence of mariculture infrastructures on each of these (see Section ‘Physical alteration of habitats and hydrodynamic regimes’), can be important in modulating shellfish grazing, phytoplankton sinking, and phytoplankton community composition – for example reduced vertical mixing favours motile dinoflagellates, while nonmotile phytoplankton such as diatoms sink below the euphotic zone and are more easily intercepted by grazers (Lucas *et al.* 2016). The influence of selective filter feeding by shellfish on plankton community structure, including HABs species, is relatively poorly understood (Newell 2004; Petersen *et al.* 2008; Lucas *et al.* 2016). Simple size selection for nano-sized plankton and above (>4 µm) and higher filtration rates in the warmer summer months may serve to reinforce seasonal succession from nano- to pico-plankton dominated communities (Newell 2004). Sensing of food particles and their surface chemistry have been suggested to play a role in selective filtering of nutritious plankton in preference to detrital and mineral particles (Ward & Shumway 2004; Espinosa *et al.* 2009; Yahel *et al.* 2009). Phycotoxins, particularly paralytic shellfish toxins (PSTs) as well as other toxin classes (e.g. NSTs and ASTs), are capable of inducing valve closure and/or reducing filtration rate in bivalves, as

well as impairing growth and reproduction and inhibiting byssus production (Burkholder 1998; Landsberg 2002; Manfrin *et al.* 2012). Nevertheless, some bivalves show preferential uptake of harmful algal cells. This has been shown in the laboratory in five bivalve species (Bay scallop, Eastern oyster, Northern quahog, softshell clam (*Mya arenaria*) and the blue mussel (*Mytilus edulis*). All bivalves, with the exception of softshell clam, ejected intact cells of three HAB species (*Prorocentrum minimum* (PST and DST), *Alexandrium fundyense* (PST) and *Heterosigma akashiwo* (NST)) in their faeces or pseudo-faeces. Only oysters exposed to *H. akashiwo* showed partial or complete valve closure and reduction in filtration rate. These results confirm that feeding responses of bivalves in the presence of HABs can be highly species-specific. Furthermore, clearance of HABs from the water by bivalves may simply result in the transfer of intact/live cells to the sediment, from which they could be resuspended (Hégaret *et al.* 2007).

Environmental impacts of mariculture and contribution to HAB risk

Long-term time-series data are required to demonstrate the influence of finfish, shellfish and/or macro-algal mariculture on HAB risk as recognised in the Science Plan of the international programme on HABs (GlobalHAB, 2017). Accumulating evidence from China, which has the longest running, largest and highest concentration of mariculture in the world, indicates that the frequency and extent of HABs has been increasing concurrently with the industry growth since 1960 (Wang *et al.* 2008; Lu *et al.* 2014; Wartenberg *et al.* 2017). The occurrence of HAB events in China increased sharply in 2009 with ~80 episodes, covering > 15,000 km² of China’s coastline in just one year. The increasing trend, however, also follows increasing urbanisation of coastal fringes (Liu & Su 2017). Potential environmental effects of mariculture are listed in Table 3, and the tendencies for these effects to promote HAB formation and impact (either directly or indirectly) are discussed in sections ‘Nutrient emission versus assimilation’, ‘Chemical treatments used to control pathogens and parasites’, ‘Escapees and introduction of invasive and/or harmful species’, ‘Physical alteration of habitats and hydrodynamic regimes’ and ‘Transmission of HAB species and alteration in the abundance and composition of plankton communities’.

Organic and inorganic nutrient emission versus assimilation

Nutrient emissions from mariculture operations are predicted to increase substantially due to industry expansion (up to sixfold by 2050). The majority of these emissions

Table 3 Environmental effects of mariculture that can promote HAB risk

-
- (i) Organic and inorganic nutrient emission versus assimilation;
 - (ii) Disease and use of preventative chemical agents;
 - (iii) Escapees and genetic interactions with wild populations;
 - (iv) Physical alteration of habitats and hydrodynamic regimes;
 - (v) Increase in HAB transmission (between relay sites) or alteration of the abundance and composition of plankton communities.
-

References for (i–iv): Lovatelli *et al.* (2013), Kapetsky *et al.* (2013), Wartenberg *et al.* (2017).

References for (v): Gibbs (2004), Grant *et al.* (2007).

comprise nutrient waste, primarily from finfish (fed mariculture) and also from shellfish, released in a dissolved form directly to the water column (Bouwman *et al.* 2013). These nutrient emissions may promote the growth of harmful algal species in the vicinity of mariculture farms (Anderson *et al.* 2002; Hallegraeff 2003). However, causal linkages between fish farming and eutrophication (Pitta *et al.* 2005; Modica *et al.* 2006) and HABs (Anderson *et al.* 2008) are often not clear (Smayda 2004; Gowen *et al.* 2012). In some cases (e.g. farming of extractive shellfish), mariculture can cause net assimilation of nutrients leading to deficits (Ferreira *et al.* 2014), while elsewhere nutrient emissions may exceed local environmental assimilation capacities (Bouwman *et al.* 2013). Problems are likely to be more acute for farms with higher stocking densities (Sellner *et al.* 2003; Bouwman *et al.* 2013). Intensive bivalve cultivation can alter the nitrogen:phosphorus (N:P) nutrient stoichiometry and change the major N species to reduced forms, especially ammonia, as well as particulate organic nitrogen, and these N forms are preferred by various harmful algae – predominated by dinoflagellates (e.g. Arzul *et al.* 2001; Glibert *et al.* 2014a, but see Davidson *et al.* 2012). Conversely, diatoms have also been shown to decline as a result of nutrient excretion by bivalves (Lucas *et al.* 2016). A further concern arises because of low assimilation efficiencies (typically 30–40% for N, or less under bloom conditions), such that shellfish can become point sources of regenerated nutrients. Benthic regeneration of the accumulated faeces and decomposing feed can be significant in shallow well mixed coastal waters (Bouwman *et al.* 2013).

Disease and use of chemical treatments to control pathogens and parasites

Infections by pathogens and infestations of parasites, exacerbated by aggregations of wild fish around mariculture installations (Dempster *et al.* 2004), present a risk to human and (shell)fish health and have similar financial impacts to those for HABs (e.g. impacts of white spot virus

on shrimp farming in Southeast Asia ~6 US\$ billion/year) (Lafferty *et al.* 2015). Consequently a range of antimicrobial chemicals and pesticides are licensed for use in mariculture, specifically for finfish culture (Johnston & Santillo 2002; Read & Fernandes 2003). Cumulative environmental exposures to these chemicals can be significant in some coastal waters (Baker-Austin *et al.* 2008; Uyaguari *et al.* 2013) and may exceed environmental quality standards (EQSs), which can be as low as 1 part in 1 trillion for some highly potent compounds (Gilliom 2007; Watts *et al.* 2017). Impacts of antimicrobial chemicals on beneficial microbes and associated ecosystem services (e.g. nutrient cycling, water quality and HAB regulation) could be significant (Woolhouse & Ward 2013; Watts *et al.* 2017). Research on the impacts of chemicals on HAB regulation has been extremely limited to date and has generally focused on the effects of pesticides on HABs in freshwater systems (Relyea 2009; Beketov *et al.* 2013; Harris & Smith 2015; Staley *et al.* 2015).

Escapees and introduction of invasive and/or harmful species

Macro-algal blooms (seaweed blooms) leading to oxygen depletion, alteration of ecosystem biodiversity and production of certain toxins (Anderson 2009) have been shown to originate from open water suspended culture systems. For example, significant escapes may occur from *Porphyra* culturing spanning more than 40 000 km² in some instances in the South China Sea. Bloom-forming species including sea lettuce (*Ulva* spp.) and gutweed (*Enteromorpha* spp.) can cause major economic loss by inundating waterways and beaches, leading to widespread asphyxiation of organisms when the blooms biodegrade (Liu & Su 2017).

Physical alteration of habitats and hydrodynamic regimes

Reduced hydrodynamic flows are known to lead to reduced turbulence, which in turn tends to promote the blooming of dinoflagellate species, including HAB species (Smayda & Reynolds 2001). Mariculture structures, including longlines for shellfish and kelp and net pens for finfish can significantly change surface current speed and direction, induce down-welling, increase stratification and reduce water exchange in sheltered and enclosed bays (Zeng *et al.* 2015; Lin *et al.* 2016; Wartenberg *et al.* 2017). Expansion of suspended mariculture in Sanggou Bay reduced the average speed of currents by 40% and the average half-life of water exchange was prolonged by ~70% (Shi & Wei 2009). It is also possible that disturbance of sediments by aquaculture and fishing operations may promote the resuspension of HAB cysts.

Transmission of HAB species and alteration in the abundance and composition of plankton communities

Risks of HAB impacts may increase directly with the future expansion of mariculture, via the movement (relaying) of 'contaminated' shellfish stocks and equipment between sites (Hégaret *et al.* 2008), including from the coast to offshore and *vice versa*, or via regular aquaculture operations and ballast water transfers (Hallegraeff & Bolch 1991, 1992). Indirect impacts include alteration of the abundance and composition of plankton communities, including HAB competitors, parasites and grazers (Roth *et al.* 2008; Eckford-Soper *et al.* 2016). Over-intensification of mariculture can also lead to depletion of planktonic larvae (including finfish, shellfish and other invertebrates) and reduced food availability for wild shellfish populations (Gibbs 2004; Ferreira *et al.* 2014; Pastres *et al.* 2018), especially in regions with low primary productivity (Gibbs 2004; Grant *et al.* 2007). This may have consequences for negative feedback control of the abundance and composition of plankton communities by native filter feeders.

Detecting and forecasting HAB events

Maximising the profitability and environmental sustainability of mariculture requires surveillance monitoring and early warning systems, forecast-based financing and strong risk governance structures (FAO, IFAD, UNICEF, WFP and WHO, 2018). The following systems are outlined in sections 'In situ monitoring', 'Satellite remote sensing (Earth observation)' and 'Predictive modelling' below.

In situ monitoring

In situ monitoring for HAB species abundance and phycotoxin concentrations in (shell)fish is the principal method for 'official control' monitoring and safeguarding of food fish safety for human consumption in Europe, North America, Asia and Australasia. *In situ* monitoring is generally conducted via the collection and analysis of representative field samples; using microscopic analysis for phytoplankton identification and enumeration, and using mass spectrometric analysis for phycotoxin identification and quantitation. The use of autonomous *in situ* molecular (qPCR) and flow cytometry methods has also proved capable of real-time sensing of algal blooms (e.g. Campbell *et al.* 2013). These *in situ* devices can be located on smart buoys or underwater gliders (Davidson *et al.* 2014). Integrative solid-phase adsorption toxin tracking (SPATT) deployed in the field for the passive

sampling of algal toxins has also been validated recently, and improved enzyme-linked immuno-sorbent assay (ELISA)-based methods with lower detection limits for more toxins have become commercially available for both screening and routine monitoring purposes (Zhang & Zhang 2015).

In Europe, routine HAB monitoring (EU Directives 2000/60/EC and 2006/113/EC) is used to quantify HAB species abundance and phytotoxin levels (Higman *et al.* 2014). Shellfish toxin concentrations are evaluated against EU action levels triggering harvesting bans (ASP > 20 mg Domoic/epi-Domoic acid; PSP > 800 µg STX equivalents (eq.); Lipophilic toxins (DSP) OA/DTXs/PTXs together > 160 µg OA eq.; AZAs > 160 µg AZA eq.; YTXs > 3.75 mg YTX eq. – see Table 2 and underlying text for expansion of abbreviations), allowing for cross-border trade of aquaculture products. While individual HABs and their toxins vary in concentration on a seasonal basis, HAB events can occur year-round, as can aquaculture harvesting. Responsibility for 'official control' resides with respective statutory authorities within EU member countries and results are published online for each designated site. *In situ* HAB monitoring data can be combined with satellite imagery (Section 'Satellite remote sensing (Earth observation)') and numerical models (Section 'Predictive modelling') to give a better indication of HAB risk, as implemented in Ireland (Leadbetter *et al.* 2018). In some cases, more proactive monitoring can occur, such as in Scotland where a group of finfish farmers collectively pay for weekly satellite remote sensing observations of *Karenia mikimotoi* surface distributions (Davidson *et al.* 2016).

In the United States, both the National Oceanic and Atmospheric Administration (NOAA) and the Environmental Protection Agency (EPA) monitor for, and provide some indication of, impending HABs. In the Gulf of Mexico, a twice-weekly risk assessment is provided during the summer-autumn HAB season, based on a regular *in situ* monitoring programme (and using meteorological models, particularly to provide warning of toxic aerosol events, e.g., caused by *Karenia brevis*). The rest of the US coastline is monitored routinely for HAB events by a volunteer network; the 'National Phytoplankton Monitoring Network', sampling twice monthly. In some locations in the United States, more intensive programmes are in place, such as the SoundToxins programme which is funded by NOAA and Washington Sea Grant and monitors 31 sites on a weekly basis in Puget Sound in Washington State, or the California Harmful Algal Bloom Monitoring and Alert Program (Cal-HABMAP) funded by US Congress and the National Aeronautics and Space Administration (NASA; Kudela *et al.* 2015).

Across Southeast Asia, some countries operate a regular programme of shellfish monitoring (e.g. Japan, Indonesia, Vietnam, Korea), while other countries lack the resources to have a robust programme or initiate sampling when blooms are detected (e.g. Laos, Myanmar; Eong & Sulit 2015). In Australasia, monitoring effort varies, with frequent sampling of high risk locations in western Australia (Dias *et al.* 2015), but overall being less well sampled and leading to high instances of human poisonings (Hallegraef *et al.* 2017). In Chile and wider Latin America, after many intoxication events, a standardised sampling programme was developed across the region in 2009, although maintaining the network and regular sampling is dependent on continued resource availability (Cuellar-Martinez *et al.* 2018).

In scaling up from regional monitoring to a Global Ocean Observing System (GOOS) for HABs, it is recognised that there is no universal ‘one-size-fits-all’ solution, but that communication is key and stakeholders require affordable, easy to understand, real-time information, for example, in the form of spatial and temporal risk mapping (Anderson *et al.* 2019).

Satellite remote sensing (Earth observation)

The use of satellite remote sensing, alongside *in situ* sensing or ground truthing, has wide-scale potential for detecting increases in potential surface dwelling HAB species or high concentrations of all surface algae (reviewed by IOCCG, 2014; Davidson *et al.* 2016) in relation to fisheries and aquaculture/mariculture (IOCCG, 2009). Images of ocean colour from visible and infrared spectrum wavelengths can be correlated statistically with HABs events or in some cases the HAB species can be observed if they are spectrally distinct (<https://www.shelleye.org/index>; <https://www.s3eurohab.eu/en/>). For example, correlations have been found between ocean colour, chlorophyll and algal biomass (Sourisseau *et al.* 2016), with some correlations incorporating the use of artificial neural networks (El-Habashi *et al.* 2017). *K. mikimotoi* and *K. brevis* are both species that have spectral signatures that allow successful identification when they are present in large abundances (Kurekin *et al.* 2014; Shutler *et al.* 2015; El-Habashi *et al.* 2017). In general, HAB species that are detectable by remote sensing are those that form significant blooms of >1000 cells/mL at the sea surface or near-surface (e.g. *Karenia mikimotoi*, Kurekin *et al.* 2014; *Karenia brevis*, El-Habashi *et al.* 2017). Satellite imaging however cannot detect species that form harmful blooms at greater depths or at low densities of ~100 cells/L (e.g. *Dinophysis* spp.) (Reguera *et al.* 2014). Remote sensing techniques are also unable to detect HABs when observation of ocean colour is obscured by cloud cover (Maguire *et al.* 2016).

Predictive modelling

Early warning of the onset of HAB events over time scales of several days, and their likely movement and changing magnitude (i.e. relative to safe limits), would be highly beneficial to the mariculture industry, allowing proactive, rather than reactive, responses to minimise impacts on businesses, customer confidence or human health (Davidson *et al.* 2016). Immediate responses may include advanced (or delayed) harvesting of stock (limited by storage capacity and by supply chain logistics) or deployment of mitigation measures (Section ‘Analysis of options for mitigating HAB risk to mariculture’). Longer-term, more strategic business planning is dependent on knowing when harvesting bans imposed by HAB outbreaks are likely to be lifted, in order to better manage business operations, staffing and supply chains. HAB predictions based on readily available physical (hydrographical and meteorological) data offer a simple, tractable solution for forewarning mariculture operators in locations where these physical ‘forcing factors’ are principle drivers of HAB initiation. These physical models are generally better at predicting HAB initiation than HAB termination, but in any event forecasting is generally limited to 1 week in advance (Davidson *et al.* 2009; Cusack *et al.* 2016; Schmidt *et al.* 2018), which corresponds with general extent and accuracy of meteorological forecasting (Davidson *et al.* 2016). Furthermore, the majority of models, which are driven predominantly by meteorological and hydrographical processes, often ‘over-predict’ HAB duration (Davidson *et al.* 2016). This is reassuring for human safety, but not so appealing for businesses desperate for harvesting bans to be lifted, as soon as it is safe to do so. Hydrophysical models coupled with HAB population models, which also incorporate biological and geochemical processes, can improve HAB predictions, by taking into account life-history data and environmental and physiological optima for HAB species (Roelke & Buyukates 2001; McGillicuddy *et al.* 2005; Glibert *et al.* 2014b; Aleynik *et al.* 2016; Gillibrand *et al.* 2016). Modelling changes in trophic mode (autotrophy versus mixotrophy) (Lee *et al.* 2016) and interactions with other plankters, including HAB parasites and grazers (Lenes *et al.* 2013) can also help to improve predictions of bloom duration. However, increasing trophic complexity in community and ecosystem models can lead to reduced resolution of species-specific dynamics, including HAB population dynamics (Flynn & McGillicuddy 2018). Other trade-offs in implementing more elaborate ecosystem models include greater specificity (spatial limitation) of model predictions and increasing requirements for input data for model parameterisation, computational processing power and expert operators (Butenschön *et al.* 2016).

Combining bio-physical modelling of HABs with satellite remote sensing data has been used successfully in short-term national forecasting systems for public health and aquaculture protection in the US and EU for example (Kudela *et al.* 2015; Shutler *et al.* 2015; Davidson *et al.* 2016; Ruiz-Villarreal *et al.* 2016) with the potential for wider detection of HABs (Anderson *et al.* 2019). There is also the potential to extend forecasting of HAB events from days to several weeks or even months in advance, by tracking successional changes in plankton community composition over time, in conjunction with traditional *in situ* monitoring and real-time sensing of impending blooms (Roelke & Buyukates 2001; Campbell *et al.* 2013). Inter-annual predictions of HAB trends and the identification of hotspots prone to recurring HAB events are also highly beneficial for strategic marine spatial planning, including for new or expanding mariculture infrastructure. These longer-term predictions are more circumspect, as the biogeographical niches of different HAB genera or species are likely to shift with a changing climate and/or become more variable (Callaway *et al.* 2012; Wells *et al.* 2015; Global-HABs, 2017).

Analysis of options for mitigating HAB risk to mariculture

Options for mitigating HAB impacts to mariculture fall into three basic categories: (i) spatial and temporal planning of mariculture operations to avoid or minimise the risk of HABs; (ii) holistic environmental management options to minimise local HAB risk around mariculture farms (e.g. multi-species, multi-trophic, ecosystem-based options favouring nutrient assimilation and recycling and/or cultivation of species which are more resistant to, or less prone to accumulate, HAB toxins); (iii) direct interventions for controlling the presence or abundance of HAB species (physical, chemical, biological control options). The advantages of various options in each of these categories and their state of readiness for application in commercial mariculture are discussed below (Sections 'Spatial and temporal planning to minimise HAB risk', 'Holistic environmental management options for minimising HAB impacts' and 'Direct interventions for controlling HAB impacts').

Spatial and temporal planning to minimise HAB risk

Spatial planning for new mariculture infrastructure can be targeted to avoid HAB hotspots, while planning harvesting outside peak HAB risk periods can be implemented at already established/licensed mariculture farms, with both options being informed by existing HAB detection and forecasting systems (outlined in Section 'Detecting and forecasting HAB events'). Development of offshore sites

with significant exposure to tides, wind and wave action (Drumm 2010; Froehlich *et al.* 2017; Buck *et al.* 2018) can potentially mitigate HAB risks linked to mariculture itself e.g. elevation of nutrient levels, physical alteration of habitats and hydrodynamics and modification of local planktonic (and benthic) communities (Section 'Environmental impacts of mariculture and contribution to HAB risk'). However, HABs often originate naturally offshore (independently from anthropogenic activities) (Whyte *et al.* 2014; Davidson *et al.* 2016; Díaz *et al.* 2016; Gobler *et al.* 2017) and there is some evidence that some HAB species may present even greater risk here compared to inshore areas (Trainer *et al.* 2012). Regulatory policy for sustainable offshore aquaculture has only recently been developed in the United States (NOAA, 2016) and is not yet formulated and published in other countries or continents, such as New Zealand, Australia and Europe (Froehlich *et al.* 2017). Emerging guidelines for assuring minimal impacts from offshore mariculture on water quality and pelagic and benthic communities relate to: minimum water depths (twice the depth of mariculture infrastructure) and minimum water flow rates (>0.05 m/s) (Belle & Nash 2008; Froehlich *et al.* 2017). In such localities, the probability of ecological effects on neighbouring natural habitats diminishes significantly beyond a distance of 90 m (Froehlich *et al.* 2017). This distance also provides a nominal guideline for the proximity/density of neighbouring offshore mariculture infrastructure. However, some ecosystem models predict significant trophic interactions between large offshore installations and more distant coastal mariculture sites, indicating wide-ranging implications for nutrient budgets and biosecurity (spread of microbial pathogens). These ecological interactions have been modelled and verified for the large (15 km²) Ria Formosa Mariculture Park located >3 nm offshore from coastal sites in the Algarve region of Portugal (Ferreira *et al.* 2014). Ecological linkages between extensive mariculture installations and the periodic occurrence of HABs along the Algarve coast have yet to be established.

Holistic environmental management options for minimising HAB impacts

Holistic environmental management of HABs addressing causative factors (e.g. minimising nutrient inputs from land-based sources and from mariculture itself) or preserving habitats and ecosystem services that help regulate HABs, may be simpler, more effective and more environmentally friendly (WHO, 2003; Wells *et al.* 2019) than attempting to control HAB outbreaks directly (Section 'Direct interventions for controlling HAB impacts'). For example, nutrient enrichment can be managed through the use of 'extractive' shellfish and macro-algal species. Furthermore, restoration of coastal

habitats, for example with seagrasses that harbour algicidal bacteria (Inaba *et al.* 2019), or cultivation of seaweeds that secrete algicidal chemicals (Zerrifi *et al.* 2018), can also help mitigate against HABs. This follows Ecosystem Approaches to Fisheries and Aquaculture (EAF/EAA) (Soto & Aguilar-Manjarez 2009; FAO, 2018), which cover three main aspects: (i) minimising environmental impacts and waste; (ii) sustaining wider ecosystem functions and services; and (iii) promoting human well-being and equity among marine stakeholders.

(i) *Minimising environmental impacts and waste* – Shellfish and macro-algal culturing can have a positive influence on the regulation of HABs, either by reduction of high biomass blooms through filter feeding or via nutrient removal (Stadmark & Conley 2011; Petersen *et al.* 2014). Nutrient removal by mariculture, curbing eutrophication, in EU coastal waters alone is valued at US\$20 to 30 billion per year (Ferreira *et al.* 2009). Furthermore, mariculture reduces the exploitation of natural shellfish stocks, which can also help regulate HABs. For example, overfishing of shellfish around Long Island, USA, has coincided with the increased occurrence of *Aerococcus anophagefferens* brown tides (Glibert *et al.* 2005).

(ii) *Sustaining wider ecosystem functions and services* – Mariculture farms can provide sheltered nursery habitats for marine/estuarine organisms, with the potential to enhance local fisheries and to support biodiversity in neighbouring marine protected areas (Le Gouvello *et al.* 2017). Maintaining biodiversity is important, since impoverishment of planktonic species and reduced species succession have been correlated with increased HAB risk. In some cases, such community changes can forewarn HAB outbreaks several months before the detection of the HAB species (e.g. *Microcystis* sp.; Roelke & Buyukates 2001).

(iii) *Promoting human well-being and equity among marine stakeholders* – Marine spatial planning is required to effectively locate mariculture and fisheries conservation areas and avoid conflicts with other uses of the marine environment. To facilitate planning, environmental models can be used to assess nutrient budgets, productivity versus eutrophication risk, the risk of transmission of pathogens, pests associated with mariculture (Ferreira *et al.* 2014; Pastres *et al.* 2018) and the risk of advection of HABs to mariculture sites (Dabrowski *et al.* 2016; Paterson *et al.* 2017).

A promising approach for delivering on each of these EAA/EAF aspects, including the potential to minimise HAB risk, is integrated multi-trophic aquaculture (IMTA) (Wartenberg *et al.* 2017). IMTA employs cultureable ‘extractive’ species (e.g. suspended bivalve shellfish and macroalgae, and benthic deposit feeders) to remove/reuse waste nutrient material discarded from the culturing of ‘fed’ species (finfish and crustaceans) thereby providing a self-sustaining and more productive food web (Figure 2)

(Soto 2009; Troell *et al.* 2009; Chopin *et al.* 2012). Macroalgae can also play a direct role in inhibiting the growth of microalgae, including HAB species, through competition for nutrients (Soto 2009; Holdt & Edwards 2014), inhibitory allelopathy (Tang & Gobler 2011; Ben Gharbia *et al.* 2017; Zerrifi *et al.* 2018), and/or by reducing light penetration (Zhou *et al.* 2006; Wang *et al.* 2007; Yang *et al.* 2015).

Further developments in IMTA, including deploying aquaculture species that are less sensitive to, or less likely to accumulate, toxins from locally re-occurring HAB species, are likely to be required to maximise benefits in terms of mitigating against HAB impacts. The long-term sustainability of IMTA for mitigating HAB risk with climate change also requires further research (Wells *et al.* 2019). For example, China has some of the world’s largest and longest established IMTA systems, including a multi-trophic system established in 1996 in Sanggou Bay, Yellow Sea (Fang *et al.* 2016). Since 2010, however, Sanggou Bay has regularly experienced brown tides of *A. anophagefferens* (Kong *et al.* 2012). Coincidentally, large-scale *A. anophagefferens* brown tides extending over 3000 km² have occurred in the north western Bohai Sea each year in early summer since 2009 and have caused significant negative impacts on scallop (*Argopecten irradians*) culture (Zhang *et al.* 2012). Other HAB species including *Karenia mikimotoi* and *Prorocentrum donghaiense* also continue to form annual blooms in nearshore waters of the Yellow Sea and neighbouring East China Sea (Li *et al.* 2009), with *K. mikimotoi* causing substantial losses to mariculture from 2005 to 2015 (Liu & Su 2017).

Direct interventions for controlling HAB impacts

Physical and chemical control methods can remove HABs efficiently and are used operationally as a last resort in mariculture, but they can be costly, lack specificity to HABs, and are generally less effective in coastal situations in comparison with enclosed or semi-enclosed aquatic systems. Alternatively, biological control methods can be potentially more specific for individual HAB species, minimising impact on other non-target species, but they are more difficult to constrain in non-enclosed systems and have not progressed beyond laboratory or field trials for mariculture applications (reviewed in NOAA, 2015; Sellner & Rensel 2018; Sun *et al.* 2018; Gallardo-Rodriguez *et al.* 2019).

Physical control methods include the use of barriers or skirts, for example, around fish net pens and/or the removal of HAB cells by water column mixing, filtering, flocculation, settlement, sediment burial and dredging, or HAB cell lysis using ultrasound (Sellner & Rensel 2018). Water column mixing using water or air pumping systems leads to disruption of thermal stratification and impairment of algal buoyancy or alteration of their daily




Figure 2 Integrated multi-trophic aquaculture (IMTA). POM – particulate organic matter; DIN – dissolved inorganic nitrogen; F/P-F – faeces/pseudo-faeces. IMTA incorporating suspended filter-feeding shellfish, and benthic deposit feeding shellfish can reduce the proliferation of HABs and recycle POM (capable of fuelling HAB growth) associated with ‘fed’ species (finfish and crustaceans). Suspended macroalgae can also reduce the growth of microalgae, including HAB species, through shading, competition for nutrients (e.g. fine POM and DIN) and inhibitory allelopathy.

migration patterns, removing them from the photic zone and preventing photosynthesis. Direct cell removal from the water column can be achieved by hydrodynamic separation, centrifugation, pump filtration, plankton net trawling or membrane filtration. A measure which has proven effective for HAB control in the open sea has been the use of clays to induce bloom flocculation. As considerable quantities of clay are needed, from 100 to 400 g/m² (Park *et al.* 2013), physical resuspension of local sediments or importation on ships are a practical solutions. Subsequent flocculation, sinking and burial of HAB cells and/or cysts can be followed by dredging and physical or chemical treatment before discharging the sediments back to the removal site (NOAA, 2015; Sellner & Rensel 2018). Potential drawbacks include the removal of non-harmful algae. More efficient flocculation can be achieved by spraying the sea surface with modified clays containing inorganic (e.g. aluminium sulphate or polyaluminium chloride) or organic (e.g. polyacrylamide or chitosan) modifiers, which can be up to 100 times more efficient in adsorbing HAB cells (and other

plankters) than natural clay sediments. This enables a reduction in application levels and time windows – reducing the risk of clay build-up and helping to reduce impacts on non-blooming (non-HAB) species (reviewed in Gallardo-Rodríguez *et al.* 2019). Furthermore, modified clays have been shown to kill HAB cells (Beaulieu *et al.* 2003), adsorb and remove extracellular HAB toxins (Pierce *et al.* 2004; Seger *et al.* 2015, 2017) and particulate nutrients (Yu *et al.* 2017), and to also reduce HAB toxin accumulation in benthic filter-feeding bivalves (Yu *et al.* 2017). Consequently, they have been used in Japan (Shirota 1989) and employed as a standard method for controlling HABs in China, since 2014 (Yu *et al.* 2017). A remaining concern, preventing uptake of these physical control methods in other countries, is their lack of specificity for controlling harmful species and possible unknown impacts on other phytoplankton and the ecosystem as a whole.

More direct chemical treatments for controlling HABs include the use of natural biosurfactants, biocides or allelochemicals (e.g. biochemical extracts from macroalgae), or

the use of synthetic chemicals, including hydrogen peroxide and isolated algicidal compounds, or metallic compounds such as copper sulphate. These various chemicals (metals and organic compounds) can interfere with HAB cell survival (algicidal chemicals), growth and reproduction (algistatic chemicals) through a variety of mechanisms (NOAA, 2015; Gallardo-Rodríguez *et al.* 2019). Biochemicals are advantageous in terms of their higher diversity, biodegradability and, in some cases, specificity and potentially lower toxicity to the wider environment (Ahn *et al.* 2003). Although many effective aqueous algicidal treatments exist, few are approved for use in open marine systems, due to environmental concerns, although some have restricted use in antifouling paints and surface treatments (NOAA, 2015; Gallardo-Rodríguez *et al.* 2019). Several biocidal chemicals have been tested and approved for use in mariculture, for controlling shellfish and finfish pathogens or parasites (Johnston & Santillo 2002; Read & Fernandes 2003) and some of these may be effective in killing some HAB species.

Biological control measures include the application of microbial (viral, bacterial, fungal and/or protistan) parasites that infect HABs and play a significant role in the natural termination of major blooms (Brussaard 2004; Chambouvet *et al.* 2008; Roth *et al.* 2008; Jones *et al.* 2011; Demuez *et al.* 2015; Pokrzywinski *et al.* 2017). Algicidal and growth inhibitory bacteria and viruses have potential for controlling HABs, due to their ability to replicate rapidly and target-specific hosts (Bibak & Hosseini 2013; Sun *et al.* 2018). However, it is possible for these parasites to be too specific, rendering them unable to infect different genetic strains of HAB species, or adapt to changing environmental conditions (Sun *et al.* 2018; Gallardo-Rodríguez *et al.* 2019). Therefore, rather than using single cultured microbial species, employing a range of microbes may be more effective. Aggregates (biofilms) immobilised on substrates may be more effective in reducing HAB cell density by inhibiting HAB cell growth via nutrient uptake and allelochemical secretion, and causing cell lysis (Alex *et al.* 2014; Sun *et al.* 2018). Research is needed to quantify the release of toxins following HAB cell lysis and the potential for microbes to degrade them. Further research is also needed to isolate, purify and identify microbial allelochemicals/exudates and to demonstrate their efficacy for controlling different HAB species and genetic strains, while incurring minimal effects on non-harmful algae and other marine organisms, including cultured shellfish and finfish species (NOAA, 2015, Sun *et al.* 2018). Other potential biological interventions include selective breeding of shellfish with resistance to HAB toxins and using them as HAB biofilters and bioremediators (NOAA, 2015). Unquantified biosecurity risks for biological control measures currently prevent their operational use in controlling HABs at mariculture sites.

Conclusions and recommendations

Marine aquaculture (mariculture) is playing an increasingly important role in global food security. One of the most significant risks to mariculture expansion, both inshore and offshore, is the occurrence of Harmful Algal Blooms (HABs).

Global impacts from HABs on mariculture (due to finfish or shellfish mortality, poisoning of human consumers and preventative harvesting bans) currently amount to something in the region of 8 US\$ billion/year; however, HAB risk assessment is not a standard requirement in the planning and classification of mariculture sites. This is, in part, because HABs are natural phenomena, and because risk factors are diverse, varying greatly both spatially and temporally. For example, HABs may originate offshore, far from anthropogenic activities, and can be advected over large distances to other areas conducive for HAB development. Further research is required to guide and enable pre-emptive measures for mitigating HAB risks, including the strategic siting of mariculture infrastructure and scheduling of harvests.

Adaptive management of HAB risk, involving the prediction of HAB events and the tactical use of appropriate and approved physical, chemical and/or biological control measures, is needed as part of the sustainable development of mariculture. However, successful application requires improved understanding on the efficacy and biosafety/specificity of the available options. There is a need also for improved understanding on the interactions among physical forcing factors (meteorological and oceanographical), and chemical (nutrient) and biological (community) factors, in order to predict where and when blooms are most likely to occur. In support of this, research should exploit the widespread occurrence of HABs, which provides opportunities for comparative assessments of HAB drivers around the world, including the extent to which HAB species, their population dynamics, and community interactions show similarities in responses within comparable ecosystem types. There is considerable scope to capitalise on advances in automation and (bio)sensor (DNA, RNA, protein and metabolite)-based technologies, with applications in: real-time, *in situ* monitoring of HAB population dynamics; defining physiological processes and underlying regulatory gene networks linked to growth and/or toxin production in HAB species; and building robust, mechanistic models for predicting HAB events.

Harmful algal bloom risks are generally perceived to be higher at coastal sites, which experience nutrient enrichment from agricultural runoff and municipal effluent discharges. Winds and tides can also transport and accumulate HABs into coastal areas, including sheltered embayments, where less turbulent and warmer waters are

conducive for the growth of various HAB species. In these and other areas with low water exchange rates, mariculture itself can have a significant influence on HAB risk by affecting local water quality (e.g. nutrient eutrophication levels), hydrodynamics (artificial structures reducing water circulation) and plankton communities (e.g. through selective filter feeding by shellfish). More studies are required to quantify HAB risks against each of the above factors and their interactions and the degree to which they are influenced by different types of mariculture.

Harmful algal bloom risks associated with nutrient enrichment and eutrophication (from terrestrial sources and mariculture itself) may be mitigated by establishing mariculture sites offshore, away from the coast and/or in areas with high horizontal water exchange rates and vertical mixing. Greater understanding is required on how hydrodynamic conditions (e.g. influenced by wind, waves, tides) and bathymetry (water depth) influence dispersal versus local deposition and resuspension of nutrients and HAB propagules/cysts.

Further capacity for HAB mitigation is offered by integrated multi-trophic aquaculture (IMTA), which employs extractive bivalve shellfish and macroalgae alongside fed finfish and crustaceans, in order to recycle nutrients, thus maximising productivity and water quality simultaneously. Macroalgae (in addition to filter-feeding shellfish) can also have a direct influence on local plankton community composition and abundance – via nutrient competition, light shading and allelochemical mechanisms. Further research is required to understand how IMTA systems could be further optimised for the additional purpose of HAB attenuation, through selection of suitable, resilient finfish, shellfish and macroalgal species, and appropriate spatial deployment and stocking densities.

A key remaining question for mariculture, both inshore and offshore, is 'How will HAB risk transpire in a future warmer climate, typified by increased sea surface temperatures and water column stratification, or alternatively in a future characterised by increased atmospheric energy and more turbulent waters?' Climate change is also likely to be accompanied by HAB range extensions towards the poles. To address these issues, collaborative effort is needed that seeks to unify research themes on 'HABs, climate change and aquaculture/mariculture', as exemplified by Global-HAB, an international programme sponsored jointly by the Scientific Committee on Oceanic Research (SCOR) and the Intergovernmental Oceanographic Commission (IOC) of UNESCO.

Acknowledgements

This review was funded by the European Maritime and Fisheries Fund (EMFF) grant ENG2360: Assessing and

mitigating the future risks of harmful algal blooms (HABs) to wild fisheries and aquaculture. The authors have no industrial links. Brown and Lilley undertook the literature review, which was led by Brown. Expert contributions and peer reviews of manuscript drafts were provided by each of the co-authors: Lilley – marine ecology; Shutler – remote sensing, water quality and aquaculture; Lowe – microbiology; Artioli and Torres – hydrodynamic and ecosystem modelling; Berdalet – HABs; and Tyler – aquaculture. Shutler has co-authored reports by the International Ocean Colour Coordinating group (IOCCG), Berdalet is the acting chair of the GlobalHAB Scientific Steering Committee (www.globalhab.info), and this paper contributes to the implementation of the programme objectives concerning HABs and Aquaculture; Tyler is co-director of the Centre for Sustainable Aquaculture Futures and co-PI on RCUK's Aquaculture Research Collaborative Hub (ARCH-UK).

References

- Adams C (2017) Red tide causes economic losses to Southwest Florida industry. UF/IFAS Florida Shellfish Aquaculture Featured News. January 13, 2017. URL (accessed April 2019): <http://shellfish.ifas.ufl.edu/news/red-tide-causes-economic-losses-sw-florida-industry/>
- Adams CM, Larkin SL, Hoagland P, Sancewich B (2018) Assessing the economic consequences of harmful algal blooms: a summary of existing literature, research methods, data, and information gaps. In Shumway SE, Burkholder JM, Morton SL (eds). *Harmful Algal Blooms: A Compendium Desk Reference*. First Edition. Chapter 8, 696 pp. ©2018 JohnWiley & Sons Ltd. Published 2018 by John Wiley & Sons Ltd, Hoboken, NJ.
- Ahn CY, Joung SH, Jeon JW, Kim HS, Yoon BD, Oh HM (2003) Selective control of cyanobacteria by surfactin-containing culture broth of *Bacillus subtilis* C1. *Biotechnology Letters* **25**: 1137–1142.
- Alex D, Mathew A, Sukumaran R (2014) Esterases immobilized on aminosilane modified magnetic nanoparticles as a catalyst for biotransformation reactions. *Bioresource Technology* **167**: 547–550.
- Aleynik D, Dale AC, Porter M, Davidson K (2016) A high resolution hydrodynamic model system suitable for novel harmful algal bloom modelling in areas of complex coastline and topography. *Harmful Algae* **53**: 102–117.
- Amin SA, Hmelo LR, van Tol HM, Durham BP, Carlson LT, Heal KR *et al.* (2015) Interaction and signalling between a cosmopolitan phytoplankton and associated bacteria. *Nature* **522**: 98–101.
- Andersen NG, Hansen PJ, Engell-Sørensen K, Nørremark LH, Andersen P, Lorenzen E *et al.* (2015) Ichthyotoxicity of the microalga *Pseudochattonella farcimen* under laboratory and field conditions in Danish waters. *Diseases of Aquatic Organisms* **116**: 165–172.

- Anderson DM (2009) Approaches to monitoring, control and management of harmful algal blooms (HABs). *Ocean and Coastal Management* **52**: 342.
- Anderson DM (2012) HABs in a changing world: a perspective on harmful algal blooms, their impacts, and research and management in a dynamic era of climactic and environmental change. *Harmful Algae* **2012**: 3–17.
- Anderson DM, Kulis DM, Doucette GJ, Gallagher JC, Balech E (1994) Biogeography of toxic dinoflagellates in the genus *Alexandrium* from the north eastern United States and Canada. *Marine Biology* **120**: 467–478.
- Anderson DM, Glibert PM, Burkholder JM (2002) Harmful Algal Blooms and eutrophication: nutrient sources, composition, and consequences. *Estuaries* **25**: 704–726.
- Anderson DM, Burkholder JM, Cochlan WP, Glibert PM, Gobler CJ, Heil CA *et al.* (2008) Harmful algal blooms and eutrophication: examining linkages from selected coastal regions of the United States. *Harmful Algae* **8**: 39–53.
- Anderson DM, Cembella AD, Hallegraeff GM (2012a) Progress in understanding harmful algal blooms: paradigm shifts and new technologies for research, monitoring, and management. *Annual Reviews in Marine Science* **4**: 143–176.
- Anderson DM, Alpermann TJ, Cembella AD, Collos Y, Masseret E, Montresor M (2012b) The globally distributed genus *Alexandrium*: multifaceted roles in marine ecosystems and impacts on human health. *Harmful Algae* **14**: 10–35.
- Anderson CR, Berdalet E, Kudela RM, Cusack CK, Silke J, O'Rourke E *et al.* (2019) Scaling up from regional case studies to a global Harmful Algal Bloom observing system. *Frontiers in Marine Science* **6**: 250. <https://doi.org/10.3389/fmars.2019.00250>
- Arzul G, Seguel M, Clement A (2001) Effect of marine animal excretions on differential growth of phytoplankton species. International Council for the Exploration of the Sea (ICES). *Journal of Marine Science* **58**: 386–390.
- ASIMUTH (2014) Applied simulations and Integrated modelling for the understanding of toxic and harmful algal blooms. Final Report Summary, FP7-SPACE Project ID: 261860.
- Baker-Austin C, McArthur JV, Tuckfield RC, Najarro M, Lindell AH, Gooch J (2008) Antibiotic resistance in the shellfish pathogen *Vibrio parahaemolyticus*, isolated from the coastal water and sediment of Georgia and South Carolina, USA. *Journal of Food Protection* **71**: 2552–2558.
- Basti L, Hégaret H, Shumway SE (2018) Harmful Algal Blooms and Shellfish. In Shumway SE, Burkholder JM, Morton SL, eds. *Harmful Algal Blooms a Compendium Desk Reference*, pp. 135–191. John Wiley & Sons Inc., Hoboken, NJ.
- Beaulieu SE, Sengco MR, Anderson DM (2003) Using clay to control harmful algal blooms: deposition and resuspension of clay/algal flocs. *Harmful Algae* **4**(1): 123–138.
- Beketov MA, Kefford BJ, Schäfer RB, Liess M (2013) Pesticides reduce regional biodiversity of stream invertebrates. *Proceedings of the National Academy of Sciences USA* **110**: 11039–11043.
- Belle SM, Nash CE (2008) Better management practices for net-pen aquaculture. In: Tucker CS, Hargreaves JA (eds) *Environmental Best Management Practices for Aquaculture*, pp. 261–330. Wiley-Blackwell, Oxford, UK.
- Ben Gharbia H, Kéfi-Daly Yahia O, Cecchi P, Masseret E, Amzil Z, Herve F *et al.* (2017) New insights on the species-specific allelopathic interactions between macrophytes and marine HAB dinoflagellates. *PLoS ONE* **12**: e0187963.
- Berdalet E, Fleming LE, Gowen R, Davidson K, Hess P, Backer LC *et al.* (2016) Marine Harmful Algal Blooms, human health and wellbeing: challenges and opportunities in the 21st century. *Journal of the Marine Biology Association of the United Kingdom* **96** (Special Issue 01): 61.
- Berdalet E, Montresor M, Reguera B, Roy S, Yamazaki H, Cembella A *et al.* (2017) Harmful algal blooms in fjords, coastal embayments, and stratified systems: recent progress and future research. *Oceanography* **30**(1): 46–57.
- Bibak M, Hosseini SA (2013) Review ways to control harmful algal bloom (HAB). *World Journal of Fish and Marine Sciences* **5**: 42–44.
- Boalch GT (1979) The dinoflagellate bloom on the coast of south west England, August-September 1978. *Journal of the Marine Biological Association of the United Kingdom* **59**: 515–517.
- Bouwman L, Beusen A, Glibert PM, Overbeek C, Pawlowski M, Herrera J *et al.* (2013) Mariculture: significant and expanding cause of coastal nutrient enrichment. *Environmental Research Letters* **8**: 044026.
- Bresnan E, Davidson K, Edwards M, Fernand L, Gowen R, Hall A *et al.* (2013) Impacts of climate change on harmful algal blooms. *MCCIP Science Review* **2013**: 236–243.
- Brussaard CPD (2004) Viral control of phytoplankton populations – a review. *Journal of Eukaryotic Microbiology* **51**: 125–138.
- Buck BH, Troell MF, Krause G, Angel DL, Grote B, Chopin T (2018) State of the art and challenges for offshore integrated multi-trophic aquaculture (IMTA). *Frontiers in Marine Science* **5**: 165.
- Burkholder JM (1998) Implications of harmful microalgae and heterotrophic dinoflagellates in management of sustainable marine fisheries. *Ecological Application* **8**: S37–S62.
- Butenschön M, Clark J, Aldridge JN, Allen JI, Artioli Y, Blackford J *et al.* (2016) ERSEM 15.06: a generic model for marine biogeochemistry and the ecosystem dynamics of the lower trophic levels. *Geoscience Model Development* **9**: 1293–1339.
- Calbet A (2008) The trophic roles of microzooplankton in marine systems. *ICES Journal of Marine Science* **65**: 325–331.
- California Ocean Science Trust (2016). *Framing the science around harmful algal blooms and California fisheries: Scientific insights, recommendations and guidance for California*. 2016. California Ocean Science Trust, Oakland, CA. (accessed November 2019) Available from URL : <https://fisheries.legislature.ca.gov/sites/fisheries.legislature.ca.gov/files/u8/2016-10-25%20HABs%20and%20CA%20Fisheries%20Science%20Guidance.pdf>

- Callaway R, Shinn AP, Grenfell SE, Bron JE, Burnell G, Cook EJ *et al.* (2012) Review of climate change impacts on marine aquaculture in the UK and Ireland. *Aquatic Conservation: Marine and Freshwater Ecosystems* **22**: 389–421.
- Campbell DA, Kelly MS, Busman M, Wiggins E, Fernandes TF (2003) Impact of preparation method on gonad domoic acid levels in the scallop, *Pecten maximus* (L.). *Harmful Algae* **2**: 215–222.
- Campbell L, Henrichs DW, Olson RJ, Sosik HM (2013) Continuous automated imaging-in-flow cytometry for detection and early warning of *Karenia brevis* blooms in the Gulf of Mexico. *Environmental Science and Pollution Research* **20**: 6896–6902.
- Castberg T, Torgersen T, Aasen J, Aune T, Naustvoll L-J (2004) Diarrhetic shellfish poisoning toxins in *Cancer pagurus* Linnaeus, 1758 (Brachyura, Canceridae) in Norwegian waters. *Sarsia* **89**: 311–317.
- Chambouvet A, Morin P, Marie D, Guillou L (2008) Control of toxic marine dinoflagellate blooms by serial parasitic killers. *Science* **322**: 1254–1257.
- Chopin T, Cooper JA, Reid G, Cross S, Moore C (2012) Open-water integrated multi-trophic aquaculture: environmental biomitigation and economic diversification of fed aquaculture by extractive aquaculture. *Reviews in Aquaculture* **4**: 209–220.
- Clément A, Lincoqueo L, Saldivia M, Brito CG, Muñoz F, Fernández C *et al.* (2016) Exceptional summer conditions and HABs of *Pseudochattonella* in Southern Chile create record impacts on salmon farms. *Harmful Algae News* **53**: 1–3.
- Cochlan WP, Trainer VL, Trick CG, Wells ML, Bill BD, Eberhart B-TL (2012). *Heterosigma akashiwo* in the Salish Sea: defining growth and toxicity leading to fish kills. In: Kim HG, Reguera B, Hallegraeff G, Lee CK, Han MS, Choi JK (eds), *Harmful Algae 2012. 15th International Conference on Harmful Algae*, pp. 203–206. International Society for the study of harmful algae, Changwon, Gyeongnam, Korea.
- CTA (2013) Fish-farming: the new driver of the blue economy, Briefing n. 32. Technical Centre for Agricultural and Rural Cooperation ACP-EU (CTA). URL (accessed 25 January 2019): https://europa.eu/capacity4dev/file/15656/download?to ken=J7_MU3_d
- Cuellar-Martinez T, Ruiz-Fernández AC, Alonso-Hernández C, Amaya-Monterrosa O, Quintanilla R, Carrillo-Ovalle HL *et al.* (2018) Addressing the Problem of Harmful Algal Blooms in Latin America and the Caribbean- A Regional Network for Early Warning and Response. *Frontiers in Marine Science* **5**: 409.
- Cusack C, Dabrowski T, Lyons K, Berry A, Westbrook G, Salas R *et al.* (2016) Harmful algal bloom forecast system for SW Ireland. Part II: are operational oceanographic models useful in a HAB warning system. *Harmful Algae* **53**: 86–101.
- Dabrowski T, Lyons K, Nolan G, Berry A, Cusack C, Silke J (2016) Bloom forecast system for SW Ireland. Part I: description and validation of an operational forecasting model. *Harmful Algae* **53**: 64–76.
- Davidson K (2014) The challenges of incorporating realistic simulations of marine protists in biogeochemically based mathematical models. *Acta Protozoologica* **53**(1): 129–138.
- Davidson K, Miller PI, Wilding TA, Shutler J, Bresnan E, Kennington K *et al.* (2009) A large and prolonged bloom of *Karenia mikimotoi* in Scottish waters in 2006. *Harmful Algae* **8**: 349–361.
- Davidson K, Gowen RJ, Tett P, Bresnan E, Harrison PJ, McKinney A *et al.* (2012) Harmful algal blooms: how strong is the evidence that nutrient ratios and forms influence their occurrence? *Estuarine, Coastal and Shelf Science* **115**: 399e413.
- Davidson K, Gowen RJ, Harrison PJ, Fleming LE, Hoagland P (2014) Anthropogenic nutrients and harmful algae in coastal waters. *Journal of Environmental Management* **146**: 206–216.
- Davidson K, Baker C, Higgins C, Higman W, Swan S, Veszelovszki A *et al.* (2015) Potential threats posed by new or emerging marine biotoxins in UK waters and examination of detection methodologies used for their control: cyclic imines. *Marine Drugs* **13**: 7087–7112.
- Davidson K, Anderson DM, Mateus M, Reguera B, Silke J, Sourisseau M *et al.* (2016) Forecasting the risk of harmful algal blooms. *Harmful Algae* **53**: 1–7.
- De Silva SS, Soto D (2009) Climate change and aquaculture: potential impacts, adaptation and mitigation. In: Cochrane K, De Young C, Soto D, Bahri T (eds). *FAO Fisheries and Aquaculture Technical Paper No. 530*. FAO, Rome, Italy. Accessed 25 Aug 2019: <http://www.fao.org/3/i0994e/i0994e00.htm>
- Dempster T, Sanchez-Jerez P, Bayle-Sempere JT, Kingsford M (2004) Extensive aggregations of wild fish at coastal sea-cage fish farms. *Hydrobiologia* **525**: 245–248.
- Demuez M, González-Fernández C, Ballesteros M (2015) Algicidal microorganisms and secreted algicides: new tools to induce microalgal cell disruption. *Biotechnology Advances* **33** (8): 1615.
- Dias PJ, Munoz J, Huisman JM, McDonald JI (2015) Biosecurity monitoring of Harmful Algal Bloom (HAB) species in Western Australian waters: first confirmed record of *Alexandrium catenella* (Dinophyceae). *BioInvasions Records* **4**: 233–241.
- Diaz RJ, Rosenberg R (2008) Spreading dead zones and consequences for marine ecosystems. *Science* **321**: 926–929.
- Díaz PA, Ruiz-Villarreal M, Pazos Y, Moita T, Reguera B (2016) Climate variability and *Dinophysis acuta* blooms in an upwelling system. *Harmful Algae* **53**: 145–159.
- Dickey TD (2001) The role of new technology in advancing ocean biogeochemical research. *Oceanography* **14**: 108–120.
- Drumm A (2010) *Evaluation of the Promotion of Offshore Aquaculture Through a Technology Platform (OATP)*. Marine Institute, Galway, Ireland. pp. 46 URL (accessed 28 August 2019): www.offshoreaquaculture.com/docs/OATP_Final_Publishable_report.pdf
- Directive 2006/113/EC of the European Parliament and of the Council of 12 December 2006 on the quality required of shellfish waters <http://data.europa.eu/eli/dir/2006/113/oj>
- Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for

- Community action in the field of water policy. <http://data.europa.eu/eli/dir/2000/60/oj>
- Eckford-Soper LK, Bresnan E, Lacaze J-P, Green DH, Keith Davidson K (2016) The competitive dynamics of toxic *Alexandrium fundyense* and non-toxic *Alexandrium tamarense*: The role of temperature. *Harmful Algae* **53**: 135–144.
- El-Habashi A, Duran CM, Lovko V, Tomlinson MC, Stumpf RP, Ahmed S (2017) Satellite retrievals of *Karenia brevis* harmful algal blooms in the West Florida shelf using neural networks and impacts of temporal variabilities. *Journal of Applied Remote Sensing* **11**: 032408.
- Eong YS, Sulit VT (2015) Monitoring and identification of Harmful Algal Blooms in Southeast Asia to support SDG 14.1. *Fish for the People, Magazine of the Southeast Asian Fisheries Development Centre* **15**: 39–46.
- Espinosa EP, Perrigault M, Ward JE, Shumway SE, Allam B (2009) Implication of lectins associated with feeding organs in particle selection in the oyster *Crassostrea virginica*. *Biological Bulletin* **217**: 130–141.
- Estrada M, Berdalet E (1997) Phytoplankton in a turbulent world. In: Marrasé C, Saiz E, Redondo JM (eds), *Lectures on Plankton and Turbulence*. *Scientia Marina* **61**: 125–140
- Fang J, Zhang J, Xiao T, Huang D, Liu S (2016) Integrated multi-trophic aquaculture (IMTA) in Sanggou Bay, China. *Aquaculture Environment Interactions* **8**: 201–205.
- FAO (2012) The State of the World Fisheries and Aquaculture 2012. FAO Fisheries and Aquaculture Department, FAO. URL (accessed 25 August 2019): <http://www.fao.org/docrep/016/i2727e/i2727e.pdf>
- FAO (2015) *FAO Global Aquaculture Production Statistics Database Updated to 2013: Summary Information*. Food and Agriculture Organization of the United Nations, Rome, Italy. URL (accessed 25 August 2019): <http://www.fao.org/3/a-i4899e.pdf>
- FAO (2018) *The State of World Fisheries and Aquaculture 2018: Meeting the Sustainable Development Goals*. 200 pp. Food and Agriculture Organization of the United Nations (UN FAO), Rome, Italy. URL (accessed 25 August 2019): <http://www.fao.org/3/i9540en/i9540en.pdf>
- FAO, IFAD, UNICEF, WFP and WHO (2018) *The State of Food Security and Nutrition in the World 2018*. Building climate resilience for food security and nutrition. UN Food and Agriculture Organisation, Rome, Italy. URL (accessed 25 August 2019): <http://www.fao.org/3/i9553en/i9553en.pdf>
- Farrell H, Gentien P, Fernand L, Lunven M, Reguera B, González-Gil S *et al.* (2012) Scales characterising a high density thin layer of *Dinophysis acuta* Ehrenberg and its transport within a coastal jet. *Harmful Algae* **15**: 36–46.
- Fehling J, Green DH, Davidson K, Bolch CJ, Bates SB (2004) Domoic acid production by *Pseudo-nitzschia seriata* (Bacillariophyceae) in Scottish waters. *Journal of Phycology* **40**: 622–630.
- Ferreira JG, Sequeira A, Hawkins AJS, Newton A, Nickell T, Pastres R *et al.* (2009) Analysis of coastal and offshore aquaculture: application of the FARM model to multiple systems and shellfish species. *Aquaculture* **292**: 129–138.
- Ferreira JG, Saurel C, Lencart E, Silva JD, Nunes JP, Vazquez F (2014) Modelling of interactions between inshore and offshore aquaculture. *Aquaculture* **426–427**: 154–164.
- Flaherty KE, Landsberg JH (2011) Effects of a persistent red tide (*Karenia brevis*) bloom on community structure and species-specific relative abundance of nekton in a Gulf of Mexico estuary. *Estuaries and Coasts* **34**(2): 417–439.
- Flynn KJ, McGillicuddy DJ Jr (2018) Modeling marine Harmful Algal Blooms: current status and future prospects. In: Shumway SE, Burkholder JM, Morton SL (eds) *Harmful Algal Blooms a Compendium Desk Reference*, pp. 115–134. John Wiley & Sons Inc., Hoboken, NJ.
- Food and Agriculture Organisation of the United Nations (FAO) (2012) *The State of World Fisheries and Aquaculture 2012*. Food and Agriculture Organisation of the United Nations, Rome, Italy. URL (accessed 25 August 2019): www.fao.org/docrep/016/i2727e/i2727e00.htm
- Franklin DJ, Brussaard CPD, Berges JA (2006) What is the role and nature of programmed cell death in phytoplankton ecology? *European Journal of Phycology* **41**(1): 1–14.
- Friedman MA, Fernandez M, Backer LC, Dickey RW, Bernstein J, Schrank K *et al.* (2017) An updated review of ciguatera fish poisoning: clinical, epidemiological, environmental, and public health management. *Marine Drugs* **15**: E72.
- Froehlich HE, Gentry RR, Halpern BS (2016) Synthesis and comparative analysis of physiological tolerance and life-history growth traits of marine aquaculture species. *Aquaculture* **460**: 75–82.
- Froehlich HE, Smith A, Gentry RR, Halpern BS (2017) Offshore Aquaculture: I Know It When I See It. *Frontiers in Marine Science* **4**: 154.
- Fukuyo Y, Imai I, Kodama M, Tamai K (2002) Red tides and other harmful algal blooms in Japan. In: Taylor FJR, Trainer VL (eds), *Harmful Algal Blooms in the PICES Region of the North Pacific*. PICES Report No. 23, pp.7–20. North Pacific Marine Science Organization (PICES), Sidney, Canada.
- Furey A, O'Doherty S, O'Callaghan K, Lehane M, James KJ (2010) Azaspiracid poisoning (AZP) toxins in shellfish: toxicological and health considerations. *Toxicon* **56**(2): 173–190.
- Gachon CMM, Sime-Ngando T, Strittmatter M, Chambouvet A, Kim GH (2010) Algal diseases: spotlight on a black box. *Trends in Plant Science* **15**: 633–640.
- Gallardo-Rodríguez JJ, Astuya-Villalón A, Llanos-Rivera A, Avello-Fontalba V, Ulloa-Jofré V (2019) A critical review on control methods for harmful algal blooms. *Reviews in Aquaculture* **2019**: 661–684.
- Gentien P, Lunven M, Lazure P, Youenou A, Crassous MP (2007) Motility and autotoxicity in *Karenia mikimotoi* (Dinophyceae). *Philosophical Transactions of the Royal Society London B* **362**: 1937–1946.
- Gentry RR, Lester SE, Kappel CV, White C, Bell TW, Stevens J *et al.* (2016) Offshore aquaculture: spatial planning principles for sustainable development. *Ecology and Evolution* **7**(2): 733–743.

- Gentry RR, Froehlich HE, Grimm D, Kareiva P, Parke M, Rust M (2017) Mapping the global potential for marine aquaculture. *Nature Ecology and Evolution* **1**: 1317–1324.
- GESAMP (2001) *Group of Experts on the Scientific Aspects of Marine Pollution (GESAMP) Reports and Studies*. United Nations Environment Programme, Nairobi: 2001. Global Environment Outlook 3, EARTH SCAN, Past, present and future perspectives. Protecting the oceans from land-based activities. Land-based sources and activities affecting the quality and uses of the marine, coastal and associated freshwater environment. No. 71.11.
- Gibbs MT (2004) Interactions between bivalve shellfish farms and fishery resources. *Aquaculture* **240**: 267–296.
- Gillibrand PA, Siemerling B, Miller PI, Davidson K (2016) Individual-based modelling of the development and transport of a *Karenia mikimotoi* bloom on the North-west European continental shelf. *Harmful Algae* **53**: 118–134.
- Gilliom RJ (2007) Pesticides in the U.S. streams and groundwater. *Environmental Science and Technology* **41**: 3408–3414.
- Glibert PM, Anderson DM, Gentien P, Granéli E, Sellner KG (2005) The global, complex phenomena of harmful algal blooms. *Oceanography* **18**(2): 136–147.
- Glibert PM, Maranger R, Sobota DJ, Bouwman L (2014a) The Haber Bosch–harmful algal bloom (HB–HAB) link. *Environmental Research Letters* **9**: 105001.
- Glibert PM, Allen IJ, Artioli Y, Beusen A, Bouwman L, Harle J *et al.* (2014b) Vulnerability of coastal ecosystems to changes in harmful algal bloom distribution in response to climate change: projections based on model analysis. *Global Change Biology* **20**: 3845–3858.
- GlobalHAB (2017). *Global Harmful Algal Blooms, Science and Implementation Plan*. E Berdalet *et al.* (eds.). SCOR and IOC, Delaware and Paris, 64pp. <http://www.globalhab.info/science/globalhab-new-topic/habs-and-aquaculture>
- Gobler CJ, Berry DL, Dyhrman ST, Wilhelm SW, Salamov A, Lobanov AV *et al.* (2011) Niche of harmful alga *Aureococcus anophagefferens* revealed through ecogenomics. *Proceedings of the National Academy of Sciences USA* **108**: 4352–4357.
- Gobler CJ, Doherty OM, Hattenrath-Lehmann TK, Griffith AW, Kang Y, Litaker RW (2017) Ocean warming since 1982 has expanded the niche of toxic algal blooms in the North Atlantic and North Pacific oceans. *Proceedings of the National Academy of Sciences USA* **114**: 4975–4980.
- Gowen RJ, Tett P, Bresnan E, Davidson K, McKinney A, Milligan S *et al.* (2012) Anthropogenic nutrient enrichment and blooms of harmful micro-algae. *Oceanography and Marine Biology* **50**: 65–126.
- Grant J, Curran KJ, Guyondet TL, Tita G, Bacher C, Koutitonsky V *et al.* (2007) A box model of carrying capacity for suspended mussel aquaculture in Lagune de la Grande-Entrée, Iles-de-la-Madeleine, Québec. *Ecological Modelling* **200**: 193–206.
- Hallegraeff GM (1993) A review of harmful algal blooms and their apparent global increase. *Phycologia* **32**(2): 79–99.
- Hallegraeff GM (2003) Harmful algal blooms: a global overview. In: Hallegraeff GM, Anderson DM, Cembella AD (eds) *Manual on Harmful Marine Microalgae*, pp. 25–49. UNESCO Publishing, Paris, France.
- Hallegraeff GM, Bolch CJ (1991) Transport of toxic dinoflagellate cysts via ships' ballast water. *Marine Pollution Bulletin* **22**: 27–30.
- Hallegraeff GM, Bolch CJ (1992) Transport of diatom and dinoflagellate resting spores in ships' ballast water: implications for plankton biogeography and aquaculture. *Journal of Plankton Research* **14**: 1067–1084.
- Hallegraeff G, Bresnan E, Enevoldsen H, Schweibold L, Zingone A (2017) Call to contribute to global harmful algal bloom status reporting. *Harmful Algae News* **58**: 1–3.
- Harris TD, Smith VH (2015) Do persistent organic pollutants stimulate cyanobacterial blooms? *Inland Waters* **6**: 124–130.
- Hedberg N, Kautsky N, Kumblad L, Wikström SA (2018) *Limitations of Using Blue Mussel Farms as a Nutrient Reduction Measure in the Baltic Sea*. Report 2/2018, Baltic Sea Centre, Stockholm University. URL (accessed 10 Jan 2019): https://www.su.se/polopoly_fs/1.406424.1539853328!/menu/standard/file/BSC_Mussels_report_2_2018_lowres.pdf
- Hégaret H, Wikfors GH, Shumway SE (2007) Diverse feeding responses of five species of bivalve mollusc when exposed to three species of harmful algae. *Journal of Shellfish Research* **26**: 549–559.
- Hégaret H, Shumway SE, Wikfors GH, Pate S, Burkholder JM (2008) Potential transport of harmful algae via relocation of bivalve molluscs. *Marine Ecology Progress Series* **361**: 169–179.
- Heisler J, Gilbert P, Burkholder J, Anderson D, Cochlan W, Dennison W *et al.* (2008) Eutrophication and harmful algal blooms: a scientific consensus. *Harmful Algae* **8**: 3–13.
- Higman WA, Turner A, Baker C, Higgins C, Veszelovszki A, Davidson K (2014) Research to support the development of a monitoring programme for new or emerging marine biotoxins in shellfish in UK Waters. Food Standards Agency report FS513005, March 2014.o
- Hinder SL, Hays GC, Brooks CJ, Davies AP, Edwards M, Walne AW *et al.* (2011) Toxic marine microalgae and shellfish poisoning in the British Isles: history, review of epidemiology, and future implications. *Environ Health* **10**: 54.
- Hoagland P, Scatasta S (2006) The Economic Effects of Harmful Algal Blooms. In: Granéli E, Turner JT (eds) *Ecology of Harmful Algae. Ecological Studies (Analysis and Synthesis)*, pp. 391–402. Springer, Berlin, Heidelberg.
- Hoagland P, Anderson DM, Kaoru Y, White AW (2002) The economic effects of harmful algal blooms in the United States: estimates, assessment issues, and information needs. *Estuaries* **25**: 677–695.
- Holdt SL, Edwards MD (2014) Cost-effective IMTA: a comparison of the production efficiencies of mussels and seaweed. *Journal of Applied Phycology* **26**: 933–945.
- Huston MA, Wolverson S (2009) The global distribution of net primary production: resolving the paradox. *Ecological Monographs* **79**: 343–377.
- Inaba N, Trainer VL, Nagain S, Kojima S, Sakami T, Takagi S *et al.* (2019) Dynamics of seagrass bed microbial communities

- used to control artificial *Chattonella* blooms: a microcosm study. *Harmful Algae* **84**: 139–150.
- International Ocean-Colour Coordinating Group (IOCCG) (2009) Remote Sensing Applications in Marine Aquaculture. Forget M-H, Stuart V, Platt T (eds.), *Reports of the International Ocean-Colour Coordinating Group, No. 8*, pp. 77–87. IOCCG, Dartmouth, Canada.
- International Ocean-Colour Coordinating Group (IOCCG) (2014) Detection of Dominant Algal Blooms by Remote Sensing. Sathyendranath S (ed.), *Reports of the International Ocean-Colour Coordinating Group, No. 15*, pp. 39–68. IOCCG, Dartmouth, Canada.
- Jacobson DM, Andersen RA (1994) The discovery of mixotrophy in photosynthetic species of *Dinophysis* (Dinophyceae): light and electron microscopical observations of food vacuoles in *Dinophysis acuminata*, *D. norvegica* and two heterotrophic dinophysoid dinoflagellates. *Phycologia* **33**(2): 97–110.
- James KJ, Carey B, O'Halloran J, Van Pelt FNAM, Krabakova ZS (2010) Shellfish toxicity: human health implications of marine algal toxins. *Epidemiology and Infection* **138**: 927–940.
- Jenkinson IR, Connors PP (1980) The occurrence of the red tide organism, *Gyrodinium aureolum* Hulbert (Dinophyceae), round the south and west of Ireland in August and September, 1979. *Journal Sherkin Island* **1**: 127–146.
- Johnston P, Santillo D (2002) *Chemical Usage in Aquaculture: Implications for Residues in Market Products*. Greenpeace Research Laboratories Technical Note 06/2002, Department of Biological Sciences, University of Exeter, Exeter, UK.
- Jones KJ, Ayres P, Bullock AM, Roberts RJ, Tett P (1982) A red tide of *Gyrodinium aureolum* in sea lochs of the firth of Clyde and associated mortality of pond-reared salmon. *Journal of the Marine Biological Association UK* **62**: 771–782.
- Jones KL, Mikulski CM, Barnhorst A, Doucette GJ (2011) Comparative analysis of bacterioplankton assemblages from *Karenia brevis* bloom and non-bloom water on the west Florida shelf (Gulf of Mexico, USA) using 16S rRNA gene clone libraries. *FEMS Microbiology Ecology* **73**: 468–485.
- Kapetsky JM, Aguilar-Manjarrez J, Jenness J (2013) *A Global Assessment of Potential for Offshore Mariculture Development From a Spatial Perspective*, 181 pp. FAO Fisheries and Aquaculture Technical Paper. No. 549. FAO, Rome, Italy.
- Kim HG (2006) Mitigation and controls of HABs. In Granéli E, Turner JT (eds), *Ecology of Harmful Algae. Ecological Studies*, Vol. **189**, pp. 327–338, Springer-Verlag, Berlin, Germany.
- Kirkpatrick B, Fleming LE, Squicciarini D, Backer LC, Clark R, Abraham W *et al.* (2004) Literature review of Florida red tide: implications for human health effects. *Harmful Algae* **3**: 99–115.
- Kong F, Yu R, Zhang Q (2012) Primary analyses on the causative species of a bloom in the Sanggou Bay. *Marine Environmental Science* **31**: 824–829.
- Krimsky L, Staugler B, Hall-Scharf B, Stump K, Burton R (2018) Understanding Florida's Red Tide. SeaGrant/University of Florida/IFAS Blog URL (accessed April 2019): <https://www.fseagrant.org/news/2018/12/understanding-floridas-red-tide/>
- Kudela RM, Seeyave S, Cochlan WP (2010) The role of nutrients in regulation and promotion of harmful algal blooms in upwelling systems. *Progress in Oceanography* **85**(1–2): 122–135.
- Kudela RM, Bickel A, Carter ML, Howard MDA, Rosenfeld L (2015). The monitoring of harmful algal blooms through ocean observing: the development of the California Harmful Algal Bloom Monitoring and Alert Program in Coastal Ocean Observing Systems. In: Liu Y, Kerkering H, Weisberg RH (eds.), *Coastal Ocean Observing Systems*. pp. 58–75. Academic Press, New York, NY.
- Kurekin A, Miller PI, Van der Woerd HJ (2014) Satellite discrimination of *Karenia mikimotoi* and *Phaeocystis* harmful algal blooms in European coastal waters: merged classification of ocean colour data. *Harmful Algae* **31**: 163–176.
- Lafferty KD, Harvell CD, Conrad JM, Friedman CS, Kent ML, Kuris AM *et al.* (2015) Infectious diseases affect marine fisheries and aquaculture economics. *Annual Reviews in Marine Science* **7**: 471–496.
- Landsberg JH (2002) The effects of Harmful Algal Blooms on aquatic organisms. *Reviews in Fisheries Science* **10**(2): 113–390.
- Landsberg JH, Flewelling LJ, Naar J (2009) *Karenia brevis* red tides, brevetoxins in the food web, and impacts on natural resources: decadal advancements. *Harmful Algae* **8**: 598–607.
- Le Gouvello R *et al.* (2017) Aquaculture and marine protected areas: potential opportunities and synergies. *Aquatic Conservation: Marine and Freshwater Ecosystems* **27**(S1): 138–150.
- Leadbetter A, Silke J, Cusack C (2018) *Creating a Weekly Harmful Algal Bloom Bulletin*. Marine Institute, Galway, Ireland, 63pp. <http://hdl.handle.net/10793/1344>
- Lee KH, Jeong HJ, Kwon JE, Kang HC, Kim JH, Jang SH *et al.* (2016) Mixotrophic ability of the phototrophic dinoflagellates *Alexandrium andersonii*, *A. affine*, and *A. fraterculus*. *Harmful Algae* **59**: 67–81.
- Lefebvre KA, Tilton SC, Bammler TK, Beyer RP, Srinouanchan S, Stapleton PL *et al.* (2007) Gene expression profiles in zebrafish brain after acute exposure to domoic acid at symptomatic and asymptomatic doses. *Toxicology Science* **107** (1): 65–77.
- Lenes JM, Walsh JJ, Darrow BP (2013) Simulating cell death in the termination of *Karenia brevis* blooms: implications for predicting aerosol toxicity vectors to humans. *Marine Ecology Progress Series* **493**: 71–81.
- Li SC, Wang WX, Hsieh DP (2002) Effects of toxic dinoflagellate *Alexandrium tamarense* on the energy budgets and growth of two marine bivalves. *Marine Environmental Research* **53**(2): 145–160.
- Li J, Glibert PM, Zhou MJ, Lu SH, Lu DD (2009) Relationships between nitrogen and phosphorus forms and ratios and the development of dinoflagellate blooms in the East China Sea. *Marine Ecology Progress Series* **383**: 11–26.
- Lin J, Li C, Zhang S (2016) hydrodynamic effect of a large offshore mussel suspended aquaculture farm. *Aquaculture* **451**: 147–155.

- Lin C-H, Flynn KJ, Mitra A, Glibert PM (2018) Simulating effects of variable stoichiometry and temperature on mixotrophy in the harmful dinoflagellate *Karlodinium veneticum*. *Frontiers in Marine Science* **5**: 320. doi.org/10.3389/fmars.2018.00320.
- Liu H, Su J (2017) Vulnerability of China's nearshore ecosystems under intensive mariculture development. *Environmental Science and Pollution Research* **24**: 8957–8966.
- Loureiro S, Jauzein C, Garcés E, Collos Y, Camp J, Vaqué D (2008) The significance of organic nutrients in the nutrition of *Pseudo-nitzschia delicatissima* (Bacillariophyceae). *Journal of Plankton Research* **31**: 399–410.
- Lovatelli A, Aguilar-Manjarrez J, Soto D (2013) *Expanding Mariculture Further Offshore: Technical, Environmental, Spatial and Governance Challenges*. FAO Technical Workshop. 22–25 March 2010. Orbetello, Italy. FAO Fisheries and Aquaculture Proceedings No. 24. FAO, Rome, Italy.
- Liu D, Qi Y, Gu H, Dai X, Wang H, Gao Y *et al.* (2014) Causative species of harmful algal blooms in Chinese coastal waters. *Algalological Studies* **145**: 145–168.
- Lucas LV, Cloern JE, Thompson JK, Stacey MT, Koseff JR (2016) Bivalve grazing can shape phytoplankton communities. *Frontiers in Marine Science* **3**: 14. <https://doi.org/10.3389/fmars.2016.00014>
- Lundholm NL, Skoy J, Pocklington R, Moestrup O (1994) Domoic acid, the toxic amino acid responsible for amnesic shellfish poisoning, now in *Pseudonitzschia seriata* (Bacillariophyceae) in Europe. *Phycologia* **33**: 475–478.
- Maguire J, Cusack C, Ruiz-Villarreal M, Silke J, McElligot D, Davidson K (2016) Applied simulations and integrated modelling for the understanding of toxic and harmful algal blooms (ASIMUTH): integrated HAB forecast systems for Europe's Atlantic Arc. *Harmful Algae* **53**: 160–166.
- Manfrin C, De Moro G, Torboli V, Venier P, Pallavicini A, Gerdol M (2012) Physiological and molecular responses of bivalves to toxic dinoflagellates. *Invertebrate Survival Journal* **9**: 184–199.
- McGillicuddy DJ Jr, Anderson DM, Lynch DR, Townsend DW (2005) Mechanisms regulating large-scale seasonal fluctuations in *Alexandrium fundyense* populations in the Gulf of Maine: results from a physical–biological model. *Deep Sea Research Part II: Topical Studies in Oceanography* **52**: 2698–2714.
- Mitchell S, Rodgers H (2007) Pathology of wild and cultured fish affected by a *Karenia mikimotoi* bloom in Ireland, 2005. *Bulletin-European Association of Fish Pathologists* **27** (1): 39–42.
- Modica A, Scilipoti D, La Torre R, Manganaro A, Sara G (2006) The effect of mariculture facilities on biochemical features of suspended organic matter (southern Tyrrhenian, Mediterranean). *Estuarine Coastal and Shelf Science* **66**: 177–184.
- Montagnes DJS, Chambouvet A, Guillou L, Fenton A (2008) Responsibility of microzooplankton and parasite pressure for the demise of toxic dinoflagellate blooms. *Aquatic Microbial Ecology* **53**: 211–225.
- Munday R (2013) Is protein phosphatase inhibition responsible for the toxic effects of okadaic acid in animals? *Toxins (Basel)* **5**(2): 267–285.
- National Oceanic and Atmospheric Administration (NOAA) (2016) *Fisheries of the Caribbean, Gulf, and South Atlantic; Aquaculture*. National Marine Fisheries Service (NMFS); National Oceanic and Atmospheric Administration (NOAA); Department of Commerce, Washington, DC.
- Nayar S, Bott K (2014) Current status of global cultivated seaweed production and markets. *World Aquaculture* **45**(2): 32–37.
- Newell RIE (2004) Ecosystem influences of natural and cultivated populations of suspension-feeding bivalve molluscs: a review. *Journal of Shellfish Research* **23**(1): 51–61.
- Nishitani L, Erickson G, Chew KK (1985) Role of the parasitic dinoflagellate *Amoebophrya ceratii* in control of *Gonyaulax catenella* populations. In: Andersen DM, White AW, Baden DG (eds), *Toxic Dinoflagellates*, pp. 225–230. Elsevier, New York, NY.
- NOAA (2015) Programmatic Environmental Assessment for the prevention, control, and mitigation of Harmful Algal Blooms Program. URL (accessed April 2019): https://cdn.coastalscience.noaa.gov/page-attachments/about/pcm_hab_pea_finaldoc.pdf
- NOAA National Oceanic and Atmospheric Administration (2004) Action Plan for Harmful Algal Blooms and the Gulf of Mexico Coastal Ocean Observing System: Results from a Regional Workshop. NOAA Coastal Services Center. NOAA/CSC/20516-PUB. 27 p.
- Okaichi T (1989) Red tide problems in the Seto Inland Sea, Japan. In: Okaichi T, Anderson DM, Nemoto T (eds) *Red Tides, Biology, Environmental Science and Toxicology*, pp. 137–142. Elsevier, New York, NY.
- Paerl HW, Hall NS, Peierls BL, Rossignol KL (2014) Evolving paradigms and challenges in estuarine and coastal eutrophication dynamics in a culturally and climatically stressed world. *Estuaries and Coasts* **37**(2): 243–258.
- Park TG, Lim WA, Park YT, Lee CK, Jeong HJ (2013) Economic impact, management and mitigation of red tides in Korea. *Harmful Algae* **30**: S131–S143.
- Pastres R, Borga F, Brigolin D, Galparsoro I, Tett P (2018) Guide to the AquaSpace ToolBox AquaSpace: Ecosystem Approach to making Space for Aquaculture. EU Horizon 2020 project grant n°. 633476. 13 pp.
- Paterson R, McNeill S, Mitchell E, Adams T, Swan S, Clarke D *et al.* (2017) Environmental control of harmful dinoflagellates and diatoms in a fjordic system. *Harmful Algae* **69**: 1–17.
- Peperzak L, Poelman M (2008) Mass mussel mortality in The Netherlands after a bloom of *Phaeocystis globosa* (prymnesiophyceae). *Journal of Sea Research* **60**: 220–222.
- Peters F, Arin L, Marrasé C, Berdalet E, Sala MM (2006) Effects of small-scale turbulence on the growth of two diatoms of different size in a phosphorus-limited medium. *Journal of Marine Systems* **61**: 134–148.

- Petersen JK, Hansen JW, Laursen MB, Clausen P, Carstensen J, Conley DJ (2008) Regime shift in a coastal marine ecosystem. *Ecological Applications* **18**: 497–510.
- Petersen JK, Hasler B, Timmermann K, Nielsen P, Tørring DB, Larsen MM *et al.* (2014) Mussels as a tool for mitigation of nutrients in the marine environment. *Marine Pollution Bulletin* **82**(1–2): 137–143.
- Pickett ML (2018) Toxic ‘red tide’ algae bloom is killing Florida wildlife and menacing tourism. Guardian newspaper Tue 14 Aug 2018. URL (accessed 14 January 2019): <https://www.theguardian.com/us-news/2018/aug/13/florida-gulf-coast-red-tide-toxic-algae-bloom-killing-florida-wildlife>
- Pierce RH, Henry MS, Higham CJ, Blum P, Sengco MR, Anderson DM (2004) Removal of harmful algal cells (*Karenia brevis*) and toxin from seawater culture by clay flocculation. *Harmful Algae* **3**: 141–148.
- Pitcher GC, Calder D (2000) Harmful algal blooms of the southern Benguela Current: a review and appraisal of monitoring from 1989 to 1997. *South African Journal of Marine Science* **22**: 255–271.
- Pitcher GC, Jiménez AB, Kudela RM, Reguera B (2017) Harmful algal blooms in eastern boundary upwelling systems: a GEO-HAB Core Research Project. *Oceanography* **30**(1): 22–35.
- Pitta P, Apostolaki ET, Giannoulaki M, Karakassis I (2005) Mesoscale changes in the water column in response to fish farming zones in three coastal areas in the Eastern Mediterranean Sea. *Estuarine, Coastal and Shelf Science* **65**: 501–512.
- Pokrzywinski KL, Tilney CL, Modla S, Caplan JL, Ross J, Warner ME *et al.* (2017) Effects of the bacterial algicide IRI-160AA on cellular morphology of harmful dinoflagellates. *Harmful Algae* **62**: 127–135.
- Rabalais NN, Diaz RJ, Levin LA, Turner RE, Gilbert D, Zhang J (2010) Dynamics and distribution of natural and human-caused hypoxia. *Biogeosciences* **7**(2): 585–619.
- Raine R, O’Boyle S, O’Higgins T, White M, Patching J, Cahill B *et al.* (2001) A satellite and field portrait of a *Karenia mikimotoi* bloom off the south coast of Ireland, August 1998. *Hydrobiologia* **465**: 187–193.
- Raine R, Edwards M, Reid PC, Bresnan E, Fernand L (2008) Harmful Algal Blooms in Marine Climate Change Impacts Annual Report Card 2007–2008. In: Baxter JM, Buckley PJ, Wallace CJ (eds.), *Scientific Review*, 8pp. URL (accessed 26 August 2019): www.mccip.org.uk/arc/2007/PDF/HABs.pdf
- Ramanan R, Kim B-H, Cho D-H, Oh HM, Kim H-S (2016) Algae–bacteria interactions: evolution, ecology and emerging applications. *Biotechnology Advances* **34**(1): 14–29.
- Ramsdell JS, Zabka TS (2008) In utero DA toxicity: a fetal basis to adult disease in the California sea lion (*Zalophus californianus*). *Marine Drugs* **6**(2): 262–290.
- Read P, Fernandes T (2003) Management of environmental impacts of marine aquaculture in Europe. *Aquaculture* **226**: 139–163.
- Reguera B, Riobó P, Rodríguez F, Díaz PA, Pizarro G, Paz B *et al.* (2014) Dinophysis Toxins: causative Organisms, Distribution and Fate in Shellfish. *Marine Drugs* **12**(1): 396–461.
- Relyea RA (2009) A cocktail of contaminants: how mixtures of pesticides at low concentrations affect aquatic communities. *Oecologia* **159**: 363–376.
- Rengefors K, Karlsson I, Hansson L-A (1998) Algal cyst dormancy: a temporal escape from herbivory. *Proceedings of the Royal Society B: Biological Sciences* **265**: 1353–1358.
- Robineau B, Gagne JA, Fortier L, Cembella AD (1991) Potential impact of a toxic dinoflagellate (*Alexandrium excavatum*) bloom on survival of fish and crustacean larvae. *Marine Biology* **108**: 293–301.
- Roelke D, Buyukates Y (2001) The diversity of harmful algal bloom-triggering mechanisms and the complexity of bloom initiation. *Human and Ecological Risk Assessment* **7**: 1347–1362.
- Roilton A, Vignier J, Volety A, Shumway S, Bricelj VM, Soudant P (2018) Impacts of exposure to the toxic dinoflagellate *Karenia brevis* on reproduction of the northern quahog, *Merccenaria mercenaria*. *Aquatic Toxicology* **202**: 153–162.
- Rosetta CH, McManus GB (2003) Feeding by ciliates on two harmful algal bloom species, *Prymnesium parvum* and *Prorocentrum minimum*. *Harmful Algae* **2**: 109–126.
- Roth PB, Mikulski CM, Doucette GJ (2008) Influence of microbial interactions on the susceptibility of *Karenia* species to algicidal bacteria. *Aquatic Microbial Ecology* **50**: 251–259.
- Ruiz-Villarreal M, García-García LM, Cobas M, Díaz PA, Reguera B (2016) Modelling the hydrodynamic conditions associated with *Dinophysis* blooms in Galicia (NW Spain). *Harmful Algae* **53**: 40–52.
- S-3 EuroHAB (2019) *Project S-3 EUROHAB to Monitor Harmful Algal Blooms From Space*. EU Interreg Programme. URL (accessed 26 August 2019): <https://interreg5a-fce.eu/en/news-and-events/news/project-s-3-eurohab-to-monitor-harmful-algal-blooms-from-space/>
- Schmidt W, Evers-King HL, Campos CJA, Jones DB, Miller PI, Davidson K *et al.* (2018) A generic approach for the development of short-term predictions of *E. coli* and biotoxins in shellfish; application to a coastal bay and an estuary. *Aquaculture and Environment Interactions* **10**: 173–185.
- Seger A, Dorantes-Aranda J, Müller M, Body A, Peristyy A, Place A *et al.* (2015) Mitigating fish-killing *Prymnesium parvum* algal blooms in aquaculture ponds with clay: the importance of pH and clay type. *Journal of Marine Science and Engineering* **3**: 154–174.
- Seger A, Park T-G, Hallegraef G (2017) Assessment of the efficacy of clay flocculation in Korean fish farm waters: *Cochlodinium* cell removal and mitigation of ichthyotoxicity. *Harmful Algae* **61**: 46–55.
- Selander E, Fagerberg T, Wohlrab S, Pavia H (2012) Fight and flight in Dinoflagellates? Kinetics of simultaneous grazer-induced responses in *Alexandrium tamarense*. *Limnology and Oceanography* **57**: 58–64.
- Sellner KG, Rensel JE (2018) Prevention, control and mitigation of Harmful Algal Bloom impacts on fish shellfish and human consumers. In: Shumway SE, Burkholder JM, Morton SL (eds), *Harmful Algal Blooms a Compendium Desk Reference*, pp. 435–492. John Wiley & Sons Inc., Hoboken, NJ.

- Sellner KG, Doucette GJ, Kirkpatrick GJ (2003) Harmful algal blooms: causes, impacts and detection. *Journal of Industrial Microbiology and Biotechnology* **30**: 383–409.
- Sephton DH, Haya K, Martin JL, LeGresley MM (2007) Paralytic shellfish toxins in zooplankton, mussels, lobsters and caged Atlantic salmon, *Salmo salar*, during a bloom of *Alexandrium fundyense* off Grand Manan Island, in the Bay of Fundy. *Harmful Algae* **6**: 745–758.
- Shi J, Wei H (2009) Simulation of hydrodynamic structures in a semi-enclosed bay with dense raft-culture. *Periodical of Ocean University of China* **39**: 1181–1187.
- Shi F, McNabb P, Rhodes L, Holland P, Webb S, Adamson J *et al.* (2012) The toxic effects of three dinoflagellate species from the genus *Karenia* on invertebrate larvae and finfish. *New Zealand Journal of Marine and Freshwater Research* **46** (2): 149–165.
- Shirota A (1989) Red tide problem and countermeasures. *International Journal of Fisheries and Aquaculture* **1**: 195–223.
- Shumway SE (1995) Phycotoxin-related shellfish poisoning: bivalve molluscs are not the only vectors. *Reviews of Fishery Science* **3**: 1–31.
- Shumway SE, Barter J, Sherman-Caswell S (1990) Auditing the impact of toxic algal blooms on oysters. *Environmental Auditor* **2**: 41–56.
- Shumway SE, Burkholder JM, Morton SL (2018) *Harmful Algal Blooms A Compendium Desk Reference*. John Wiley & Sons Inc., Hoboken, NJ.
- Shutler JD, Warren MA, Miller PI, Barciela R, Mahdon R, Land PE *et al.* (2015) Operational monitoring and forecasting of bathing water quality through exploiting satellite Earth observation and models: the AlgaRisk demonstration service. *Computers & Geosciences* **77**: 87–96.
- Silke J, O'Beirn F, Cronin M (2005) *Karenia mikimotoi: An Exceptional Dinoflagellate Bloom in Western Irish Waters, Summer 2005*, Marine Environment and Health Series No. 21, 48 page. Marine Institute. (accessed November 2019). Available from URL: <https://oar.marine.ie/bitstream/handle/10793/240/No%2021%20Marine%20Environment%20and%20Health%20Series.pdf;jsessionid=DD29022691DFCCBB7EAEFBD A7FC29D54?sequence=1>
- Smayda TJ (1989) Causes and impacts of recurrent brown tide and other unusual blooms. In: Cosper EM, Carpenter EJ, Bricelej M (eds) *Novel Phytoplankton Blooms*, pp. 213–228. Springer-Verlag, New York, NY.
- Smayda TJ (1997) Harmful algal blooms: their ecophysiology and general relevance to phytoplankton blooms in the sea. *Limnology and Oceanography* **42**: 1137–1153.
- Smayda TJ (2004) Harmful algal bloom communities in Scottish coastal waters: relationship to fish farming and regional comparisons – a review. Scottish Executive Environment Group Paper 2006/3.
- Smayda TJ, Reynolds CS (2001) Community assembly in marine phytoplankton: application of recent models to harmful dinoflagellate blooms. *Journal of Plankton Research* **23**(5): 447–461.
- Smayda TJ, Trainer VL (2010) Dinoflagellate blooms in upwelling systems: seeding, variability, and contrasts with diatom bloom behaviour. *Progress in Oceanography* **85**(1–2): 92–107.
- Soliño L, Ferrer-Obiol J, Navarro-Herrero L, González-Solís J, Reis Costa PR (2019) Are pelagic seabirds exposed to amnesic shellfish poisoning toxins? *Harmful Algae* **84**: 172–180.
- Soto D (2009) *Integrated Mariculture: A Global Review*. FAO Fisheries and Aquaculture Technical Paper No. 529. FAO, Rome, Italy. 183pp. URL (accessed 29 December 2018): www.fao.org/docrep/012/i1092e/i1092e00.htm.
- Soto D, Aguilar-Manjárez J (2009) *FAO Expert Workshop on "Guidelines for the Implementation of an Ecosystem Approach to Aquaculture (EAA)"*. FAO Aquaculture Newsletter 42, pp. 8–9. FAO, Rome, Italy.
- Sourisseau M, Jegou K, Lunven M, Quere J, Gohin F, Bryere P (2016) Distribution and dynamics of two species of Dinophyceae producing high biomass blooms over the French Atlantic Shelf. *Harmful Algae* **53**: 53–63.
- Stadmark J, Conley DJ (2011) Mussel farming as a nutrient reduction measure in the Baltic Sea: consideration of nutrient biogeochemical cycles. *Marine Pollution Bulletin* **62**: 1385–1388.
- Staley ZR, Harwood VJ, Rohr JR (2015) A synthesis of the effects of pesticides on microbial persistence in aquatic ecosystems. *Critical Reviews in Toxicology* **45**: 813–836.
- Stüken A, Orr RJS, Kellmann R, Murray SA, Neilan BA, Kjetill SJ (2011) Discovery of nuclear-encoded genes for the neurotoxin saxitoxin in dinoflagellates. *PLoS ONE* **6**: e0020096.
- Sun R, Sun P, Zhang J, Esquivel-Elizondo S, Wu Y (2018) Microorganisms-based methods for harmful algal blooms control. *Annual Reviews in Bioresource Technology* **248**(Pt. B): 12–20.
- Svensen MBS, Andersen NR, Hansen PJ, Steffensen JF (2018) Effects of Harmful Algal Blooms on fish: Insights from *Prymnesium parvum*. *Fishes* **3**: 11.
- Tang YZ, Gobler CJ (2011) The green macroalga, *Ulva lactuca*, inhibits the growth of seven common harmful algal bloom species via allelopathy. *Harmful Algae* **10**: 480–488.
- Toth GB, Norén F, Selander E, Pavia H (2004) Marine dinoflagellates show induced life-history shifts to escape parasite infection in response to water-borne signals. *Proceedings of the Royal Society: B* **271**: 733–738.
- Touzet N, Davidson K, Romain P, McCoy GR, Amzil Z, Maher M *et al.* (2010) Co-occurrence of the West European (Gr. III) and North American (Gr. I) ribotypes of *Alexandrium tamarense* (Dinophyceae) in Shetland, Scotland. *Protist* **161**: 370–384.
- Trainer VL, Yoshida T (2014) Proceedings of the Workshop on Economic Impacts of Harmful Algal Blooms on Fisheries and Aquaculture. *PICES Scientific Reports* **47**: 85.
- Trainer VL, Bates SS, Lundholm N, Thessen AE, Cochlan WP, Adams NG *et al.* (2012) *Pseudo-nitzschia* physiological ecology, phylogeny, toxicity, monitoring and impacts on ecosystem health. *Harmful Algae* **14**: 271–300.

- Troell M, Joyce A, Chopin T, Neori A, Buschmann AH, Fang J-G (2009) Ecological engineering in aquaculture – potential for integrated multi-trophic aquaculture (IMTA) in marine offshore systems. *Aquaculture* **297**: 1–9.
- Turner JT, Tester PA (1997) Toxic marine phytoplankton, zooplankton grazers, and pelagic food webs. *Limnology and Oceanography* **42**: 1203–1214.
- Turner AD, Higgins C, Davidson K, Veszelszki A, Payne D, Hungerford J *et al.* (2015) Potential threats posed by new or emerging marine biotoxins in UK Waters and examination of detection methodology used in their control: Brevetoxins. *Marine Drugs* **13**: 1224–1254.
- Twiner MJ, Rehmann N, Hess P, Doucette GJ (2008) Azaspiracid Shellfish Poisoning: a review on the chemistry, ecology, and toxicology with an emphasis on human health impacts. *Marine Drugs* **6**(2): 39–72.
- UN (2017) The 2017 Revision of World Population Prospects. URL (accessed 31 December 2018): <https://population.un.org/wpp/>
- Uyaguari M, Scott G, Norman RS (2013) Abundance of class 1-3 integrons in South Carolina estuarine ecosystems under high and low levels of anthropogenic influence. *Marine Pollution Bulletin* **15**: 77–84.
- Wang Y, Yu Z, Song X, Tang X, Zhang S (2007) Effects of macroalgae *Ulva pertusa* (Chlorophyta) and *Gracilaria lemaneiformis* (Rhodophyta) on growth of four species of bloom-forming dinoflagellates. *Aquatic Botany* **86**: 139–147.
- Wang S, Tang D, He F, Fukuyo Y, Azanza RV (2008) Occurrences of Harmful Algal Blooms (HABs) associated with ocean environments in the South China Sea. *Hydrobiologia* **596**: 79–93.
- Ward JE, Shumway SE (2004) Separating the grain from the chaff: particle selection in suspension- and deposit-feeding bivalves. *Journal of Experimental Marine Biology and Ecology* **300**: 83–130.
- Wartenberg R, Feng L, Jun WuJ, Mak YL, Chan LL, Telfer TC *et al.* (2017) The impacts of suspended mariculture on coastal zones in China and the scope for Integrated Multi-Trophic Aquaculture. *Ecosystem Health and Sustainability* **3**: 1340268.
- Watkins SM, Reich A, Fleming LE, Hammond R (2008) Neurotoxic Shellfish Poisoning. *Marine Drugs* **6**: 431–455.
- Watts JEM, Schreier HJ, Lanska L, Hale MS (2017) The rising tide of antimicrobial resistance in aquaculture: sources, sinks and solutions. *Marine Drugs* **15**: 158.
- Wells ML, Trick CG, Cochlan WP, Hughes P, Trainer VL (2005) Domoic Acid: the synergy of iron, copper and the toxicity of diatoms. *Limnol. Oceanography* **50**: 1908–1917.
- Wells ML, Trainer VL, Smayda TJ, Karlson BS, Trick CG, Kudela RM *et al.* (2015) Harmful algal blooms and climate change: learning from the past and present to forecast the future. *Harmful Algae* **49**: 68–93.
- Wells ML, Karlson B, Wulff A, Kudela R, Trick C, Asnaghi V *et al.* (2019). Future HAB science: Directions and challenges in a changing climate. *Harmful Algae*, available online, doi.org/10.1016/j.hal.2019.101632.
- WHO (2003) Algae and cyanobacteria in coastal and estuarine waters in Guidelines for Safe Recreational Water Environments, pp. 128–136. URL (accessed 16 April 2017): <http://apps.who.int/iris/bitstream/10665/42591/1/9241545801.pdf>
- Whyte C, Swan S, Davidson K (2014) Changing wind patterns linked to unusually high *Dinophysis* blooms around the Shetland Islands, Scotland. *Harmful Algae* **39**: 365–373.
- Wiese M, D’Agostino PM, Mihali TK, Moffitt MC, Neilan BA (2010) Neurotoxic Alkaloids: Saxitoxin and its analogs. *Marine Drugs* **8**: 2185–2211.
- Wilson WH, Tarran GA, Schroeder D, Cox M, Oke J, Malin G (2002) Isolation of viruses responsible for the demise of an *Emiliania huxleyi* bloom in the English Channel. *Journal of the Marine Biological Association UK* **82**: 369–377.
- Wolrhaf S (2013) Characterization of grazer-induced responses in the marine dinoflagellate *Alexandrium tamarense*. PhD Dissertation, Universität Bremen, Germany.
- Woolhouse MEJ, Ward MJ (2013) Sources of Antimicrobial Resistance. *Science* **341**: 1460–1461.
- Yahel G, Marie D, Beninger PG, Eckstein S, Genin A (2009) In situ evidence for pre-capture qualitative selection in the tropical bivalve *Lithophaga simplex*. *Aquatic Biology* **6**: 235–246.
- Yang Y, Chai Z, Wang Q, Chen W, He Z, Jiang S (2015) Cultivation of seaweed *Gracilaria* in Chinese coastal waters and its contribution to environmental improvements. *Algal Research* **9**: 236–244.
- Yu Z, Song X, Cao X, Liu Y (2017) Mitigation of harmful algal blooms using modified clays: theory, mechanisms, and applications. *Harmful Algae* **69**: 48–64.
- Zeng D, Huang D, Qiao X, He Y, Zhang T (2015) Effect of suspended kelp culture on water exchange as estimated by in situ current measurement in Sanggou Bay, China”. *Journal of Marine Systems* **149**: 14–24.
- Zerriffi SEA, El Khalloufi F, Oudra B, Vasconcelos V (2018) Seaweed bioactive compounds against pathogens and microalgae: potential uses on pharmacology and Harmful Algae Bloom control. *Marine Drugs* **16** (2): E55.
- Zhang C, Zhang J (2015) Current techniques for detecting and monitoring algal toxins and causative harmful algal blooms. *Journal of Environmental Analytical Chemistry* **2**: 123.
- Zhang QC, Qiu LM, Yu RC, Kong FZ, Wang YF, Yan T *et al.* (2012) Emergence of brown tides caused by *Aureococcus anophagefferens* Hargraves *et Sieburth* in China. *Harmful Algae* **19**: 117–124.
- Zhou Y, Yang H, Hu H, Liu Y, Mao Y, Zhou H *et al.* (2006) Bioremediation potential of the macroalga *Gracilaria lemaneiformis* (Rhodophyta) integrated into fed fish culture in coastal waters of North China”. *Aquaculture* **252**: 264–276.