



Citation for published version:

Jackson, DR, Fuller-Rowell, TJ, Griffin, DJ, Griffith, MJ, Kelly, CW, Marsh, DR & Walach, MT 2019, 'Future Directions for Whole Atmosphere Modeling: Developments in the Context of Space Weather', *Space Weather*, vol. 17, no. 9, pp. 1342-1350. <https://doi.org/10.1029/2019SW002267>

DOI:

[10.1029/2019SW002267](https://doi.org/10.1029/2019SW002267)

Publication date:

2019

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

(C) 2019 American Geophysical Union. The final published version is available online via: <https://doi.org/10.1029/2019SW002267>

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Space Weather

COMMENTARY

10.1029/2019SW002267

Key Points:

- We have reached a paradigm shift, where any self-respecting space weather model of the upper atmosphere now needs to have some representation of the lower atmosphere
- Further model developments are required in several key areas, including dynamical cores and the improved representation of gravity waves
- A road map of future actions is presented to ensure good progress continues to be made; this includes the development of a multi-model verification strategy

Correspondence to:

D. R. Jackson,
david.jackson@metoffice.gov.uk

Citation:






Jackson, D. R., Fuller-Rowell, T. J., Griffin, D. J., Griffith, M. J., Kelly, C. W., Marsh, D. R., & Walach, M.-T. (2019). Future directions for whole atmosphere modeling: Developments in the context of space weather. *Space Weather*, 17. <https://doi.org/10.1029/2019SW002267>

Received 31 MAY 2019

Accepted 24 AUG 2019

Accepted article online 30 AUG 2019

Future Directions for Whole Atmosphere Modeling: Developments in the Context of Space Weather

David R. Jackson¹ , Tim J. Fuller-Rowell², Dan J. Griffin¹, Matthew J. Griffith³ , Christopher W. Kelly⁴ , Daniel R. Marsh^{4,5} , and Maria-Theresia Walach⁶ 

¹Met Office, Exeter, UK, ²CIRES, University of Colorado, Boulder, CO, USA, ³Department of Mathematical Sciences, University of Bath, Bath, UK, ⁴School of Earth and Environment, University of Leeds, Leeds, UK, ⁵National Center for Atmospheric Research, Boulder, CO, USA, ⁶Department of Physics, Lancaster University, Lancaster, UK

Abstract Coupled Sun-to-Earth models represent a key part of the future development of space weather forecasting. With respect to predicting the state of the thermosphere and ionosphere, there has been a recent paradigm shift; it is now clear that any self-respecting model of this region needs to include some representation of forcing from the lower atmosphere, as well as solar and geomagnetic forcing. Here we assess existing modeling capability and set out a road map for the important next steps needed to ensure further advances. These steps include a model verification strategy, analysis of the impact of nonhydrostatic dynamical cores, and a cost-benefit analysis of model chemistry for weather and climate applications.

Plain Language Summary Numerical models that comprehensively simulate the region between the Sun and the Earth represent a key part of the future development of space weather forecasting. With respect to predicting the Earth's upper atmosphere, there has been a recent paradigm shift; it is now clear that any self-respecting model of this region needs to include some representation of impacts from below (the lower atmosphere) as well as from above (solar variability and the effects of solar wind fluctuations). Here we assess existing modeling capability and set out a road map for the important next steps needed to ensure further advances. These steps include a strategy for checking the accuracy of the models, an analysis of the impact of methods chosen to represent upper atmosphere dynamics, and an assessment of the relative benefits of comprehensive (but expensive) and simplified (but inexpensive) model representations of upper atmosphere chemistry.

1. Introduction

We are at the stage in the development of operational space weather forecasts where individual models of components of the Sun-to-Earth domain (including the ionosphere and the thermosphere) are beginning to be coupled together. Such a coupled modeling system, constrained by assimilation of near real time observations, has the potential to provide considerably better forecasts than currently available. It is clear that representing the impact of, for example, a coronal mass ejection, across the whole Sun-to-Earth domain can potentially improve forecasts in the ionosphere. The potential for improved forecasts has already been demonstrated for parts of the Sun-to-Earth system. For example, coupling a global magnetosphere model with an inner magnetosphere drift physics model considerably improves forecasts of geomagnetic storms (Liemohn et al., 2018) and improved representation of the thermosphere leads to improved ionospheric evolution (e.g., Chartier et al., 2013). In addition, there is a strong connection between the lower atmosphere state and the ionosphere that was highlighted initially by Immel et al. (2006) and demonstrated in later modeling studies (e.g., Pedatella et al., 2016). Furthermore, data assimilation (DA) schemes are already used for operational ionosphere models (e.g., Schunk et al., 2016), and experimental systems show that assimilation can improve model initial conditions in the thermosphere (e.g., Murray et al., 2015), the magnetosphere (e.g., Merkin et al., 2016), and the heliosphere (e.g., Lang & Owenst, 2019).

However, it is also becoming increasingly apparent that, in addition to correctly specifying this space weather forcing, thermosphere and ionosphere forecasts can also benefit from an accurate representation of coupling from within and below. The motivation for a whole atmosphere model (i.e., a model that extends from the ground up to the exobase) is thus twofold:

1. Recent research (e.g., Chartier et al., 2013, 2016; Hsu et al., 2014) has shown that no matter how accurately one represents the current ionospheric state, the quality of the subsequent ionospheric forecasts crucially depends on the ability to also represent the thermosphere and its evolution.
2. Both the ionosphere and thermosphere are sensitive to forcing from the lower atmosphere. The seminal paper by Immel et al. (2006) indicated connections between tidal patterns in the lower thermosphere and the *F* region ionosphere and noted that the tidal structure was linked to patterns of convection in the equatorial troposphere. Furthermore, numerous papers (e.g., Goncharenko, Chau, et al., 2010, Goncharenko, Coster, et al., 2010; Liu & Roble, 2002; McDonald et al., 2018; Pedatella et al., 2012) have shown how planetary wave forcing, specifically via stratospheric sudden warmings (SSWs), can affect lower thermospheric tides and thus the ionosphere.

Akmaev (2011) reviewed whole atmosphere models at a time when these models were quite new and our understanding of the links between the lower and upper atmosphere was developing. A Whole Atmosphere Modelling Workshop was held in Tres Cantos, Spain in June 2018 and a strong consensus emerged: the need to have some representation of the lower atmosphere in space weather models of the upper atmosphere. This is highly significant for the continued development of whole atmosphere models. In this commentary we review existing models, how their building blocks can be further developed, and how we can use observations (via DA and verification) to confront the model simulations and potentially produce improved forecasts.

2. Existing Models

There are three current whole atmosphere space weather models:

1. The Whole Atmosphere Model (WAM; Akmaev et al., 2008; Fuller-Rowell et al., 2008) is based on the U.S. National Weather Service Numerical Weather prediction model and extends from the surface to around 600 km. It is being combined with a separate ionosphere model Ionosphere Plasmasphere Electrodynamics (Maruyama et al., 2016) to produce a coupled model of the ionosphere and neutral atmosphere. WAM represents both the mean state and tides in the thermosphere well (e.g., Lieberman et al., 2013, show good agreement with diurnal and time mean Challenging Mini Satellite Payload winds). The pattern of changes seen in ionospheric vertical plasma drift and Total Electron Content (TEC; which occur in response to SSW forcing from below) agrees well with observations (e.g., Wang et al., 2014).
2. The Whole Atmosphere Community Climate Model with thermosphere and ionosphere extension (WACCM-X; Liu et al., 2010; Liu et al., 2018) is focused primarily on climate time scales (in contrast to WAM, which is focused on weather forecast time scales). With a comparable altitude range to WAM, it has a much more detailed representation of neutral and ion chemistry. Liu et al. (2018) report that in WACCM-X the amplitudes and seasonal variations of atmospheric tides in the mesosphere and lower thermosphere (MLT), equatorial ionosphere anomaly structures and storm time ionospheric behavior are all in good agreement with observations.
3. The Ground to topside model of the Atmosphere and Ionosphere for Aeronomy (GAIA) combines neutral atmosphere, ionospheric, and electrodynamic models. The neutral model covers the entire atmosphere from the Earth's surface up to the top of the thermosphere and contains a comprehensive range of physical parametrizations (e.g., Fujiwara & Miyoshi, 2010). Jin et al. (2012) show the ability of GAIA to model the impact of an SSW on migrating tides and the associated ionospheric response, with in general good agreement shown with Sounding of the Atmosphere using Broadband Emission and Constellation Observing System for Meteorology, Ionosphere, and Climate observations.

For clarification, weather models focus on short forecast time scales (often less than 10 days) and use as fine a resolution as possible in order to represent meteorological features such as weather fronts. Since forecast quality will depend on initial conditions, weather models must be initialized using DA. Coupling to other models (such as an ocean model) is usually not required on forecast time scales, and the need to run quickly in near real time precludes the use of such coupled models, and it is necessary to use fast, less-complex representations of physics and chemistry. Climate models are run for long forecast time scales such as annual or multidecadal periods and so generally have coarser resolutions than weather models. Coupling to comprehensive models of the Earth system (chiefly ocean and atmospheric chemistry models) is required to

represent long-term variability and climate change. For the specific case of whole atmosphere models, WAM and WACCM do not completely meet the description given above (e.g., WACCM can run at a finer horizontal resolution than WAM), but the WAM chemistry scheme is simple and designed for fast weather forecasts, whereas the WACCM chemistry scheme is considerably more complex, and it can be coupled to an ocean model. This enables WACCM to be used in activities like the Coupled Model Intercomparison Project 5, studying, for example, climate change from 1850 (Marsh et al., 2013) and climate impacts associated with long-term ozone change (Eyring et al., 2013).

3. Building Blocks for Better Models

3.1. Dynamics—Gravity Waves and Dynamical Formulation

The representation of gravity waves is very important for accurate modeling of the thermosphere. They are the prime driver of the middle atmosphere circulation and affect tidal amplitudes and thus can influence the mechanisms connecting the lower atmosphere with the thermosphere and ionosphere (see, e.g., Yiğit et al., 2016). Furthermore, accurate simulation of medium and small-scale traveling ionospheric disturbances (MSTIDs) and associated ionospheric plasma bubbles that impact precision application of Global Navigation Satellite System data require the ability to represent subgrid-scale gravity waves in whole atmosphere models. This information on MSTIDs could be input into existing tools for estimating Global Navigation Satellite System positioning error from TIDs (e.g., Lejeune et al., 2012). Gravity waves also play an important role in the transport of chemical constituents, which is discussed in more detail later.

Liu et al. (2014) ran a fine-resolution ($0.25^\circ \times 0.25^\circ$ horizontal, 0.1 scale height vertical) version of WACCM to demonstrate the simulation and impact of gravity waves up to around 100 km. However, it is not clear whether such resolutions are needed at higher levels in the thermosphere. Miyoshi et al. (2018) showed that a GAIA simulation with a resolution of $1^\circ \times 1^\circ$ produces fluctuations in electron density with length scales less than around 1,000 km and periods of less than around 2 hr, which are in good agreement with observations and which are not seen in a coarser resolution ($2.5^\circ \times 2.5^\circ$) simulation. The fluctuations reported by Miyoshi et al. are attributed to TIDs that are excited by secondary gravity waves. These waves typically have horizontal wavelengths of around 100 km to several 1,000s of kilometers (Vadas & Crowley, 2010). This also appears consistent with Gardner and Schunk (2011), who indicated observed gravity waves in the thermosphere typically have horizontal scales of around 100–500 km. Furthermore, at altitudes above around 110-km molecular viscosity and thermal conduction strongly influence gravity wave filtering and dissipation, as opposed to winds and wave breaking lower in the atmosphere (see, e.g., Vadas & Fritts, 2005). Accordingly, lower atmosphere gravity wave parametrization schemes may not be appropriate in the thermosphere. Schemes that specifically focus on parameterizing gravity waves in the thermosphere (e.g., Yiğit et al., 2008) could be adopted for coarse horizontal resolution whole atmosphere model simulations.

Presently, WAM, WACCM-X, and GAIA use hydrostatic dynamical cores. The dynamical core solves the governing fluid and thermodynamic equations in the model on resolved scales, while parametrizations represent subgrid-scale processes and other processes not included in the dynamical core such as radiative transfer (Thuburn, 2008). Certainly for some applications, such as satellite drag, the hydrostatic approximation appears adequate (see, e.g., Bruinsma et al., 2018), but there is still a need to identify the impact on model results that may arise from nonhydrostatic processes. For some applications that require accurate representation of the wave fluctuations (such as radio wave propagation in the bottomside F region for HF applications), the hydrostatic approximation may be inappropriate in the thermosphere, and adoption of nonhydrostatic (non-H) dynamical cores appears to be a logical next step. The hydrostatic approximation breaks in the presence of large vertical accelerations (e.g., Curry & Webster, 1998), and using a non-H dynamical core may affect the modeled gravity wave spectrum, particularly when applied at fine horizontal resolution. High-frequency waves with horizontal wavelength less than $4\pi H$ (where H is scale height) should be treated nonhydrostatically (Akmaev, 2011). For example, Eckermann et al. (2016) showed observations of gravity waves that had propagated from the surface to the lower thermosphere with vertical velocities of several tens of meters per second. They concluded that these waves must be nonhydrostatic, since if they were hydrostatic, they would have broken in the troposphere or lower stratosphere rather than propagating higher. Therefore, selection of a non-H dynamical core can affect the modeled gravity wave spectrum in the MLT, and thus the simulation of MSTIDs. A fine horizontal resolution is required to represent such

waves in the first place, and, given that whole atmosphere models currently have resolutions of ~100 to 200 km, the case for using non-H cores at such resolutions is not yet well made. Three new whole atmosphere models are being developed, which use non-H cores: the Navy Global Environmental Model (NAVGEN; e.g., McCormack et al., 2017), the Met Office Extended Unified Model (UM) and WAM, where the current dynamical core is being replaced with the Geophysics Fluid Dynamics Laboratory Finite-Volume on a Cubed-Sphere (FV3) non-H core (Ullrich et al., 2017). In addition, Borchert et al. (2018) report on work to extend the ICOSahedral Non-hydrostatic general circulation model up to 150-km altitude. NAVGEN and the UM have the option to switch between hydrostatic and non-H formulations, and both these models could play key roles in evaluating the importance of non-H cores in whole atmospheric models.

There can also be issues with the robustness of non-H dynamical cores in the thermosphere. Griffin and Thuburn (2018) showed that the UM required the addition of molecular viscosity and diffusion in order to realistically stabilize artificial wave growth, as this viscosity has a significant damping effect in the thermosphere. Another challenge arises above the turbopause (around 105 km) where diffusive separation means that air parcels are no longer turbulently mixed and the molecular weight of a species determines its dynamical evolution. Therefore, ideally, each species should have its own set of dynamical equations that need to be solved. The molecular diffusion is also affected by variable gravity, which in turn modifies atmospheric scale heights. Thus, there is a need to reformulate the dynamical core to properly model the individual species, as well as a need to add a correction to the thermal equation.

3.2. Radiation and Chemistry

Accurate radiation and chemistry schemes are needed throughout the whole atmosphere model domain, most obviously in the MLT where the radiation scheme calculates the absorption of solar radiation that drive the large rise in temperature with height there. This means that radiation schemes need to include the far ultraviolet, extreme ultraviolet, and soft X-ray spectral ranges that are usually ignored in lower atmosphere models. In the MLT, heating from exothermic reactions becomes important (especially during polar night) and must be accounted for to correctly simulate the thermal structure. Quenching of $O(^1D)$ is a large source of heating throughout the MLT, above 100-km ion reactions, and reactions involving atomic nitrogen are significant sources of heat, and below 100 km O_x and HO_x reactions are the dominant producers of chemical heating (Marsh et al., 2007). In addition, above the midmesosphere, local thermodynamic equilibrium (LTE) schemes need to be replaced by non-LTE formulations, since both near infrared heating and infrared cooling are overestimated by the LTE schemes. The Fomichev non-LTE parametrization (Fomichev & Blanchet, 1995; Fomichev et al., 1998; Ogibalov & Fomichev, 2003) is the only scheme currently available for Earth GCMs. Its formulation is based on recent atmospheric conditions, and it lacks the adaptability to be used for climate change experiments. The UM's radiation scheme is being extended to include far ultraviolet and extreme ultraviolet wavelengths. The scheme is highly flexible, with the option of being run using different spectral resolutions. In future it could be further modified to include a more comprehensive representation of non-LTE heating, possibly based on a scheme developed for Mars (López-Valverde & López-Puertas, 1994), which potentially represents a considerable improvement on the Fomichev scheme. Since the scheme is also publically available, it could be a highly important community resource for future collaborative whole atmosphere model development.

While only relatively few major chemical reactions are sufficient to adequately represent the large rise in temperature in the MLT (Marsh et al., 2007), other challenges remain. Below 85 km the atmospheric chemistry is dominated by compounds, and above 100 km by ion chemistry. Particularly interesting chemistry exists in between, where atoms including highly reactive hydrogen and oxygen atoms are in abundance, with maximum mixing ratios observed at around 85 and 90–95 km, respectively (Plane et al., 2015). WACCM simulations of metal layers originating from the ablation of meteoroids in the MLT give good model agreement with data at midlatitudes but show worse agreement at high latitudes. For example, for Fe chemistry Feng et al. (2013) show that the model significantly overestimates winter Fe and underestimates summer Fe compared to observations from three Antarctic ground-based lidars. This implies that the model vertical transport of chemical species may be significantly underestimated. A possible issue is that global models cannot capture transport associated with small-scale gravity waves, and adding diffusion terms to account for this does help with reducing the large bias. Observations of MLT chemistry are sparse, and thus, there is great scope for new observations to significantly improve our knowledge of the interaction

between chemistry and transport. For example, recent observations made by the Atmospheric Chemistry Experiment indicate nitrous oxide (N_2O) is being produced in the MLT (Sheese et al., 2016). N_2O is a precursor of odd nitrogen (NO_x), which destroys stratospheric ozone. A new chemical source of N_2O has been successfully added to WACCM by Kelly et al. (2018). Model simulations were able to capture the observed N_2O layer and well replicate seasonal variations near the poles. Recent studies have also highlighted the importance of radiation and chemistry schemes working together to produce the strong NO cooling, which is observed in the immediate aftermath of geomagnetic storm time thermospheric heating (e.g. Knipp et al., 2017).

3.3. Ionosphere and Electrodynamics

The coupling between the thermosphere and ionosphere is important, as mentioned above, in ensuring a more accurate evolution of the ionospheric state. Fang et al. (2013) performed an intercomparison of a range of ionospheric models. It is clear that the thermosphere/ionosphere coupling was modeled better when the models employed a fully consistent representation of the electrodynamics. This led to the development of the Ionosphere Plasmasphere Electrodynamics model, which includes the following requirements: It represents the ionosphere globally with similar resolution to the neutral atmospheric model (WAM) it is coupled to; it uses self-consistent electrodynamics for quiet and storm time dynamo processes; it uses a coupling infrastructure.

Also important is an accurate representation of the electric field and its variation. There are limitations with current empirical electric field models, such as those developed by Heelis et al. (1982) and Weimer (2005). These are climatological in nature, but more observations are required to capture the electric field variability. The introduction of Super Dual Auroral Radar Network (SuperDARN) data crucially adds extra observations poleward of 40° geomagnetic latitude (as well as providing observations at lower latitudes), and the deviation of SuperDARN high-latitude electric fields from the average ionospheric state shows the importance of accounting for the prior evolution of the ionospheric state. M.-T. Walach (presentation available at [http://www.research.lancs.ac.uk/portal/en/activities/characterising-and-understanding-temporal-variability-in-ionospheric-flows-using-superdarn-data\(21f8f287-e085-4418-8a1c-387d597ef2f0\).html](http://www.research.lancs.ac.uk/portal/en/activities/characterising-and-understanding-temporal-variability-in-ionospheric-flows-using-superdarn-data(21f8f287-e085-4418-8a1c-387d597ef2f0).html)) used SuperDARN data to show that greater solar wind corresponds to greater variability in convection and is currently investigating the drivers of this variability in more detail. Use of SuperDARN observations in the Canadian Ionosphere and Atmosphere Model (Martynenko et al., 2014) allows detailed features in the plasma density distribution to be reproduced, especially in the topside ionosphere at high latitudes. Data from the Assimilative Mapping of Ionospheric Electrodynamics can be used to assimilate multiple data sources (SuperDARN) for testing in whole atmosphere models. The electric field model chosen also influences modeled Joule heating, and it is important to continue to confront empirical model-based estimates with observations (e.g., Billett et al., 2018).

3.4. Observations for DA and Model Verification

DA is important in attempting to ensure the model state is constrained to be close to the true atmospheric state and has been applied extensively in WACCM-X, WAM, and NAVGEM. DA in WACCM-X is done using an ensemble Kalman filter while the NAVGEM DA system is a hybrid of 4D-Var and an ensemble Kalman Filter. The ensemble Kalman Filter (Evensen, 1994) is a combination of a Kalman Filter (which evolves the state and estimate covariance as new observations arrive) and Monte Carlo estimation methods (the full estimate covariance matrix is explicitly evolved using an ensemble—sample of evolved states). The NAVGEM system has been shown to add a lot of value in the thermosphere. As an example, in Figure 1 the observed Wave Number 4 structure in TEC is best reproduced when the NAVGEM model thermosphere is forced by 3-hourly analyses; forcing by 6-hourly analyses is less accurate. A major challenge is that the models cover a large altitude range, so waves can grow exponentially, and to maintain model stability with DA, more damping is often added to deal with spurious small-scale waves. A consequence of this approach is that while model dynamics and chemical transport are improved, it is at the cost of the tidal amplitudes being too weak. To add to the challenge in the upper atmosphere, data are sparse, and processes act on shorter time scales than in the lower atmosphere. Provision of considerably more near real-time observations of the upper atmosphere, particularly of the thermosphere, is vital if we are to exploit DA in order to produce improved model forecasts.




Figure 1. (a) Jet Propulsion Laboratory (JPL) global ionospheric map of total electron content (TEC) on 12 January 2010 shown at constant local time of 13:00 LT. (b) /NOGAPS-ALPHA simulation of TEC. (c) Navy Global Environmental Model (NAVEM) simulation of TEC. The simulated TEC is scaled by a factor of 0.7 (from McDonald et al., 2018). WACCM = Whole Atmosphere Community Climate Model.

To compound the lack of observations, the instruments that produce many of the upper atmosphere observations used in the DA schemes (e.g., Sounding of the Atmosphere using Broadband Emission, and MLS, the Microwave Limb Sounder) are well past their nominal mission lifetimes, and no follow-on programs are planned. Furthermore, these instruments only observe up to the lower thermosphere and observations higher in the thermosphere are extremely sparse. The QB50 Cubesat project (e.g., Gill et al., 2013) focused on the building and launching of instruments to measure thermospheric neutral density, but with little or no attention given to coordination and reception of data. However, the constellation of Cubesats used could be a pathfinder for a future operational observations system, with the critical proviso that this constellation would need to be underpinned by associated systems for near real time data reception and cross-calibration of data. In addition, new data from the Global-scale Observations of the Limb and Disk mission will help address the paucity of thermospheric data. The planned assimilation of Global-scale Observations of the Limb and Disk O/N₂ observations into WAM could test the assumption that temperature is a key variable for the initialization of upper atmosphere models. Since O/N₂ plays a key role as a diagnostic of thermospheric transport, it is possible that future DA schemes could instead use O/N₂ as a primary control variable.

Model verification using existing data has proved invaluable. However, there is a need for a consistent model verification strategy, and in particular community-wide agreement on which metrics to compare—this could include basic seasonal variability, tide amplitudes and variability, TEC, and the magnitude of the solar

semidiurnal migrating tide. An important consideration is to understand which observations are trusted and therefore should be used to validate model output, and there are benefits in an Intergovernmental Panel on Climate Change style model intercomparison, and a cooperative approach. An example is Coupled Model Intercomparison Project 5 (Taylor et al., 2012), in which an agreed set of experiments addressing major gaps in understanding was run using multiple models, and output data were formatted in a common way and made freely available via data portals. Empirical models may not be ideal for use as a level of comparison, and we suggest the employment of a more general model comparison system, for example, as implemented in the International Land Model Benchmarking Project (Collier et al., 2018).

4. Future Research Directions and Activities

Based on the discussions throughout the workshop, the following road map for future collaboration was agreed:

1. Compare existing hydrostatic models to understand impacts of dynamical formulation (also interactions with chemistry, the ionosphere, and radiation)
2. Comparison of non-H and hydrostatic dynamical cores to assess impact of non-H cores (and whether non-H is even needed at coarser resolution)
3. Assess numerical cost/benefit of comprehensive chemistry schemes designed for climate applications (e.g., WACCM) against simpler schemes designed for near-real time operational use (e.g., as used in WAM)
4. Development of a verification strategy and methodology, which is required to underpin the above three actions. Clearly, it makes sense to make links with other activities to guide our future actions. These include the Committee on Space Research International Space Weather Action Team and the Community Coordinated Modeling Center Space Weather Modeling Capabilities Assessment (Scherliess et al., 2019).

Of course, other issues that were discussed at the workshop (such as near real time availability of observations and DA) are very important, but the first focus here is on assessment and developing the whole atmosphere models themselves.

There was a further suggestion that the joint development of parametrizations would be

incredibly useful in unifying parametrization strategy across multiple models. The International Space Science Institute has a good setup for accomplishing verification with data, and this setting would be helpful for deciding a verification strategy. To monitor progress, it was also agreed to organize a follow up workshop in mid-2020.

Acknowledgments

Acknowledgements We thank Fabrizio Sassi (Naval Research Laboratory) for supplying Figure 1. No data were generated for use in this manuscript.

References

- Akmaev, R. A. (2011). Whole atmosphere modeling: Connecting terrestrial and space weather. *Reviews of Geophysics*, *49*, RG4004. <https://doi.org/10.1029/2011RG000364>
- Akmaev, R. A., Fuller-Rowell, T. J., Wu, F., Forbes, J. M., Zhang, X., Anghel, A. F., et al. (2008). Tidal variability in the lower thermosphere: Comparison of Whole Atmosphere Model (WAM) simulations with observations from TIMED. *Geophysical Research Letters*, *35*, L03810. <https://doi.org/10.1029/2007GL032584>
- Billett, D. D., Grocott, A., Wild, J. A., Walach, M.-T., & Kosch, M. J. (2018). Diurnal variations in global Joule heating morphology and magnitude due to neutral winds. *Journal of Geophysical Research: Space Physics*, *123*, 2398–2411. <https://doi.org/10.1002/2017JA025141>
- Borchert, S., Zhou, G., Baldauf, M., Schmidt, H., Zängl, G., & Reinert, D. (2018). The upper-atmosphere extension of the ICON general circulation model. *Geoscientific Model Development Discussion*, 1–50. <https://doi.org/10.5194/gmd-2018-289>
- Bruinsma, S., Sutton, E., Solomon, S. C., Fuller-Rowell, T., & Fedrizzi, M. (2018). Space weather modeling capabilities assessment: Neutral density for orbit determination at low Earth orbit. *Space Weather*, *16*, 1806–1816. <https://doi.org/10.1029/2018SW002027>
- Chartier, A. T., Jackson, D. R., & Mitchell, C. N. (2013). A comparison of the effects of initializing different thermosphere-ionosphere model fields on storm time plasma density forecasts. *Journal of Geophysical Research: Space Physics*, *118*, 7329–7337. <https://doi.org/10.1002/2013JA019034>
- Chartier, A. T., Matsuo, T., Anderson, J. L., Collins, N., Hoar, T. J., Lu, G., et al. (2016). Ionospheric data assimilation and forecasting during storms. *Journal of Geophysical Research: Space Physics*, *121*, 764–778. <https://doi.org/10.1002/2014JA020799>
- Collier, N., Hoffman, F. M., Lawrence, D. M., Keppel-Aleks, G., Koven, C. D., Riley, W. J., et al. (2018). The International Land Model Benchmarking (ILAMB) system: Design, theory, and implementation. *Journal of Advances in Modeling Earth Systems*, *10*, 2731–2754. <https://doi.org/10.1029/2018MS001354>
- Curry, J. A., & Webster, P. J. (1998). *Thermodynamics of atmospheres & oceans, International Geophysics Series* (Vol. 65). London: Academic Press.

- Eckermann, S. D., Broutman, D., Ma, J., Doyle, J. D., Pautet, P., Taylor, M. J., et al. (2016). Dynamics of orographic gravity waves observed in the mesosphere over the Auckland Islands during the Deep Propagating Gravity Wave Experiment (DEEPWAVE). *Journal of the Atmospheric Sciences*, *73*(10), 3855–3876. <https://doi.org/10.1175/JAS-D-16-0059.1>
- Evensen, G. (1994). Sequential data assimilation with a nonlinear quasi-geostrophic model using Monte Carlo methods to forecast error statistics. *Journal of Geophysical Research*, *99*, 10,143–10,162. <https://doi.org/10.1029/94JC00572>
- Eyring, V., Arblaster, J. M., Cionni, I., Sedláček, J., Perlwitz, J., Young, P. J., et al. (2013). Long-term ozone changes and associated climate impacts in CMIP5 simulations. *Journal of Geophysical Research: Atmospheres*, *118*, 5029–5060. <https://doi.org/10.1002/jgrd.50316>
- Fang, T.-W., Anderson, D., Fuller-Rowell, T. J., Akmaev, R., Codrescu, M., Millward, G., et al. (2013). Comparative studies of theoretical models in the equatorial ionosphere. In J. D. Huba (Ed.), *Modeling the Ionosphere-Thermosphere*, Geophys. Monogr. Ser. (Vol. 201, pp. 133–144). Washington, DC: American Geophysical Union.
- Feng, W., Marsh, D. R., Chipperfield, M. P., Janches, D., Höffner, J., Yi, F., & Plane, J. M. C. (2013). A global atmospheric model of meteoric iron. *Journal of Geophysical Research: Atmospheres*, *118*, 9456–9474. <https://doi.org/10.1002/jgrd.50708>
- Fomichev, V. I., & Blanchet, J.-P. (1995). Development of the new CCC/GCM radiation model for extension into the Middle Atmosphere. *Atmosphere-Ocean*, *33*(3), 513–529. <https://doi.org/10.1080/07055900.1995.9649543>
- Fomichev, V. I., Blanchet, J.-P., & Turner, D. S. (1998). Matrix parameterization of the 15 μm CO₂ band cooling in the middle and upper atmosphere for variable CO₂ concentration. *Journal of Geophysical Research*, *103*(D10), 11,505–11,528. <https://doi.org/10.1029/98JD00799>
- Fujiwara, H., & Miyoshi, Y. (2010). Morphological features and variations of temperature in the upper thermosphere simulated by a whole atmosphere GCM. *Annales de Geophysique*, *28*(2), 427–437. <https://doi.org/10.5194/angeo-28-427-2010>
- Fuller-Rowell, T. J., Akmaev, R. A., Wu, F., Anghel, A. F., Maruyama, N., Anderson, D. N., et al. (2008). Impact of terrestrial weather on the upper atmosphere. *Geophysical Research Letters*, *35*, L09808. <https://doi.org/10.1029/2007GL032911>
- Gardner, L. C., & Schunk, R. W. (2011). Large-scale gravity wave characteristics simulated with a high-resolution global thermosphere-ionosphere model. *Journal of Geophysical Research*, *116*, A06303. <https://doi.org/10.1029/2010JA015629>
- Gill, E., Sundaramoorthy, P., Bouwmeester, J., Zandbergen, B., & Reinhard, R. (2013). Formation flying within a constellation of nanosatellites: The QB50 mission. *Acta Astronautica*, *82*(1), 110–117. <https://doi.org/10.1016/j.actaastro.2012.04.029>
- Goncharenko, L., Chau, J., Liu, H.-L., & Coster, A. J. (2010). Unexpected connections between the stratosphere and ionosphere. *Geophysical Research Letters*, *37*, L10101. <https://doi.org/10.1029/2010GL043125>
- Goncharenko, L., Coster, A. J., Chau, J., & Valladares, C. (2010). Impact of sudden stratospheric warmings on equatorial ionization anomaly. *Journal of Geophysical Research*, *115*, A00G07. <https://doi.org/10.1029/2010JA015400>
- Griffin, D. J., & Thuburn, J. (2018). Numerical effects on vertical wave propagation in deep-atmosphere models. *Quarterly Journal of The Royal Meteorological Society*, *144*(711), 567–580.
- Heelis, R. A., Lowell, J. K., & Spiro, R. W. (1982). A model of the high-latitude ionospheric convection pattern. *Journal of Geophysical Research*, *87*(A8), 6339–6345. <https://doi.org/10.1029/JA087iA08p06339>
- Hsu, C. T., Matsuo, T., Wang, W., & Liu, J. Y. (2014). Effects of inferring unobserved thermospheric and ionospheric state variables by using an Ensemble Kalman Filter on global ionospheric specification and forecasting. *Journal of Geophysical Research: Space Physics*, *119*, 9256–9267. <https://doi.org/10.1002/2014JA020390>
- Immel, T. J., Sagawa, E., England, S. L., Henderson, S. B., Hagan, M. E., Mende, S. B., et al. (2006). The control of equatorial ionospheric morphology by atmospheric tides. *Geophysical Research Letters*, *33*, L15108. <https://doi.org/10.1029/2006GL026161>
- Jin, H., Miyoshi, Y., Pancheva, D., Mukhtarov, P., Fujiwara, H., & Shinagawa, H. (2012). Response of migrating tides to the stratospheric sudden warming in 2009 and their effects on the ionosphere studied by a whole atmosphere-ionosphere model GAIA with COSMIC and TIMED/SABER observations. *Journal of Geophysical Research*, *117*, A10323. <https://doi.org/10.1029/2012JA017650>
- Kelly, C. W., Chipperfield, M. P., Plane, J. M. C., Feng, W., Sheese, P. E., Walker, K. A., & Boone, C. D. (2018). An explanation for the nitrous oxide layer observed in the mesopause region. *Geophysical Research Letters*, *45*, 7818–7827. <https://doi.org/10.1029/2018GL078895>
- Knipp, D. J., Pette, D. V., Kilcommons, L. M., Isaacs, T. L., Cruz, A. A., Mlynczak, M. G., et al. (2017). Thermospheric nitric oxide response to shock-led storms. *Space Weather*, *15*, 325–342. <https://doi.org/10.1002/2016SW001567>
- Lang, M., & Owens, M. J. (2019). A variational approach to data assimilation in the solar wind. *Space Weather*, *17*, 59–83. <https://doi.org/10.1029/2018SW0018>
- Lejeune, S., Wautelet, G., & Warnant, R. (2012). Ionospheric effects on relative positioning within a dense GPS network. *GPS Solutions*, *16*(1), 105–116. <https://doi.org/10.1007/s10291-011-0212-1>
- Lieberman, R. S., Akmaev, R. A., Fuller-Rowell, T. J., & Doornbos, E. (2013). Thermospheric zonal mean winds and tides revealed by CHAMP. *Geophysical Research Letters*, *40*, 2439–2443. <https://doi.org/10.1002/grl.50481>
- Liemohn, M., Ganushkina, N. Y., De Zeeuw, D. L., Rastaetter, L., Kuznetsova, M., Welling, D. T., et al. (2018). Real-time SWMF at CCMC: Assessing the Dst output from continuous operational simulations. *Space Weather*, *16*, 1583–1603. <https://doi.org/10.1029/2018SW001953>
- Liu, H.-L., Bardeen, C. G., Foster, B. T., Lauritzen, P., Liu, J., Lu, G., et al. (2018). Development and validation of the Whole Atmosphere Community Climate Model with thermosphere and ionosphere extension (WACCM-X 2.0). *Journal of Advances in Modeling Earth Systems*, *10*, 381–402. <https://doi.org/10.1002/2017MS001232>
- Liu, H.-L., Foster, B. T., Hagan, M. E., McInerney, J. M., Maute, A., Qian, L., et al. (2010). Thermosphere extension of the Whole Atmosphere Community Climate Model. *Journal of Geophysical Research*, *115*, A12302. <https://doi.org/10.1029/2010JA015586>
- Liu, H.-L., McInerney, J. M., Santos, S., Lauritzen, P. H., Taylor, M. A., & Pedatella, N. M. (2014). Gravity waves simulated by high-resolution Whole Atmosphere Community Climate Model. *Geophysical Research Letters*, *41*, 9106–9112. <https://doi.org/10.1002/2014GL062468>
- Liu, H.-L., & Roble, R. G. (2002). A study of a self-generated stratospheric sudden warming and its mesospheric–lower thermospheric impacts using the coupled TIME-GCM/CCM3. *Journal of Geophysical Research*, *107*(D23), 4695. <https://doi.org/10.1029/2001JD001533>
- López-Valverde, M. A., & López-Puertas, M. (1994). A non-local thermodynamic equilibrium radiative transfer model for infrared emissions in the atmosphere of Mars. 1: Theoretical basis and nighttime populations of vibrational levels. *Journal of Geophysical Research*, *99*(E6), 13,093–13,115. <https://doi.org/10.1029/94JE00635>
- Marsh, D. R., Garcia, R. R., Kinnison, D. E., Boville, B. A., Sassi, F., Solomon, S. C., & Matthes, K. (2007). Modeling the whole atmosphere response to solar cycle changes in radiative and geomagnetic forcing. *Journal of Geophysical Research*, *112*, D23306. <https://doi.org/10.1029/2006JD008306>

- Marsh, D. R., Mills, M. J., Kinnison, D. E., & Lamarque, J.-F. (2013). Climate Change from 1850 to 2005 Simulated in CESM1(WACCM). *Journal of Climate*, *26*(19), 7372–7391. <https://doi.org/10.1175/JCLI-D-12-00558.1>
- Martynenko, O. V., Fomichev, V. I., Semeniuk, K., Beagley, S. R., Ward, W. E., McConnell, J. C., & Namgaladze, A. A. (2014). Physical mechanisms responsible for forming the 4-peak longitudinal structure of the 135.6 nm ionospheric emission: First results from the Canadian IAM. *Journal of Atmospheric and Solar - Terrestrial Physics*, *120*, 51–61. <https://doi.org/10.1016/j.jastp.2014.08.014>
- Maruyama, N., Sun, Y.-Y., Richards, P. G., Middlecoff, J., Fang, T.-W., Fuller-Rowell, T. J., et al. (2016). A new source of the midlatitude ionospheric peak density structure revealed by a new Ionosphere-Plasmasphere model. *Geophysical Research Letters*, *43*, 2429–2435. <https://doi.org/10.1002/2015GL067312>
- McCormack, J. P., Hoppel, K., Kuhl, D., de Wit, R., Stober, G., Espy, P., et al. (2017). Comparison of mesospheric winds from a high-altitude meteorological analysis system and meteor radar observations during the boreal winters of 2009–2010 and 2012–2013. *Journal of Atmospheric and Solar - Terrestrial Physics*, *154*, 132–166. <https://doi.org/10.1016/j.jastp.2016.12.007>
- McDonald, S. E., Sassi, F., Tate, J., McCormack, J., Kuhl, D. D., Drob, D. P., et al. (2018). Impact of non-migrating tides on the low latitude ionosphere during a sudden stratospheric warming event in January 2010. *Journal of Atmospheric and Solar-Terrestrial Physics*, *171*, 188–200. <https://doi.org/10.1016/j.jastp.2017.09.012>
- Merkin, V. G., Kondrashov, D., Ghil, M., & Anderson, B. J. (2016). Data assimilation of low-altitude magnetic perturbations into a global magnetosphere model. *Space Weather*, *14*, 165–184. <https://doi.org/10.1002/2015SW001330>
- Miyoshi, Y., Jin, H., Fujiwara, H., & Shinagawa, H. (2018). Numerical study of traveling ionospheric disturbances generated by an upward propagating gravity wave. *Journal of Geophysical Research: Space Physics*, *123*, 2141–2155. <https://doi.org/10.1002/2017JA025110>
- Murray, S. A., Henley, E. M., Jackson, D. R., & Bruinsma, S. L. (2015). Assessing the performance of thermospheric modeling with data assimilation throughout solar cycles 23 and 24. *Space Weather*, *13*, 220–232. <https://doi.org/10.1002/2015SW001163>
- Ogibalov, V. P., & Fomichev, V. I. (2003). Parameterization of solar heating by the near IR CO₂ bands in the mesosphere. *Advances in Space Research*, *32*(5), 759–764. [https://doi.org/10.1016/S0273-1177\(03\)80069-8](https://doi.org/10.1016/S0273-1177(03)80069-8)
- Pedatella, N. M., Fang, T.-W., Jin, H., Sassi, F., Schmidt, H., Chau, J. L., et al. (2016). Multimodel comparison of the ionosphere variability during the 2009 sudden stratosphere warming. *Journal of Geophysical Research: Space Physics*, *121*, 7204–7225. <https://doi.org/10.1002/2016JA022859>
- Pedatella, N. M., Liu, H.-L., Richmond, A. D., Maute, A., & Fang, T.-W. (2012). Simulations of solar and lunar tidal variability in the mesosphere and lower thermosphere during sudden stratosphere warmings and their influence on the low-latitude ionosphere. *Journal of Geophysical Research*, *117*, A08326. <https://doi.org/10.1029/2012JA017858>
- Plane, J. M. C., Feng, W., & Dawkins, E. C. M. (2015). The mesosphere and metals: Chemistry and changes. *Chemical Reviews*, *115*(10), 4497–4541. <https://doi.org/10.1021/cr500501m>
- Scherliess, L., Tsagouri, I., Yizengaw, E., Bruinsma, S., Shim, J. S., Coster, A., & Retterer, J. M. (2019). The International Community Coordinated Modeling Center space weather modeling capabilities assessment: Overview of ionosphere/thermosphere activities. *Space Weather*, *17*, 527–538. <https://doi.org/10.1029/2018SW002036>
- Schuck, R. W., Scherliess, L., Eccles, V., Gardner, L. C., Sojka, J. J., Zhu, L., et al. (2016). Space weather forecasting with a Multimodel Ensemble Prediction System (MEPS). *Radio Science*, *51*, 1157–1165. <https://doi.org/10.1002/2015RS005888>
- Sheese, P. E., Walker, K. A., Boone, C. D., Bernath, P. F., & Funke, B. (2016). Nitrous oxide in the atmosphere: First measurements of a lower thermospheric source. *Geophysical Research Letters*, *43*, 2866–2872. <https://doi.org/10.1002/2015GL067353>
- Taylor, K. E., Stouffer, R. J., & Meehl, G. A. (2012). An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, *93*(4), 485–498. <https://doi.org/10.1175/BAMS-D-11-00094.1>
- Thuburn, J. (2008). Some conservation issues for the dynamical cores of NWP and climate models. *Journal of Computational Physics*, *227*(7), 3715–3730. <https://doi.org/10.1016/j.jcp.2006.08.016>
- Ullrich, P. A., Jablonowski, C., Kent, J., Lauritzen, P. H., Nair, R., Reed, K. A., et al. (2017). DCMIP2016: a review of non-hydrostatic dynamical core design and intercomparison of participating models. *Geoscientific Model Development*, *10*(12), 4477–4509. <https://doi.org/10.5194/gmd-10-4477-2017>
- Vadas, S. L., & Crowley, G. (2010). Sources of the traveling ionospheric disturbances observed by the ionospheric TIDDBIT sounder near Wallops Island on October 30, 2007. *Journal of Geophysical Research*, *115*, A07324. <https://doi.org/10.1029/2009JA015053>
- Vadas, S. L., & Fritts, D. C. (2005). Thermospheric responses to gravity waves: Influences of increasing viscosity and thermal diffusivity. *Journal of Geophysical Research*, *110*, D15103. <https://doi.org/10.1029/2004JD005574>
- Wang, H., Akmaev, R. A., Fang, T.-W., Fuller-Rowell, T. J., Wu, F., Maruyama, N., & Iredell, M. D. (2014). First forecast of a sudden stratospheric warming with a coupled whole-atmosphere/ionosphere model IDEA. *Journal of Geophysical Research: Space Physics*, *119*, 2079–2089. <https://doi.org/10.1002/2013JA019481>
- Weimer, D. (2005). Improved ionospheric electrodynamic models and application to calculating Joule heating rates. *Journal of Geophysical Research*, *110*, A05306. <https://doi.org/10.1029/2004JA010884>
- Yiğit, E., Aylward, A. D., & Medvedev, A. S. (2008). Parameterization of the effects of vertically propagating gravity waves for thermosphere general circulation models: Sensitivity study. *Journal of Geophysical Research*, *113*, D19106. <https://doi.org/10.1029/2008JD010135>
- Yiğit, E., Koucká Knížová, P., Georgieva, K., & Ward, W. (2016). A review of vertical coupling in the Atmosphere–Ionosphere system: Effects of waves, sudden stratospheric warmings, space weather, and of solar activity. *Journal of Atmospheric and Solar-Terrestrial Physics*, *141*, 1–12. <https://doi.org/10.1016/j.jastp.2016.02.011>