

# THE JOURNAL OF PHYSICAL CHEMISTRY B

Subscriber access provided by Brought to you by ST ANDREWS UNIVERSITY LIBRARY

B: Liquids, Chemical and Dynamical Processes in Solution, Spectroscopy in Solution

## Efficient Singlet Oxygen Photogeneration by Zinc Porphyrin-Dimers Upon One- and Two-Photon Excitation

Leszek Mateusz Mazur, Thomas J. Roland, Stephanie Leroy-Lhez, Vincent Sol, Marek Samoc, Ifor D.W. Samuel, and Katarzyna Matczyszyn

*J. Phys. Chem. B*, **Just Accepted Manuscript** • DOI: 10.1021/acs.jpcc.8b12561 • Publication Date (Web): 05 Mar 2019

Downloaded from <http://pubs.acs.org> on March 12, 2019

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



ACS Publications

is published by the American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# Efficient Singlet Oxygen Photogeneration by Zinc Porphyrin-Dimers Upon One- and Two-Photon Excitation.

*Leszek M. Mazur<sup>a</sup>, Thomas Roland<sup>b</sup>, Stephanie Leroy-Lhez<sup>c</sup>, Vincent Sol<sup>c</sup>, Marek*

*Samoc<sup>a</sup>, Ifor D.W. Samuel<sup>b</sup>, Katarzyna Matczyszyn<sup>\*a</sup>*

<sup>a</sup> Advanced Materials Engineering and Modelling Group, Faculty of Chemistry, Wrocław

University of Science and Technology, Wybrzeże Wyspińskiego 27, 50-370 Wrocław,

Poland

<sup>b</sup> Organic Semiconductor Centre, SUPA, School of Physics and Astronomy, University

of St Andrews, Fife, KY16 9SS, UK

<sup>c</sup> Laboratoire PEIRENE EA 7500, Faculté des Sciences et Techniques 123, Avenue

Albert Thomas, Limoges cedex 87060, France

## ABSTRACT

The development of photodynamic therapy at depth requires photosensitizers which have both sufficient quantum yield for singlet oxygen generation and strong two-photon absorption. Here we show that this can be achieved by conjugated linkage of zinc porphyrins to make dimers. We determined the quantum yield of generation of  $^1\text{O}_2$ ,  $\phi_\Delta$ , by measuring emission at 1270 nm using a near infra-red streak camera and found it to increase from 15% for a single porphyrin unit to 27-47% for the dimers with a conjugated linker. Then, we measured the spectra of two-photon absorption cross section,  $\sigma_2$ , by a focus-tunable Z-scan method, which allows for nondestructive investigation of light-sensitive materials. We observed a strong enhancement of the two-photon absorption coefficient in the dimers, especially those with an alkyne linker. These results lead to an excellent figure of merit for two-photon production of singlet oxygen (expressed by the product  $\sigma_2 \times \phi_\Delta$ ) in the porphyrin dimers, of around 3700 GM, which is very promising for applications involving treatment of deep tumors by photodynamic therapy.

## 1. INTRODUCTION

1  
2  
3  
4 Photodynamic therapy (PDT) is a widely killing *Propionibacterium acnes*),  
5  
6  
7 used cancer treatment, particularly for the periodontology<sup>6</sup>, treatment of viral lesions<sup>7</sup>  
8  
9  
10 treatment of superficial cancers such as and age-related macular degeneration<sup>8</sup>.  
11  
12  
13 many skin cancers<sup>1</sup>. PDT involves the use In order to efficiently photogenerate  
14  
15  
16 of light in combination with a singlet oxygen, a material should fulfill a  
17  
18  
19 photosensitizer to cause a photochemical few criteria. It needs to absorb excitation  
20  
21  
22 reaction, leading to the generation of light and have a triplet energy level higher  
23  
24  
25 reactive oxygen species (ROS)<sup>2</sup>, which than 397 kJ/mol to allow for efficient  
26  
27  
28 then kill neighboring tumor cells. Whilst energy transfer to ground state oxygen  
29  
30  
31 most often used to kill cancer cells, the and hence the generation of singlet  
32  
33  
34 reactive oxygen species can also be used oxygen. Moreover, it should have high  
35  
36  
37 to kill bacteria thereby providing a triplet quantum yield and long triplet state  
38  
39  
40  
41  
42 powerful alternative to antibiotics in the lifetime<sup>2</sup>. Despite these requirements,  
43  
44  
45 management of topical infections<sup>3, 4</sup>. For there are many materials capable of  
46  
47  
48 example, in addition to cancers, PDT has photogenerating singlet oxygen, they can  
49  
50  
51  
52  
53 been used in the treatment of acne<sup>5</sup> (by be grouped into: (1) the organic dyes and  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 aromatics<sup>9</sup>; (2) the porphyrins, chlorins, wavelengths in the biological  
4  
5  
6 phthalocyanines, and related transparency windows of tissue at  
7  
8  
9  
10 macrocycles<sup>10</sup>; (3) semiconductors<sup>11</sup>; and 650-950 nm and 1000-1250 nm<sup>14</sup>. As only  
11  
12  
13  
14 (4) metal nanoparticles<sup>12, 13</sup>. Among these few photosensitizers have a linear  
15  
16  
17 photosensitizers, porphyrin derivatives absorption band in this wavelength  
18  
19  
20 attracted significant attention and some of region<sup>15</sup>, two-photon excitation in which  
21  
22  
23  
24 them are already in clinical use. two longer wavelength photons are  
25  
26  
27

28 Photodynamic therapy has been widely simultaneously absorbed is a natural way  
29  
30  
31 used to kill tumor cells, which are located to achieve the desired excitation energy  
32  
33  
34 close to skin surface, due to the fact that with longer wavelength light that can  
35  
36  
37 the photosensitizers are excited with penetrate further into tissue. The  
38  
39  
40 wavelengths which are strongly absorbed photosensitizers which aim to be used in  
41  
42  
43 and scattered by the tissue. PDT could two-photon excited PDT should be  
44  
45  
46 treat a much wider range of conditions if characterized by high quantum yield of  
47  
48  
49 could work at greater depth which could singlet oxygen generation,  $\phi_{\Delta}$ , and large  
50  
51  
52 be achieved by using excitation two-photon absorption cross-sections.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Typical porphyrin derivatives<sup>16-18</sup>, even consider this figure of merit when  
5  
6 those accepted for clinical use like normalized by molecular weight i.e.  
7  
8 Photofrin<sup>®</sup>, Verteporfin<sup>®</sup> or Foscan<sup>®</sup> are  $\sigma_2 \times \phi_{\Delta} / M$ , marked as  $(\sigma_2 \times \phi_{\Delta})^*$  to facilitate  
9  
10 indeed efficient singlet oxygen generators comparison of large and small candidate  
11  
12 but suffer from low two-photon absorption photosensitizers. In the case of  
13  
14 cross-sections, on the order of 1 GM<sup>19, 20</sup>. porphyrins, in order to maximize the  
15  
16 In that context, there is a need to design parameter  $\sigma_2 \times \phi_{\Delta}$ , insertion of a metal ion  
17  
18 and synthesize new molecules with leads to a trade-off as it causes reduction  
19  
20 excellent figure of merit for two-photon of the singlet oxygen quantum yield<sup>21</sup> and  
21  
22 production of singlet oxygen (expressed increase of the 2PA cross-section.  
23  
24 by the product  $\sigma_2 \times \phi_{\Delta}$ ). It is also relevant to  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 In the field of nonlinear optics, the been shown that centrosymmetric  
43  
44 general rules for molecular design to quadrupolar compounds have larger  
45  
46 maximize 2PA cross-section exist in the values of  $\sigma_2$  than their dipolar  
47  
48 literature and are supported by both counterparts<sup>22</sup>. In addition, introducing  
49  
50 theoretical and experimental work. It has strong electron donating (D) or electron  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


1  
2  
3 withdrawing (A) groups to yield the or higher order oligomers<sup>17, 23, 24</sup>. In the  
4 structures like D-A-D or A-D-A further present work we use metal containing  
5  
6 structures like D-A-D or A-D-A further present work we use metal containing  
7  
8 improves 2PA properties<sup>23</sup>. Finally, the porphyrins and explore the effect of  
9  
10 process of simultaneous absorption of two increasing electron delocalization by  
11  
12 photons requires strong overlap of the conjugated links between macrocycles. In  
13  
14  $\pi$ -orbitals over a large system, making it particular, we explore the photogeneration  
15  
16 sensitive to molecular conformation of singlet oxygen and two-photon  
17  
18 changes, which may alter the electronic absorption in a series of zinc porphyrins,  
19  
20 coupling. either being a monomer (compound 1), or  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

34  
35 The general structure-property rules can a dimer, linked by ethyne (2) or di-ethyne  
36  
37 be applied to porphyrin based link (3) or by a single bond (4), as depicted  
38  
39 compounds. It has been shown that 2PA in Scheme 1.  
40  
41  
42  
43  
44


45  
46 properties may be tuned by metal There are numerous papers  
47  
48 insertion, adding electron donating or investigating either the two-photon  
49  
50 withdrawing groups and expanding the absorption or singlet oxygen  
51  
52  $\pi$ -conjugated system by creating dimers photogeneration properties of porphyrin  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 based compounds, but not so many where  
4  
5  
6  
7 the combination of these two effects is  
8  
9  
10 presented<sup>25-27</sup>. Our goal was to carry out  
11  
12  
13 interdisciplinary research covering the  
14  
15  
16 nonlinear optical properties and singlet  
17  
18 oxygen photogeneration, thus providing  
19  
20  
21 strong support to the results and  
22  
23  
24 conclusions concerning the structure-  
25  
26  
27 property optimization. Singlet oxygen  
28  
29  
30 generation was studied using an infra-red  
31  
32  
33  
34  
35  
36  
37

streak camera. The two-photon absorption  
properties were investigated by the focus-  
tunable Z-scan method, which has the  
important advantage over alternative  
methods of being quick, thereby avoiding  
photodegradation which is a particular  
consideration in the measurement of  
materials intended to cause  
photochemistry.







**Scheme 1.** Structures of investigated compounds.

## 2. EXPERIMENTAL SECTION

The molecules used in this work were synthesized and purified according to the protocols described in<sup>28</sup>. The molecules were dissolved in chloroform (Sigma-Aldrich) and quartz cuvettes were filled with the solutions for the photophysical measurements. UV-vis absorption spectra were measured using a Perkin Elmer Lambda 1050 UV/Vis/NIR

spectrophotometer. The concentration of the solution was adjusted to reach optical densities suitable for the PL and emission from singlet oxygen experiments. Steady-state PL spectra were recorded using an automated spectrofluorometer (Fluorolog, Horiba Jobin-Yvon), with a 450 W Xenon lamp excitation source. The solution singlet oxygen quantum yield (SOQY) ( $\phi_{\Delta}$ )

values were determined using PHAROS regenerative amplifier (output tetraphenylporphyrin (H<sub>2</sub>TPP) from Fisher Scientific in chloroform solution as a standard reference because of its well-known SOQY of 55%<sup>29</sup>. The direct emission of singlet oxygen at 1270 nm was detected by a NIR streak camera from Hamamatsu Photonics. The camera was working in Photon Counting mode and the experiment was automatically stopped after 50 000 frames for each sample. In order to prevent the degradation, the samples solutions were placed in a rotating cell, placed in such a way that the exciting beam was shifted from the center.

The beam from a Light Conversion

PHAROS regenerative amplifier (output wavelength 1028 nm, pulse duration 290 fs, repetition rate 5 kHz) was directed to an optical parametric amplifier (OPA, a Light Conversion OPHEUS-F). The OPA generated 828 nm, which was further frequency doubled to obtain 414 nm used to excite the samples. The output pulse duration was less than 100 fs. The quantum yield of singlet oxygen generation may be determined using the modified method used by Tanielian and Heinrich<sup>30</sup> and using equation 1<sup>21</sup>:

$$\phi_{\Delta x} = \phi_{\Delta r} \frac{A_r(\lambda_r) I(\lambda_r) D_x \tau_r}{A_x(\lambda_x) I(\lambda_x) D_r \tau_s} \quad (1)$$

1  
2  
3 where  $\phi_{\Delta}$  stands for the quantum yield of  
4  
5  
6  
7 singlet oxygen generation, the subscripts  
8  
9  
10  $x$  and  $r$  refer to the compound to be  
11  
12  
13 characterized and to the reference  
14  
15  
16 solutions, respectively.  $\lambda$  is the excitation  
17  
18  
19 wavelength,  $A(\lambda)$  is the absorbance,  $I(\lambda)$  is  
20  
21  
22 the relative intensity of the excitation light,  
23  
24  
25  
26  
27  $\tau$  is the lifetime of singlet oxygen emission  
28  
29  
30  
31 in the sample and reference solvent, and  
32  
33  
34  
35  $D$  is the integrated area under the  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
emission spectra.

As, the samples and reference solutions  
are excited at the same conditions and  
they are dissolved in chloroform, the  
above equation can be simplified to:

$$\phi_{\Delta x} = \phi_{\Delta r} \frac{A_r(\lambda_r) D_x}{A_x(\lambda_x) D_r} \quad (2)$$

The two-photon absorption cross-sections were determined by f-scan method described in<sup>31</sup>. The laser system used for this study was a Quantronix Integra Ti:Sapphire regenerative amplifier (output wavelength: 800 nm; pulse duration: 130 fs; repetition rate: 1 kHz) that pumped an optical parametric amplifier (OPA, a Quantronix Palitra) to provide tunable excitation. The sample solution was prepared by dissolving the corresponding chromophore in chloroform at concentration of 0.5% (w/w) and placed in a glass cell with 1 mm optical path-

length. The measurements were carried out in a relative manner, calibrating all the data against f-scans carried out on a fused silica plate and taking into account the nonlinear signals obtained from a cell containing pure solvent.

### 3. RESULTS AND DISCUSSION

#### Steady-state electronic absorption.

The absorption and photoluminescence spectra of the investigated compounds were measured in dilute  $\text{CHCl}_3$  solutions with  $1 \times 10^{-6}$  mol/L concentration. Maximum absorption wavelengths and maximum emission wavelengths are listed in Table 1. The absorption spectra are displayed in Figure 1. For the monomeric





porphyrin, compound **1**, (Figure 1a), two absorption bands are clearly visible. The Soret band is located at 414 nm and so called Q-band at 543 nm. For the dimers, the Q-bands are intensified they show bathochromic shift compared to molecule **1**. Moreover, both Soret and Q-bands are much broader with additional peaks visible in the absorption spectra for compounds **2-4**. It is worth to mention that the stronger Q-band absorption is responsible for enhanced 2PA properties in the NIR. Fluorescence occurs from the Q-band, whichever band is excited. The fluorescence bands of compounds **2** and **3** show a bathochromic shift with respect

1  
2  
3 to compound **1** (Figure 1a-c). The as  $B_x(0-0)$  transition. There are also other  
4 bathochromic shift of the absorption and  $B_x$  and  $B_y$  transitions which span a wide  
5  
6 fluorescence bands can be explained by region to the blue from the  $B_x(0-0)$   
7 the decreased HOMO-LUMO gap due to transition. In the case of compound **4** the  
8  $\pi$ -delocalization between the neighboring molecule is no longer planar due to steric  
9 hindrance, and so the dimerization does  
10 not lead to increased electron  
11 delocalization. There is an almost equally  
12 intense split in the Soret absorption and Q-  
13 band emission of **4** (Figure 1d), which can  
14 be explained by the excitonic coupling  
15 between the two parallel strong dipole  
16 transitions of each porphyrin ring<sup>28, 32</sup>. The  
17 magnitude of the absorption band split is  
18 equal to 2002  $\text{cm}^{-1}$  and the emission band  
19 split is equal to 854  $\text{cm}^{-1}$ .  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The crystallographic structure analysis performed by another group yielded the torsional angle of the porphyrin dimer linked by a single bond to be  $69.1(2)^\circ$ <sup>33</sup>.

The torsional angle of the dimers linked by ethyne and di-ethyne bonds were

determined by DFT calculations to be zero, which means that the molecules are planar<sup>34</sup>, which enhances both 2PA and singlet oxygen photogeneration due to faster intersystem crossing ( $S_1-T_1$ ), compared to twisted conformers<sup>35</sup>.



1  
2  
3  
4 **Figure 1.** Normalized absorption (black curves) and steady-state fluorescence spectra  
5  
6  
7 (red curves) of investigated porphyrins. Figures a) to d) correspond to molecules **1** to **4**,  
8  
9  
10 respectively.


### 15 **Singlet oxygen detection.**

16  
17  
18 The singlet oxygen quantum yield,  $\phi_{\Delta}$ , emission. The spectra obtained for sample  
19  
20 was determined by recording direct **4** and the reference are shown in Figure 2.  
21  
22 emission from singlet oxygen centered at Finally, the spectra were integrated to  
23  
24 1270 nm by a NIR streak camera. The obtain the  $D$  factors necessary to calculate  
25  
26 transition from first excited singlet state the quantum yield of singlet oxygen  
27  
28  $^1\Delta_g$  to ground state being triplet,  $^3\Sigma_g^-$ , is photogeneration. The values of  $\phi_{\Delta}$  of all  
29  
30 spin forbidden<sup>2</sup>. This is the reason for the investigated compounds are summarized  
31  
32 emission lifetime on the order of  $\mu$ s, which in Table 1. The least efficient is the  
33  
34 strongly depends on the solvent<sup>21</sup>. monomeric unit (**1**). The dimerization  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 The signals were integrated from 8 to leads to enhanced properties of singlet  
50  
51  
52 50  $\mu$ s to obtain the spectra of  $^1\text{O}_2$  oxygen photogeneration, regardless the  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 type of the link. One can notice that the  
4  
5  
6  
7 value of  $\phi_{\Delta}$  for compound **4**, which is not  
8  
9  
10 conjugated due to steric hindrance, is  
11  
12 approximately doubled, compared to **1**.  
13  
14  
15 On the other hand, compounds **2** and **3**,  
16  
17  
18 which have large  $\pi$ -delocalization and are  
19  
20  
21 close to planar, are characterized with  
22  
23  
24 much larger  $\phi_{\Delta}$  being equal to 35 and  
25  
26  
27 47 %. These findings are in good  
28  
29  
30 agreement with other works on porphyrins  
31  
32  
33 dimers, e.g. with porphyrin units linked by  
34  
35  
36 butadiyne bond linked to TIPS-protected  
37  
38  
39 ethynyl groups<sup>36</sup>. The determined  $\phi_{\Delta}$  of  
40  
41  
42 such porphyrin dimer in DCM was  
43  
44  
45 reported to be 37 %, which means that the  
46  
47  
48 structural design of compounds **2** and **3**

are more efficient in terms of singlet  
oxygen production.



**Figure 2.** Singlet oxygen emission spectra of **4** and H<sub>2</sub>TPP, obtained by integrating the streak camera images from 8 to 50  $\mu$ s.

**Table 1.** Steady-state photophysical properties of the investigated compounds (**1-4**) and their singlet oxygen quantum yields. The positions of the most intensive bands are given in bold.

<b>porphyrins</b>	$\lambda_{\text{abs}}$ (nm)	$\lambda_{\text{em}}$ (nm)	$\phi_{\Delta}$ (%)
-------------------	-----------------------------	----------------------------	---------------------



	Soret band	Q-band		
<b>1</b>	393 <b>414</b>	<b>542</b>	554 585 <b>637</b> 700	15
<b>2</b>	420 441 <b>450</b> 482	566 622 <b>673</b>	627 <b>681</b> 751	35

**Two-photon absorption cross-section spectra.**


The spectra of two-photon absorption cross-section may be obtained by many methods, two-photon excited fluorescence (TPEF)<sup>37</sup> and fs Z-scan<sup>38-41</sup> being the most common nowadays. They strongly differ from each other, as in TPEF one detects fluorescence spectrum at various wavelengths and compares the integrated PL with a standard reference with known PLQY and  $\sigma_2$ . In Z-scan one

<b>3</b>	414 431 444 <b>481</b>	562 <b>681</b>	599 654 <b>706</b> 785	47
<b>4</b>	417 <b>455</b>	538 <b>558</b> 600	567 <b>624</b> <b>658</b>	27

measures the changes of sample transmission as a function of the sample position in Z direction and fit the data recorded on so called Open Aperture curve as described in the literature<sup>38, 41</sup> to determine the value of two-photon absorption cross-section at the specific wavelength. In order to obtain reliable data by these two aforementioned techniques, the samples have to be perfectly photostable.

1  
2  
3  
4       Upon photoexcitation with one- and       the f-scan technique, instead of translating  
5  
6  
7       two-photons the singlet oxygen which is       the sample, the focal point is shifted by an  
8  
9  
10       generated by the investigated porphyrins       electrically controlled lens<sup>31</sup>. Combined  
11  
12  
13  
14       causes their degradation. It is important to       with improved data acquisition technique,  
15  
16  
17       mention that each photosensitizer       this method reduces the time necessary  
18  
19  
20  
21       molecule can typically produce  $10^3$ - $10^5$        for a single scan from a few minutes to  
22  
23  
24       molecules of  $^1\text{O}_2$  before being degraded       several seconds, thus limiting the  
25  
26  
27  
28       through photobleaching by  $^1\text{O}_2$  or by some       possibility of photodegradation. While the  
29  
30  
31       other process<sup>2</sup>. Thus, it is crucial to       vast majority of published papers  
32  
33  
34  
35       minimize the time of irradiation especially       described only the singlet oxygen  
36  
37  
38       with high power laser used to study the       photogeneration process, or two-photon  
39  
40  
41  
42       nonlinear effects. Hence we use the so-       absorption properties of the studied  
43  
44  
45       called “f-scan” technique which is a       photosensitizers, in this work both are  
46  
47  
48       modification of the standard Z-scan       treated with equal attention and detail,  
49  
50  
51  
52       method, to determine the spectra of       allowing for more reliable conclusions.  
53  
54  
55  
56       two-photon absorption cross-section. In

The spectra of two-photon absorption cross-sections of investigated porphyrins obtained by f-scan technique are shown in Figure 3. They are overlaid with linear absorption spectra, shown as red curves, plotted vs. twice the wavelength. For all porphyrins, both the monomer (1) and dimers (2-4) the maximum of two-photon absorption cross-sections is within first biological window which exists at wavelengths between 650 nm and 950 nm<sup>14</sup>.



1  
2  
3 **Figure 3.** Spectra of two-photon absorption cross-sections of the investigated porphyrins  
4  
5  
6  
7 (1-4). The spectra were overlaid with linear absorption (red curves) plotted vs. twice the  
8  
9  
10 wavelength. Figures a) to d) correspond to compounds 1 to 4.

16 In case of the monomer 1, the maximum means that it is favorable for the  
17 value of  $\sigma_2$  is around 1000 GM at 725 nm, molecules to be excited to higher vibronic  
18 as shown in Figure 3. a). One can see that state upon 2PA excitation, compared to  
19 the band of 2PA is split into two, even 1PA process. Despite the fact that linear  
20 though the linear absorption shows only absorption of 4 is split into two equally  
21 one peak at 414 nm. It is caused by intense bands, the 2PA spectrum is  
22 different selection rules for one- and different. One can see that the band at  
23 two-photon absorption, as described in the 725 nm is much more intensive than that at  
24 literature<sup>22</sup>. It is worth noticing that the high 875 nm. The maximum value of  $\sigma_2$ ,  
25 energy 2PA band at 725 nm is observed measured at 725 nm is approximately  
26 for all the compounds and has the highest doubled compared to those of compound  
27 value of  $\sigma_2$  among other 2PA bands. It 1 and is equal to 2500 GM. In the studies

1  
2  
3 of two-photon absorption cross-sections but its position remains the same, the  
4  
5  
6 various normalization factors are used, wavelength of maximum  $\sigma_2$  is equal to  
7  
8  
9 among which the normalization to molar 725 nm. However, the low energy band,  
10  
11 mass is the most popular. If the maximum which is located at 875 nm for other  
12  
13  $\sigma_2$  is divided by molar mass, a similar investigated compounds, is  
14  
15 value is obtained, which means that such bathochromically shifted to 1050 nm.  
16  
17 structural change does not impose neither Similarly to compound 1, there is a dip in  
18  
19 cooperation nor deleterious effects on the the two-photon absorption cross-section  
20  
21 multiphoton absorption properties of the spectra at twice the wavelength of  
22  
23 investigated compounds 1 and 4. In case maximum 1PA band. The 2PA spectrum  
24  
25 of compounds 2 and 3, the spectra of of compound 3 is presented in Figure 3. c).  
26  
27 two-photon absorption cross-section show The higher energy band is  
28  
29 more complex structures, similarly to their bathochromically shifted by 50 nm to  
30  
31 1PA spectra. For compound 2 775 nm and the value of  $\sigma_2$  is around 8000  
32  
33 (Figure 3. b), the higher energy 2PA band GM being slightly lower than for  
34  
35 is broader than for compounds 1 and 4, compound 2. For both compounds 2 and  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

3, the maximum value of two-photon absorption cross-section is enhanced few times, while keeping the molar mass similar to compound 4. The results prove that a simple modification of the linker, from single bond (4) to ethyne (2) and di-ethyne (3) links between the zinc porphyrins cores cause cooperation effect, enhancing the  $\sigma_{2^*}$  figure of merit.

The values of maximum two-photon absorption cross-section and the figure of merit are listed in Table 2. The dimers of zinc porphyrins, especially connected by ethyne and di-ethyne link, are much more effective two-photon photosensitizers than tetraphenyl porphyrin, or other porphyrin derivatives that are commercially used for PDT.

**Table 2.** Two-photon absorption cross-section, merit factors and two-photon excited oxygen sensitization properties of the investigated porphyrins (1-4).

	$\lambda_{2PA}^{max}$ (nm)	$\sigma_2^{max}$ (GM)	$\sigma_2^{max*}$ (GM·mol/g)	$\sigma_2^{max} \cdot \phi_{\Delta}$ (GM)	$\sigma_2^{max} \cdot \phi_{\Delta}^*$ (GM·mol/g)	Two-photon excited oxygen sensitization enhancement factor <sup>[c]</sup>
<b>1</b>	725	1030	1.4	155	0.21	21
<b>2</b>	725	8900	6.1	3115	2.15	433
<b>3</b>	775	7950	5.4	3737	2.53	519
<b>4</b>	725	2500	1.7	675	0.45	94
H <sub>2</sub> TPP	790	12 <sup>[a]</sup>	0.02	7.2 <sup>[b]</sup>	0.01	1

<sup>[a]</sup> taken from reference <sup>42</sup>

1  
2  
3 [b] taken from reference <sup>27</sup>  
4

5 [c] Enhancement factor:  $\sigma_2^{max} \cdot \phi_{\Delta}$  of the compound normalized to that of H<sub>2</sub>TPP.  
6  
7

8 It was previously shown e.g. by TPEF conditions in the three-level system and  
9 measurements<sup>17</sup>, that dimerization of dramatic enhancement of the excited-  
10 porphyrin units leads to enhanced 2PA state singlet-singlet transition due to linear  
11 cross-sections and enhanced two-photon elongation of the  $\pi$ -conjugated system.  
12 production of singlet oxygen. The  $\sigma_2$  Insertion of side groups, being either  
13 determined for the porphyrin dimer linked electron acceptors (A) or donors (D)  
14 by ethyne and di-ethyne bonds were on influence the  $\sigma_2$  in a way that two-photon  
15 the order of 8200 and 5500 GM, which is absorption cross-section is the lowest for  
16 similar to the values found in the current neutral porphyrins, higher for the  
17 work for similar porphyrin derivatives. The porphyrins extended with A and the  
18 cooperative effect was explained by a highest for compounds with D groups<sup>43</sup>.  
19 combination of several factors, including Even though the vast majority of  
20 strong enhancement of the lowest one- molecules targeted to be used in 2PA PDT  
21 photon Q-transition, better resonance are based on the porphyrin structure,  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 there are some alternative approaches photogeneration properties in a series of  
4 with reasonable figures of merit. One zinc porphyrins, either in a monomer or  
5 example is a macromolecular dimer form. The dimers were connected  
6 photosensitizer, which was obtained via by single bond, or by ethyne, or di-ethyne  
7 the covalent anchoring of several link. We observed strong enhancement of  
8 molecular PS based on a the obtained quantum yields for  
9 dibromobenzene (DBB) core onto a water- two-photon excited singlet oxygen  
10 soluble polymer chain based on generation (expressed by the product  
11 poly(N-acryloylmorpholine). It resulted in  $\sigma_2^{max} \phi_{\Delta}$ ) in the porphyrin dimer  
12 large 2PA cross-section up to 2600 GM at connected by di-ethyne link, around  
13 740 nm and was proved to be an efficient 3700 GM, which is around 500 times more  
14 candidate to cause melanoma cells death efficient than tetraphenyl porphyrin. It can  
15 upon photo irradiation<sup>44</sup>. be seen that in case of zinc porphyrin

#### 48 4. CONCLUSIONS

49 In this work, we investigated two-photon high two-photon absorption cross-section  
50 absorption and singlet oxygen simultaneously enhances the properties of  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4 singlet oxygen photogeneration. Thus, in  
5  
6  
7 order to obtain a material capable of  
8  
9  
10 efficient production of singlet oxygen upon  
11  
12  
13 two-photon excitation, one should  
14  
15  
16  
17 increase the conjugation length, so that  
18  
19  
20  
21  $\pi$ -delocalization is maximized, while  
22  
23  
24 maintaining planar structure, as shown for  
25  
26  
27  
28 compounds **2** and **3**. The results are very  
29  
30  
31 promising for applications involving  
32  
33  
34  
35 treatment of deep tumors by  
36  
37  
38 photodynamic therapy.  
39  
40  
41

## 42 AUTHOR INFORMATION

### 43 Corresponding Author

44  
45  
46 katarzyna.matczyszyn@pwr.edu.pl;  
47  
48 idws@st-andrews.ac.uk  
49

### 50 ORCID

51  
52  
53 Leszek M. Mazur: 0000-0002-4562-0743  
54  
55  
56  
57  
58  
59  
60

Vincent Sol: 0000-0003-0175-0156

Marek Samoc: 0000-0002-5404-2455

Ifor D.W. Samuel: 0000-0001-7821-7208

Katarzyna Matczyszyn: 0000-0001-8578-  
8340

## Conflict of interest statement

There are no conflicts to declare.

## ACKNOWLEDGEMENTS

The authors gratefully acknowledge the support from National Science Centre (Poland) under Opus program DEC-2013/09/B/ST5/03417, the EPSRC of the UK through grant EP/L017008/1 and the Conseil Régional du Limousin in France for financial support. The work was financed by a statutory activity subsidy

1  
2  
3 from the Polish Ministry of Science and  
4  
5  
6  
7 Higher Education for the Faculty of  
8  
9  
10 Chemistry of Wroclaw University of  
11  
12  
13 Science and Technology. IDWS  
14  
15  
16 acknowledges a Royal Society Wolfson  
17  
18  
19 Research Merit Award. We thank  
20  
21  
22 the European Union as this work is a part  
23  
24  
25 of the ITN-EJD-2017 project POLYTHEA  
26  
27  
28 which has received funding from the  
29  
30  
31 European Union's Horizon 2020 research  
32  
33  
34 and innovation programme under the  
35  
36  
37 Marie Sklodowska-Curie grant agreement  
38  
39  
40 n°764837.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

## 52 REFERENCES:

53 1. Brown, S. B.; Brown, E. A.; Walker, I.,  
54 The Present and Future Role of Photodynamic  
55 Therapy in Cancer Treatment. *Lancet Oncol.* **2004**,  
56 *5*, 497-508.

- 57  
58  
59  
60
2. DeRosa, M. C.; Crutchley, R. J.,  
Photosensitized Singlet Oxygen and Its  
Applications. *Coord. Chem. Rev.* **2002**, *233–234*,  
351-371.
3. Yin, R.; Agrawal, T.; Khan, U.; Gupta, G.  
K.; Rai, V.; Huang, Y.-Y.; Hamblin, M. R.,  
Antimicrobial Photodynamic Inactivation in  
Nanomedicine: Small Light Strides against Bad  
Bugs. *Nanomedicine* **2015**, *10*, 2379-2404.
4. Tardivo, J. P.; Adami, F.; Correa, J. A.;  
Pinhal, M. A. S.; Baptista, M. S., A Clinical Trial  
Testing the Efficacy of PDT in Preventing  
Amputation in Diabetic Patients. *Photodiagn.*  
*Photodyn. Ther.* **2014**, *11*, 342-350.
5. Bissonnette, R., Treatment of Acne with  
Photodynamic Therapy. *G. Ital. Dermatol.*  
*Venerol.* **2011**, *146*, 445-456.
6. Kumar, V.; Sinha, J.; Verma, N.; Nayan,  
K.; Saimbi, C.; Tripathi, A., Scope of  
Photodynamic Therapy in Periodontics. *Indian J.*  
*Dent. Res.* **2015**, *26*, 439-442.
7. Kharkwal, G. B.; Sharma, S. K.; Huang,  
Y.-Y.; Dai, T.; Hamblin, M. R., Photodynamic  
Therapy for Infections: Clinical Applications.  
*Lasers Surg. Med.* **2011**, *43*, 755-767.
8. Boscia, F.; Furino, C.; Sborgia, L.;  
Reibaldi, M.; Sborgia, C., Photodynamic Therapy  
for Retinal Angiomatous Proliferations and  
Pigment Epithelium Detachment. *Am. J.*  
*Ophthalmol.* **2004**, *138*, 1077-1079.
9. Redmond, R. W.; Gamlin, J. N., A  
Compilation of Singlet Oxygen Yields from  
Biologically Relevant Molecules. *Photochem.*  
*Photobiol.* **1999**, *70*, 391-475.
10. Murtinho, D.; Pineiro, M.; Pereira, M. M.;  
Rocha Gonsalves, A. M. d. A.; Arnaut, L. G.;  
Miguel, M. d. G.; Burrows, H. D., Novel  
Porphyrins and a Chlorin as Efficient Singlet  
Oxygen Photosensitizers for Photooxidation of  
Naphthols or Phenols to Quinones. *J. Chem. Soc.,*  
*Perkin Trans. 2* **2000**, 2441-2447.
11. Yamamoto, Y.; Imai, N.; Mashima, R.;  
Konaka, R.; Inoue, M.; Dunlap, W. C., Singlet  
Oxygen from Irradiated Titanium Dioxide and  
Zinc Oxide. *Methods Enzymol* **2000**, *319*, 29-37.
12. Chen, C.-W.; Hsu, C.-Y.; Lai, S.-M.; Syu,  
W.-J.; Wang, T.-Y.; Lai, P.-S., Metal Nanobullets  
for Multidrug Resistant Bacteria and Biofilms.  
*Adv. Drug Del. Rev.* **2014**, *78*, 88-104.
13. Huang, C.-C.; Liu, T.-M., Controlled Au-  
Polymer Nanostructures for Multiphoton Imaging,

- Prodrug Delivery, and Chemo-Photothermal Therapy Platforms. *ACS Appl. Mater. Interfaces* **2015**, *7*, 25259-25269.
14. Smith, A. M.; Mancini, M. C.; Nie, S., Bioimaging: Second Window for in vivo Imaging. *Nat. Nanotechnol.* **2009**, *4*, 710-711.
15. Habermeyer, B.; Guillard, R., Some Activities of Porphyrin Illustrated by the Applications of Porphyrinoids in PDT, PIT and PDI. *Photochem. Photobiol. Sci.* **2018**, *17*, 1675-1690.
16. Goyan, R. L.; Cramb, D. T., Near-Infrared Two-Photon Excitation of Protoporphyrin IX: Photodynamics and Photoproduct Generation. *Photochem. Photobiol.* **2000**, *72*, 821-827.
17. Drobizhev, M.; Stepanenko, Y.; Dzenis, Y.; Karotki, A.; Rebane, A.; Taylor, P. N.; Anderson, H. L., Extremely Strong near-IR Two-Photon Absorption in Conjugated Porphyrin Dimers: Quantitative Description with Three-Essential-States Model. *J. Phys. Chem. B* **2005**, *109*, 7223-7236.
18. Nowak-Król, A.; Wilson, C. J.; Drobizhev, M.; Kondratuk, D. V.; Rebane, A.; Anderson, H. L.; Gryko, D. T., Amplified Two-Photon Absorption in Trans-A2B2-Porphyrins Bearing Nitrophenylethynyl Substituents. *ChemPhysChem* **2012**, *13*, 3966-3972.
19. Balaz, M.; Collins, H. A.; Dahlstedt, E.; Anderson, H. L., Synthesis of Hydrophilic Conjugated Porphyrin Dimers for One-Photon and Two-Photon Photodynamic Therapy at NIR Wavelengths. *Org. Biomol. Chem.* **2009**, *7*, 874-888.
20. Karotki, A.; Khurana, M.; Lepock, J. R.; Wilson, B. C., Simultaneous Two-Photon Excitation of Photofrin in Relation to Photodynamic Therapy. *Photochem. Photobiol.* **2006**, *82*, 443-452.
21. Mathai, S.; Smith, T. A.; Ghiggino, K. P., Singlet Oxygen Quantum Yields of Potential Porphyrin-Based Photosensitisers for Photodynamic Therapy. *Photochem. Photobiol. Sci.* **2007**, *6*, 995-1002.
22. Pawlicki, M.; Collins, H. A.; Denning, R. G.; Anderson, H. L., Two-Photon Absorption and the Design of Two-Photon Dyes. *Angew. Chem. Int. Ed.* **2009**, *48*, 3244-3266.
23. Wilkinson, J. D.; Wicks, G.; Nowak-Król, A.; Lukasiewicz, L. G.; Wilson, C. J.; Drobizhev, M.; Rebane, A.; Gryko, D. T.; Anderson, H. L., Two-Photon Absorption in Butadiyne-Linked Porphyrin Dimers: Torsional and Substituent Effects. *J. Mater. Chem. C* **2014**, *2*, 6802-6809.
24. Kim, K. S.; Lim, J. M.; Osuka, A.; Kim, D., Various Strategies for Highly-Efficient Two-Photon Absorption in Porphyrin Arrays. *J. Photochem. Photobiol., C* **2008**, *9*, 13-28.
25. Yao, D.; Hugues, V.; Blanchard-Desce, M.; Mongin, O.; Paul-Roth, C. O.; Paul, F., Dendritic Molecular Assemblies for Singlet Oxygen Generation: Meso-Tetraphenylporphyrin-Based Biphotonic Sensitizers with Remarkable Luminescence. *New J. Chem.* **2015**, *39*, 7730-7733.
26. McIlroy, S. P.; Cló, E.; Nikolajsen, L.; Frederiksen, P. K.; Nielsen, C. B.; Mikkelsen, K. V.; Gothelf, K. V.; Ogilby, P. R., Two-Photon Photosensitized Production of Singlet Oxygen: Sensitizers with Phenylene-Ethynylene-Based Chromophores. *J. Org. Chem.* **2005**, *70*, 1134-1146.
27. Yao, D.; Zhang, X.; Triadon, A.; Richy, N.; Mongin, O.; Blanchard-Desce, M.; Paul, F.; Paul-Roth, C. O., New Conjugated Meso-Tetrafluorenylporphyrin-Cored Derivatives as Fluorescent Two-Photon Photosensitizers for Singlet Oxygen Generation. *Chem. - Eur. J.* **2017**, *23*, 2635-2647.
28. Jiblaoui, A.; Baudequin, C.; Chaleix, V.; Ducourthial, G.; Louradour, F.; Ramondenc, Y.; Sol, V.; Leroy-Lhez, S., An Easy One-Pot Desilylation/Copper-Free Sonogashira Cross-Coupling Reaction Assisted by Tetra-Butylammonium Fluoride (TBAF): Synthesis of Highly  $\pi$ -Conjugated Porphyrins. *Tetrahedron* **2013**, *69*, 5098-5103.
29. Poon, C. T.; Zhao, S. S.; Wong, W. K.; Kwong, D. W. J., Synthesis, Excitation Energy Transfer and Singlet Oxygen Photogeneration of Covalently Linked N-Confused Porphyrin-Porphyrin and Zn(II) Porphyrin Dyads. *Tetrahedron Lett.* **2010**, *51*, 664-668.
30. Tanielian, C.; Heinrich, G., Effect of Aggregation on the Hematoporphyrin-Sensitized Production of Singlet Molecular Oxygen. *Photochem. Photobiol.* **1995**, *61*, 131-135.
31. Kolkowski, R.; Samoc, M., Modified Z - Scan Technique Using Focus-Tunable Lens. *J. Opt.* **2014**, *16*, 125202.
32. L. Anderson, H., Building Molecular Wires from the Colours of Life: Conjugated

- 1  
2  
3 Porphyrin Oligomers. *Chem. Commun.* **1999**,  
4 2323-2330.
- 5 33. Senge Mathias, O.; Pinteá, M.; Ryan  
6 Aoife, A., Synthesis and Crystal Structure of a  
7 Meso-Meso Directly Linked Bisporphyrin. *Z.*  
8 *Naturforsch., B: Chem. Sci.* **2011**, *66*, 553-558.
- 9 34. Peeks, M. D.; Neuhaus, P.; Anderson, H.  
10 L., Experimental and Computational Evaluation of  
11 the Barrier to Torsional Rotation in a Butadiyne-  
12 Linked Porphyrin Dimer. *Phys. Chem. Chem.*  
13 *Phys.* **2016**, *18*, 5264-5274.
- 14 35. Kuimova, M. K.; Balaz, M.; Anderson, H.  
15 L.; Ogilby, P. R., Intramolecular Rotation in a  
16 Porphyrin Dimer Controls Singlet Oxygen  
17 Production. *J. Am. Chem. Soc.* **2009**, *131*, 7948-  
18 7949.
- 19 36. Schmitt, J.; Heitz, V.; Jenni, S.; Sour, A.;  
20 Bolze, F.; Ventura, B.,  $\pi$ -Extended Porphyrin  
21 Dimers as Efficient near-Infrared Emitters and  
22 Two-Photon Absorbers. *Supramol. Chem.* **2017**,  
23 *29*, 769-775.
- 24 37. Deiana, M.; Mettra, B.; Mazur, L. M.;  
25 Andraud, C.; Samoc, M.; Monnereau, C.;  
26 Matczyszyn, K., Two-Photon Macromolecular  
27 Probe Based on a Quadrupolar Anthracenyl  
28 Scaffold for Sensitive Recognition of Serum  
29 Proteins under Simulated Physiological  
30 Conditions. *ACS Omega* **2017**, *2*, 5715-5725.
- 31 38. Van Stryland, E. W.; Sheik-Bahae, M., Z-  
32 Scan Measurements of Optical Nonlinearities. In  
33 *Characterization Techniques and Tabulations for*  
34 *Organic Nonlinear Materials*, Kuzyk, M. G.; Dirk,  
35 C., Eds. Marcel Dekker: 1998; pp 655-692.
- 36 39. Gao, B.; Mazur, L. M.; Morshedi, M.;  
37 Barlow, A.; Wang, H.; Quintana, C.; Zhang, C.;  
38 Samoc, M.; Cifuentes, M. P.; Humphrey, M. G.,  
39 Exceptionally Large Two- and Three-Photon  
40 Absorption Cross-Sections by OPV  
41 Organometalation. *Chem. Commun.* **2016**, *52*,  
42 8301-8304.
- 43 40. Lin, T.-C.; Chien, W.; Mazur, L. M.; Liu,  
44 Y.-Y.; Jakubowski, K.; Matczyszyn, K.; Samoc,  
45 M.; Amini, R. W., Two- and Three-Photon  
46 Absorption Properties of Fan-Shaped Dendrons  
47 Derived from 2,3,8-Trifunctionalized  
48 Indenoquinoxaline Units: Synthesis and  
49 Characterization. *J. Mater. Chem. C* **2017**, *5*, 8219-  
50 8232.
- 51 41. Sheik-Bahae, M.; Said, A. A.; Wei, T. H.;  
52 Hagan, D. J.; Stryland, E. W. V., Sensitive  
53 Measurement of Optical Nonlinearities Using a  
54 Single Beam. *IEEE J. Quantum Electron.* **1990**,  
55 *26*, 760-769.
- 56 42. Makarov, N. S.; Drobizhev, M.; Rebane,  
57 A., Two-Photon Absorption Standards in the 550–  
58 1600 nm Excitation Wavelength Range. *Opt.*  
59 *Express* **2008**, *16*, 4029-4047.
- 60 43. Koszelewski, D.; Nowak-Krol, A.;  
Drobizhev, M.; Wilson, C. J.; Haley, J. E.; Cooper,  
T. M.; Romiszewski, J.; Gorecka, E.; Anderson, H.  
L.; Rebane, A.; Gryko, D. T., Synthesis and Linear  
and Nonlinear Optical Properties of Low-Melting  
 $\pi$ -Extended Porphyrins. *J. Mater. Chem. C* **2013**,  
*1*, 2044-2053.
44. Cepraga, C.; Marotte, S.; Ben Daoud, E.;  
Favier, A.; Lanoë, P.-H.; Monnereau, C.; Baldeck,  
P.; Andraud, C.; Marvel, J.; Charreyre, M.-T.;  
Leverrier, Y., Two-Photon Photosensitizer–  
Polymer Conjugates for Combined Cancer Cell  
Death Induction and Two-Photon Fluorescence  
Imaging: Structure/Photodynamic Therapy  
Efficiency Relationship. *Biomacromolecules*  
**2017**, *18*, 4022-4033.

