

University of Dundee

An ancestral non-proteolytic role for presenilin proteins in multicellular development of the social amoeba dictyostelium discoideum

Ludtmann, Marthe H.R.; Otto, Grant P.; Schilde, Christina; Chen, Zhi Hui; Allan, Claire Y.; Brace, Selina

Published in:
Journal of Cell Science

DOI:
[10.1242/jcs.140939](https://doi.org/10.1242/jcs.140939)

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Ludtmann, M. H. R., Otto, G. P., Schilde, C., Chen, Z. H., Allan, C. Y., Brace, S., ... Williams, R. S. B. (2014). An ancestral non-proteolytic role for presenilin proteins in multicellular development of the social amoeba dictyostelium discoideum. *Journal of Cell Science*, 127(7), 1576-1584. <https://doi.org/10.1242/jcs.140939>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

An ancestral non-proteolytic role for presenilin proteins in multicellular development of the social amoeba *Dictyostelium discoideum*

Marthe H. R. Ludtmann^{1,*‡}, Grant P. Otto^{1,‡}, Christina Schilde², Zhi-Hui Chen², Claire Y. Allan³, Selina Brace⁴, Philip W. Beesley¹, Alan R. Kimmel⁵, Paul Fisher³, Richard Killick⁶ and Robin S. B. Williams^{1,§}

ABSTRACT

Mutations in either of two presenilin genes can cause familial Alzheimer's disease. Presenilins have both proteolysis-dependent functions, as components of the γ -secretase complex, and proteolysis-independent functions in signalling. In this study, we investigate a conserved function of human presenilins in the development of the simple model organism *Dictyostelium discoideum*. We show that the block in *Dictyostelium* development caused by the ablation of both *Dictyostelium* presenilins is rescued by the expression of human presenilin 1, restoring the terminal differentiation of multiple cell types. This developmental role is independent of proteolytic activity, because the mutation of both catalytic aspartates does not affect presenilin ability to rescue development, and the ablation of nicastrin, a γ -secretase component that is crucial for proteolytic activity, does not block development. The role of presenilins during *Dictyostelium* development is therefore independent of their proteolytic activity. However, presenilin loss in *Dictyostelium* results in elevated cyclic AMP (cAMP) levels and enhanced stimulation-induced calcium release, suggesting that presenilins regulate these intracellular signalling pathways. Our data suggest that presenilin proteins perform an ancient non-proteolytic role in regulating intracellular signalling and development, and that *Dictyostelium* is a useful model for analysing human presenilin function.

KEY WORDS: *Dictyostelium*, γ -secretase, Presenilin

INTRODUCTION

To date, more than 170 different mutations in presenilin 1 (*PSEN1*) and 13 mutations in presenilin 2 (*PSEN2*) genes are known to give rise to a familial form of Alzheimer's disease (FAD) (De Strooper and Annaert, 2010). The protein products of the mammalian *PSEN* genes are components of an aspartate protease complex, γ -secretase (De Strooper, 2003), which is responsible for the regulated, intramembranous cleavage of a

number of type 1 transmembrane proteins (Wolfe, 2006), including the Notch receptors (Huppert et al., 2000) and the amyloid precursor protein (APP). It is from APP that β -amyloid (A β), the likely causative agent of both the familial and sporadic forms of Alzheimer's disease, is derived (De Strooper et al., 1998; Herreman et al., 2000). Prior to incorporation into the γ -secretase complex, the presenilin 1 protein undergoes autoproteolytic cleavage into two parts (Chávez-Gutiérrez et al., 2008). Presenilin proteins have also been shown to have a non-proteolytic function as a scaffold for the regulation of glycogen synthase kinase 3 β (GSK3 β)-dependent β -catenin phosphorylation (Kang et al., 1999; Kang et al., 2002). Finally, presenilin proteins have also been implicated in altered calcium signalling (Tu et al., 2006) through an unknown mechanism. Understanding the discrete roles of presenilin proteins thus remains an important goal in understanding basic cell function and the progression of Alzheimer's disease (De Strooper and Annaert, 2010).

Although research into presenilin protein function has often focussed on presenilin-knockout mice (Schaeffer et al., 2011), it has been difficult to determine the precise function of presenilins because the loss of both presenilins leads to embryonic lethality. A range of alternative non-mammalian models have been employed to examine the function of presenilin proteins, including *Physcomitrella patens* (Khandelwal et al., 2007), *Caenorhabditis elegans* (Calahorra and Ruiz-Rubio, 2011), *Danio rerio* (van Tijn et al., 2011) and *Drosophila melanogaster* (Coen et al., 2012). However, the social amoeba *Dictyostelium discoideum* is the simplest model organism that possesses two presenilin proteins as well as the other three components of the γ -secretase complex (Boeckeler and Williams, 2007; McMains et al., 2010). *Dictyostelium* has been extensively used in a range of motility (Janetopoulos and Firtel, 2008), developmental (Loomis and Shaulsky, 2011) and biomedical studies (Chang et al., 2012; Francione and Fisher, 2011; Ludtmann et al., 2011; Myre et al., 2011; Terbach et al., 2011), and has a number of experimental advantages over existing models (Williams et al., 2006). In this study, we employ the advantages of *Dictyostelium* to explore the role of the human presenilin 1 protein (PSEN1) in development. We generated an isogenic *Dictyostelium* strain lacking both presenilin A and presenilin B genes (*psenA*⁻/*psenB*⁻), and found that development was blocked at the aggregation stage prior to morphogenesis. We show that either human PSEN1 or *Dictyostelium* PsenB can rescue the developmental phenotype of this mutant. Presenilin proteins that are mutated at the catalytic aspartic acid residues retain the ability to rescue the developmental block, demonstrating that the proteolytic activity of presenilin is not required for phenotypic rescue. A conserved functional role for

¹Centre for Biomedical Sciences, School of Biological Sciences, Royal Holloway University of London TW20 0EX, UK. ²School of Life Sciences, University of Dundee, Dundee DD1 5EH, UK. ³Faculty of Science, Technology and Engineering, La Trobe University, Bundoora VIC 3086, Australia. ⁴Centre for Ecology and Evolution, School of Biological Sciences, Royal Holloway University of London, Egham, Surrey TW20 0EX, UK. ⁵NIDDK, National Institutes of Health, Bethesda, MD 20892, USA. ⁶Institute of Psychiatry, King's College London, London SE5 8AF, UK.

*Present address: UCL Institute of Neurology, London WC1N 3BG, UK.

[‡]These authors contributed equally to this work

[§]Author for correspondence (robin.williams@rhul.ac.uk)

Received 23 August 2013; Accepted 29 December 2013

the human protein in *Dictyostelium* provides new insights into the ancient function(s) of presenilin proteins.

RESULTS

Dictyostelium presenilins share substantial similarity with human presenilins

The social amoeba *Dictyostelium discoideum* represents one of the earliest branches from the common ancestor of all eukaryotes, and thus provides a useful means for understanding the ancestral eukaryotic genome (Eichinger et al., 2005). It is also a simple widely used model organism for understanding developmental signalling and is increasingly being used to investigate the role of disease-related proteins (Williams et al., 2006). The *Dictyostelium* genome sequence has identified orthologues of a range of proteins associated with human diseases and conditions (Eichinger et al., 2005), including two presenilin proteins (PsenA and PsenB). Phylogenetic analysis reveals that they are more closely related (sister) to the monophyletic animal presenilin clade than to the distinct monophyletic plant presenilin clade (Fig. 1A). Furthermore, PsenA and PsenB (Fig. 1A) are similar in size and structure to the two human homologues and, like their human counterparts, they contain the conserved catalytic aspartic acid residues for proteolytic activity (Wolfe, 2006) and the GxDG motif for γ -secretase integration (Fig. 1B). A comparison of the predicted transmembrane domains reveals that these regions are 43–50% identical in the *Dictyostelium* and human proteins. Additionally, more than half (67/112) of the residues in either PSEN1 or PSEN2 that are mutated in FAD are conserved in either *Dictyostelium* PsenA or PsenB (supplementary material Fig. S1) [see also fig. S2 in McMains et al. (McMains et al., 2010)]. The sequence similarity and predicted common structure between the human and *Dictyostelium* presenilin proteins suggests a possible conservation of their function in these distantly related species.

Presenilins function redundantly in development

Presenilin proteins have previously been shown to play a key role in *Dictyostelium* development (McMains et al., 2010). In wild-type cells, starvation triggers the aggregation of ~100,000 individual amoebae that progress through a series of multicellular morphological stages over a 24-h period,

ultimately giving rise to a fruiting body comprising spores held within a sorus suspended above the substratum by a stalk composed of dead vacuolated cells (Fig. 2) (Williams et al., 2006). To test whether PsenA and PsenB act redundantly, we ablated each *Dictyostelium* gene individually using homologous integration of a knockout cassette (Faix et al., 2004) (supplementary material Fig. S2). The ablation of either *Dictyostelium* gene individually did not alter this basic developmental morphology under our conditions (supplementary material Fig. S3A), in contrast to earlier reports showing a developmental block following *psenB* ablation (McMains et al., 2010), which might be accounted for by a difference in parental strain or by differing experimental conditions. However, the removal of both genes (generating a line hereafter referred to as *psenA*⁻/*psenB*⁻ cells) caused a strong block in development (Fig. 2). The *psenA*⁻/*psenB*⁻ cells aggregated to complete early development, forming a rounded structure attached to the substratum, but morphogenesis halted at the beginning of tip formation. This supports the role previously suggested for *Dictyostelium* presenilins in development (McMains et al., 2010) and demonstrates that the presenilin homologues have redundant functions, consistent with what is observed in mammals (Feng et al., 2004; Kim et al., 2011).

Human PSEN1 rescues the defective development of *psenA*⁻/*psenB*⁻ cells

To investigate whether *Dictyostelium* and human presenilin proteins have conserved functions, we expressed the PsenB and PSEN1 proteins tagged with an N-terminal GFP (Veltman et al., 2009) in the *psenA*⁻/*psenB*⁻ genetic background. Expression of either PsenB or PSEN1 reversed the morphological block in development in this mutant (Fig. 2), resulting in the formation of wild-type fruiting bodies composed of both a stalk and a fully formed sorus (Fig. 2). We also expressed the human PSEN2 protein tagged with an N-terminal GFP, and found that development was partially rescued, so that it progressed beyond the block at the tipped mound observed in *psenA*⁻/*psenB*⁻ cells, to the early culminant (Fig. 2). Because PSEN2 expression did not fully complement development in the *psenA*⁻/*psenB*⁻ genetic background, our subsequent analyses were restricted to human




Fig. 1. The *Dictyostelium* PsenA and PsenB proteins show structural similarity to human presenilins. (A) Bayesian-derived phylogeny of presenilin amino-acid-sequence data from animal and plant species and *D. discoideum*. Human SPPLA2A was used as the outgroup. Numbers adjacent to nodes represent Bayesian posterior probability values. Scale bar: the number of substitutions per site. (B) Schematic of both *Dictyostelium* presenilin proteins showing the putative orientation in the membrane, cytosolic and transmembrane regions, the conserved catalytic residues and the proline-alanine-leucine (PAL) sequence.




Fig. 2. *Dictyostelium* presentilin proteins redundantly control multicellular development that is complemented by expression of human presentilin 1. Following starvation, wild-type (WT) cells undergo development over a 24-h period, leading to the formation of fruiting bodies; left, aerial view at low magnification; right, side view at higher magnification. Ablation of both presentilin genes ($psenA^-/psenB^-$) in one cell line gives rise to a block in fruiting-body formation, leading to a small round structure lacking a stalk. Overexpression of GFP-tagged *Dictyostelium* PsenB or human PSEN1 in these cells restores development, whereas PSEN2 expression partially rescues development. Scale bars: 1 mm.

PSEN1. This developmental complementation by PSEN1 in this model organism suggests that the function of PSEN1 is conserved from humans to *Dictyostelium*.

Our previous study also identified altered development in *Dictyostelium* lacking presentilin activity (McMains et al., 2010), so we assessed whether the *Dictyostelium* and human presentilin proteins could complement these defects in stalk- and spore-cell production and developmental-marker expression (Fig. 3). Stalk-cell formation was induced in low-density cultures in the presence of differentiation induction factor-1 (DIF-1) (Fig. 3A)

(Williams et al., 1999). The ablation of either *psenA* or *psenB* had no effect on DIF-1-induced stalk-cell formation, whereas the $psenA^-/psenB^-$ mutant, although still able to form stalk cells, showed a significant 31% reduction in stalk-cell production ($P < 0.05$), mirroring the developmental phenotype of each mutant. We also analysed the inhibition of this developmental pathway by cAMP, which leads to a glycogen-synthase-kinase-A (GSKA)-dependent block in stalk-cell production (GSKA is the *Dictyostelium* homologue of mammalian GSK3 β) (Williams et al., 1999). The ablation of either *Dictyostelium* presentilin gene had no effect on cAMP-mediated inhibition of stalk-cell differentiation compared with wild-type cells, whereas the presentilin double-null mutant showed a significant 84% increase in cAMP-dependent stalk-cell inhibition compared with wild-type cells ($P < 0.004$), which is consistent with the absence of a stalk in the fruiting body after 24 h. Furthermore, complementing the $psenA^-/psenB^-$ mutant with PsenB and PSEN1 restored stalk-cell production to levels close to that achieved by wild-type cells either in the presence or absence of cAMP.

Because *Dictyostelium* presentilin proteins have previously been shown to regulate spore production in monolayer assays (McMains et al., 2010), we also assessed whether expressing the *Dictyostelium* and human proteins complemented this aspect of multicellular development (Fig. 3B). The ablation of either *psenA* or *psenB* caused a 40% reduction in spore production during multicellular development. However, the $psenA^-/psenB^-$ mutant showed a complete loss of spore formation. The expression of either PsenB or PSEN1 restored spore production to the level of single knockouts, confirming the functionality of the human presentilin protein in *Dictyostelium* development. Together with the effects on stalk-cell differentiation and the observed developmental arrest at the initiation of tip formation, our results indicate that presentilin proteins are required not for cell-type choice but for differentiation along both the stalk and spore pathways.

Finally, as an independent verification of developmental complementation, we used quantitative transcriptional analysis (qPCR) to monitor the expression of developmental and cell-type markers in the $psenA^-/psenB^-$ mutant and following complementation with the *Dictyostelium* and human proteins (Fig. 3C). The cAMP receptor 1 (*cARI*) gene is expressed in wild-type cells in early development (Saxe, III et al., 1991), whereas a threefold increase in expression in early (6 h) development and elevated expression in mid-late (12–24 h) development was observed in the $psenA^-/psenB^-$ mutant compared with wild-type cells. In addition, the pre-stalk-cell marker *ecmA* (Harwood, 2008) showed a highly significant reduction in expression in the $psenA^-/psenB^-$ mutant in early and mid-development, whereas the pre-spore-cell marker *psA* (Harwood, 2008) showed only a small elevation in the double mutant compared with wild-type cells after 24 h of development. Complementation with PsenB returned marker expression to wild-type levels. Complementation with PSEN1 only partially rescued marker expression, with levels not reaching those of either wild-type cells or $psenA^-/psenB^-$ cells complemented with PsenB, suggesting a reduced efficacy of the human protein in *Dictyostelium*. Also, expression of either human or *Dictyostelium* proteins caused an elevation of late (24 h) stalk-cell-marker expression and mid-late (12–24 h) spore-cell-marker expression, suggesting a common controlling role for both presentilin proteins in terminal differentiation. These data suggest that the




Fig. 3. Expression of PsenB or PSEN1 can rescue the block in multiple differentiation pathways resulting from the ablation of presenilin activity in the *psenA*⁻/*psenB*⁻ mutant. (A) Stalk-cell production is unaffected by ablation of either presenilin individually but is reduced when both presenilins are removed and is restored by expression of PsenB or PSEN1. (B) Spore production is reduced by ablation of either presenilin gene individually, is blocked in *psenA*⁻/*psenB*⁻ mutants and is restored by expression of PsenB or PSEN1. Insets show light microscopy images of the cell type analysed in each panel. (C) The expression of developmentally regulated genes (0, 6, 12 and 24 h post-starvation), using quantitative transcriptional analysis for the early developmental marker *cAR1* and the cell-type-specific marker genes (pre-stalk, *ecmA*; pre-spore, *psA*) in wild-type and in *psenA*⁻/*psenB*⁻ cells, and following complementation with PsenB or PSEN1. Increased *cAR1* expression and reduced *ecmA* expression caused by the loss of both presenilin genes is restored by the expression of PsenB or PSEN1. All data are presented as the mean \pm standard deviation. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

differentiation of pre-stalk and pre-spore cells in the mound preceding tip formation is not dramatically impaired by the loss of presenilin (but is enhanced by the ectopic expression of either the *Dictyostelium* or human presenilin protein). By contrast, the further differentiation to mature stalk and spore cells is blocked, as is morphogenesis beyond the beginning of tip formation. These data also confirm the functionality of the human presenilin 1 protein in stalk- and spore-cell differentiation in *Dictyostelium*.

The presenilins are predominantly localised at the endoplasmic reticulum in *Dictyostelium*

We next examined the subcellular localisation of the expressed *Dictyostelium* and human proteins in the *psenA*⁻/*psenB*⁻ mutant. In mammals, presenilins are localised on the endoplasmic reticulum (ER) (Walter et al., 1996) and nuclear membrane (Li et al., 1997). In *Dictyostelium*, presenilin subcellular localisation was determined by fixing cells expressing GFP-tagged versions of PsenB and PSEN1 to reveal that both proteins were located on a fine interlaced network of membrane structures resembling the ER and nuclear envelope (Fig. 4). Colocalisation with an ER-specific marker, calnexin (Müller-Taubenberger et al., 2001), confirmed that both the human and *Dictyostelium* presenilin proteins localise to the ER and nuclear envelope in *D. discoideum*. These data once again confirm the similar behaviour of human and *Dictyostelium* presenilin proteins.

The proteolytic activity of presenilins is dispensable for development

We next investigated the molecular mechanism of presenilin function in *Dictyostelium*. The human presenilin proteins have been ascribed either a proteolytic function within the γ -secretase complex, dependent upon two crucial catalytic aspartic-acid residues in the protein (Yamasaki et al., 2006), or a structural role related to the docking of intracellular signalling components

(Fluhrer et al., 2004; Tesco and Tanzi, 2000; Twomey and McCarthy, 2006). The *Dictyostelium* presenilin proteins have been shown to proteolytically process recombinant human APP in an identical manner to APP processing in mammalian cells (McMains et al., 2010). This is probably due to γ -secretase activity because the proteolytic ability is lost upon ablation of the γ -secretase components nicastrin (Ncstn) and anterior pharynx defective 1 (Aph1) (McMains et al., 2010). To establish a role for γ -secretase proteolytic activity in *Dictyostelium* development, we mutated the catalytic aspartic-acid residues to alanines in both *Dictyostelium* and human proteins [D348A/D394A in PsenB or D257A/D385A in PSEN1 (Tesco and Tanzi, 2000); see supplementary material Fig. S1] and examined whether these mutant presenilin proteins could complement the defective fruiting-body formation of *psenA*⁻/*psenB*⁻ cells. Expression of either the *Dictyostelium* or human double aspartic-acid-mutated protein rescued *Dictyostelium* morphological development, enabling wild-type fruiting bodies to form (Fig. 5A). Furthermore, ablation of the *ncstn* gene, which is necessary for γ -secretase proteolytic ability (Edbauer et al., 2003), had no effect on fruiting-body development (Fig. 5A). As an independent verification of the normal developmental morphology that we observed in these experiments, we measured spore production in the complemented *psenA*⁻/*psenB*⁻ cells or the *ncstn*⁻ mutant and confirmed that spore production was similar to the wild-type cells (Fig. 5B). These results suggest that presenilin proteins do not require proteolytic activity, presumably as part of the γ -secretase complex, to function in *Dictyostelium* development.

The proteolytic activity of mammalian presenilins depends on their PEN2-dependent autoproteolytic cleavage at the intracellular loop, giving rise to a 34-kDa N-terminal and a 22-kDa C-terminal fragment (CTF) (Ahn et al., 2010; Haass and De Strooper, 1999). We therefore probed protein extracts from *psenA*⁻/*psenB*⁻ cells transformed with GFP-tagged wild-type PSEN1 and PSEN2 by western blotting, to ascertain whether




Fig. 4. The *Dictyostelium* and human presenilins localise to the ER in *Dictyostelium* cells. The GFP tag on PsenB and PSEN1 proteins indicates that both proteins colocalise with the ER marker calnexin (labelled with a specific antibody). Blue, 4',6-diamidino-2-phenylindole (DAPI) staining. Scale bar: 10 μ m.

cleavage occurred. This analysis showed that a band of 20–25 kDa corresponding to the CTF was detected for both proteins (Fig. 5C), indicating that endoproteolysis of both the human presenilins occurs in *Dictyostelium*. This result suggests that these human proteins can form an active protease complex together with the remaining *Dictyostelium* γ -secretase components.

Presenilins function in cAMP signalling and calcium homeostasis in *Dictyostelium*

We then investigated two signalling pathways that might be controlled by presenilin activity during *Dictyostelium* development. A role for presenilins in cAMP signalling was likely, based upon two observations: first, *psenA*⁻/*psenB*⁻ cells do not show streaming during early development (where a visible

trail of aggregating cells is seen leading to a mound; supplementary material Movie 1 and supplementary material Fig. S3B) – a phenotype associated with altered cAMP signalling (Garcia and Parent, 2008; Veltman and van Haastert, 2008); and second, the mutant cells also show increased sensitivity to cAMP-induced stalk-cell inhibition in late development (Fig. 3A). Therefore, we assessed intracellular cAMP levels during early development in wild-type cells, the *psenA*⁻/*psenB*⁻ mutant and double-mutant cells complemented with PsenB or PSEN1 (Fig. 6A). No significant difference was found between the different strains during growth ($t=0$ h) or early development ($t=4$ h). However, at $t=8$ h, the *psenA*⁻/*psenB*⁻ mutant showed significantly higher cAMP levels than wild-type cells ($P<0.001$). This was rescued following complementation with PsenB and




Fig. 5. *Dictyostelium* fruiting-body development does not require the proteolytic activity of presenilins or a complete γ -secretase complex. (A) Expression of *Dictyostelium* PsenB^{D348A/D394A} or human PSEN1^{D257A/D385A}, lacking the key catalytic aspartic-acid residues that are necessary for proteolytic activity, rescued *Dictyostelium* development in *psenA*⁻/*psenB*⁻ cells. The ablation of nicastrin (*ncstr*⁻) had no effect on *Dictyostelium* fruiting-body (developmental) morphology. Images show low-magnification aerial view (left) and high-magnification side view (right). Scale bars: 1mm. (B) Spore production in *psenA*⁻/*psenB*⁻ cells is restored by expression of either *Dictyostelium* or human presenilins mutated at the two catalytic aspartic-acid residues and is unaffected by the ablation of nicastrin (*ncstr*⁻). WT, wild type. Results show means \pm s.e.m. Inset shows a light microscopy image of the cell type analysed. (C) Both human presenilins, when expressed in *Dictyostelium* *psenA*⁻/*psenB*⁻ cells, undergo endoproteolysis to yield a single band of 20–25 kDa, corresponding to the C-terminal fragment, as demonstrated by western analysis. Molecular-mass markers in kilodaltons are indicated on the left; -, untransformed *psenA*⁻/*psenB*⁻ control; BD6, mouse-blastocyst-like cells showing processing of endogenous mouse presenilins. The antibodies used for western blotting (WB) are indicated below the blots.




Fig. 6. Loss of presenilin proteins in *Dictyostelium* alters cAMP and calcium levels. (A) cAMP levels in cells during growth ($t=0$ h) and early development ($t=4$ and 8 h) in wild-type (WT) cells and *psenA*⁻/*psenB*⁻ cells with and without complementation (rescue) with PsenB or PSEN1. Data are derived from five independent experiments and show a statistically significant increase in cAMP in *psenA*⁻/*psenB*⁻ and the PSEN1-rescue cells at 8 h only. (B) Schematic representation of the calcium-response recording prior to and upon 50 nM cAMP stimulation; A, the time until calcium response after cAMP stimulation; B, the length of the calcium response. The magnitude of the cAMP-induced calcium release is also shown. Data for A and B are presented in supplementary material Fig. S4. (C) The magnitude of the calcium response in wild-type and presenilin-mutant cells. The magnitude of the calcium response is significantly increased in *psenA*⁻, *psenB*⁻ and *psenA*⁻/*psenB*⁻ cells when compared to wild-type cells. Values shown are means \pm s.e.m, $n=5$. * $P<0.05$, ** $P<0.01$, *** $P<0.001$.

reduced following complementation with PSEN1, and both proteins rescued the lack of streaming observed in early development (supplementary material Fig. S3B). These data demonstrate a role for the *Dictyostelium* presenilin proteins in regulating cAMP signalling, an effect that has also been seen in mammals (Müller et al., 2011; Wang et al., 2011). Once again, these results demonstrate that the *Dictyostelium* and human proteins have overlapping functions and perform a conserved role in *Dictyostelium* development.

Because both human and *Dictyostelium* presenilin proteins are localised to the ER and presenilin ablation has been shown to alter calcium signalling in mammals (Tu et al., 2006), we also analysed whether calcium levels were affected in the presenilin mutants. In these experiments, cells expressing apoaequorin were loaded with coelenterazine and calcium levels were measured by luminescence in resting cells and following cAMP stimulation (Fig. 6). This analysis showed that the resting calcium levels, the cAMP-dependent time until calcium response and the duration of the calcium response were not significantly different in the single- and double-presenilin-null cells when compared to wild-type cells (supplementary material Fig. S4). However, the magnitude of the cAMP-induced calcium response was 2–2.5-fold greater in *psenA*⁻, *psenB*⁻ and *psenA*⁻/*psenB*⁻ cells ($P<0.05$, $P<0.01$ and $P<0.01$, respectively) when compared to the wild-type cells (Fig. 6C). These results suggest a role for presenilin proteins in *Dictyostelium* calcium homeostasis.

***Dictyostelium* presenilins can cleave mammalian substrates**

The conserved function for the human presenilin 1 protein in *Dictyostelium* development prompted us to examine whether the reverse is true; that is, whether *Dictyostelium* presenilins can function in mammalian cells. We tested the ability of presenilin proteins to function in the human γ -secretase complex to cleave Notch-1 within the transmembrane domain, as measured by the activation of a construct containing a Notch-and-CBF1-dependent luciferase reporter gene (Hooper et al., 2006). We transfected

mouse-blastocyst-derived cells lacking all copies of both presenilin genes [BD8 cells (Donoviel et al., 1999)] with constructs expressing either PsenA, PsenB (both with and without FLAG tags) or human PSEN1, an S2-cleaved but membrane-tethered form of human Notch-1 (Δ EN1) and the CBF1-luc reporter construct. Both *Dictyostelium* presenilin proteins showed significant activity in their ability to participate in γ -secretase cleavage of the S3 cleavage site in Δ EN1, and to induce reporter activation (Fig. 7), indicating that the *Dictyostelium* presenilin proteins are functionally active in mammalian cells, presumably as part of the γ -secretase complex. These results are in agreement with our previous demonstration of the cleavage of an N-terminally truncated human APP in *Dictyostelium* (McMains et al., 2010).

DISCUSSION

There is a pressing need for new models in presenilin research owing to the lethality of double-presenilin-null mutants in mammals (Shen et al., 1997), which makes research in the absence of background presenilin activity highly problematic. Alternative non-mammalian model systems (Calahorra and Ruiz-Rubio, 2011; Coen et al., 2012; Khandelwal et al., 2007; van Tijn et al., 2011) have been used to provide some insight into presenilin function, but studies in these models are complicated by difficulties in gene ablation or in the overexpression of tagged proteins in stable isogenic cell lines to enable cell-signalling and biochemistry studies. This study in *Dictyostelium*, in combination with our previous work (McMains et al., 2010), thus demonstrates the experimental benefits provided by this novel model for the analysis of presenilin function.

We have addressed the conservation of function of the *Dictyostelium* and human presenilins in development by analysing their ability to rescue the block mid-way through development (at the mound stage) observed in a *psenA*⁻/*psenB*⁻ mutant (Fig. 2). The ability of PSEN1 to efficiently restore morphological development strongly supports the conclusion that




Fig. 7. The *Dictyostelium* presenilins are functional in mammalian cells. The PsenA and PsenB proteins were expressed in mouse-blastocyst-derived BD8 cells that have both *psen1* and *psen2* ablated. γ -secretase activity was assessed by the fold increase in luminescence resulting from cleavage of the substrate Δ EN1 (a membrane-tethered Notch1 lacking the extracellular domain) by transfected human presenilin 1 (WT) or *Dictyostelium* presenilin proteins (PsenA and PsenB) with or without a C-terminal FLAG tag. Results show means \pm s.e.m. Data represent two to three independent experiments carried out in duplicate ($n=4-6$). * $P<0.05$, ** $P<0.01$, *** $P<0.001$.

the human and *Dictyostelium* presenilins share a conserved and evolutionarily ancient function. In addition, the ability of PsenB to rescue the development of the double-null mutant suggests that there is functional redundancy between the *Dictyostelium* presenilins, even though their starkly different expression profiles (McMains et al., 2010), with *psenB* expressed highly in growth and *psenA* expressed highly during later development, would suggest specific functions for each presenilin. The reverse is also true: APP expressed ectopically in *Dictyostelium* is cleaved efficiently by the endogenous γ -secretase complex and in a similar way as in mammalian cell lines (McMains et al., 2010), and we show that *Dictyostelium* presenilins can cleave mammalian Notch 1 when they are expressed in *Psen1*⁻/*Psen2*⁻ mouse-blastocyst-like cells (Fig. 7). Thus, it appears that human presenilin can be incorporated into the *Dictyostelium* γ -secretase complex and vice versa, and that both of these complexes are proteolytically active. Additionally, the shared cellular localisation [on the ER and nuclear envelope (Walter et al., 1996)] in both species and a shared function in regulating intracellular (cAMP) signalling (Müller et al., 2011; Wang et al., 2011) and calcium homeostasis (Berridge, 2011) further support a common function for presenilins in mammals and *Dictyostelium* (Figs 4,6). These observations together suggest that presenilins might have a conserved function throughout multiple kingdoms of life and across millennia of evolution.

Despite a conserved function of the *Dictyostelium* and human presenilin proteins in a range of cellular processes, the physiological function of the proteins in the development of the two species appears to be different. Mutating the catalytic aspartic-acid residues of either PsenB or PSEN1 did not remove the ability of these proteins to rescue the morphological defect of the *psenA*⁻/*psenB*⁻ mutant during development (Fig. 5A). Also, ablation of nicastrin, which completely blocks γ -secretase proteolytic activity in mammalian models (Edbauer et al., 2003), did not block *Dictyostelium* development, unlike the strong effect observed when both *psenA* and *psenB* are disrupted. Thus, these data showing that presenilin proteolytic activity is dispensable for development suggest an archaic structural role for presenilin proteins in this amoeba, which is independent of

proteolytic function. In support of this, a study in the moss *Physcomitrella patens* showed that both human presenilin 1 and the PSEN1^{D385A} variant were able to rescue the growth defect observed in the moss presenilin-null mutant (Khandelwal et al., 2007). However, it appears that certain growth and other developmental aspects in *Dictyostelium* require the other components of the γ -secretase complex (McMains et al., 2010). Our data thus provide evidence for a highly conserved structural role for presenilin proteins in development.

The conserved cellular role for human presenilin 1 that we have shown in *Dictyostelium* will enable a more thorough analysis of the cellular function of the wild-type protein in isogenic cell lines. Multiple gram weights of identical cells can be employed for biochemical analysis in this model, and a range of genetic approaches [including suppression screens and pharmacogenetics (Williams, 2005)] can be employed to better understand wild-type presenilin function in cells. *Dictyostelium* can also be used to analyse the >170 presenilin mutations that give rise to FAD (Parks and Curtis, 2007) in relation to basic cellular functions (such as calcium regulation) and in relation to γ -secretase activity [we can monitor the cleavage of human APP in *Dictyostelium*, giving rise to a potential change in the A β 40:A β 42 ratio associated with Alzheimer's disease progression (McMains et al., 2010)]. This new presenilin model could provide an important versatile system for future research on FAD and drug-development studies in translational research.

MATERIALS AND METHODS

Phylogenetic and structural analysis

MEGA4 was employed to align protein sequence data (Kumar et al., 2008), with further edits and the removal of highly variable regions conducted in Se-ALV2.0 (Rambaut, 1996). The phylogenetic relationships within this dataset were estimated by Bayesian analyses, using human signal-peptide-peptidase-like-2A (SPLLA2; Q8TCT8) as the outgroup, because presenilins and this protein belong to different subgroups of the peptidase A22 family (A22A and A22B, respectively). Bayesian trees were constructed in MrBayes 3.2 (Ronquist and Huelsenbeck, 2003) with gamma distribution, proportion of invariable sites and implementation of the 'aamodelpr=mixed prior' that allowed selection for the optimum substitution model. Two independent runs using four chains (three heated, one cold) were run for 1×10^6 generations, with sampling every 1×10^3 generations and a burn-in period of 250 trees. Nodal support was determined by approximate posterior probabilities performed in MrBayes. Stabilisation and convergence between runs was assessed using Tracer 1.5 (Rambaut and Drummond, 2009). Protein structural analysis was based upon the human protein structure (Parks and Curtis, 2007).

Presenilin-mutant cell lines and cell-type-differentiation analysis

Presenilin- and nicastrin-null mutants were generated by homologous integration using the Cre-Lox system (Faix et al., 2004). Wild-type *psenB* and human *psen1* (isoform 2) full-length cDNAs were ligated into pDM317 and pDM448 N-terminal-GFP expression constructs under control of the *act15* promoter (Veltman et al., 2009). Wild-type human *psen2* (isoform 1) with a *D. discoideum* codon bias was synthesised by MWG-Biotech (Germany) and ligated into the pDM448 N-terminal-GFP expression construct. Stalk-cell development was conducted as described previously (Williams et al., 1999). Spore production was assessed by the lysis of mature fruiting bodies (24 h) in 0.1% NP40 or 0.3% Triton X-100 and counting spores with a haemocytometer.

cAMP assays

Cells were grown in HL-5 medium containing selection antibiotics where appropriate. Cells were developed on non-nutrient (NN) agar (0.68 g/l KH₂PO₄, 0.89 g/l Na₂HPO₄•2H₂O pH 6.5), 15 g/l agar], and samples

were harvested in PB (0.68 g/l KH_2PO_4 , 0.89 g/l $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$ pH 6.5) at 0, 4 and 8 h of development. 5×10^7 cells were resuspended in 200 μl of PB and were lysed with 300 μl 3.5% (v/v) HClO_4 . The lysates were neutralised by the addition of an equal volume of 50% saturated KHCO_3 and 200 μl of cAMP-assay buffer (4 mM EDTA in 150 mM sodium phosphate pH 7.5). The lysates were centrifuged for 5 min at 3000 g to precipitate protein and KClO_4 . cAMP was assayed in 20 μl and 40 μl of the supernatant fraction by isotope-dilution assay, using purified protein kinase A regulatory subunit (PKA-R) from beef muscle as a cAMP-binding protein and $[2,8\text{-}^3\text{H}]\text{cAMP}$ as the competitor. The data represent the mean and s.e.m. of five experiments performed in triplicate. The data from individual measurements were not normally distributed and significant differences between datasets were therefore estimated by Kruskal–Wallis ANOVA on ranks.

Calcium assays

Calcium homeostasis in *Dictyostelium* cells was measured by using the calcium-sensitive aequorin approach in which strains were transformed with an apoaequorin-expressing plasmid (pPROF120) that allows real-time assay of cytosolic free Ca^{2+} levels (Allan and Fisher, 2009). Briefly, 10^8 cells were incubated in 5 ml of MES development buffer [10 mM 2-(*N*-morpholino)ethanesulfonic acid (MES, pH 6.2), 10 mM KCl, 0.25 mM CaCl_2] containing coelenterazine-*h* (0.5 $\mu\text{g}/\text{ml}$ dissolved in 20% w/v Pluronic F-127) for 7 h in shaking suspension, followed by washing with MES development buffer to remove residual coelenterazine-*h*. The coelenterazine-*h* allows *in vivo* reconstitution of the functional photoprotein, which upon calcium binding produces luminescence that can be detected with a photometer. In order to measure calcium influx upon cAMP stimulation, the total possible light emission was determined to normalise the aequorin luminescence signals. These values allowed calculation of an *in vitro* calcium concentration-effect curve upon 1 μM cAMP stimulation. All measurements were carried out in a New Brunswick ATP Photometer, as described previously (Allan and Fisher, 2009), and analysed using the R statistical package (R Core Team, 2012, <http://www.R-project.org/>).

Immunofluorescence and western analysis

Monoclonal antibodies against calnexin were kindly provided by Annette Müller-Taubenberger (Müller-Taubenberger et al., 2001). For immunofluorescence labelling, cell lines expressing the *D. discoideum* or human genes were grown overnight in low-fluorescence medium, fixed, probed with anti-calnexin antibodies and labelled with goat anti-mouse-IgG (NEB 4409s; Alexa Fluor 555; Ipswich, MA). Rabbit monoclonal anti-presenilin-1 D39D1 antibody (#5643) was obtained from Cell Signaling Technology (Danvers, MA) and anti-presenilin-2 (EP1515Y) antibody (ab51249) was obtained from Abcam (Cambridge, MA) and used at 1:1000 for western analysis.

Quantitative RT-PCR

Total RNA was extracted from *Dictyostelium* using the High Pure RNA Isolation kit (11828665001; Roche; Welwyn Garden City, UK) and cDNA was amplified using the First Strand cDNA Synthesis Kit (K1612; Fermentas; Loughborough, UK). Real-time amplification with SYBR Green (Sigma, S4438) was performed in a Rotor-Gene 6000 (Qiagen Ltd, Manchester, UK). Triplicate samples were collected at each time point with two RT-PCR technical replicates and the level of transcription was quantified using the $2^{-\Delta\Delta\text{CT}}$ method (Livak and Schmittgen, 2001). PCR primer pairs were designed using Primer3 (Rozen and Skaletsky, 2000).

Notch cleavage assay

An S2-cleaved form of human Notch 1 that requires γ -secretase-dependent S3 cleavage was used in a CBF1-reporter-gene assay as described previously (Hooper et al., 2006). The CBF1 reporter contains four tandem consensus DNA-binding sites in the promoter driving luciferase expression (a gift from Gerry Weinmaster, UCLA Medical School, CA). Presenilin-dependent generation of the Notch intracellular domain is required to activate CBF1-dependent transcription. Full-length

Dictyostelium presenilin was expressed from the pFLAG-CMV5a vector (Sigma-Aldrich St Louis, MO).

Acknowledgements

G.P.O. (funded by the Dr Hadwen Trust) and R.S.B.W. did not participate in experiments involving animals, or cells or tissues from animals or from human embryos. The Dr Hadwen Trust is the UK leading medical research charity that funds and promotes exclusively human-relevant research that encourages the progress of medicine with the replacement of the use of animals in research.

Competing interests

The authors declare no competing interests.

Author contributions

The project was conceived and designed by R.S.B.W., with additional experimental design input by M.H.R.L. M.H.R.L. and G.P.O. performed the majority of the experiments. C.S. and Z.-H.C. performed the cAMP experiments. C.A. and M.H.R.L. performed the calcium experiments. R.K. and M.H.R.L. performed the mammalian experiments. S.B. performed the phylogenetic analysis. P.W.B., A.R.K., P.F. and R.K. contributed reagents, materials or analysis tools and provided manuscript comments. R.S.B.W., M.H.R.L. and G.P.O. wrote the paper. G.P.O. did not participate in experiments involving animals, or cells or tissues from animals or from human embryos.

Funding

This work was supported by the Alzheimer's Society UK (R.K.); by the Intramural Research Programs of the National Institutes of Health; and the National Institute of Diabetes and Digestive and Kidney Diseases (A.R.K.); and by an Alzheimer's Research UK PhD studentship to R.S.B.W. M.H.R.L. received a Central Research Fund and Helen Shackleton travel scholarship. G.P.O. is funded by the Dr Hadwen Trust (DHT). Deposited in PMC for release after 12 months.

Supplementary material

Supplementary material available online at <http://jcs.biologists.org/lookup/suppl/doi:10.1242/jcs.140939/-DC1>

References

- Ahn, K., Shelton, C. C., Tian, Y., Zhang, X., Gilchrist, M. L., Sisodia, S. S. and Li, Y. M. (2010). Activation and intrinsic gamma-secretase activity of presenilin 1. *Proc. Natl. Acad. Sci. USA* **107**, 21435–21440.
- Allan, C. Y. and Fisher, P. R. (2009). *In vivo* measurements of cytosolic calcium in *Dictyostelium discoideum*. *Methods Mol. Biol.* **571**, 291–308.
- Berridge, M. J. (2011). Calcium signalling and Alzheimer's disease. *Neurochem. Res.* **36**, 1149–1156.
- Boeckeler, K. and Williams, R. S. B. (2007). *Dictyostelium* as a biomedical model. *Encyclopedia of Life Sciences*. **2007**, 1–6.
- Calahorra, F. and Ruiz-Rubio, M. (2011). *Caenorhabditis elegans* as an experimental tool for the study of complex neurological diseases: Parkinson's disease, Alzheimer's disease and autism spectrum disorder. *Invert. Neurosci.* **11**, 73–83.
- Chang, P., Orabi, B., Deranieh, R. M., Dham, M., Hoeller, O., Shimshoni, J. A., Yagen, B., Bialer, M., Greenberg, M. L., Walker, M. C. et al. (2012). The antiepileptic drug valproic acid and other medium-chain fatty acids acutely reduce phosphoinositide levels independently of inositol in *Dictyostelium*. *Dis. Model. Mech.* **5**, 115–124.
- Chávez-Gutiérrez, L., Tolia, A., Maes, E., Li, T., Wong, P. C. and de Strooper, B. (2008). Glu(332) in the Nicastrin ectodomain is essential for gamma-secretase complex maturation but not for its activity. *J. Biol. Chem.* **283**, 20096–20105.
- Coen, K., Flannagan, R. S., Baron, S., Carraro-Lacroix, L. R., Wang, D., Vermeire, W., Michiels, C., Munck, S., Baert, V., Sugita, S. et al. (2012). Lysosomal calcium homeostasis defects, not proton pump defects, cause endo-lysosomal dysfunction in PSEN-deficient cells. *J. Cell Biol.* **198**, 23–35.
- De Strooper, B. (2003). Aph-1, Pen-2, and Nicastrin with Presenilin generate an active gamma-Secretase complex. *Neuron* **38**, 9–12.
- De Strooper, B. and Annaert, W. (2010). Novel research horizons for presenilins and γ -secretases in cell biology and disease. *Annu. Rev. Cell Dev. Biol.* **26**, 235–260.
- De Strooper, B., Saftig, P., Craessaerts, K., Vanderstichele, H., Guhde, G., Annaert, W., Von Figura, K. and Van Leuven, F. (1998). Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature* **391**, 387–390.
- Donoviel, D. B., Hadjantonakis, A. K., Ikeda, M., Zheng, H., Hyslop, P. S. and Bernstein, A. (1999). Mice lacking both presenilin genes exhibit early embryonic patterning defects. *Genes Dev.* **13**, 2801–2810.
- Edbauer, D., Winkler, E., Regula, J. T., Pesold, B., Steiner, H. and Haass, C. (2003). Reconstitution of gamma-secretase activity. *Nat. Cell Biol.* **5**, 486–488.
- Eichinger, L., Pachebat, J. A., Glöckner, G., Rajandream, M. A., Sugang, R., Berriman, M., Song, J., Olsen, R., Szafrański, K., Xu, Q. et al. (2005). The genome of the social amoeba *Dictyostelium discoideum*. *Nature* **435**, 43–57.

- Faix, J., Kreppel, L., Shaulsky, G., Schleicher, M. and Kimmel, A. R. (2004). A rapid and efficient method to generate multiple gene disruptions in *Dictyostelium discoideum* using a single selectable marker and the Cre-loxP system. *Nucleic Acids Res.* **32**, e143.
- Feng, R., Wang, H., Wang, J., Shrom, D., Zeng, X. and Tsien, J. Z. (2004). Forebrain degeneration and ventricle enlargement caused by double knockout of Alzheimer's presenilin-1 and presenilin-2. *Proc. Natl. Acad. Sci. USA* **101**, 8162–8167.
- Fuhrer, R., Friedlein, A., Haass, C. and Walter, J. (2004). Phosphorylation of presenilin 1 at the caspase recognition site regulates its proteolytic processing and the progression of apoptosis. *J. Biol. Chem.* **279**, 1585–1593.
- Francione, L. M. and Fisher, P. R. (2011). Heteroplasmic mitochondrial disease in *Dictyostelium discoideum*. *Biochem. Pharmacol.* **82**, 1510–1520.
- Garcia, G. L. and Parent, C. A. (2008). Signal relay during chemotaxis. *J. Microsc.* **231**, 529–534.
- Haass, C. and De Strooper, B. (1999). The presenilins in Alzheimer's disease – proteolysis holds the key. *Science* **286**, 916–919.
- Harwood, A. J. (2008). *Dictyostelium* development: a prototypic Wnt pathway? *Methods Mol. Biol.* **469**, 21–32.
- Herreman, A., Serneels, L., Annaert, W., Collen, D., Schoonjans, L. and De Strooper, B. (2000). Total inactivation of gamma-secretase activity in presenilin-deficient embryonic stem cells. *Nat. Cell Biol.* **2**, 461–462.
- Hooper, C., Tavassoli, M., Chapple, J. P., Uwanogho, D., Goodyear, R., Melino, G., Lovestone, S. and Killick, R. (2006). TAp73 isoforms antagonize Notch signalling in SH-SY5Y neuroblastomas and in primary neurones. *J. Neurochem.* **99**, 989–999.
- Huppert, S. S., Le, A., Schroeter, E. H., Mumm, J. S., Saxena, M. T., Milner, L. A. and Kopan, R. (2000). Embryonic lethality in mice homozygous for a processing-deficient allele of Notch1. *Nature* **405**, 966–970.
- Janetopoulos, C. and Firtel, R. A. (2008). Directional sensing during chemotaxis. *FEBS Lett.* **582**, 2075–2085.
- Kang, D. E., Soriano, S., Frosch, M. P., Collins, T., Naruse, S., Sisodia, S. S., Leibowitz, G., Levine, F. and Koo, E. H. (1999). Presenilin 1 facilitates the constitutive turnover of beta-catenin: differential activity of Alzheimer's disease-linked PS1 mutants in the beta-catenin-signaling pathway. *J. Neurosci.* **19**, 4229–4237.
- Kang, D. E., Soriano, S., Xia, X., Eberhart, C. G., De Strooper, B., Zheng, H. and Koo, E. H. (2002). Presenilin couples the paired phosphorylation of beta-catenin independent of axin: implications for beta-catenin activation in tumorigenesis. *Cell* **110**, 751–762.
- Khandelwal, A., Chandu, D., Roe, C. M., Kopan, R. and Quatran, R. S. (2007). Moonlighting activity of presenilin in plants is independent of gamma-secretase and evolutionarily conserved. *Proc. Natl. Acad. Sci. USA* **104**, 13337–13342.
- Kim, J., Chang, A., Dudak, A., Federoff, H. J. and Lim, S. T. (2011). Characterization of nectin processing mediated by presenilin-dependent γ -secretase. *J. Neurochem.* **119**, 945–956.
- Kumar, S., Nei, M., Dudley, J. and Tamura, K. (2008). MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences. *Brief. Bioinform.* **9**, 299–306.
- Li, J., Xu, M., Zhou, H., Ma, J. and Potter, H. (1997). Alzheimer presenilins in the nuclear membrane, interphase kinetochores, and centrosomes suggest a role in chromosome segregation. *Cell* **90**, 917–927.
- Livak, K. J. and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402–408.
- Loomis, W. F. and Shaulsky, G. (2011). Developmental changes in transcriptional profiles. *Dev. Growth Differ.* **53**, 567–575.
- Ludtmann, M. H., Boeckeler, K. and Williams, R. S. (2011). Molecular pharmacology in a simple model system: implicating MAP kinase and phosphoinositide signalling in bipolar disorder. *Semin. Cell Dev. Biol.* **22**, 105–113.
- McMains, V. C., Myre, M., Kreppel, L. and Kimmel, A. R. (2010). *Dictyostelium* possesses highly diverged presenilin/gamma-secretase that regulates growth and cell-fate specification and can accurately process human APP: a system for functional studies of the presenilin/gamma-secretase complex. *Dis. Model. Mech.* **3**, 581–594.
- Müller, M., Cárdenas, C., Mei, L., Cheung, K. H. and Foskett, J. K. (2011). Constitutive cAMP response element binding protein (CREB) activation by Alzheimer's disease presenilin-driven inositol trisphosphate receptor (InsP3R) Ca²⁺ signaling. *Proc. Natl. Acad. Sci. USA* **108**, 13293–13298.
- Müller-Taubenberger, A., Lupas, A. N., Li, H., Ecke, M., Simmeth, E. and Gerisch, G. (2001). Calreticulin and calnexin in the endoplasmic reticulum are important for phagocytosis. *EMBO J.* **20**, 6772–6782.
- Myre, M. A., Lumsden, A. L., Thompson, M. N., Wasco, W., MacDonald, M. E. and Gusella, J. F. (2011). Deficiency of huntingtin has pleiotropic effects in the social amoeba *Dictyostelium discoideum*. *PLoS Genet.* **7**, e1002052.
- Parks, A. L. and Curtis, D. (2007). Presenilin diversifies its portfolio. *Trends Genet.* **23**, 140–150.
- Rambaut, A. (1996). Se-ALV2.0a11: sequence alignment editor.
- Rambaut, A. and Drummond, A. J. (2009). Tracer v1.5.
- Ronquist, F. and Huelsenbeck, J. P. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574.
- Rozen, S. and Skaletsky, H. (2000). Primer3 for general users and for biologist programmers. *Methods Mol. Biol.* **132**, 365–386.
- Saxe, C. L., III, Johnson, R., Devreotes, P. N. and Kimmel, A. R. (1991). Multiple genes for cell surface cAMP receptors in *Dictyostelium discoideum*. *Dev. Genet.* **12**, 6–13.
- Schaeffer, E. L., Figueiro, M. and Gattaz, W. F. (2011). Insights into Alzheimer disease pathogenesis from studies in transgenic animal models. *Clinics (Sao Paulo)* **66 Suppl.** **1**, 45–54.
- Shen, J., Bronson, R. T., Chen, D. F., Xia, W., Selkoe, D. J. and Tonegawa, S. (1997). Skeletal and CNS defects in Presenilin-1-deficient mice. *Cell* **89**, 629–639.
- Terbach, N., Shah, R., Kelemen, R., Klein, P. S., Gordienko, D., Brown, N. A., Wilkinson, C. J. and Williams, R. S. (2011). Identifying an uptake mechanism for the antiepileptic and bipolar disorder treatment valproic acid using the simple biomedical model *Dictyostelium*. *J. Cell Sci.* **124**, 2267–2276.
- Tesco, G. and Tanzi, R. E. (2000). GSK3 beta forms a tetrameric complex with endogenous PS1-CTF/NTF and beta-catenin. Effects of the D257/D385A and FAD-linked mutations. *Ann. N. Y. Acad. Sci.* **920**, 227–232.
- Tu, H., Nelson, O., Bezprozvanny, A., Wang, Z., Lee, S. F., Hao, Y. H., Serneels, L., De Strooper, B., Yu, G. and Bezprozvanny, I. (2006). Presenilins form ER Ca²⁺ leak channels, a function disrupted by familial Alzheimer's disease-linked mutations. *Cell* **126**, 981–993.
- Twomey, C. and McCarthy, J. V. (2006). Presenilin-1 is an unprimed glycogen synthase kinase-3beta substrate. *FEBS Lett.* **580**, 4015–4020.
- van Tijn, P., Kamphuis, W., Marlatt, M. W., Hol, E. M. and Lucassen, P. J. (2011). Presenilin mouse and zebrafish models for dementia: focus on neurogenesis. *Prog. Neurobiol.* **93**, 149–164.
- Veltman, D. M. and van Haastert, P. J. (2008). The role of cGMP and the rear of the cell in *Dictyostelium* chemotaxis and cell streaming. *J. Cell Sci.* **121**, 120–127.
- Veltman, D. M., Akar, G., Bosgraaf, L. and Van Haastert, P. J. (2009). A new set of small, extrachromosomal expression vectors for *Dictyostelium discoideum*. *Plasmid* **61**, 110–118.
- Walter, J., Capell, A., Grünberg, J., Pesold, B., Schindzielorz, A., Prior, R., Podlisny, M. B., Fraser, P., Hyslop, P. S., Selkoe, D. J. et al. (1996). The Alzheimer's disease-associated presenilins are differentially phosphorylated proteins located predominantly within the endoplasmic reticulum. *Mol. Med.* **2**, 673–691.
- Wang, D., Yuen, E. Y., Zhou, Y., Yan, Z. and Xiang, Y. K. (2011). Amyloid beta peptide-(1–42) induces internalization and degradation of beta2 adrenergic receptors in prefrontal cortical neurons. *J. Biol. Chem.* **286**, 31852–31863.
- Williams, R. S. B. (2005). Pharmacogenetics in model systems: defining a common mechanism of action for mood stabilisers. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **29**, 1029–1037.
- Williams, R. S. B., Eames, M., Ryves, W. J., Viggars, J. and Harwood, A. J. (1999). Loss of a prolyl oligopeptidase confers resistance to lithium by elevation of inositol (1,4,5) trisphosphate. *EMBO J.* **18**, 2734–2745.
- Williams, R. S., Boeckeler, K., Gräf, R., Müller-Taubenberger, A., Li, Z., Isberg, R. R., Wessels, D., Soll, D. R., Alexander, H. and Alexander, S. (2006). Towards a molecular understanding of human diseases using *Dictyostelium discoideum*. *Trends Mol. Med.* **12**, 415–424.
- Wolfe, M. S. (2006). The gamma-secretase complex: membrane-embedded proteolytic ensemble. *Biochemistry* **45**, 7931–7939.
- Yamasaki, A., Eimer, S., Okochi, M., Smialowska, A., Kaether, C., Baumeister, R., Haass, C. and Steiner, H. (2006). The GxGD motif of presenilin contributes to catalytic function and substrate identification of gamma-secretase. *J. Neurosci.* **26**, 3821–3828.