



University of Dundee

New 8-nitroquinolinone derivative displaying submicromolar in vitro activities against both *Trypanosoma brucei* and *cruzi*

Pedron, Julien; Boudot, Clotilde; Brossas, Jean-yves; Pinault, Emilie; Bourgeade-Delmas, Sandra; Sournia-Saquet, Alix

Published in:
ACS Medicinal Chemistry Letters

DOI:
[10.1021/acsmchemlett.9b00566](https://doi.org/10.1021/acsmchemlett.9b00566)

Publication date:
2020

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Pedron, J., Boudot, C., Brossas, J., Pinault, E., Bourgeade-Delmas, S., Sournia-Saquet, A., Boutet-Robinet, E., Destere, A., Tronnet, A., Bergé, J., Bonduelle, C., Deraeve, C., Pratviel, G., Stigliani, J-L., Paris, L., Mazier, D., Corvaisier, S., Since, M., Malzert-Freon, A., ... Verhaeghe, P. (2020). New 8-nitroquinolinone derivative displaying submicromolar in vitro activities against both *Trypanosoma brucei* and *cruzi*. *ACS Medicinal Chemistry Letters*, 11(4), 464-472. <https://doi.org/10.1021/acsmchemlett.9b00566>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

New 8-Nitroquinolinone Derivative Displaying Submicromolar *in Vitro* Activities against Both *Trypanosoma brucei* and *cruzi*

Julien Pedron,⁺ Clotilde Boudot,⁺ Jean-Yves Brossas, Emilie Pinault, Sandra Bourgeade-Delmas, Alix Sournia-Saquet, Elisa Boutet-Robinet, Alexandre Destere, Antoine Tronnet, Justine Bergé, Colin Bonduelle, Céline Deraeve, Geneviève Pratviel, Jean-Luc Stigliani, Luc Paris, Dominique Mazier, Sophie Corvaisier, Marc Since, Aurélie Malzert-Fréon, Susan Wyllie, Rachel Milne, Alan H. Fairlamb, Alexis Valentin, Bertrand Courtioux, and Pierre Verhaeghe*

Cite This: <https://dx.doi.org/10.1021/acsmmedchemlett.9b00566>

Read Online

ACCESS |


Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: An antikinoplastid pharmacomodulation study was conducted at position 6 of the 8-nitroquinolin-2(1H)-one pharmacophore. Fifteen new derivatives were synthesized and evaluated *in vitro* against *L. infantum*, *T. brucei brucei*, and *T. cruzi*, in parallel with a cytotoxicity assay on the human HepG2 cell line. A potent and selective 6-bromo-substituted antitrypanosomal derivative **12** was revealed, presenting EC₅₀ values of 12 and 500 nM on *T. b. brucei* trypomastigotes and *T. cruzi* amastigotes respectively, in comparison with four reference drugs (30 nM ≤ EC₅₀ ≤ 13 μM). Moreover, compound **12** was not genotoxic in the comet assay and showed high *in vitro* microsomal stability (half life >40 min) as well as favorable pharmacokinetic behavior in the mouse after oral administration. Finally, molecule **12** (E° = -0.37 V/NHE) was shown to be bioactivated by type 1 nitroreductases, in both *Leishmania* and *Trypanosoma*, and appears to be a good candidate to search for novel antitrypanosomal lead compounds.

KEYWORDS: *Trypanosoma brucei brucei*, *Trypanosoma cruzi*, 8-nitroquinolin-2(1H)-ones, redox potentials, NTR1



Human African trypanosomiasis (HAT),¹ Chagas diseases (CD),² and visceral leishmaniasis (VL)³ are infectious diseases caused by unicellular flagellated kinetoplastid parasites belonging to the *Trypanosoma* and *Leishmania* genera. From an epidemiological point of view, the World Health Organization (WHO) estimates that millions of people are at risk of contracting HAT, CD, and VL. These three diseases are also estimated to be responsible for about 30 000 annual deaths, with the caveat that these numbers may be underestimated due to the difficulty in accessing some rural areas and the unspecific symptoms during the early stage of these infections.⁴ Although these diseases are lethal if untreated, there are very few efficient and safe drugs available, each of them presenting various drawbacks for the patient (toxicity, emergence of resistances, mode of administration, treatment cost). Thus because of the lack of consideration and investments from the pharmaceutical industry, the WHO classified HAT, CD, and VL among “neglected tropical diseases” (NTDs).⁵

In this context, nitroaromatic derivatives play a key role in the fight against kinetoplastids (Figure 1). Beyond Nifurtimox and Benznidazole (antitrypanosomal drugs), nitroaromatics are fully part of the research efforts of the Drugs for Neglected Diseases initiative (DNDi). Thus among the few novel chemical entities in clinical trials against these infections,⁶ the development of fexinidazole (an orally available 5-

nitroimidazole derivative that received an agreement from the European Medicines Agency (EMA) in the treatment of HAT in 2018 and that is in phase 2 against CD),^{7–9} delamanid (a derivative that is marketed against tuberculosis and that was studied against VL),¹⁰ and DNDi-0690 (a 2-nitroimidazooxazine derivating from delamanid and that has recently entered phase-I clinical trials against VL) illustrate well the strong potential of nitroaromatic molecules in the search for novel antitrypanosomatids.¹¹

The antiparasitic mechanism of action of these molecules involves parasitic nitroreductases (NTRs). NTRs contain a flavin as a cofactor and catalyze the one- or two-electron reduction of nitrodrugs into cytotoxic electrophilic metabolites such as nitroso and hydroxylamine derivatives, which are able to form covalent adducts with DNA or proteins.¹² Two NTRs were found in *Leishmania*: a mitochondrial NTR1 catalyzing a two-electron reduction¹³ and a cytosolic NTR2 catalyzing a one-electron reduction.¹⁴ In *Trypanosoma*, only a mitochondrial type 1 NTR is expressed.^{15,16} It is important to note that

Received: December 2, 2019

Accepted: February 6, 2020

Published: February 6, 2020




Figure 1. Structures of nitroaromatic drugs displaying antikinoplastid activity.





Figure 2. Antitrypanosomatid profile of previously reported 8-nitroquinolin-2(1*H*)-one hit compounds.

Scheme 1. Synthesis of Compounds 1–8 from *p*-Trifluoromethylaniline



these enzymes are absent from mammalian cells, an important feature for developing selective antikinoplastid molecules. Moreover, depending on the chemical structure of nitrated derivatives, the selectivity for NTRs could be modulated. For instance, fexinidazole is selectively bioactivated by NTR1 in both *Leishmania* and *Trypanosoma*; meanwhile, delamanid is only bioactivated by NTR2 in *Leishmania*.¹⁴

In this context, by studying the antiparasitic potential of the 8-nitroquinoline scaffold,¹⁷ our research team reported a new antileishmanial pharmacophore: 8-nitroquinolin-2(1*H*)-one.¹⁸ After pharmacomodulation studies at position 4,^{19,20} an electrochemistry-guided work highlighted the importance of an intramolecular hydrogen bond between the nitro group and the lactam function.²¹ This work also led to the identification of a new 3-bromosubstituted antikinoplastid hit compound derivative (**Hit 1**, **Figure 2**), which is bioactivated by type I

NTRs, in both *L. donovani* and *T. brucei brucei* (*T. b. brucei*).²¹ Recently, by introducing a *p*-carboxyphenyl functionality at position 3 of the pharmacophore, we reported a new selective antitrypanosomal hit compound (**Hit 2**, **Figure 2**).²²

We herein present a medicinal chemistry work that greatly improved the antitrypanosomal activity of the current series via the synthesis of novel derivatives bearing electron-withdrawing groups at position 6 of the quinolinone scaffold.

To increase the oxidant character of the series and facilitate its bioactivation by nitroreductases, 15 derivatives (**1–15**) were prepared, bearing various electron-withdrawing groups (EWGs) at position 6. As presented in **Scheme 1**, eight compounds bearing a trifluoromethyl group at position 6 of the quinolinone scaffold were synthesized according to a previously reported protocol.²³ In brief, 3-ethoxyacryloyl chloride was synthesized in two steps by the saponification

Scheme 2. Synthesis of Compounds 9–15





Figure 3. Redox potentials of synthesized nitroaromatic compounds determined by cyclic voltammetry.

of commercial 3,3'-diethoxyethylpropionate followed by a reaction with thionyl chloride. This acyl chloride was then reacted with *p*-trifluoromethylaniline to afford, after a cyclization step in H_2SO_4 (98%), compound **1** in 42% yield (a). Molecule **1** was then either nitrated at position 8, using classical conditions, to afford compound **2** in 89% yield (b) or selectively halogenated at position 3 to generate derivative **3** or **4** in 48 and 54% yield, respectively (c and d). The *O*-methylation of compound **2** was performed with a protocol using a mixture of methyl iodide and sodium hydride to prepare derivative **5** in 69% yield (e). Nitrated molecule **2** could also undergo chlorination at position 3, affording compound **6** in 88% yield (f), or react with NaBrO_3 in refluxing HBr , according to a previously reported protocol,²⁴ to afford halogenated derivative **7** in 94% yield (g). The nitro group of compound **7** was finally reduced to afford the amino-derivative **8**, considered as a negative control, using SnCl_2 in 68% yield (h).

The synthesis pathway used for preparing derivatives bearing a halogen atom at position 6 of the quinolinone ring is presented in Scheme 2. It starts from 8-nitroquinolin-2(1*H*)-

one,²¹ which was reacted with bromine in refluxing acetic acid, saturated with sodium acetate, leading to a mixture of both 6-bromo and 3,6-dibromo derivatives **9** and **10** in 9 and 33% yield, respectively (a). Dichloro analogue **11** was also prepared from 8-nitroquinolin-2(1*H*)-one by reacting with 5 equiv of sodium chlorate in refluxing hydrochloric acid for 24 h (b). By using the same reaction, decreasing the amount of sodium chlorate to 3 equiv, and limiting the reaction time to 45 min, 3-chloro-8-nitroquinolin-2(1*H*)-one was formed (c).²¹ This compound could be brominated at position 6 by reacting with an excess of NaBrO_3 in refluxing HBr , leading to **12** in good yield (d). Taking into account that we previously reported the benefit of introducing a *p*-carboxyphenyl moiety at position 3 of the scaffold toward antitrypanosomal activity,²² we finally engaged dibromo-derivative **10** into a Suzuki–Miyaura reaction with *p*-(methoxycarbonyl)phenylboronic acid and then saponified the resulting ester coupling product to afford carboxylic acid **13** (e). Finally, from brominated compound **12**, derivatives **14** and **15** were obtained through a Sonogashira coupling reaction (f).

Table 1. Antikinetoplastid Activities and Cytotoxicity of Compounds 1–15




















Cpd	Structure	<i>L. infantum</i> axenic amastigote EC ₅₀ (μM)	<i>T. brucei brucei</i> trypomastigotes EC ₅₀ (μM)	Cytotoxicity HepG2 CC ₅₀ (μM) & Selectivity index regarding <i>T. b. brucei</i>
Phar.		15.5 +/-0.5	23.4 +/-5.7	164 +/-28 7
Hit 1		7.1 +/-1.5	1.9 +/-0.4	92 +/-13 48
A ^[27]		16 +/-0.9	1.2 +/-0.2	85 +/-4 71
Hit 2		> 100 ^a	1.5 +/- 0.2	120+/-7 80
1		> 100 ^a	> 50 ^a	> 100 ^a ND
2		18.5 +/- 3.9	1.7 +/- 0.9	41.3 +/- 3.9 24
3		> 100 ^a	> 50 ^a	> 100 ^a ND
4		> 100 ^a	> 50 ^a	> 100 ^a ND
5		> 100 ^a	20.9 +/- 14.5	> 100 ^a >5
6		9.2 +/- 1.5	0.57 +/- 0.25	31.4 +/- 3.3 55
7		3.9 +/- 1.5	0.2 +/- 0.09	41.3 +/- 3.9 206
8		> 100 ^a	8.7 +/- 3.2	> 100 ^a >11
9		11.6 +/- 2.5	0.07 +/- 0.01	27.9 +/- 6.0 399
10		6.5 +/- 1.8	0.05 +/- 0.008	16.6 +/- 1.2 332
11		6.2 +/- 0.6	0.1 +/- 0.04	24.3 +/- 0.9 243
12		6.7 +/- 0.8	0.012 +/- 0.002	18.1 +/- 2.5 1508
13		8.0 +/- 1.5	0.2 +/- 0.02	60.4 +/- 14.4 302
14		8.9 +/- 0.6	1.0 +/- 0.09	30.9 +/- 2.1 31

Table 1. continued

Cpd	Structure	<i>L. infantum</i> axenic amastigote EC ₅₀ (μM)	<i>T. brucei brucei</i> trypomastigotes EC ₅₀ (μM)	Cytotoxicity HepG2 CC ₅₀ (μM) & Selectivity index regarding <i>T. b. brucei</i>
15		10.1 +/- 1.0	0.85 +/- 0.1	32.7 +/- 0.4 38
	Doxorubicin ^b	-	-	0.2 +/- 0.02
	Amphotericin B ^c	0.06 +/- 0.001	-	5.5 +/- 0.25
	Miltefosine ^c	0.8 +/- 0.2	-	85 +/- 8.8
	Fexinidazole ^{c,d,e}	3.4 +/- 0.8	0.4 +/- 0.2	> 200 ^a > 500
	Suramin ^d	-	0.03 +/- 0.009	> 100 ^a > 3333
	Eflornithine ^d	-	13.3 +/- 2.1	> 100 ^a > 8

^aEC₅₀ or CC₅₀ value was not reached at the highest tested concentration. ^bDoxorubicin was used as a cytotoxic reference drug. ^cAmphotericin B, miltefosine, and fexinidazole were used as antileishmanial reference drugs. ^dFexinidazole, suramin, and eflornithine were used as anti-*Trypanosoma brucei* reference drugs.

Electrochemistry. Because the compounds were expected to be nitroreductase substrates, we studied the influence of the introduction of EWGs at position 6 of the pharmacophore toward redox potentials. Thus redox potentials (couple R-NO₂/RNO₂^{•-}) of the synthesized compounds were measured by cyclic voltammetry in DMSO (one-electron reversible reduction). Redox potentials ranged between -0.36 and -0.75 V/NHE (Figure 3). In comparison with the original pharmacophore, the introduction of a bromine atom at position 6 (compound 9) led to a significant increase in the E° value from -0.54 to -0.40 V. The same effect was observed when a bromine atom was introduced at position 6 of hit 2 (compound 13), with an increase in the E° value of +0.16 V. It is important to note that introducing a bromine atom has a higher influence on the redox potential at position 6 (compound 9) compared with position 3 (hit 1): +0.14 versus +0.09 V. The influence of the CF₃ group toward the redox potential of the scaffold was the same as that of the bromine atom: +0.15 V (comparison between the pharmacophore and compound 2). Then, when two bromine atoms were combined at positions 3 and 6 (compound 10), a slight increase of +0.03 V was observed (no additive effect) in comparison with 6-bromo-derivative 9. Surprisingly, by comparing compounds 7 and 10–12, there was almost no effect of the variation of the EWG at position 3 or 6 toward the redox potential values, with all compounds ranging from -0.36 to -0.37 V. Finally, by comparing compound 2 with its O-methylated derivative 5, it appeared that the redox potential value was drastically lower (-0.36 V shift). This shift is not only due to the donating mesomeric effect of the methoxy group but also due to the disappearance of the intramolecular hydrogen bond between the nitro group and the lactam function.²¹




Compound Evaluation. The cytotoxicity of all compounds was first assessed on the human HepG2 cell line using doxorubicin as a positive control (Table 1). Compared with this reference drug (CC₅₀ = 0.2 μM), the series showed little (CC₅₀ = 17 μM) to no (CC₅₀ > 100 μM) cytotoxicity. Non-nitrated molecules 1, 3, and 4, O-methylated derivative 5, and 8-amino-derivative 8, considered as negative controls, displayed high CC₅₀ values, >100 μM. The results show that the

introduction of a bromine atom or a CF₃ group at position 6 of the nitrated scaffold leads to derivatives (2 and 9) that are four to five times more cytotoxic (CC₅₀ = 28–41 μM) than the 8-nitroquinolinone pharmacophore (CC₅₀ = 164 μM). When introducing an additional halogen atom at position 3 of the scaffold (6, 7, 10–12), the cytotoxicity remains close to the ones of 6-monosubstituted derivatives (CC₅₀ = 17–41 μM). Nevertheless, introducing a *p*-carboxyphenyl functionality at position 3 of the scaffold (compound 13) results in a three-fold decrease in the cytotoxicity (CC₅₀ = 60 μM) in comparison with the 3,6-dibromo derivative 10 (CC₅₀ = 17 μM).

All synthesized molecules were screened *in vitro* for their activity against *Leishmania infantum* axenic amastigotes, and their EC₅₀ values were compared with the ones of three reference drugs (amphotericin B, miltefosine, and fexinidazole) and antikinoplastid hit 1 (Table 1). Among the tested compounds, only compound 7 (E° = -0.36 V) showed a weak antileishmanial activity, displaying an EC₅₀ value of 3.9 μM in addition to a selectivity index superior to 10. Derivatives of 7 without a nitro group at position 8 of the scaffold (1, 3, 4, and 8) were inactive toward *L. infantum* as well as the O-methylated analogue 5 (E° = -0.75 V), confirming both the key role of the nitro group toward antiparasitic activity and the importance of the interaction between the nitro group and the lactam function to confer suitable redox potentials for displaying antileishmanial properties in this chemical series.²¹ Note that, contrary to hit 2, which was not active against *L. infantum* (EC₅₀ > 100 μM), the 6-bromo derivative 13 was active (EC₅₀ = 8 μM).

All molecules were also screened *in vitro* for their antitrypanosomal activity against *T. b. brucei* trypomastigotes, and their EC₅₀ values were compared with the ones of three reference drugs, suramin, eflornithine, and fexinidazole, and with antitrypanosomal hit 2. Among the 15 tested molecules, 7, 9, 10, 11, 12, and 13 displayed both submicromolar activity (12 ≤ EC₅₀ ≤ 200 nM) and good selectivity indices (SI > 200) in comparison with suramin (EC₅₀ = 30 nM), eflornithine (EC₅₀ = 13 μM), fexinidazole (EC₅₀ = 400 nM), and hit 2 (EC₅₀ = 1.5 μM). Molecule 12 was the most active and selective in the series: EC₅₀ = 12 nM and SI = 1508. Compounds without a nitro group at position 8 of the scaffold

Table 2. *In Vitro* Pharmacokinetic, Physicochemical, and Toxicological Properties of Compounds 7, 12, and 13

			
Microsomal stability (min)	> 40	> 40	> 40
Albumin binding assay (%)	99.4	99.0	> 99
BBB PAMPA assay (nm/s)	450.3 ± 170.5	405.1 ± 69.3	12.7 ± 0.9
cLogD ^a at pH = 7.4	3.24	2.97	0.44
Thermodynamic solubility at pH = 7.4 (μM)	-	21.7 +/-4.1	-
Comet assay at 1 and 10 μM	-	Negative	-

^aCalculated with MarvinSketch (ChemAxon).

appeared to be inactive (compounds 1, 3, and 4), and amino-derivative 8 was quite less active ($EC_{50} = 9 \mu\text{M}$). It is important to note that the introduction of a bromine atom at position 6 of the scaffold seems crucial to afford highly active antitrypanosomal molecules, with compound 9 being 334 times more active than the 8-nitroquinolin-2(1*H*)-one pharmacophore and 27 times more active than its 3-bromoposition isomer (**hit 1**). This key effect of the bromine atom at position 6 of the scaffold also appears when comparing the activities of 3-chloro-6-bromo hit-compound 12 with its 3-chloro-analogue **A**: 12 is 100 times more active than **A**. When comparing molecules 2 and 9, it appeared that the CF_3 group does not provide the same level of improvement in activity. Introducing an additional halogen atom at position 3 of hit molecule 9, leading to dibromo-derivative 10 and dihalo-derivative 12, afforded the most active compounds in the series ($EC_{50} = 12\text{--}50 \text{ nM}$). A comparison between compound 12 and 3,6-dichlorosubstituted derivative 11 showed that the replacement of the bromine atom by a chlorine atom at position 6 was not in favor of antitrypanosomal activity. During pharmacomodulation studies at position 3 of the scaffold, **hit 2** was identified as a promising antitrypanosomal hit with an EC_{50} of $1.5 \mu\text{M}$.²² When a bromine atom is introduced at position 6 of **hit 2**, leading to compound 13, the antitrypanosomal activity is improved, reaching an EC_{50} value of 200 nM.

Thus the molecule presenting the best antitrypanosomal profile in the series, regarding both the activity and the selectivity, was compound 12. To better evaluate its antitrypanosomal potential, an *in vitro* evaluation against intracellular *T. cruzi* amastigotes was carried out using benzimidazole and fexinidazole as positive controls (Table 1). Quite interestingly, despite being about 10 times more cytotoxic than reference drugs, molecule 12 displayed a high activity ($EC_{50} = 0.5 \mu\text{M}$) against *T. cruzi*, identical to the one of benzimidazole ($EC_{50} = 0.5 \mu\text{M}$) and better than the one of fexinidazole ($EC_{50} = 3 \mu\text{M}$).

To investigate more deeply the potential of compounds 7, 12, and 13 as the most active antikinoplastid compounds of the studied series, an *in vitro* pharmacokinetic evaluation was performed, including a microsomal stability assay (female mouse microsomes), a human albumin binding assay, and a PAMPA assay using a blood–brain barrier (BBB) model (Table 2). All compounds presented good microsomal stability ($T_{1/2} > 40 \text{ min}$) and strong binding to human albumin ($\geq 99\%$). However, the development of antitrypanosomal agents requires that molecules cross the BBB, as the second stage of HAT is meningoencephalitic. A PAMPA assay on a BBB model was then performed.^{25,26} Compounds 7 and 12 diffused through the lipid bilayer with Pe values significantly higher than the positive control (Pe = 450.3 and 405.1 nm/s, respectively), whereas compound 13 displayed a low Pe value (Pe = 12.7 nm/s), indicating a limited crossing of the BBB, hindering the further development of 13. This last result is not surprising because electrostatic repulsions between the negatively charged carboxylic acid group of this molecule and the negatively charged membrane constituents are probably responsible for its low permeability across the BBB in the PAMPA model. Moreover, 13 has a molecular weight approaching 400 g/mol, which could also contribute to limiting its diffusion across the BBB.²⁷ Unfortunately, the methyl ester precursor or 13 could not be evaluated *in vitro* because of a lack of aqueous solubility. The ability of hit-compound 12 to cross the BBB in the PAMPA assay is in accordance with its central nervous system multiparameter optimization (CNS MPO) score²⁸ that has a calculated value of 4.35. Finally, the aqueous solubility of hit-compound 12 was measured with a thermodynamic solubility assay at physiological pH and was determined at $21.7 \mu\text{M}$ (Table 2).

To check the antikinoplastid mechanism of the bio-activation of compounds 7, 12 and 13, their activities were evaluated *in vitro* toward *Leishmania donovani* promastigote strains overexpressing leishmanial nitroreductases (NTR1 or NTR2). EC_{50} values were determined and compared with the ones obtained on a wild-type strain to evaluate the

bioactivation of these nitroaromatic compounds (Table 3). As reported for hit 1,²¹ compounds 7 and 12 are selectively

Table 3. Evaluation of the Bioactivation of Compounds 7, 12, and 13 by Leishmanial and Trypanosomal NTRs

<i>Leishmania donovani</i> promastigotes EC ₅₀ (μM)			
compound	wild type	NTR1 overexpressing	NTR2 overexpressing
Hit 1	5.9 ± 0.2	0.47 ± 0.02	4.6 ± 0.12
7	4.1 ± 0.2	1.2 ± 0.05	4.9 ± 0.3
12	4.7 ± 0.09	0.3 ± 0.01	2.9 ± 0.08
13	2.0 ± 0.2	1.3 ± 0.1	2.1 ± 0.1
<i>Trypanosoma brucei brucei</i> trypomastigotes EC ₅₀ (μM)			
compound	wild type	NTR overexpressing	
Hit 1	17.7 ± 1.0	3.9 ± 0.1	
Hit 2	5.4 ± 0.12	4.2 ± 0.2	
7	6.7 ± 0.4	0.9 ± 0.03	
12	0.3 ± 0.06	0.02 ± 0.008	
13	4.0 ± 0.2	0.1 ± 0.009	
Nifurtimox	1.9 ± 0.05	0.6 ± 0.05	

bioactivated by leishmanial NTR1, with EC₅₀ values being 3.5 and 16 times lower on the strain overexpressing the NTR than on the wild type. Contrary to 7 and 12, compound 13 did not demonstrate significant bioactivation by NTR1 or by NTR2 in *Leishmania*. The same experiments were carried out with compounds 7, 12, and 13 on *T. b. brucei* strains overexpressing the type 1 trypanosomal NTR, comparing the EC₅₀ values with the ones determined on the wild-type strain (Table 3). All tested compounds were bioactivated by the type 1 nitroreductase of *T. b. brucei*, being, respectively, 7, 15, and 40 times more active against the strain overexpressing the NTR than against the wild type. It is interesting to note that hit 2 ($E^\circ = -0.56$ V) did not seem to be bioactivated by *T. b. brucei* NTR, whereas its 6-brominated derivative 13 ($E^\circ = -0.40$ V) was. Finally, hit-compound 12 also appeared to be six times more active than nifurtimox against *T. b. brucei*, underlining the potential of this molecule as an antitrypanosomal candidate.

Even though many safe nitroaromatic molecules are today on the market (for example, antiprotozoal 5-nitroimidazoles such as metronidazole, several antihypertensive dihydropyridines like nicardipine, some benzodiazepines like loprazolam, anti-Parkinson's catechol *O*-methyltransferase (COMT) inhibitors like entacapone, antiandrogens like flutamide, anti-inflammatory nimesulide, immunomodulating azathioprine, anticoagulant acenocoumarol, anthelmintic niclosamide, etc.) or in clinical trials (for instance, delamanid), drugs including a nitroaromatic moiety suffer from a bad reputation as possible mutagenic or genotoxic molecules, which narrows their pharmaceutical development.²⁹ The Ames test is the most commonly used assay to evaluate the mutagenic potential of compounds, using *Salmonella typhimurium* strains. However, these bacteria possess NTR, making the Ames test generally unsuitable for the evaluation of most of nitroaromatic compounds, with the genotoxicity being attributed to the reduced metabolites that are bioactivated by these bacterial enzymes (absent in human cells).³⁰ Thus it is accepted that the comet assay or the micronucleus assay are more pertinent methods to evaluate the genotoxic potential of nitroaromatic compounds. For example, fexinidazole is positive in the Ames test but negative in the micronucleus assay.³¹ A comet assay was then performed on the HepG2 cell line with hit compound

12, using methylmethanesulfonate (MMS) as a positive control (see the Supporting Information). Compound 12 was not genotoxic after 2 or 72 h of incubation or at 1 or 10 μM, concentrations chosen to be lower than its CC₅₀ on HepG2 cell line (CC₅₀ = 18 μM).

Regarding tolerance in the mouse, a once daily oral dose of compound 12, at 25 or 50 mg/kg, was administrated to eight mice (four mice in each group). These two doses were well tolerated with just a little apathy during 1 h for all mice receiving the dose of 50 mg/kg. The administration of a repeated dose of 25 mg/kg once daily for 5 days was well tolerated without any adverse effect noted. The no-observed-adverse-effect level (NOAEL)³¹ in mice was then set at 25 mg/kg/day. After euthanasia, no lesions were found on the different organs (kidney, liver, brain, heart, and lung).

Finally, the *in vivo* pharmacokinetic parameters were determined in the mouse, after oral administration at 25 mg/kg in a mixture of 5% Tween 80 and 95% of 0.5% carboxymethylcellulose in water, by using the QuEChERS method.³² Monolix Lixoft software was used to analyze the data by a noncompartmental model to fit pharmacokinetics parameters.³³ As presented in the Supporting Information (Figure S17), hit-compound 12 showed good pharmacokinetic properties: 12 is highly and rapidly absorbed after oral administration and shows good exposure. Its plasmatic half life (2.9 h) is higher than the one of the drug fexinidazole (0.8 h).³¹

Conclusions. Via an antitrypanosomal pharmacomodulation study of the 8-nitroquinolin-2(1*H*)-one scaffold, by introducing an EWG at positions 3 and 6 of the quinolinone ring, a novel potent antitrypanosomal hit-compound (12) was identified, displaying high activities toward *T. b. brucei* (EC₅₀ = 12 nM) and *T. cruzi* (EC₅₀ = 0.5 μM) in comparison with eflornithine (EC₅₀ = 13 μM), suramin (EC₅₀ = 30 nM), benznidazole (EC₅₀ = 0.5 μM), and fexinidazole (EC₅₀ = 0.4 and 3.0 μM). Compound 12 displays good *in vitro* pharmacokinetic parameters (good microsomal stability and BBB permeability), is rapidly absorbed after oral administration, and is well tolerated in the mouse at 25 mg/kg. Nitroaromatic hit compound 12 is selectively bioactivated by type 1 NTR in both *Leishmania* and *Trypanosoma*. An electrochemistry study showed the strong influence of the introduction of an EWG in the benzenic part of the pharmacophore toward the redox potentials and the corresponding antitrypanosomal activities. Finally, the lack of genotoxicity of compound 12 in the comet assay is an additional favorable element so as to consider further development. Compound 12 meets all criteria for an antitrypanosomal hit, as defined by Katsuno and coworkers.³⁴ *In vivo* studies on the mice model of trypanosomiasis are now planned to investigate the potential of this compound to become an antitrypanosomal lead.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsmmedchemlett.9b00566>.

Materials and methods, additional analysis of hit compounds 7, 12, and 13 as well as electrochemistry data, *in vitro* pharmacokinetics (PK) data, and *in vivo* PK data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Pierre Verhaeghe – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France; orcid.org/0000-0003-0238-2447; Phone: +33 561 333 236; Email: pierre.verhaeghe@lcc-toulouse.fr

Authors

Julien Pedron – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France

Clotilde Boudot – Université de Limoges, UMR INSERM 1094, Faculté de Pharmacie, 87025 Limoges, France

Jean-Yves Brossas – AP-HP, Groupe Hospitalier Pitié-Salpêtrière, Service de Parasitologie Mycologie, 75013 Paris, France

Emilie Pinault – Université de Limoges, BISCEM Mass Spectrometry Platform, CBRS, F-87025 Limoges, France; orcid.org/0000-0001-8811-2661

Sandra Bourgeade-Delmas – UMR 152 PharmaDev, Université de Toulouse, IRD, UPS, 31077 Toulouse, France

Alix Sournia-Saquet – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France

Elisa Boutet-Robinet – Toxalim, Université de Toulouse, INRA, ENVT, INP-Purpan, UPS, 31077 Toulouse, France; orcid.org/0000-0002-5223-4568

Alexandre Destere – Department of Pharmacology, Toxicology and Pharmacovigilance, CHU Limoges, France, INSERM, UMR 1248, University of Limoges, F-87025 Limoges, France

Antoine Tronnet – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France

Justine Bergé – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France

Colin Bonduelle – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France; orcid.org/0000-0002-7213-7861

Céline Deraeve – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France

Geneviève Pratiel – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France

Jean-Luc Stigliani – LCC-CNRS, Université de Toulouse, CNRS, UPS, 31077 Toulouse, France

Luc Paris – AP-HP, Groupe Hospitalier Pitié-Salpêtrière, Service de Parasitologie Mycologie, 75013 Paris, France

Dominique Mazier – CIMI-Paris, Sorbonne Université, 75013 Paris, France

Sophie Corvaisier – Centre d'Etudes et de Recherche sur le Médicament de Normandie (CERMN), Normandie Université, 14032 Caen, France

Marc Since – Centre d'Etudes et de Recherche sur le Médicament de Normandie (CERMN), Normandie Université, 14032 Caen, France

Auréli Malzert-Fréon – Centre d'Etudes et de Recherche sur le Médicament de Normandie (CERMN), Normandie Université, 14032 Caen, France; orcid.org/0000-0001-6023-1861

Susan Wyllie – University of Dundee, School of Life Sciences, Division of Biological Chemistry and Drug Discovery, Dundee DD1 5EH, United Kingdom; orcid.org/0000-0001-8810-5605

Rachel Milne – University of Dundee, School of Life Sciences, Division of Biological Chemistry and Drug Discovery, Dundee DD1 5EH, United Kingdom

Alan H. Fairlamb – University of Dundee, School of Life Sciences, Division of Biological Chemistry and Drug Discovery,

Dundee DD1 5EH, United Kingdom; orcid.org/0000-0001-5134-0329

Alexis Valentin – UMR 152 PharmaDev, Université de Toulouse, IRD, UPS, 31077 Toulouse, France

Bertrand Courtioux – Université de Limoges, UMR INSERM 1094, Faculté de Pharmacie, 87025 Limoges, France

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsmchemlett.9b00566>

Author Contributions

*J.P. and C.B. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Paul Sabatier University and the Région Occitanie are acknowledged for financial support. A.H.F., S.W., and R.M. are supported by funding from the Wellcome Trust (WT105021). C. Piveteau from Institut Pasteur de Lille, C. Bijani from Laboratoire de Chimie de Coordination de Toulouse, and C. Claparols, N. Martins-Froment, V. Bourdon, and E. Leroy from the Institut de Chimie de Toulouse are also acknowledged for their support. We also thank Dr. J.-B. Woillard for the *in vivo* pharmacokinetic analysis, Mr. F.-L. Sauvage for his help with the mass spectrometry analysis, and Dr. Ismail Belfquih for his assistance with the culture of Vero cells and *T. cruzi*.

■ REFERENCES

- (1) Büscher, P.; Cecchi, G.; Jamonneau, V.; Priotto, G. Human african trypanosomiasis. *Lancet* **2017**, *390*, 2397–2405.
- (2) Pérez-Molina, J. A.; Molina, I. Chagas disease. *Lancet* **2018**, *391*, 82–94.
- (3) Burza, S.; Croft, S. L.; Boelaert, M. Leishmaniasis. *Lancet* **2018**, *392*, 951–970.
- (4) World Health Organization. *Neglected Tropical Diseases*. https://www.who.int/neglected_diseases/diseases/en/ (accessed November 1, 2019).
- (5) Molyneux, D.; Savioli, L.; Engels, D. Neglected tropical diseases: progress towards addressing the chronic pandemic. *Lancet* **2017**, *389*, 312–325.
- (6) Drugs for Neglected Diseases Initiative (DNDi). *DNDi R&D Portfolio June 2019*. <https://www.dndi.org/diseases-projects/portfolio/> (accessed November 1, 2019).
- (7) Wyllie, S.; Patterson, S.; Stojanovski, L.; Simeons, F. R. C.; Norval, S.; Kime, R.; Read, K. D.; Fairlamb, A. H. The anti-trypanosome drug fexinidazole shows potential for treating visceral leishmaniasis. *Sci. Transl. Med.* **2012**, *4*, 119re1.
- (8) Patterson, S.; Fairlamb, A. H. Current and future prospects of nitro-compounds as drugs for trypanosomiasis and leishmaniasis. *Curr. Med. Chem.* **2019**, *26*, 4454–4475.
- (9) Mesu, V. K. B. K.; Kalonji, W. M.; Bardonneau, C.; Mordt, O. V.; Blesson, S.; Simon, F.; Delhomme, S.; Bernhard, S.; Kuziena, W.; Lubaki, J. F.; Vuvu, S. L.; Ngima, P. N.; Mbembo, H. M.; Ilunga, M.; Bonama, A. K.; Heradi, J. A.; Solomo, J. L. L.; Mandula, G.; Badibabi, L. K.; Dama, F. R.; Lukula, P. K.; Tete, D. N.; Lumbala, C.; Scherrer, B.; Strub-Wourgaft, N.; Tarral, A. Oral fexinidazole for late-stage African Trypanosoma brucei gambiense trypanosomiasis: a pivotal multicentre, randomised, non-inferiority trial. *Lancet* **2018**, *391*, 144–154.
- (10) Patterson, S.; Wyllie, S.; Norval, S.; Stojanovski, L.; Simeons, F. R. C.; Auer, J. L.; Osuna-Cabello; Read, K. D.; Fairlamb, A. H. The anti-tubercular drug delamanid as a potential oral treatment for visceral leishmaniasis. *eLife* **2016**, *5*, No. e09744.

- (11) Ang, C. W.; Jarrad, A. M.; Cooper, M. A.; Blaskovich, M. A. T. Nitroimidazoles: Molecular Fireworks That Combat a Broad Spectrum of Infectious Diseases. *J. Med. Chem.* **2017**, *60*, 7636–7657.
- (12) Patterson, S.; Wyllie, S. Nitro drugs for the treatment of trypanosomatid diseases: past, present, and future prospects. *Trends Parasitol.* **2014**, *30*, 289–298.
- (13) Wyllie, S.; Patterson, S.; Fairlamb, A. H. Assessing the essentiality of leishmania donovani nitroreductase and its role in nitro drug activation. *Antimicrob. Agents Chemother.* **2013**, *57*, 901–906.
- (14) Wyllie, S.; Roberts, A. J.; Norval, S.; Patterson, S.; Foth, B. J.; Berriman, M.; Read, K. D.; Fairlamb, A. H. Activation of bicyclic nitro-drugs by a novel nitroreductase (NTR2) in *Leishmania*. *PLoS Pathog.* **2016**, *12*, No. e1005971.
- (15) Hall, B. S.; Bot, C.; Wilkinson, S. R. Nifurtimox activation by trypanosomal type I nitroreductases generates cytotoxic nitrile metabolites. *J. Biol. Chem.* **2011**, *286*, 13088–13095.
- (16) Hall, B. S.; Wilkinson, S. R. Activation of Benznidazole by trypanosomal type I nitroreductases results in glyoxal formation. *Antimicrob. Agents Chemother.* **2012**, *56*, 115–123.
- (17) Verhaeghe, P.; Rathelot, P.; Rault, S.; Vanelle, P. Convenient preparation of original vinylic chlorides with antiparasitic potential in quinolone series. *Lett. Org. Chem.* **2006**, *3*, 891–897.
- (18) Paloque, L.; Verhaeghe, P.; Casanova, M.; Castera-Ducros, C.; Dumètre, A.; Mbatchi, L.; Hutter, S.; Kraiem-M'Rabet, M.; Laget, M.; Remusat, V.; Rault, S.; Rathelot, P.; Azas, N.; Vanelle, P. Discovery of a new antileishmanial hit in 8-nitroquinoline series. *Eur. J. Med. Chem.* **2012**, *54*, 75–86.
- (19) Kieffer, C.; Cohen, A.; Verhaeghe, P.; Hutter, S.; Castera-Ducros, C.; Laget, M.; Remusat, V.; M'Rabet, M. K.; Rault, S.; Rathelot, P.; Azas, N.; Vanelle, P. Looking for new antileishmanial derivatives in 8-nitroquinolin-2(1H)-one series. *Eur. J. Med. Chem.* **2015**, *92*, 282–294.
- (20) Kieffer, C.; Cohen, A.; Verhaeghe, P.; Paloque, L.; Hutter, S.; Castera-Ducros, C.; Laget, M.; Rault, S.; Valentin, A.; Rathelot, P.; Azas, N.; Vanelle, P. Antileishmanial pharmacomodulation in 8-nitroquinolin-2(1H)-one series. *Bioorg. Med. Chem.* **2015**, *23*, 2377–2386.
- (21) Pedron, J.; Boudot, C.; Hutter, S.; Bourgeade-Delmas, S.; Stigliani, J.-L.; Sournia-Saquet, A.; Moreau, A.; Boutet-Robinet, E.; Paloque, L.; Mothes, E.; Laget, M.; Vendier, L.; Pratviel, G.; Wyllie, S.; Fairlamb, A. H.; Azas, N.; Courtioux, B.; Valentin, A.; Verhaeghe, P. Novel 8-nitroquinolin-2(1H)-ones as NTR-bioactivated antikinetoplastid molecules: synthesis, electrochemical and SAR study. *Eur. J. Med. Chem.* **2018**, *155*, 135–152.
- (22) Pedron, J.; Boudot, C.; Bourgeade-Delmas, S.; Sournia-Saquet, A.; Paloque, L.; Rastegari, M.; Abdoulaye, M.; El-Kashef, H.; Bonduelle, C.; Pratviel, G.; Wyllie, S.; Fairlamb, A. H.; Courtioux, B.; Verhaeghe, P.; Valentin, A. Antitrypanosomal pharmacomodulation at position 3 of the 8-nitroquinolin-2(1H)-one scaffold using pallado-catalyzed cross coupling reactions. *ChemMedChem* **2018**, *13*, 2217–2228.
- (23) Zaragoza, F.; Stephensen, H.; Peschke, B.; Rimvall, K. 2-(4-alkylpiperazin-1-yl)quinolines as a new class of imidazole-free histamine H₃ receptor antagonists. *J. Med. Chem.* **2005**, *48*, 306–311.
- (24) O'Brien, N. J.; Brzozowski, M.; Wilson, D. J. D.; Deady, L. W.; Abbott, B. M. Synthesis and biological evaluation of substituted 3-anilinoquinolin-2(1H)-ones as PDK1 inhibitors. *Bioorg. Med. Chem.* **2014**, *22*, 3781–3790.
- (25) Di, L.; Kerns, E. H.; Bezar, I. F.; Petusky, S. L.; Huang, Y. Comparison of blood-brain barrier permeability assays: in situ brain perfusion, MDR1-MDCKII and PAMPA-BBB. *J. Pharm. Sci.* **2009**, *98*, 1980–1991.
- (26) Jourdan, J. P.; Since, M.; El Kihel, L.; Lecoutey, C.; Corvaisier, S.; Legay, R.; Sopkova-de Oliveira Santos, J.; Cresteil, T.; Malzert-Fréon, A.; Rochais, C.; Dallemagne, P. Benzylphenylpyrrolizones with anti-amyloid and radical scavenging effects, potentially useful in Alzheimer's disease treatment. *ChemMedChem* **2017**, *12*, 913–916.
- (27) Van De Waterbeemd, H.; Camenisch, G.; Folkers, G.; Chretien, J. R.; Raevsky, O. A. Estimation of blood-brain barrier crossing of drugs using molecular size and shape, and H-bonding descriptors. *J. Drug Target* **1998**, *6*, 151–165.
- (28) Wager, T. T.; Hou, X.; Verhoest, P. R.; Villalobos, A. Moving beyond rules: the development of a central nervous system multiparameter optimization (CNS MPO) approach to enable alignment of druglike properties. *ACS Chem. Neurosci.* **2010**, *1*, 435–449.
- (29) Nepali, K.; Lee, H.-Y.; Liou, J.-P. Nitro-Group-Containing Drugs. *J. Med. Chem.* **2019**, *62*, 2851–2893.
- (30) Rosenkranz, E. J.; McCoy, E. C.; Mermelstein, R.; Rosenkranz, H. S. Evidence for the existence of distinct nitroreductases in *Salmonella typhimurium*: roles in mutagenesis. *Carcinogenesis* **1982**, *3*, 121–123.
- (31) Torrelee, E.; Bourdin Trunz, B.; Tweats, D.; Kaiser, M.; Brun, R.; Mazué, G.; Bray, M. A.; Pécoul, B. Fexinidazole - a new oral nitroimidazole drug candidate entering clinical development for the treatment of sleeping sickness. *PLoS Neglected Trop. Dis.* **2010**, *4*, No. e923.
- (32) Westland, J.; Dorman, F. QuEChERS extraction of benzodiazepines in biological matrices. *J. Pharm. Anal.* **2013**, *3*, 509–517.
- (33) Pkanalix. *Data Processing and Calculation Rules*. <http://pkanalix.lixoft.com/calculation-rules/> (accessed September 15, 2019).
- (34) Katsuno, K.; Burrows, J. N.; Duncan, K.; van Huijsduijnen, R. H.; Kaneko, T.; Kita, K.; Mowbray, C. E.; Schmatz, D.; Warner, P.; Slingsby, B. T. Hit and lead criteria in drug discovery for infectious diseases of the developing worlds. *Nat. Rev. Drug Discovery* **2015**, *14*, 751–758.