

**Manuscript version: Author's Accepted Manuscript**

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

**Persistent WRAP URL:**

<http://wrap.warwick.ac.uk/134007>

**How to cite:**

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk).

Understanding suicide risk in autistic adults: comparing the Interpersonal Theory of Suicide  
in autistic and non-autistic samples

*Abstract*

This study explored whether the Interpersonal Theory of suicide informs our understanding of high rates of suicidality in autistic adults. Autistic and non-autistic adults (n=695, mean age 41.7 years, 58% female) completed an online survey of self-reported thwarted belonging, perceived burden, autistic traits, suicidal capability, trauma, and lifetime suicidality. Autistic people reported stronger feelings of perceived burden, thwarted belonging and more lifetime trauma than non-autistic people. The hypothesised interaction between burdensomeness and thwarted belonging were observed in the non-autistic group but not in the autistic group. In both groups autistic traits influenced suicidality through burdensomeness/thwarted belonging. Promoting self-worth and social inclusion are important for suicide prevention and future research should explore how these are experienced and expressed by autistic people.

## *Introduction*

Research reports higher suicidality rates amongst adults diagnosed with an autism spectrum condition (ASC) than the general population, but research has yet to explain *how* the unique challenges facing autistic adults lead to suicidality (Cassidy & Rodgers, 2017; Segers & Rawana, 2014). A recent population study reported a nine-fold increase in death by suicide amongst autistic adults<sup>1</sup> (Hirvikoski et al., 2016) and higher rates of suicide ideation have been noted in autistic samples than general population, psychotic and student samples (Cassidy et al., 2014). Autistic adults more frequently experience general population suicide risk factors, such as abuse (Storch et al., 2013) and depression (Cassidy et al., 2014), but also report greater impact of less commonly reported risk factors, such as camouflaging (masking autistic characteristics to fit in) or coping with life without required support (Cassidy, Bradley, Shaw, & Baron-Cohen, 2018). However, studies to date have not applied theoretically based empirical models and so do not provide the detailed insight into suicide mechanisms required to design interventions that meet the specific needs of autistic people (Cassidy & Rodgers, 2017; Franklin et al., 2017; Joiner, 2005; Segers & Rawana, 2014; Van Orden et al., 2010). The current study, thus, explores for the first time how a widely cited theory of suicide, the Interpersonal Theory of Suicide (Joiner, 2005), applies to suicide in autistic adults and whether this is different from non-autistic adults.

(FIGURE 1 ABOUT HERE)

---

<sup>1</sup>This study refers to adults without intellectual disability who are reported to be at greatest risk of suicidal behaviour (Hirvikoski 2016).

The ITS proposes that three proximal risk factors for suicide, shown in Figure 1, lead to a lethal or near lethal suicide attempt: 1) strong feelings of ‘thwarted belonging’ (absence of reciprocal relationships with family, friends and society) and 2) ‘perceived burdensomeness’ (a belief that others are better off without oneself) (Joiner, 2005) are proposed to jointly lead to a desire for suicide. A suicide attempt will only be made if an individual has 3) ‘suicidal capability’, a biological change in the body’s pain and fear systems that allows rather than inhibits a suicide attempt. The theory suggests this stable trait develops in response to lifetime exposure to painful and frightening experiences (Chu et al., 2017; Joiner, 2005; Van Orden et al., 2010). The ITS has been tested in a range of populations (Anestis, J. C. et al., 2018; Bryan, Hernandez, Allison, & Clemans, 2013; Cole et al., 2013; Czyz, Berona, & King, 2015; Heelis, Graham, & Jackson, 2016; Miller, Esposito-Smythers, & Leichtweis, 2016; Wilson, Kowal, Henderson, McWilliams, & Péloquin, 2013; Wolford-Clevenger et al., 2017) with broad support for the theory reported in a meta-analysis of 122 studies (Chu et al., 2017), and its associations have under-pinned evidence-based efforts to increase identification of suicide risk (Ribeiro, Bodell, Hames, Hagan, & Joiner, 2013) and treatments (Joiner Jr, Van Orden, Witte, & Rudd, 2009a). Our earlier study reported that, in a non-clinical sample, thwarted belonging and perceived burden were associated with suicidality and that this association was not attenuated by high levels of autistic traits (Pelton & Cassidy, 2017). Further, an online survey of 98 autistic adults, reported that thwarted belongingness and perceived burdensomeness were associated with social dissatisfaction and current suicidal ideation (Dow et al., 2019). These studies were small in scale, and concluded with the recommendation that future research should assess the hypotheses of the ITS in a sample of autistic adults with a matched comparison group of non-autistic people. This is the aim of the current study.

The ITS may be of particular relevance to autistic people: its emphasis on the fundamental human needs of social self-worth and inclusion are conducive with views in the autistic community that the social marginalisation of autistic people should be central to understanding high rates of suicidality (Milton & Moon, 2012). The clinical diagnosis of an ASC comprises social, relationship and sensory differences, narrow interests and repetitive behaviours (APA 2013). Participatory research has reported that poor understanding and acceptance of such differences is associated with reduced social belonging, independence and quality of life and increased mental health difficulties for autistic people (Cage, Di Monaco, & Newell, 2018; Crane, Laura, Adams, Harper, Welch, & Pellicano, 2019; Mason et al., 2018). Autistic people are more likely to report behaviours indicative of thwarted belonging, such as loneliness (Haertl, Callahan, Markovics, & Sheppard, 2013; Hickey, Crabtree, & Stott, 2018; Müller, Schuler, & Yates, 2008), family stress, childhood maltreatment and intimate partner violence (Griffiths et al., 2019) than people who are non-autistic. ‘Social camouflaging’ (masking autistic mannerisms to ‘fit in’) (Hull et al., 2017), and the ‘double empathy problem’ (expectation that autistic people communicate social preferences and emotional states according to non-autistic norms) (Milton, 2012) reduce reciprocity in social interactions and could confer additional risk for thwarted belonging that feels hopeless and permanent (Joiner, 2005). Autistic adults frequently report experiences indicative of perceived burdensomeness, such as higher levels of unemployment (National Autistic Society 2016), incarceration (Fazio, Pietz, & Denney, 2012), homelessness (Stone, 2019), physical illness (Cashin, Buckley, Trollor, & Lennox, 2016; Hirvikoski et al., 2016) and lower self-esteem (Williamson, Craig, & Slinger, 2008) than non-autistic people. Research reports associations between unmet support needs, burdensomeness, poor mental health and suicidality (Camm-Crosbie, Bradley, Shaw, Baron-Cohen, & Cassidy, 2018; Cassidy et al.,

2018). Overall, this could suggest that autistic people experience stronger feelings of thwarted belonging and perceived burden than people who are not autistic.

The ITS proposes that suicidal capability develops in a dose-response relationship with lifetime experience of painful and frightening experiences (Joiner, 2005; Van Orden et al., 2010). Research reports shockingly high rates of trauma amongst autistic people (Griffiths et al., 2019): autistic children report higher rates of maltreatment (McDonnell et al., 2019), removal from their biological birth family (Green, Leadbitter, Kay, & Sharma, 2016), bullying (Mañano, Normand, Salvas, Moullec, & Aimé, 2016) school discipline (including exclusion), police contact and psychiatric hospitalisation (Humphrey, 2008; Turcotte, Shea, & Mandell, 2018). Autistic adults are more likely to be victims of hate crime (Beadle-Brown et al., 2014; Chaplin & M., 2018), sexual victimization (Brown-Lavoie, Vecili, & Weiss, 2014; Weiss & Fardella, 2018), incarceration (Fazio et al., 2012) and engage in non-suicidal self-injury (Cassidy et al., 2018; Maddox, Trubanova, & White, 2017). Overall, this suggests that autistic people experience more painful and frightening traumatic life events, which the ITS hypothesises suggests higher rates of suicidal capability than non-autistic people.

The ITS also proposes that gender differences in suicidal behaviour results from gender differences in the prevalence of thwarted belonging, perceived burden and suicidal capability. Amongst non-autistic people, men are reported to more frequently die by suicide because they are more likely to develop suicidal capability through violent sports, fighting or aggression. Women are more likely to think about suicide as they attach greater importance to social connections making them more vulnerable to thwarted belonging (Joiner, 2005). However, in contrast to the general population, autistic women have been reported more likely to die by suicide than autistic men (Hirvikoski et al., 2016). Autistic women are reported to be more socially motivated than autistic men (Lai, Lombardo, Auyeung,

Chakrabarti, & Baron-Cohen, 2015; Sedgewick, Crane, Hill, & Pellicano, 2019) but struggle to find genuine social reciprocity which could make them more vulnerable to thwarted belonging (Hull et al., 2017; Tint et al., 2018). Autistic women present distinct autistic characteristics which reduce likelihood of diagnosis (Duvekot et al., 2017; Dworzynski, Ronald, Bolton, & Happé, 2012; Evans, Boan, Bradley, & Carpenter, 2018; Ratto et al., 2018) typically associated with poor access to services (Cassidy et al., 2014), employment problems (Cassidy et al., 2014), and poor mental health (Hull et al., 2017; Trubanova, Donlon, Kreiser, Ollendick, & White, 2014), which could contribute to feelings of burdensomeness. Combined unmet support and social needs could explain higher incidence of mistreatment, such as abusive or coercive sexual relationships (Ohlsson Gotby, Lichtenstein, Långström, & Pettersson, 2018), maladaptive coping strategies such as controlled eating (Bargiela, Steward, & Mandy, 2016; Lai et al., 2015) or non-suicidal self-injury (Cassidy et al., 2018; Maddox et al., 2017) than autistic men. Overall, this could suggest that autistic women are more likely to experience thwarted belonging, perceived burden, lifetime trauma and suicidal capability than autistic men.

However, the Interpersonal Theory of Suicide argues that, in addition to being present, it is the *interaction* of theory constructs that provides the unique conjunction consistent with the rarity of suicide (Joiner, 2005; Van Orden et al., 2010). Thus, the theory proposes that the interaction of perceived burden and thwarted belonging are significantly associated with the presence of suicidal desire. Empirically, this interaction has been observed in a range of non-autistic groups, such as American (Van Orden, Witte, Gordon, Bender, & Joiner, 2008) and Chinese undergraduates (Zhang, Lester, Zhao, & Zhou, 2013), community samples (Campos & Holden, 2016), military personnel (Shelef, Fruchter, Mann, & Yacobi, 2014) and American Indians (O'Keefe et al., 2014) with a meta-analysis of 58 studies finding that the interaction of perceived burden and thwarted belonging added unique

variance in suicidal ideation beyond the main effects of each construct (Chu et al., 2017). However, our earlier study suggested that high levels of autistic traits could lead to under-reporting feelings of thwarted belonging, in line with well-documented reports of alexithymia (difficulty in expressing internal emotional states) in autistic people (Bird & Cook, 2013; Pelton & Cassidy, 2017). Furthermore, research has suggested that alternative risk factors such as social camouflaging (Cassidy et al., 2018), social support (Hedley, Uljarević, Wilmot, Richdale, & Dissanayake, 2017) or unmet support needs (Cassidy et al., 2018) may play a more significant role in the development of suicidal thoughts and behaviours in autistic people than non-autistic people. Overall, this could suggest that the model operates differently in autistic people. Thus, this study will explore for the first time whether the hypothesised interaction of perceived burden and thwarted belonging is associated with suicidal ideation in autistic adults and whether this is similar in non-autistic adults.

The ITS represents a novel innovation in suicide theory as it argues that risk factors for suicide ideation are distinct from those for suicide attempt. The construct of suicidal capability is proposed to uniquely enable a suicide attempt when present alongside the temporary cognitive states of thwarted belonging and perceived burden. Thus, suicidal capability theoretically doesn't exert a main effect on suicidal behaviour but, the three-way interaction of thwarted belonging, perceived burden and suicidal capability will transform suicidal ideation into a lethal or near lethal suicide attempt (Joiner, 2005; Van Orden et al., 2010). This three-way interaction has been upheld in samples of firefighters (Chu, Buchman-Schmitt, Hom, Stanley, & Joiner Jr, 2016) and military personnel (Anestis, M. D., Khazem, Mohn, & Green, 2015), but it has received less empirical attention than the two-way interaction described above: the same meta-analysis (Chu et al., 2017) included only 13 studies of the three-way interaction and reported that it was only significantly associated with suicide attempt when suicide attempt history was measured continuously (Chu et al., 2017).



Factors that trigger transition from suicidal thoughts to behaviours are of great concern in autistic people given reports that autistic people are more likely to attempt suicide (Chen, Pan, Lan, Hsu, Huang, Su, Li, Lin, Wei, Chen, & Bai, 2017), select lethal methods for a suicide attempt yet be unknown to clinical services (Kato et al., 2013) and die by suicide (Hirvikoski et al., 2016) than non-autistic people. Commentators have speculated on the reasons for this suggesting, for example, that autistic people may relate differently to death than non-autistic people or may be more likely to perseverate on death (Gillberg, 2002; Lai, J., Rhee, & Nicholas, 2017). However, empirical research to date has not yet explored how distinct risk factors may lead to suicide attempt versus ideation in autistic people. Thus, this study will explore, for the first time, whether the theoretical three-way interaction of thwarted belonging, perceived burden and suicidal capability is significantly associated with suicide attempt in autistic and non-autistic people.

Furthermore, the ITS argues that suicidal capability is developed through opponent process theory: repeated painful and frightening experiences ultimately render future painful and frightening experiences as bringing emotional and/ or physical pain relief (Joiner, 2005; Solomon & Corbit, 1974). As outlined above, autistic people are more likely to experience lifetime trauma compared to non-autistic people but, in addition, autistic people have been reported to experience a more profound effect of trauma due to absent social support, differences in pain and emotional communication and cognitive rigidity (Kerns, Newschaffer, & Berkowitz, 2015; Rumball, 2018). The functional model of suicidal capability argues that the development of suicidal capability depends on innate vulnerabilities in interaction with the environment (Smith & Cukrowicz, 2010). Thus, differences in pain (Rattaz, Dubois, & Baghdadli, 2016), emotional processing and expression, high trait anxiety and sensory sensitivities (Mazefsky et al., 2013) could suggest that suicidal capability is more quickly developed in autistic people. Thus, this study will explore whether the pathway from

traumatic life events to suicidal thoughts and behaviours is through suicidal capability and compare whether this is similar to non-autistic people.

Finally, the ITS proposes that distal factors, including oft-cited suicide risk factors - such as mental health difficulties - lead to suicidal behaviour because they increase the likelihood that an individual will experience thwarted belonging, perceived burden (Davidson, Wingate, Grant, Judah, & Mills, 2011; Silva, Ribeiro, & Joiner, 2015; Van Orden et al., 2010). In clinical and non-clinical samples autistic characteristics are independently associated with suicidality beyond other factors, such as psychiatric disorder (Pelton & Cassidy, 2017; Upthegrove et al., 2018), previous suicide attempts (Chen et al., 2017), depression, anxiety, unemployment and satisfaction with living arrangements (Cassidy et al., 2018). Understanding *how* autistic traits associate with lifetime suicidality is vital to inform suicide prevention interventions: autistic traits are not modifiable but support can be provided to reduce the negative social impacts (Joiner Jr, Van Orden, Witte, & Rudd, 2009b; Milton & Moon, 2012). Our earlier study reported that in a non-clinical young adult sample the association between autistic traits and lifetime suicidality was mediated by perceived burden and thwarted belonging suggesting (Pelton & Cassidy, 2017). This study will seek to replicate this finding and compare whether a similar pathway exists for autistic people.

### *The current study*

In summary, the current study aimed to fill a gap in previous research by examining how the ITS explains suicide in autistic people and comparing this to non-autistic people. This study will explore (i) prevalence rates of ITS constructs in autistic and non-autistic people, (ii) whether hypothesised interactions between ITS variables are present in autistic and non-autistic groups (iii) putative pathways from distal factors to suicidal thoughts and

behaviours through theory constructs are similar in autistic and non-autistic people.

Specifically, this study hypothesises that,

- (i) Given more frequent reports of loneliness, unmet support needs and trauma, autistic people will report stronger feelings of thwarted belonging and perceived burdensomeness, higher suicidal capability and more frequent lifetime trauma than non-autistic people.
- (ii) Given gender differences in the autistic female phenotype, autistic women will report stronger feelings of thwarted belonging and perceived burden, higher suicidal capability and more frequent lifetime trauma than autistic men.
- (iii) Given broad published support, the respective interactions in ITS variables in association with suicidal ideation and suicide attempt will be observed in the non-autistic group but - given suggested differences in suicidality in autistic people - these may not be observed in the autistic group.
- (iv) In both groups, given the proposed development of suicidal capability in response to lifetime trauma there will be a significant pathway from lifetime trauma through suicidal capability to suicidality but – given reported more profound effect of trauma – the pathway from lifetime trauma to suicidal capability will be stronger in autistic people.
- (v) In both groups, the previously reported pathways from autistic traits to suicidality (Pelton & Cassidy, 2017) through perceived burden and thwarted belonging will be observed.

### *Methods*

#### *Participants*

Participants were 695 autistic (64.6% female, mean age= 41.9, 18-90 years) and non-autistic people (58.7% female, mean age = 41.3, 18-73 years) described in Table 1. Autistic participants self-reported a diagnosis of ASC from a trained clinician. Mean AQ-S scores in the autistic group were consistent with published clinical cut off score of >65 indicative of ASC (Hoekstra et al., 2011a), with AQ-S scores significantly lower in the non-autistic compared to the autistic group ( $t(636.98)=-31, p<.001$ ).

There was no significant difference in age ( $t(672.7)=-.55, p=.59$ ) or proportion of female participants ( $X^2(1)=2.53, p=.11$ ) between the autistic and non-autistic group. Significantly more non-autistic people reported being in full time employment ( $X^2(1)=14.47, p<.001$ ) and holding a post-graduate degree ( $X^2(1)=7.30, p<.01$ ) than autistic people. Autistic people were more likely to report at least one additional neurodevelopmental condition ( $X^2(1)=55.45, p<.001$ ) and at least one mental health condition ( $X^2(1)=106.32, p<.001$ ) than non-autistic people.

Non-autistic people reported significantly lower lifetime suicidality ( $t(656.98)=-14.52, p<.001$ ) than autistic people. Reported rates of past suicide attempt were higher in the autistic (38.3%) than the non-autistic group (10.5%), our earlier study of 8.6% (Pelton & Cassidy, 2017) and the UK Adult Psychiatric Morbidity study (6.7%) (McManus, Bebbington, Jenkins, & Brugha, 2016).

The autistic group was recruited through the Cambridge Autism Research Database, the Autistica Discover network, local and national autism organisations and social media. The non-autistic group was recruited through the Cambridge Psychology Database with

opportunity sampling, Coventry University psychology research participation scheme, social media channels and suicide-focussed research websites to balance group size, mean age and gender frequency with the autistic group. Adverts clearly stated that the study would ask about suicidal thoughts and behaviours but did not specify inclusion criteria based on these.

### *Materials*

*Demographics:* participants indicated age, gender, living status, employment status, mental health, autism diagnoses and the presence of any other neurodevelopmental disorders.

**Thwarted belonging** and **perceived burden** were measured using *The Interpersonal Needs Questionnaire 10 (INQ-10)*. The INQ 10 is a 10 item validated scale containing two subscales for distinct but related interpersonal constructs that represent the desire to die: ‘thwarted belonging’ and ‘perceived burdensomeness’ (Van Orden, Cukrowicz, Witte, & Joiner, 2012). It contains statements such as ‘These days the people in my life would be better off if I were gone’ and ‘these days I think the people in my life wish they could be rid of me’ which are responded to on a 7-point scale from ‘strongly agree’ to ‘strongly disagree’. Our steering group of autistic adults advised choosing the INQ-10, with comparable validity (Hill et al., 2015) over the INQ-15 to avoid frustration from questions that could be perceived as similar by autistic people. They also advised changing the instructions to clearly communicate the intended meaning to autistic people without disadvantaging non-autistic people: *‘Please read the items below. Click on the option that best describes how you have been feeling. Where the questionnaire refers to ‘these days’ please consider how you have been feeling in general over the past two weeks’* (original wording in Appendix). The scale author confirmed that change was acceptable prior to administration. (Burdensomeness subscale Cronbach’s alpha = .93, Thwarted belonging Cronbach’s alpha=.92)

**Capability for Suicide** was measured using *Acquired Capability for Suicide Scale – Fearlessness of Death (ACSS-FAD)* (Ribeiro, Witte, Van Orden, Selby, Gordon, Bender, & Joiner, 2014a). The ACSS-FAD is a validated 7-item scale measuring a sense of fearlessness around the concept of death. Scale items include ‘the prospect of my own death arouses anxiety in me’ and ‘I am not at all afraid to die’. Responses are given on a five point Likert scale with scores from 0 to 4 with higher scores indicating a higher capability for suicide (Ribeiro, Witte, Van Orden, Selby, Gordon, Bender, & Joiner Jr, 2014). Cronbach’s alpha =.85.

**Traumatic life events** were measured using *Vulnerability Experience Quotient (VEQ)* (Griffiths et al., 2019). The VEQ is a 60-item scale which has been developed through participatory methods with autistic adults to reflect adverse life experiences across 10 themes, such as childhood maltreatment, non-suicidal self-injury, bullying and victimisation as a child or adult and discrimination. This scale reflects many of the themes of Painful and Provocative Events scale designed for use in the ITS but has been developed and validated to reflect the experiences of autistic people (Griffiths et al., 2019). The response scale has been amended to give an indication of frequency as this is hypothesised to enable transition from suicide ideation to attempt. A simplified scale of ‘not applicable’, ‘never’, ‘once’ or ‘more than once’ assessed frequency without repeated need for counting instances of traumatic life events. This response scale has been recommended in a recent psychometric evaluation of the Painful and Provocative Events Scale (Brown, Roush, Marshall, Mitchell, & Cukrowicz, 2018) and was confirmed with the scale author prior to administration. Cronbach’s alpha=.93

**Lifetime suicidal thoughts and behaviours** were measured by *Suicide Behaviours*

*Questionnaire – Revised (SBQ-R), item 1*: The first item of the SBQ-R measures lifetime suicidal ideation and/ or attempt with the question ‘Have you ever thought about or attempted to kill yourself?’. There are six possible responses from ‘never’ to ‘I have attempted to kill myself and really hoped to die’ (Osman et al., 2001). Participant responses are categorised in four subgroups from non-suicidal to suicide attempt. Item 1 was employed in our previous study (Pelton & Cassidy, 2017) and has been reported to demonstrate comparable measurement properties amongst autistic and non-autistic adults (Cassidy et al., 2018)

**Autistic characteristics** were measured using *Autism Quotient Short Form (AQ-S)*: The AQ-S is a 28-item subset of the full 50 item Autism Quotient based upon DSM-IV criteria for autism (APA 2000). The scale includes items such as ‘it does not upset me if my daily routine is disturbed’ and ‘I find it easy to work out what someone is thinking or feeling’ with a four item response scale from 1 “definitely agree” to 4 “definitely disagree (Hoekstra et al., 2011b). It is scored using the full four-point Likert scale giving a total range from 28-112 and a cut off of 65 for potential consideration of a clinical diagnosis of autism. Scores are highly correlated with the full AQ-50 and the AQ-S is reported to demonstrate the same latent traits in autistic and non-autistic people (Murray, Booth, McKenzie, Kuenssberg, & O’Donnell, 2014). Cronbach’s alpha = .95.

## **Procedure**

Participants were invited to complete a survey of online self-report measure of self-report questionnaires using Qualtrics. The dataset from which participants were drawn also includes possibly autistic people and four-week repeated measures of anxiety and depression. Analyses planned with these data are described in the discussion. Research reports that people feel more able to disclose potentially sensitive information about stigmatised

behaviours such as suicide and self-harm online than in situations where they cannot remain anonymous (Michaels, Chu, Silva, Schulman, & Joiner, 2015; Nock et al., 2008).

Participants indicated informed consent to participate via an online form. They were warned of the content of questions in each section, prompted to take breaks and given information about support services throughout the survey. The study materials and scales were designed with the assistance of autistic adults (one male, one female) who suggested modifications to instructions to clarify meaning, suggested substitutions of scales to maximise clarity for autistic people, suggested amendments to online instructions and guidance to reduce risk to ensure that the study was equally accessible for autistic and non-autistic people. The study received ethical approval from Coventry University Psychology Ethics Committee and was approved by the scientific advisory group at the Autism Research Centre, University of Cambridge.

(TABLE 1 ABOUT HERE)



## Results

### *Analytic Approach*

Data were analysed in SPSS version 25. Boxplots were examined for univariate outliers and multivariate outliers were identified using Mahalanobis distance. Non-normal distribution of burdensomeness in the non-autistic group was addressed with square root transformation: kurtosis reduced from 5.22 to 2.14 and extreme outliers came within normal distribution. Independent samples t-tests and three-way ANOVA were used to explore differences in group means (thwarted belonging, perceived burden, suicidal capability, lifetime trauma). Multinomial logistic regression explored whether the ITS hypothesised interactions were associated with past suicide ideation and suicide attempt. Simple linear regressions explored the association between ITS constructs and suicidal thoughts and behaviours. Mediation analysis explored the pathways from distal factors to suicidality through ITS constructs using ordinary least squares path analysis (PROCESS version 3.3 model 4) and employing bootstrapping for non-normal distribution. PROCESS model 59 explored whether mediation models were moderated by autism diagnosis.

*Hypothesis (i) autistic people report stronger feelings of thwarted belonging, perceived burden, higher suicidal capability and more prevalent lifetime trauma than non-autistic people.*

As shown in Table 1, autistic people reported significantly higher burdensomeness ( $t(635.09)=-11.33, p<.001$ ), thwarted belonging ( $t(652.28)=-16.43, p<.001$ ) and traumatic life events ( $t(648.83)=-17.46, p<.001$ ) than non-autistic people, but there was no significant difference in the mean score for suicidal capability between groups.

*Hypothesis (ii) autistic women report stronger feelings of thwarted belonging, perceived burden, higher suicidal capability and more frequent lifetime trauma than autistic men.*

Three-way ANOVA reported there was no significant difference in mean scores for thwarted belonging ( $F(2,337)=.55$ ,  $p=.575$ ), perceived burdensomeness ( $F(2,337)=.27$ ,  $p=.762$ ), suicidal capability ( $F(2,337)=2.87$ ,  $p=.058$ ) and traumatic life events ( $F(2,331)=1.21$ ,  $p=.332$ ) between autistic men and autistic women and those who identify as other gender. As this finding was against the direction of our hypothesis, moderation analysis explored whether the association between any predictor variables (burden, capability, thwarted belonging and traumatic life events) and lifetime suicidality was moderated by gender. We found no significant moderation. Furthermore, in this sample, there was no significant difference in reported lifetime suicidality between autistic men and autistic women ( $F(2,337)=2.08$ ,  $p=.126$ ). Thus, data from autistic men and women were analysed together in subsequent analyses.

In the non-autistic group<sup>2</sup>, independent t-tests showed that non-autistic men reported significantly higher suicidal capability ( $t(325)=2.02$ ,  $p=.044$ ) and significantly lower perceived burdensomeness ( $t(296.74)=-2.24$ ,  $p=.026$ ) than non-autistic women. There was no significant difference in thwarted belonging ( $t(328)=1.02$ ,  $p=.27$ ) between non-autistic men and non-autistic women.

---

<sup>2</sup> The 'other' gender group were not included as  $n = 2$ .

*Hypothesis (iii)(a) The interaction of thwarted belonging and perceived burden is significantly associated with suicidal thoughts.*

A multinomial logistic regression was undertaken with the categorical outcome variable SBQ-R item 1 (frequencies shown in Table 1) and the predictors thwarted belonging, perceived burden and their interaction term.

As shown in Table 2, in the non-autistic group, participants who reported significantly lower scores of thwarted belonging (Wald  $X^2(1)=18.16$ ,  $p<.001$ ), and perceived burden (Wald  $X^2=10.06$ ,  $p=.002$ ) were more likely to endorse the ‘no past suicidality’ response option versus ‘past suicidal thoughts’ and their interaction was significant within the regression (Wald  $X^2=10.67$ ,  $p=.001$ ).

(TABLE 2 ABOUT HERE)

In the autistic group, participants who reported higher perceived burden (Wald  $X^2(1)=4.13$ ,  $p=.04$ ) and thwarted belonging (Wald  $X^2(1)=6.25$ ,  $p=.01$ ) were more likely to endorse ‘past suicide attempt’ than ‘past suicidal thoughts’ and the interaction between thwarted belonging and perceived burden was not significant within the model. The model explained only 10% of the variance in the autistic group versus 31% in the non-autistic group.

*Hypothesis (iii)(b): the three-way interaction of thwarted belonging, perceived burden and suicidal capability will be significantly associated with past suicide attempt.*

A further multinomial logistic regression was undertaken with the categorical outcome variable SBQ-R item 1 (frequency responses shown in Table 1) and the predictors were

thwarted belonging, perceived burden and suicidal capability, the two-way interaction (belong\*burden) and their three-way interaction (belong\*burden\*capability).

As shown in Table 3, in the non-autistic group, participants who reported significantly lower levels of suicidal capability (Wald  $X(1)^2=5.98$ ,  $p=.014$ ) were more likely to endorse past suicidal thoughts versus past suicide attempt. The three-way interaction of perceived burden, thwarted belonging and suicidal capability was not significant within the model.

In the autistic group, as shown in Table 3, participants who reported significantly lower levels of suicidal capability (Wald  $X^2=18.89$ ,  $p<.001$ ) were more likely to endorse ‘past suicidal thoughts’ versus ‘past suicide attempt’ and the three-way interaction of thwarted belonging, perceived burden and suicidal capability was not significant within the model. In the autistic group, the model explained only 15% of the variance versus 34% in the non-autistic group.

(TABLE 3 ABOUT HERE)

*Hypothesis (iv): the association between lifetime trauma and suicidality will be significantly mediated by suicidal capability in both groups. In the autistic group the association between trauma and suicidal capability will be significantly strengthened.*

Simple linear regressions, shown in Table 4, reported that autistic traits, perceived burden, thwarted belonging, suicidal capability and traumatic life events were significantly associated with lifetime suicidality.

(TABLE 4 ABOUT HERE)

Simple mediation analysis was undertaken with the predictor, lifetime trauma, the mediator, suicidal capability and the outcome variable, suicidality. There was a direct effect of trauma on lifetime suicidality independent of suicidal capability in autistic ( $c'=.02$ ,  $p<.001$ ) and non-autistic ( $c'=.037$ ,  $p<.001$ ) groups and a weak but significant indirect effect of trauma on lifetime suicidality through suicidal capability in the autistic group ( $ab=.0001$ , 95% CI [.0001, .002]). As shown in Figure 2, moderation analyses reported an attenuated association between lifetime trauma and lifetime suicidality in the autistic group ( $VEQ*diagnosis=-.016$ ,  $SE=.004$ ,  $p<.001$ ) and no significant moderation of other mediation pathway by autism diagnosis.

(FIGURE 2 ABOUT HERE)

*Hypothesis (v): the association between autistic traits and suicidality will be significantly mediated by perceived burden and thwarted belonging in autistic and non-autistic groups.*

Simple mediation analysis was undertaken with the predictor autistic traits, the mediator thwarted belonging and the outcome variable suicidality. As shown in Figures 3 and 4, results indicated a significant indirect effect of autistic traits on lifetime suicidality through thwarted belonging in both autistic ( $ab=.008$ , 95% CI [.004, .012]) and non-autistic groups ( $ab=.019$ , 95% CI [.014, .025]). Moderation analysis revealed that the pathway from thwarted belonging and lifetime suicidality was significantly attenuated in the autistic group ( $belonging*autism\ diagnosis=-.027$ ,  $SE=.010$ ,  $p=.008$ ) as shown in Figure 5 but no other mediation pathway was significantly moderated by autism diagnosis.

(FIGURE 3 ABOUT HERE)

(FIGURE 4 ABOUT HERE)

(FIGURE 5 ABOUT HERE)

A further simple mediation analysis was undertaken with the predictor autistic traits, the mediator perceived burden and the outcome variable suicidality. As shown in Figures 6 and 7, there was a significant indirect effect of autistic traits on lifetime suicidality through perceived burden in both autistic ( $ab=.004$ , 95% CI [.002, .007]) and non-autistic ( $ab=.011$ , 95% CI [.008, .015]) groups. Moderation analyses reported that the pathway from perceived burden to lifetime suicidality was significantly attenuated in the autistic group ( $burdensomeness*autism\_diagnosis=-.048$ ,  $SE=.099$ ,  $p<.001$ ) as shown in Figure 8, but no other pathway was significantly moderated by autism diagnosis.

(FIGURE 6 ABOUT HERE)

(FIGURE 7 ABOUT HERE)

(FIGURE 8 ABOUT HERE)

## **Discussion**

This study aimed, for the first time, to explore whether the associations of the Interpersonal Theory of Suicide are informative for understanding and addressing high rates of suicidality in autistic adults. Autistic adults reported stronger feelings of thwarted belonging, perceived burdensomeness and traumatic life events than non-autistic adults. Autistic men and women reported similar levels of perceived burden, thwarted belonging, traumatic life events and suicidal capability. By contrast, non-autistic men reported higher suicidal capability and weaker feelings of perceived burden than non-autistic women. The proposed interaction between thwarted belonging and perceived burden was present in the

non-autistic group but not the autistic group. In both groups, participants endorsing lower suicidal capability were more likely to report suicide ideation versus attempt. In both groups, there was a direct effect of trauma on lifetime suicidality and autistic traits led to feelings of thwarted belonging and perceived burden, which led to lifetime suicidality. Finally, the pathway from each of thwarted belonging and perceived burden and lifetime trauma to lifetime suicidality was significantly attenuated in the autistic group. Overall, these results add to the limited body of knowledge describing the development of suicidal thoughts and behaviours in autistic people (Cassidy et al., 2018; Hedley, Uljarević, Foley, Richdale, & Trollor, 2018a) as well as extending previous explorations of the influence of autistic traits in the ITS (Pelton & Cassidy, 2017) and how the ITS works in autistic people (Dow et al., 2019).

Our findings echo a well-established body of research reporting higher rates of lifetime suicidality in autistic than non-autistic adults (Chen et al., 2017; Hirvikoski et al., 2016; Kirby et al., 2019). In line with the ITS hypotheses and the hypotheses of this study, this was accompanied by higher levels of thwarted belonging and perceived burden in the autistic group than in the non-autistic group (Joiner, 2005; Van Orden et al., 2010). This finding is consistent with research suggesting that social isolation and burdensomeness may contribute to poor mental health, reduced quality of life and suicidality in autistic adults (Camm-Crosbie et al., 2018; Cassidy et al., 2018; Hedley et al., 2017; McConachie et al., 2018). Autistic people also reported significantly higher rates of traumatic life events than people who were non-autistic, consistent with other recent research (Fuld, 2018; Rumball, 2018; Taylor & Gotham, 2016) and with research suggesting that trauma contributes to suicidality (Storch et al., 2013). Against our expectations there was no significant difference in suicidal capability between the autistic and non-autistic groups.

Against recent research findings, this study reported no significant difference in reported lifetime suicidality between autistic women and men. Recent population studies have reported significantly higher rates of suicidality amongst autistic women than men (Hirvikoski et al., 2016; Kirby et al., 2019), a trend noticeable for its opposite pattern to the non-autistic population. However, this study echoes the findings of another study in our group, which also found no significant difference in suicidality between autistic women and men (Cassidy et al., 2018). Against our hypotheses, there was no significant difference in mean scores on any other study variable between autistic women, men and other gender. This is in contrast to the non-autistic group where the hypotheses of the ITS (Joiner, 2005) were largely upheld with women reporting higher burdensomeness and men reporting higher suicidal capability. This suggests that suicidal behaviour in autistic people may not reflect expected gender-based patterns and processes observed in the non-autistic population. Clinicians should be aware that, in particular, autistic women may be at increased risk if they are assumed to be at lower risk due to female gender. However, these findings are preliminary and future research should explore whether, and how, gender may influence suicidality in autistic adults.

We examined, for the first time, the hypothesised interactions of the ITS in autistic and non-autistic adults. In the non-autistic group, the hypothesis of the ITS regarding suicide ideation was largely upheld: participants reporting lower perceived burden and thwarted belonging were more likely to report no past suicidal behaviour versus suicide ideation and the interaction between the variables was significant. In the autistic group, by contrast, the ITS was not supported: participants reporting stronger feelings of thwarted belonging and perceived burden were more likely to report suicide attempt than suicide ideation or no past



suicidal behaviour. This suggests that there may be differences in how suicidal thoughts and behaviours develop in autistic people and, worryingly, that increases in feelings of thwarted belonging and burdensomeness may lead to suicide attempt rather than ideation. This is in line with commentary and anecdotal report that autistic people may transition to suicide attempt more quickly than people who are not autistic (Kato et al., 2013). Clinicians should be aware that suicidal thoughts and behaviours may develop differently in autistic people and, in the absence of validated care pathways and measures, undertake individualised, detailed risk assessments. Furthermore, there was no hypothesised interaction between thwarted belonging and perceived burden. The ITS proposes that perceived burden and thwarted belonging are related but distinct constructs but our results could suggest that autistic differences – possibly theory of mind (Baron-Cohen, Leslie, & Frith, 1985) or cognitive rigidity - could render them more separate in autistic than non-autistic people. This is in line with our earlier study, which suggested that differences in theory of mind could reduce the expression of these constructs (Pelton & Cassidy, 2017). Overall, this suggests that future research should continue to explore how risk factors for suicide may be different in autistic people as well as considering whether the expression of such states may present differently.

We found only partial support for the suicidal capability construct in both groups: participants reporting significantly lower suicidal capability were more likely to report suicide ideation rather than suicide attempt suggesting that suicidal capability is uniquely associated with suicide attempt. However, the three-way interaction of thwarted belonging, perceived burden and suicidal capability was not significant in either group. This is in line with recent meta-analysis which reported similar findings when suicide attempt was measured dichotomously as in this study (Chu et al., 2017). This is also in line with suicide theorists who agree with Joiner's assertion that a change in the relationship with fear of death and with pain is necessary to transition from suicide ideation to attempt but have argued that

other factors are also significant. For example, the Integrated Motivational-Volitional model describes suicidal capability as only one of many possible volitional moderators that may provide the bridge from suicidal intention to attempt (O'Connor, 2011; O'Connor, Cleare, Eschle, Wetherall, & Kirtley, 2016). This suggests an urgent need for research to explore distinct risk factors for suicide attempt in autistic and non-autistic people. Future research should be theoretically rigorous, explore the applicability of other suicide models and explore alternative pathways to confirm direction of causality. This could also include inductive theoretical research taking as its starting point individual accounts of autistic people, and comparing those against current theoretical accounts to build a valid theory. However, all interpretations of these findings regarding the hypothesised interactions should be considered in the context that, in the autistic group, the variables together accounted for around half the variance compared to the non-autistic group. This is discussed below.

In both groups, trauma was significantly directly associated with lifetime suicidality. This finding is in line with research in non-autistic people reporting significant negative life outcomes, including mental health difficulties and suicidality resulting from trauma (Luukkonen, Räsänen, Hakko, & Riala, 2009). However, this does not fully explain *how* trauma is associated with lifetime suicidality as the direct path does not support the ITS hypothesis that painful and provocative events lead to a change in the body's fear and pain system (suicidal capability) that enables suicidal behaviour. This could suggest that the measure of suicidal capability employed in this study does not truly represent the changes hypothesised to result from trauma (Joiner, 2005). Within this study, we used the most widely validated measure of suicidal capability – the ACSS-FAD - however, this represents a single construct – a reduced fear of death – and the scale author has already concluded that this may be too narrow to measure the changes that lead to suicidality (Ribeiro, Witte, Van Orden, Selby, Gordon, Bender, & Joiner, 2014b). Clinicians should thus be aware that those

with traumatic backgrounds are at increased risk but we don't yet have measures to clearly detail why. Future research should explore whether alternative suicide theories are able to provide detailed empirical evidence regarding the proposed faster transition to suicide attempt amongst autistic people and associated higher rates of death by suicide. An alternative approach would be to take as a starting point the lived experience of those who experience these difficulties and use these to build an inductive theory.

In the autistic group, there was also an indirect effect of trauma on suicidality through suicidal capability but detailed analysis of this finding found this did not support our proposal of a strengthened pathway from trauma to suicidal capability in autistic people. By contrast, we found a significant attenuation of the association between traumatic life events and lifetime suicidality which suggests that trauma (at least sometimes) occurs in the absence of suicidal behaviour for autistic people. Given such high rates of trauma in autistic people, this could suggest that these are everyday experiences for autistic people, against which they become resilient, rather than rare experiences associated with suicidality. This could, however, also reflect the use of our 'once' or 'more than once' rating scale that could under-report trauma for autistic people: the original Painful and Provocative Events Scale designed for non-autistic people includes a range of categorical up to 20 incidents of each painful event (Brown, Roush, Marshall, Mitchell, & Cukrowicz, 2018). It is vital that future research continues to explore how trauma contributes to suicidality alongside continuing research to understand of how trauma impacts autistic people. Suicide prevention efforts should continue to address the enforcement of statutory protection for autistic people with a view to reducing trauma.

The findings of our earlier study (Pelton & Cassidy, 2017), and our hypothesised pathways from autistic traits to suicidality through both perceived burden and thwarted belonging, were upheld. This suggests that feelings of social disconnection and reduced self-

worth associated with autistic traits provide the bridge to suicidal behaviour. This is in line with research linking autistic traits to increased loneliness and social disconnection in autistic and non-autistic populations (Hedley, Uljarević, Wilmot, Richdale, & Dissanayake, 2018; White & Roberson-Nay, 2009), and in line with research suggesting these concepts may link to suicidality (Hedley, Uljarević, Foley, Richdale, & Trollor, 2018b). This supports the ITS model that, in any population group, distal risk factors contribute to suicidality via the two proximal risk factors thwarted belonging and perceived burden (Christensen, Batterham, Mackinnon, Donker, & Soubelet, 2014; Cole et al., 2013; Corbin, 2017; Horton et al., 2016; Miller et al., 2016; Pennings, Finn, Houtsma, Green, & Anestis, 2017; Van Orden et al., 2008). This also suggests that these two concepts may represent modifiable constructs that can be targeted in suicide prevention programmes, thus suicide prevention programmes that promote social inclusion and feelings of self-worth are important in autistic and non-autistic people.

However, we also reported attenuated associations between each of perceived burden, thwarted belonging and lifetime suicidality in the autistic group compared to the non-autistic group. This could, similar to the case of trauma, suggest that these are everyday experiences for autistic people, rather than experiences that differentiate those experiencing suicidal thoughts and behaviours from those who are not. Similar findings have been reported in clinical groups: a study of individuals with a first episode psychosis reported that the concepts of the ITS resonated with the experience of psychosis regardless of suicidality (Heelis et al., 2016). An alternative interpretation, however, is that both thwarted belonging and burdensomeness could be proximal risk factors for suicide in autistic people yet may be experienced and expressed differently by autistic adults. This could suggest that the INQ measure, designed for use in the general population, doesn't capture the same concepts for autistic people. This is in line with a wide body of research reporting the need for measures,

which accurately describe and capture emotional and psychological processes in autistic adults (Cassidy et al., 2018; Cassidy, Bradley, Bowen, Wigham, & Rodgers, 2018; Gotham, Unruh, & Lord, 2015). Indeed, our design group already identified problematic items: for example, ‘These days, I feel like I belong’ is confusing in its lack of specificity whereas ‘These days I often feel like an outsider at social gatherings’ could describe everyday autistic life rather than indicate elevated suicidal feelings. In the case of burdensomeness this attenuation of association could support our earlier assertion that the second latent construct of burdensomeness (Van Orden et al., 2010) described as self-hatred, poor self-esteem and agitation could present in autistic people even if differences in theory of mind reduce the ability to consider oneself a burden on others (Pelton & Cassidy, 2017). This is possibly a more likely explanation, as recent qualitative research has identified burdensomeness as a significant theme in autistic quality of life and mental health difficulties (Camm-Crosbie et al., 2018; Crane, L., Adams, Harper, Welch, & Pellicano, 2017). Future research should seek to understand the experience of thwarted belonging and burdensomeness from the perspective of autistic people and identify the specific experiences, if any, which could indicate a desire to die. However, as a first step, future research should explore potential measurement differences in the INQ-10 and ACSS-FAD between autistic and non-autistic people.

This study has several strengths: it is the first large-scale study to compare suicidal behaviour in autistic and non-autistic samples within a well-validated theory of suicidal behaviour. This responds to repeated calls within the research literature (Segers & Rawana, 2014) and within the autistic community for theoretically based suicide research in autistic adults (Cassidy & Rodgers, 2017). Thus, it explores an under-researched area and provides a foundation for broader theoretical modelling work within the academic and practitioner communities. This foundation is vital to respond to calls from within the autistic community to explore other specified priorities, including, but not limited to identification of suicidal

thoughts and behaviours, appropriate adaptation of interventions and designing and delivering crisis services that facilitate help-seeking (Cassidy et al 2019).

It is also important to acknowledge this study's limitations. This study included a single outcome variable, lifetime suicidality, whereas the ITS specifies current suicidal ideation, attempt and death by suicide as its outcomes. The key focus of this study was to examine differences between the autistic and non-autistic groups, thus, an outcome variable was selected with known measurement similarities across autistic and non-autistic people (Cassidy et al., 2018). This study has also not explored the role of anxiety and depression and complex mental health difficulties in the autistic group, and the ITS makes specific hypotheses regarding how mental health difficulties impact suicidal behaviour, which should be urgently explored in future research. This study has relied on self-reported autism diagnosis due to sample size required but future research could employ a diagnostic test. Finally, mediation analyses are based upon cross-sectional data so limitations on directions of causality are acknowledged. Future analyses are planned with repeated measures data, which will also explore the role of depression and anxiety. Finally, the study has not explored the potential influence of demographic variables such as age, living or marital status. There is an urgent need to incorporate these into future research studies but these were beyond the scope of this paper.

In summary, this study represents the first large scale study to compare the associations of the ITS in autistic and non-autistic samples. This study provides evidence that perceptions of burdensomeness, reduced social belonging and exposure to traumatic life events are significantly associated with lifetime suicidality in autistic adults and addressing these is vital to reduce suicide rates. However, this study also highlights the importance of understanding how these feelings are experienced and communicated by autistic people and ensuring that our current measures and clinical practices capture these. This study also

highlights the fact that a model of suicidal behaviour that works for autistic people may need to be tailored to reflect distinct experiences, communication and social preferences of autistic people. Public policy should urgently address rates of stigmatising and abusive traumatic experiences of autistic people. Overall, improving inclusion and self-worth and reducing trauma could reduce death by suicide in autistic and non-autistic people.

## Appendix

Original wording instructions of the Interpersonal Needs Questionnaire:

“The following questions ask you to think about yourself and other people. Please respond to each question by using your own current beliefs and experiences, NOT what you think is true in general, or what might be true for other people. Please base your responses on how you’ve been feeling recently. Use the rating scale to find the number that best matches how you feel and circle that number. There are no right or wrong answers: we are interested in what you think and feel.”



## Figure Captions

*Figure 1.* Pathways of the Interpersonal Theory of Suicide to be tested in this study

*Figure 2.* A visual representation of the conditional effect of traumatic life events on lifetime suicidality amongst autistic and non-autistic adults at low, moderate and high frequency of traumatic life events

*Figure 3.* Simple mediation model for the influence of autistic traits on suicidality through feelings of thwarted belonging in autistic adults

*Figure 4.* Simple mediation model for the influence of autistic traits on suicidality through feelings of thwarted belonging in non-autistic adults

*Figure 5.* A visual representation of the conditional effect of thwarted belonging (M) on lifetime suicidality (Y) in autistic (W=0) and non-autistic (W=1) at low, moderate and high levels thwarted belonging

*Figure 6.* Simple mediation model for the influence of autistic traits on suicidality through feelings of burdensomeness in autistic adults

*Figure 7.* Simple mediation model for the influence of autistic traits on suicidality through feelings of burdensomeness in non-autistic adults

*Figure 8.* A visual representation of the conditional effect of burdensomeness on lifetime suicidality in autistic and non-autistic adults amongst those experiencing low, moderate and high feelings of burdensomeness

Figure 1 top




Figure 2 top






Figure 3 top



1. AQS=Autism Quotient short form, 2. INQTB=Interpersonal Needs Questionnaire 10 Thwarted belonging subscale, 3. SBQ-R=Item 1 Suicidal Behaviours Questionnaire revised

Figure 4 top



1. AQS=Autism Quotient short form, 2. INQTB=Interpersonal Needs Questionnaire 10 Thwarted belonging subscale, 3. SBQ-R=item 1 Suicidal Behaviours Questionnaire revised

Figure 5 top






Figure 6 top



1. AQS=Autism Quotient short form, 2. INQP=Interpersonal Needs Questionnaire 10 Perceived burden subscale, 3. SBQ-R=item 1 Suicidal Behaviours Questionnaire revised

Figure 7 top



Direct effect ( $c'$ ) = .004,  $p = .26$   
 Indirect effect ( $ab$ ) = .011, 95% CI [.008, .015]

1. AQS=Autism Quotient short form, 2. INQP=Interpersonal Needs Questionnaire 10 Perceived burden subscale, 3. SBQ-R=item 1 Suicidal Behaviours Questionnaire revised



Figure 8 top




Table 1. Participant characteristics and descriptive statistics

	Non-autistic group			Autistic group		
Participant characteristics	Male (n=118)	Female (n=219)	Other gender (n=2)	Male (n=124)	Female (n=206)	Other gender (n=20)
Age mean (SD)	43.68 (18.47)	40.0 (13.95)	24.5 (3.54)	44.65 (13.82)	41.13 (13.29)	33.05 (10.56)
Range (min-max)	72 (18-90)	56 (18-74)	5 (22-27)	52 (18-70)	55 (18-73)	36 (18-54)
N (%) endorsing postgraduate degree	48 (40.7%)	103 (47%)	0	48 (38.7%)	67 (32.5%)	7 (35%)
Living with support n (%)	21 (17.8%)	21 (9.6%)	1 (50%)	42 (33.9%)	43(20.9%)	5 (25%)
Employed full time n (%)	55 (46.6%)	86 (39.3%)	1 (50%)	40 (32.3%)	55 (26.7%)	5 (25%)
Diagnosed with neurodevelopmental condition endorsing n=yes (%)	14 (11.9%)	14 (5.9%)	1 (50%)	36 (29%)	62 (30.1%)	8 (40%)
Diagnosed with at least one co-occurring mental health condition n=yes (%)	26 (22%)	75 (34.2%)	1 (50%)	76 (61.3%)	148 (71.8%)	16 (80%)
Descriptive statistics mean (sd)	Male (n=118)	Female (n=219)	Other gender (n=2)	Male (n=124)	Female)(n=206)	Other gender (n=20)
Burdensomeness (INQ10 PB)	6.19 (12.49)	7.25 (5.11)	22 (8.49)	14.61 (7.65)	14.85 (8.20)	16.05 (9.05)
Thwarted belonging (INQ10 TB)	17.18 (8.06)	16.16 (7.91)	31.5 (4.95)	26.43 (6.70)	25.61 (6.93)	26.15 (5.67)
Suicidal capability (ACSSFAD)	17.23 (6.87)	15.61 (6.91)	15.5 (16.26)	17.38 (6.63)	15.77 (7.58)	18.65 (5.64)
Traumatic life events (VEQ)	17.80 (12.65)	22.37 (15.66)	51.5 (31.82)	42.67 (18.46)	45.80 (17.81)	43.75 (14.78)
Mean autistic traits (AQ-S)	63.45 (12.49)	59.64 (12.55)	76.5 (7.78)	89.62 (12.04)	91.85 (11.15)	89.75 (9.00)
Mean lifetime suicidality (SBQ-R)	2.00 (.90)	2.16 (1.01)	3 (1.41)	3.01 (.90)	3.19 (.82)	3.30 (.73)
Breakdown of past suicidal behaviour by type n (%)	Non-autistic combined male, female and other gender			Autistic combined male, female and other gender		
No past suicidal thoughts or behaviours	106 (31.7)			16 (4.7)		
Past suicidal thoughts	120 (35.9)			55 (16.1)		
Past suicide plans	73 (21.9)			140 (40.9)		
Past suicide attempt	35 (10.5)			131 (38.3)		

Table 2: Logistic regression exploring the association between thwarted belonging, perceived burden and their interaction and past reported suicidal thoughts and behaviours in autistic and non-autistic adults

	Non-autistic adults				Autistic adults			
		95%CI for Odds ratio				95% CI for Odds ratio		
	<b>b(SE)</b>	<b>Lower</b>	<b>Odds ratio</b>	<b>Upper</b>	<b>b(SE)</b>	<b>Lower</b>	<b>Odds ratio</b>	<b>Upper</b>
<b>Non-suicidal versus suicide ideation</b>								
<b>Intercept</b>	8.57 (2.23)				.41 (.90)			
<b>Belong</b>	-.39 (.092)*	.57	.68	.81*	-.06 (.06)	.85	.95	1.06
<b>Burden</b>	-3.01 (.95)*	.01	.05	.32*	-.05 (.04)	.87	.95	1.04
<b>Belong*burden</b>	.117 (.036)*	1.05	1.12	1.21*				
<b>Suicide plan versus suicide ideation</b>								
<b>Intercept</b>	-3.39 (1.82)							
<b>Belong</b>	.029 (.09)	.87	1.03	1.22	-.58 (.60)			
<b>Burden</b>	.936 (.67)	.69	2.55	9.39	.02 (.03)	.97	1.02	1.07
<b>Belong*burden</b>	-.007 (.03)	.94	.99	1.05	.05 (.03)	1.00	1.05	1.11
<b>Suicide attempt versus suicide ideation</b>								
<b>Intercept</b>	-2.65 (2.24)				.			
<b>Belong</b>	-.09 (.10)	.74	.91	1.11	.64			
<b>Burden</b>	.28 (.79)	.28	1.32	6.17	.05 (.03)*	1.0	1.05	1.11
<b>Belong*burden</b>	.04 (.03)	.97	1.03	1.10	.07 (.03)*	1.01	1.07	1.31
Note $R^2=.31$ (Cox & Snell), $.34$ (Nagelkerke). Model $X^2(9)=128.29$ , $p<.001$					Note $R^2=.10$ (Cox & Snell), $.11$ (Nagelkerke). Model $X^2(6)=36.67$ , $p<.001$ . * $p<.001$ .			

Table 3 Logistic regression exploring the association between thwarted belonging, perceived burden, suicidal capability and their interactions and past reported suicidal thoughts and behaviours in autistic and non-autistic adults

	Non-autistic people				Autistic people			
		95%CI for Odds ratio				95%CI for Odds ratio		
	b(SE)	Lower	Odds ratio	Upper	b(SE)	Lower	Odds ratio	Upper
<b>Non-suicidal versus suicide attempt</b>								
Intercept	7.91 (1.47)*				2.78 (1.14)			
Belong	-.22 (.06)**	.72	.80	.90	-.11 (.06)**	.81	.90	.99
Burden	-.61 (17)*	.39	.54	.76	-.12 (.04)**	.82	.89	.97
Suicidal capability	-.05 (.04)	.89	.95	1.02	-.04 (.04)	.89	.96	1.04
Belong*burden	.02 (.01)**	1.01	1.02	1.03				
<b>Suicide ideation versus suicide attempt</b>								
Intercept	4.23 (2.4)				3.32 (.79)			
Belong	.08 (.11)	.90	.91	1.12	-.05 (.03)**	.90	.95	1.00
Burden	-.36 (.82)	.71	.92	1.17	-.07 (.82)**	.88	.93	.99
Suicidal capability	-.08 (.03) *	.87	.87	.98	-.11 (.03)*	.86	.90	.94
Belong*burden	-.03 (.03)	.99	.99	1.01				
<b>Suicide plan versus suicide attempt</b>								
Intercept	-.12 (2.33)				1.53 (.62)			
Belong	.11 (.10)	.93	1.03	1.14	-.03 (.02)	.94	.97	1.00
Burden	.58 (72)	.87	1.06	1.31	-.02 (.02)	.94	.98	1.03
Suicidal capability	.04 (.03)	.92	.98	1.05	.03 (.02)	.94	.97	1.01
Belong*burden	-.04 (.03)	.99	.99	1.00				
	Note $R^2=.34$ (Cox & Snell), $.36$ (Nagelkerke). Model $X^2(12)=136.46$ , $p<.001$ * $p<.001$ ** $p<.05$				Note $R^2=.15$ (Cox & Snell), $.17$ (Nagelkerke). Model $X^2(9)=56.88$ , $p<.001$ * $p<.001$ ** $p<.05$			

*Table 4.* Simple linear regressions showing associations between lifetime suicidality and all predictor variables (thwarted belonging, perceived burden, capability, traumatic life events and autistic traits) in autistic and non-autistic groups

	<b>Autistic group</b>			<b>Non-autistic group</b>		
Predictor	R <sup>2</sup>	F statistic (df)	p value	R <sup>2</sup>	F statistic (df)	p value
Thwarted belong (INQTB)	.078	29.74 (1,340)	<.001	.211	90.293 (1,332)	<.001
Perceived burden (INQPB)	.063	23.86 (1,340)	<.001	.227	97.476 (1,332)	<.001
Capability (ACSS)	.037	9.77 (1,341)	<.001	.010	4.489 (1,330)	.035
Traumatic life events (VEQ)	.190	103.07 (1,335)	<.001	.349	173.856 (1,322)	<.001
Autistic traits (AQS)	.042	15.35 (1,330)	<.001	.040	14.412 (1,317)	<.001

#### Compliance with ethical standards

All procedures performed in studies involving human participants were in accordance with the ethical standards of Coventry University (ref p61841) and with the 1964 Helsinki declaration and its later amendments or comparable ethical approval standards. Informed consent was obtained from all individual participants included in the study. This article does not contain any studies with animals performed by any of the authors.

The authors declare that they have no conflict of interest.

## References

- Anestis, J. C., Finn, J. A., Gottfried, E. D., Hames, J. L., Bodell, L. P., Hagan, C. R., . . . Joiner, T. E. (2018). Burdensomeness, belongingness, and capability: Assessing the Interpersonal–Psychological theory of suicide with MMPI-2-RF scales. *Assessment, 25*(4), 415-431. doi:10.1177/1073191116652227
- Anestis, M. D., Khazem, L. R., Mohn, R. S., & Green, B. A. (2015). Testing the main hypotheses of the interpersonal–psychological theory of suicidal behavior in a large diverse sample of united states military personnel. *Comprehensive Psychiatry, 60*, 78-85.
- Bargiela, S., Steward, R., & Mandy, W. (2016). The experiences of late-diagnosed women with autism spectrum conditions: An investigation of the female autism phenotype. *Journal of Autism and Developmental Disorders, 46*(10), 3281-3294.
- Baron-Cohen, S., Leslie, A. M., & Frith, U. (1985). Does the autistic child have a “theory of mind”? *Cognition, 21*(1), 37-46.
- Beadle-Brown, J., Richardson, L., Guest, C., Malovic, A., Bradshaw, J., & Himmerich, J. (2014). Living in fear: Better outcomes for people with learning disabilities and autism.
- Bird, G., & Cook, R. (2013). Mixed emotions: The contribution of alexithymia to the emotional symptoms of autism. *Translational Psychiatry, 3*(7), e285. doi:10.1038/tp.2013.61
- Brown, S. L., Roush, J. F., Marshall, A. J., Mitchell, S. M., & Cukrowicz, K. C. (2018). A psychometric investigation of the painful and provocative events scale: Moving forward. *Archives of Suicide Research, , 1-16*.
- Brown-Lavoie, S. M., Vecili, M. A., & Weiss, J. A. (2014). Sexual knowledge and victimization in adults with autism spectrum disorders. *Journal of Autism and Developmental Disorders, 44*(9), 2185-2196.

- Bryan, C. J., Hernandez, A. M., Allison, S., & Clemans, T. (2013). Combat exposure and suicide risk in two samples of military personnel. *Journal of Clinical Psychology, 69*(1), 64-77.
- Cage, E., Di Monaco, J., & Newell, V. (2018). Experiences of autism acceptance and mental health in autistic adults. *Journal of Autism and Developmental Disorders, 48*(2), 473-484.
- Camm-Crosbie, L., Bradley, L., Shaw, R., Baron-Cohen, S., & Cassidy, S. (2018). 'People like me don't get support': Autistic adults' experiences of support and treatment for mental health difficulties, self-injury and suicidality. *Autism, , 136236131881605*.  
doi:10.1177/1362361318816053
- Campos, R. C., & Holden, R. R. (2016). Testing a theory-based model of suicidality in a community sample. *OMEGA-Journal of Death and Dying, 74*(2), 119-137.
- Cashin, A., Buckley, T., Trollor, J. N., & Lennox, N. (2016). A scoping review of what is known of the physical health of adults with autism spectrum disorder. *Journal of Intellectual Disabilities, 22*(1), 96-108. doi:10.1177/1744629516665242
- Cassidy, S., Bradley, L., Bowen, E., Wigham, S., & Rodgers, J. (2018). Measurement properties of tools used to assess depression in adults with and without autism spectrum conditions: A systematic review. *Autism Research, 11*(5), 738-754. doi:10.1002/aur.1922
- Cassidy, S., Bradley, L., Shaw, R., & Baron-Cohen, S. (2018). Risk markers for suicidality in autistic adults. doi:10.17863/CAM.33147
- Cassidy, S., Bradley, L., Shaw, R., Bowen, E., Glod, M., Baron-Cohen, S., & Rodgers, J. Measurement properties of the suicide behaviours questionnaire - revised in autistic adults. Paper presented at the *3rd Suicide and Self-Harm Early and Mid-Career Researchers' Forum*,
- Cassidy, S., Bradley, P., MRCPsych, Robinson, J., DClinPsy, Allison, C., PhD, McHugh, M., BSc, & Baron-Cohen, S., Prof. (2014). Suicidal ideation and suicide plans or attempts in adults with



- asperger's syndrome attending a specialist diagnostic clinic: A clinical cohort study. *Lancet Psychiatry, The, 1*(2), 142-147. doi:10.1016/S2215-0366(14)70248-2
- Cassidy, S., & Rodgers, J. (2017). Understanding and prevention of suicide in autism. *Lancet Psychiatry, The, 4*(6), e11. doi:10.1016/S2215-0366(17)30162-1
- Chaplin, & M. (2018). Autism spectrum disorder and hate crime. *Advances in Autism, 4*(1), 30-36. doi:10.1108/AIA-08-2017-0015
- Chen, M., Pan, T., Lan, W., Hsu, J., Huang, K., Su, T., . . . Chen, T. (2017). Risk of suicide attempts among adolescents and young adults with autism spectrum disorder: A nationwide longitudinal follow-up study. *The Journal of Clinical Psychiatry, 78*(9), e1174-e1179.
- Chen, M., Pan, T., Lan, W., Hsu, J., Huang, K., Su, T., . . . Bai, Y. (2017). Risk of suicide attempts among adolescents and young adults with autism spectrum disorder: A nationwide longitudinal follow-up study. *The Journal of Clinical Psychiatry, 78*(9), e1174-e1179. doi:10.4088/JCP.16m11100
- Christensen, H., Batterham, P. J., Mackinnon, A. J., Donker, T., & Soubelet, A. (2014). Predictors of the risk factors for suicide identified by the interpersonal-psychological theory of suicidal behaviour. *Psychiatry Research, 219*(2), 290-297. doi:10.1016/j.psychres.2014.05.029
- Chu, C., Buchman-Schmitt, J. M., Hom, M. A., Stanley, I. H., & Joiner Jr, T. E. (2016). A test of the interpersonal theory of suicide in a large sample of current firefighters. *Psychiatry Research, 240*, 26-33.
- Chu, C., Buchman-Schmitt, J. M., Stanley, I. H., Hom, M. A., Tucker, R. P., Hagan, C. R., . . . Ringer, F. B. (2017). The interpersonal theory of suicide: A systematic review and meta-analysis of a decade of cross-national research. *Psychological Bulletin, 26*
- Cole, A., Wingate, L. R., Shish, M. L., Tucker, R. P., Hollingsworth, D. W., & Victoria M. O'Keefe. (2013). Burdensomeness, depression, and suicide in a sample of american-indian college students.

*Ethnicity and Inequalities in Health and Social Care*, 6(2), 77-86. doi:10.1108/EIHSC-10-2013-0026

Corbin, J. M. (2017). *Do high burdensomeness and low belongingness interact with an individual's sense of acquired capability to increase risk of suicidal behavior within an inpatient population?*. (2016-58394-103). Retrieved from <http://search.ebscohost.com/login.aspx?direct=true&db=psych&AN=2016-58394-103&site=ehost-live>

Crane, L., Adams, F., Harper, G., Welch, J., & Pellicano, E. (2017). Know your normal: Mental health in young autistic adults. *Ambitious about Autism*,

Crane, L., Adams, F., Harper, G., Welch, J., & Pellicano, E. (2019). 'Something needs to change': Mental health experiences of young autistic adults in England. *Autism*, 23(2), 477-493. doi:10.1177/1362361318757048

Czyz, E. K., Berona, J., & King, C. A. (2015). A prospective examination of the Interpersonal-Psychological theory of suicidal behavior among psychiatric adolescent inpatients. *Suicide and Life-Threatening Behavior*, 45(2), 243-259.

Davidson, C., Wingate, L., Grant, D., Judah, M., & Mills, A. (2011). Interpersonal suicide risk and ideation: The influence of depression and social anxiety. *Journal of Social and Clinical Psychology*, 30(8), 842-855.

Dow, D., Morgan, L., Hooker, J. L., Michaels, M. S., Joiner, T. E., Woods, J., & Wetherby, A. M. (2019). Anxiety, depression, and the interpersonal theory of suicide in a community sample of adults with autism spectrum disorder. *Archives of Suicide Research*, , 1-18.

Duvekot, J., van der Ende, J., Verhulst, F. C., Slappendel, G., van Daalen, E., Maras, A., & Greaves-Lord, K. (2017). Factors influencing the probability of a diagnosis of autism spectrum disorder in girls versus boys. *Autism*, 21(6), 646-658.

Dworzynski, K., Ronald, A., Bolton, P., & Happé, F. (2012). How different are girls and boys above and below the diagnostic threshold for autism spectrum disorders? *Journal of the American Academy of Child & Adolescent Psychiatry*, 51(8), 788-797.

Evans, S. C., Boan, A. D., Bradley, C., & Carpenter, L. A. (2018). Sex/gender differences in screening for autism spectrum disorder: Implications for evidence-based assessment. *Journal of Clinical Child and Adolescent Psychology : The Official Journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division 53*, , 1-15.  
doi:10.1080/15374416.2018.1437734

Fazio, R. L., Pietz, C. A., & Denney, R. L. (2012). An estimate of the prevalence of autism spectrum disorders in an incarcerated population. *Open Access Journal of Forensic Psychology*, 4, 69-80.

Franklin, J. C., Ribeiro, J. D., Fox, K. R., Bentley, K. H., Kleiman, E. M., Huang, X., . . . Nock, M. K. (2017). Risk factors for suicidal thoughts and behaviors: A meta-analysis of 50 years of research. *Psychological Bulletin*, 143(2), 187-232. doi:10.1037/bul0000084

Fuld, S. (2018). Autism spectrum disorder: The impact of stressful and traumatic life events and implications for clinical practice. *Clinical Social Work Journal*, 46(3), 210-219.  
doi:10.1007/s10615-018-0649-6

Gillberg, C. (2002). *A guide to asperger syndrome* Cambridge University Press.

Gotham, K., Unruh, K., & Lord, C. (2015). Depression and its measurement in verbal adolescents and adults with autism spectrum disorder. *Autism*, 19(4), 491-504. doi:10.1177/1362361314536625

Green, J., Leadbitter, K., Kay, C., & Sharma, K. (2016). Autism spectrum disorder in children adopted after early care breakdown. *Journal of Autism and Developmental Disorders*, 46(4), 1392-1402.

Griffiths, S., Allison, C., Kenny, R., Holt, R., Smith, P., & Baron-Cohen, S. (2019). The vulnerability experiences quotient (VEQ): A study of vulnerability, mental health and life satisfaction in autistic adults. *Autism Research*, (000), 1-13. doi:10.1002/aur.2162

- Haertl, K., Callahan, D., Markovics, J., & Sheppard, S. S. (2013). Perspectives of adults living with autism spectrum disorder: Psychosocial and occupational implications. *Occupational Therapy in Mental Health, 29*(1), 27-41.
- Hedley, D., Uljarević, M., Foley, K., Richdale, A., & Trollor, J. (2018a). Risk and protective factors underlying depression and suicidal ideation in autism spectrum disorder. *Depression and Anxiety,*
- Hedley, D., Uljarević, M., Foley, K., Richdale, A., & Trollor, J. (2018b). Risk and protective factors underlying depression and suicidal ideation in autism spectrum disorder. *Depression and Anxiety, 35*(7), 648-657.
- Hedley, D., Uljarević, M., Wilmot, M., Richdale, A., & Dissanayake, C. (2017). Brief report: Social support, depression and suicidal ideation in adults with autism spectrum disorder. *Journal of Autism and Developmental Disorders, 47*(11), 3669-3677.
- Hedley, D., Uljarević, M., Wilmot, M., Richdale, A., & Dissanayake, C. (2018). Understanding depression and thoughts of self-harm in autism: A potential mechanism involving loneliness. *Research in Autism Spectrum Disorders, 46,* 1-7.
- Heelis, R., Graham, H., & Jackson, C. (2016). A preliminary test of the interpersonal psychological theory of suicidal behavior in young people with a first episode of psychosis. *Journal of Clinical Psychology, 72*(1), 79-87. doi:10.1002/jclp.22233
- Hickey, A., Crabtree, J., & Stott, J. (2018). 'Suddenly the first fifty years of my life made sense': Experiences of older people with autism. *Autism, 22*(3), 357-367.
- Hill, R. M., Rey, Y., Marin, C. E., Sharp, C., Green, K. L., & Pettit, J. W. (2015). Evaluating the interpersonal needs questionnaire: Comparison of the reliability, factor structure, and predictive validity across five versions. *Suicide and Life-Threatening Behavior, 45*(3), 302-314. doi:10.1111/sltb.12129

- Hirvikoski, Mittendorfer-Rutz, E., Boman, M., Larsson, H., Lichtenstein, P., & Bolte, S. (2016).  
 Premature mortality in autism spectrum disorder. *The British Journal of Psychiatry : The Journal of Mental Science*, 208(3), 232. Retrieved from  
<http://kipublications.ki.se/Default.aspx?queryparsed=id:133077842>
- Hoekstra, R. A., Vinkhuyzen, A. A., Wheelwright, S., Bartels, M., Boomsma, D. I., Baron-Cohen, S., . . . van der Sluis, S. (2011a). The construction and validation of an abridged version of the autism-spectrum quotient (AQ-short). *Journal of Autism and Developmental Disorders*, 41(5), 589-596.
- Hoekstra, R. A., Vinkhuyzen, A. A., Wheelwright, S., Bartels, M., Boomsma, D. I., Baron-Cohen, S., . . . van der Sluis, S. (2011b). The construction and validation of an abridged version of the autism-spectrum quotient (AQ-short). *Journal of Autism and Developmental Disorders*, 41(5), 589-596.
- Horton, S. E., Hughes, J. L., King, J. D., Kennard, B. D., Westers, N. J., Mayes, T. L., & Stewart, S. M. (2016). Preliminary examination of the interpersonal psychological theory of suicide in an adolescent clinical sample. *Journal of Abnormal Child Psychology*, 44(6), 1133-1144.  
 doi:10.1007/s10802-015-0109-5
- Hull, L., Petrides, K. V., Allison, C., Smith, P., Baron-Cohen, S., Lai, M., & Mandy, W. (2017). "Putting on my best normal": Social camouflaging in adults with autism spectrum conditions. *Journal of Autism and Developmental Disorders*, 47(8), 2519-2534.
- Humphrey, N. (2008). Including pupils with autistic spectrum disorders in mainstream schools. *Support for Learning*, 23(1), 41-47. doi:10.1111/j.1467-9604.2007.00367.x
- Joiner Jr, T. E., Van Orden, K. A., Witte, T. K., & Rudd, M. D. (2009a). *The interpersonal theory of suicide: Guidance for working with suicidal clients*. American Psychological Association.
- Joiner Jr, T. E., Van Orden, K. A., Witte, T. K., & Rudd, M. D. (2009b). *The interpersonal theory of suicide: Guidance for working with suicidal clients*. American Psychological Association.
- Joiner, T. E. (2005). *Why people die by suicide*. Cambridge, Mass: Harvard Univ. Press.

- Kato, K., Mikami, K., Akama, F., Yamada, K., Maehara, M., Kimoto, K., . . . Matsumoto, H. (2013). Clinical features of suicide attempts in adults with autism spectrum disorders. *General Hospital Psychiatry, 35*(1), 50-53. doi:10.1016/j.genhosppsych.2012.09.006
- Kerns, C. M., Newschaffer, C. J., & Berkowitz, S. J. (2015). Traumatic childhood events and autism spectrum disorder. *Journal of Autism and Developmental Disorders, 45*(11), 3475-3486.
- Kirby, A. V., Bakian, A. V., Zhang, Y., Bilder, D. A., Keeshin, B. R., & Coon, H. (2019). A 20-year study of suicide death in a statewide autism population. *Autism Research, 12*(4), 658-666.  
doi:10.1002/aur.2076
- Lai, J., Rhee, E., & Nicholas, D. (2017). Suicidality in autism spectrum disorder: A commentary. *Advances in Neurodevelopmental Disorders, 1*(3), 190-195. doi:10.1007/s41252-017-0018-4
- Lai, Lombardo, M. V., Auyeung, B., Chakrabarti, B., & Baron-Cohen, S. (2015). Sex/gender differences and autism: Setting the scene for future research. *Journal of the American Academy of Child & Adolescent Psychiatry, 54*(1), 11-24. doi://doi.org/10.1016/j.jaac.2014.10.003
- Luukkonen, A., Räsänen, P., Hakko, H., & Riala, K. (2009). Bullying behavior is related to suicide attempts but not to self-mutilation among psychiatric inpatient adolescents. *Psychopathology, 42*(2), 131-138.
- Maddox, B. B., Trubanova, A., & White, S. W. (2017). Untended wounds: Non-suicidal self-injury in adults with autism spectrum disorder. *Autism, 21*(4), 412-422.
- Maïano, C., Normand, C. L., Salvat, M., Moullec, G., & Aimé, A. (2016). Prevalence of school bullying among youth with autism spectrum disorders: A systematic review and Meta-Analysis. *Autism Research, 9*(6), 601-615.
- Mason, D., McConachie, H., Garland, D., Petrou, A., Rodgers, J., & Parr, J. R. (2018). Predictors of quality of life for autistic adults. *Autism Research, 11*(8), 1138-1147. doi:10.1002/aur.1965

Mazefsky, C. A., Herrington, J., Siegel, M., Scarpa, A., Maddox, B. B., Scahill, L., & White, S. W. (2013).

The role of emotion regulation in autism spectrum disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 52(7), 679-688.

McConachie, H., Mason, D., Parr, J. R., Garland, D., Wilson, C., & Rodgers, J. (2018). Enhancing the

validity of a quality of life measure for autistic people. *Journal of Autism and Developmental Disorders*, 48(5), 1596-1611.

McDonnell, C. G., Boan, A. D., Bradley, C. C., Seay, K. D., Charles, J. M., & Carpenter, L. A. (2019).

Child maltreatment in autism spectrum disorder and intellectual disability: Results from a population-based sample. *Journal of Child Psychology and Psychiatry*, 60(5), 576-584.

doi:10.1111/jcpp.12993

McManus, S., Bebbington, P., Jenkins, R., & Brugha, T. (2016). Mental health and wellbeing in

England: Adult psychiatric morbidity survey 2014. A survey carried out for NHS Digital by NatCen social research and the department of health sciences, university of Leicester.

Michaels, M. S., Chu, C., Silva, C., Schulman, B. E., & Joiner, T. (2015). Considerations regarding

online methods for suicide-related research and suicide risk assessment. *Suicide and Life-Threatening Behavior*, 45(1), 10-17. doi:10.1111/sltb.12105

Miller, A. B., Esposito-Smythers, C., & Leichtweis, R. N. (2016). A short-term, prospective test of the

interpersonal psychological theory of suicidal ideation in an adolescent clinical sample. *Suicide and Life-Threatening Behavior*, 46(3), 337-351. doi:10.1111/sltb.12196

Milton, D. (2012). On the ontological status of autism: The 'double empathy problem'. *Disability &*

*Society*, 27(6), 883-887. doi:10.1080/09687599.2012.710008

Milton, D., & Moon, L. (2012). The normalisation agenda and the psycho-emotional disablement of

autistic people. Retrieved from <https://kar.kent.ac.uk/62638>

- Müller, E., Schuler, A., & Yates, G. B. (2008). Social challenges and supports from the perspective of individuals with asperger syndrome and other autism spectrum disabilities. *Autism: The International Journal of Research and Practice*, 12(2), 173-190. doi:10.1177/1362361307086664
- Murray, A. L., Booth, T., McKenzie, K., Kuenssberg, R., & O'Donnell, M. (2014). Are autistic traits measured equivalently in individuals with and without an autism spectrum disorder? an invariance analysis of the autism spectrum quotient short form. *Journal of Autism and Developmental Disorders*, 44(1), 55-64.
- Nock, M. K., Borges, G., Bromet, E. J., Cha, C. B., Kessler, R. C., & Lee, S. (2008). Suicide and suicidal behavior. *Epidemiologic Reviews*, 30(1), 133-154.
- O'Connor, R. C. (2011). The integrated motivational-volitional model of suicidal behavior. *Crisis: The Journal of Crisis Intervention and Suicide Prevention*, 32(6), 295-298. doi:10.1027/0227-5910/a000120
- O'Connor, R. C., Cleare, S., Eschle, S., Wetherall, K., & Kirtley, O. J. (2016). The integrated Motivational-Volitional model of suicidal behavior: An update. *The International Handbook of Suicide Prevention*, , 220-240.
- Ohlsson Gotby, V., Lichtenstein, P., Långström, N., & Pettersson, E. (2018). Childhood neurodevelopmental disorders and risk of coercive sexual victimization in childhood and adolescence – a population-based prospective twin study. *Journal of Child Psychology and Psychiatry*, 59(9), 957-965. doi:10.1111/jcpp.12884
- O'Keefe, V. M., Wingate, L. R., Tucker, R. P., Rhoades-Kerswill, S., Sligh, M. L., & Davidson, C. L. (2014). Interpersonal suicide risk for american indians: Investigating thwarted belongingness and perceived burdensomeness. *Cultural Diversity and Ethnic Minority Psychology*, 20(1), 61.



- Osman, A., Bagge, C. L., Gutierrez, P. M., Konick, L. C., Kopper, B. A., & Barrios, F. X. (2001). The suicidal behaviors questionnaire-revised (SBQ-R): Validation with clinical and nonclinical samples. *Assessment, 8*(4), 443-454.
- Pelton, M. K., & Cassidy, S. A. (2017). Are autistic traits associated with suicidality? A test of the interpersonal-psychological theory of suicide in a non-clinical young adult sample. *Autism Research, 10*(11), 1891-1904. doi:10.1002/aur.1828
- Pennings, S. M., Finn, J., Houtsma, C., Green, B. A., & Anestis, M. D. (2017). Posttraumatic stress disorder symptom clusters and the interpersonal theory of suicide in a large military sample. *Suicide and Life-Threatening Behavior, 47*(5), 538-550. doi:10.1111/sltb.12317
- Rattaz, C., Dubois, A., & Baghdadli, A. (2016). How do people with autism spectrum disorders (ASD) experience pain? *An introduction to pain and its relation to nervous system disorders / editors, anna A. battaglia* (pp. 295-315) Wiley-Blackwell. Retrieved from <https://hal.univ-rennes2.fr/hal-01885009>
- Ratto, A. B., Kenworthy, L., Yerys, B. E., Bascom, J., Wieckowski, A. T., White, S. W., . . . Ollendick, T. H. (2018). What about the girls? sex-based differences in autistic traits and adaptive skills. *Journal of Autism and Developmental Disorders, 48*(5), 1698-1711.
- Ribeiro, J. D., Bodell, L. P., Hames, J. L., Hagan, C. R., & Joiner, T. E. (2013). An empirically based approach to the assessment and management of suicidal behavior. *Journal of Psychotherapy Integration, 23*(3), 207-221. doi:10.1037/a0031416
- Ribeiro, J. D., Witte, T. K., Van Orden, K. A., Selby, E. A., Gordon, K. H., Bender, T. W., & Joiner Jr, T. E. (2014). Fearlessness about death: The psychometric properties and construct validity of the revision to the acquired capability for suicide scale. *Psychological Assessment, 26*(1), 115.
- Ribeiro, J. D., Witte, T. K., Van Orden, K. A., Selby, E. A., Gordon, K. H., Bender, T. W., & Joiner, J., Thomas E. (2014a). Fearlessness about death: The psychometric properties and construct

validity of the revision to the acquired capability for suicide scale. *Psychological Assessment*, 26(1), 115-126. doi:10.1037/a0034858

Ribeiro, J. D., Witte, T. K., Van Orden, K. A., Selby, E. A., Gordon, K. H., Bender, T. W., & Joiner, T. E. J. (2014b). Fearlessness about death: The psychometric properties and construct validity of the revision to the acquired capability for suicide scale. *Psychological Assessment*, 26(1), 115-126. doi:10.1037/a0034858; 10.1037/a0034858.supp (Supplemental)

Rumball, F. (2018). A systematic review of the assessment and treatment of posttraumatic stress disorder in individuals with autism spectrum disorders. *Review Journal of Autism and Developmental Disorders*, , 1-31.

Sedgewick, F., Crane, L., Hill, V., & Pellicano, E. (2019). Friends and lovers: The relationships of autistic and neurotypical women. *Autism in Adulthood*, 1(2), 112-123. doi:10.1089/aut.2018.0028

Segers, M., & Rawana, J. (2014). What do we know about suicidality in autism spectrum disorders? A systematic review. *Autism Research*, 7(4), 507-521. doi:10.1002/aur.1375

Shelef, L., Fruchter, E., Mann, J. J., & Yacobi, A. (2014). Correlations between interpersonal and cognitive difficulties: Relationship to suicidal ideation in military suicide attempters. *European Psychiatry*, 29(8), 498-502.

Silva, C., Ribeiro, J. D., & Joiner, T. E. (2015). Mental disorders and thwarted belongingness, perceived burdensomeness, and acquired capability for suicide. *Psychiatry Research*, 226(1), 316-327. doi:10.1016/j.psychres.2015.01.008

Smith, P., & Cukrowicz, K. (2010). Capable of suicide: A functional model of the acquired capability component of the Interpersonal-Psychological theory of suicide. *Suicide and Life-Threatening Behavior*, 40(3), 266-274. doi:10.1521/suli.2010.40.3.266

- Solomon, R. L., & Corbit, J. D. (1974). An opponent-process theory of motivation: I. temporal dynamics of affect. *Psychological Review*, *81*(2), 119.
- Stone, B. (2019). 'The domino effect': Pathways in and out of homelessness for autistic adults. *Disability & Society*, , 1-6.
- Storch, E. A., Sulkowski, M. L., Nadeau, J., Lewin, A. B., Arnold, E. B., Mutch, P. J., . . . Murphy, T. K. (2013). The phenomenology and clinical correlates of suicidal thoughts and behaviors in youth with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, *43*(10), 2450-2459. doi:10.1007/s10803-013-1795-x
- Taylor, J. L., & Gotham, K. O. (2016). Cumulative life events, traumatic experiences, and psychiatric symptomatology in transition-aged youth with autism spectrum disorder. *Journal of Neurodevelopmental Disorders*, *8*(1), 28. doi:10.1186/s11689-016-9160-y
- Tint, A., Hamdani, Y., Sawyer, A., Desarkar, P., Ameis, S. H., Bardikoff, N., & Lai, M. (2018). Wellness efforts for autistic women. *Current Developmental Disorders Reports*, *5*(4), 207-216.
- Trubanova, A., Donlon, K., Kreiser, N. L., Ollendick, T. H., & White, S. W. (2014). Under-identification of ASD in females: A case series illustrating the unique presentation of ASD in young adult females. *Scandinavian Journal of Child and Adolescent Psychiatry and Psychology*, *2*(2), 66-76.
- Turcotte, P., Shea, L. L., & Mandell, D. (2018). School discipline, hospitalization, and police contact overlap among individuals with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, *48*(3), 883-891.
- Upthegrove, R., Abu-Akel, A., Chisholm, K., Lin, A., Zahid, S., Pelton, M., . . . Wood, S. J. (2018). Autism and psychosis: Clinical implications for depression and suicide. *Schizophrenia Research*, *195*, 80-85. doi:10.1016/j.schres.2017.08.028
- Van Orden, K. A., Cukrowicz, K. C., Witte, T. K., & Joiner, T. E. (2012). Thwarted belongingness and perceived burdensomeness: Construct validity and psychometric properties of the

interpersonal needs questionnaire. *Psychological Assessment*, 24(1), 197-215.

doi:10.1037/a0025358

Van Orden, K. A., Witte, T. K., Cukrowicz, K. C., Braithwaite, S. R., Selby, E. A., & Joiner Jr, T. E. (2010).

The interpersonal theory of suicide. *Psychological Review*, 117(2), 575-600.

Van Orden, K. A., Witte, T. K., Gordon, K. H., Bender, T. W., & Joiner, T. E. J. (2008). Suicidal desire

and the capability for suicide: Tests of the interpersonal-psychological theory of suicidal

behavior among adults. *Journal of Consulting and Clinical Psychology*, 76(1), 72-83.

doi:10.1037/0022-006X.76.1.72

Weiss, J. A., & Fardella, M. A. (2018). Victimization and perpetration experiences of adults with

autism. *Frontiers in Psychiatry*, 9, 203. doi:10.3389/fpsy.2018.00203/full

White, S. W., & Roberson-Nay, R. (2009). Anxiety, social deficits, and loneliness in youth with autism

spectrum disorders. *Journal of Autism & Developmental Disorders*, 39(7), 1006-1013.

doi:10.1007/s10803-009-0713-8

Williamson, S., Craig, J., & Slinger, R. (2008). Exploring the relationship between measures of self-

esteem and psychological adjustment among adolescents with asperger syndrome. *Autism :*

*The International Journal of Research and Practice*, 12(4), 391-402.

doi:10.1177/1362361308091652 [doi]

Wilson, K. G., Kowal, J., Henderson, P. R., McWilliams, L. A., & Pélouquin, K. (2013). Chronic pain and

the interpersonal theory of suicide. *Rehabilitation Psychology*, 58(1), 111.

Wolford-Clevenger, C., Brem, M. J., Elmquist, J., Florimbio, A. R., Smith, P. N., & Stuart, G. L. (2017). A

test of the interpersonal-psychological theory of suicide among arrested domestic violence

offenders. *Psychiatry Research*, 249, 195-199.

Zhang, J., Lester, D., Zhao, S., & Zhou, C. (2013). Suicidal ideation and its correlates: Testing the interpersonal theory of suicide in chinese students. *Archives of Suicide Research, 17*(3), 236-241.