

City Research Online

City, University of London Institutional Repository

Citation: Sehmi, R., Smith, N., Nguyen, A. and McManus, S. ORCID: 0000-0003-2711-0819 (2019). Trends in long-term prescribing of dependence forming medicines. London, UK: PHRC/NatCen.

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: https://openaccess.city.ac.uk/id/eprint/23651/

Link to published version:

Copyright and reuse: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

City Research Online: http://openaccess.city.ac.uk/ publications@city.ac.uk/



Trends in long-term prescribing of dependence forming medicines





Sehmi R, Nguyen A, McManus S, Smith N. (2019) *Trends in long-term prescribing of dependence forming medicines*. London: PHRC/NatCen.

This is independent research commissioned and funded by the National Institute for Health Research (NIHR) Policy Research Programme (PHPEHF50/14). The views expressed are those of the authors and not necessarily those of the NIHR, the Department of Health and Social Care or its arm's length bodies, or other Government Departments. Information about the wider programme of the Public Health Research Consortium (PHRC) is available from http://phrc.lshtm.ac.uk/. Study summary available at: www.natcen.ac.uk/dependence

Cover image by Rosie Rodgers.



NatCen Social Research 35 Northampton Square London EC1V 0AX T 020 7250 1866

www.natcen.ac.uk

A Company Limited by Guarantee Registered in England No.4392418 A Charity registered in England and Wales (1091768) and Scotland (SC038454) This project was carried out in compliance with ISO20252.

NatCen Social Research 2

Contents

| Exe | ecutive summary | 1 |
|-----|--|------|
| 1 | Introduction | 14 |
| 1.1 | Background | . 14 |
| 1.2 | New analyses in this report | . 14 |
| 1.3 | Objectives | . 15 |
| 2 | Methods | 17 |
| 2.1 | Data extraction | . 17 |
| 2.2 | Estimating length of individual prescriptions | . 17 |
| 2.3 | Estimating length of continuous prescribing periods | . 17 |
| 2.4 | Base population | . 18 |
| 2.5 | Analysis | . 18 |
| 2.6 | Exclusion criteria | . 19 |
| 2.7 | Missing data | . 19 |
| 2.8 | Over-the-counter sales | . 19 |
| 3 | Trends in DFM prescribing | 20 |
| 3.1 | Proportion of people prescribed to | . 20 |
| 3.2 | Mean average length of prescribing | . 20 |
| 3.3 | Prescriptions exceeding thresholds | . 20 |
| 3.4 | Continuous prescribing periods exceeding thresholds | . 21 |
| | 3.4.1 Benzodiazepines | |
| | 3.4.2 Z-drugs | |
| | 3.4.3 Opioids | |
| 4 | 3.4.4 GABAergic medicines | |
| 4 | Trends in gabapentin and pregabalin prescribing | |
| 4.1 | Items prescribed in England 2008-2018 | |
| 4.2 | Average length of prescriptions and continuous prescribing periods | |
| 4.3 | Prescriptions exceeding thresholds | |
| 4.4 | Continuous prescribing periods exceeding thresholds | |
| | 4.4.1 Gabapentin | |
| | 4.4.2 Pregabalin | 30 |
| 5 | Characteristics associated with DFM prescribing in | |
| exc | ess of 6 months | 31 |
| 5.1 | Variation in continuous prescribing by age-group | . 31 |

| | 5.1.1 Benzodiazepines | 31 |
|------------|--|-------|
| | 5.1.2 Z-drugs | |
| | 5.1.3 Opioids | |
| 5.2 | 5.1.4 GABAergic medicines Variation in continuous prescribing by gender | |
| 5.3 | Variation in continuous prescribing by area level deprivation | |
| 5.4 | Characteristics associated with long-term prescribing in 2014 | |
| . . | 5.4.1 Gender variations in long-term prescribing | |
| | 5.4.2 Age-group variations in long-term prescribing | 36 |
| | 5.4.3 Area level deprivation variations in long-term prescribing | |
| | 5.4.4 Regional variations in long-term prescribing | 36 |
| 6 | Characteristics associated with gabapentin and | |
| pre | gabalin prescribing in excess of 6 months | 41 |
| 6.1 | Variation in continuous prescribing by age-group | 41 |
| 6.2 | Variation in continuous prescribing by gender | |
| 6.3 | Variation in continuous prescribing by area level deprivation | |
| 6.4 | Characteristics associated with long-term prescribing in 2014 | |
| | 6.4.1 Gender variations in long-term prescribing | |
| | 6.4.3 Area level deprivation variations in long-term prescribing | |
| | 6.4.4 Regional variations in long-term prescribing | |
| 7 | Time trends in sales of codeine- and dihydrocod | eine- |
| con | taining medications | 47 |
| 8 | Conclusions | 48 |
| 9 | References | 49 |
| App | pendix A. Data tables | 51 |
| App | pendix B. Evaluation of published estimates of | |
| dep | endence on prescribed medicine | 91 |
| App | pendix C. Estimating the minimum proportion of | the |
| pop | oulation prescribed DFM long-term in 2014 | 95 |
| App | pendix D. Clinical Practice Research Datalink | 96 |
| App | pendix E. Medicines included in the analysis | 100 |
| | | |

Figures

| Figure 1: Proportion of prescriptions exceeding 30 days by DFM type 21 |
|---|
| Figure 2: Benzodiazepines: Proportion of continuous prescribing periods exceeding thresholds |
| Figure 3: Benzodiazepines: Proportion of continuous prescribing periods exceeding 6 and 12 months |
| Figure 4: Z-drugs: Proportion of continuous prescribing periods exceeding thresholds |
| Figure 5: Z-drugs: Proportion of continuous prescribing periods exceeding 6 and 12 months |
| Figure 6: Opioids: Proportion of continuous prescribing periods exceeding thresholds |
| Figure 7: Opioids: Proportion of continuous prescribing periods exceeding 6 and 12 months |
| Figure 8: GABAergic medicines: Proportion of continuous prescribing periods exceeding thresholds |
| Figure 9: GABAergic medicines: Proportion of continuous prescribing periods exceeding 6 and 12 months |
| Figure 10: Number of GABAergic medicine items dispensed in 2008-2018 26 |
| Figure 11: Average number of days of prescriptions by GABAergic medicine type |
| Figure 12: Average number of days of continuous prescribing periods by GABAergic medicine type |
| Figure 13: Proportion of prescriptions exceeding 30 days by GABAergic medicine type |
| Figure 14: Gabapentin: Proportion of continuous prescribing periods exceeding thresholds |
| Figure 15: Pregabalin: Proportion of continuous prescribing periods exceeding thresholds |
| Figure 16: Pregabalin: Proportion of continuous prescribing periods exceeding 6 and 12 months |
| Figure 17: Benzodiazepines: Proportion to exceed continuous prescribing threshold of 6 months, by age |
| Figure 18: Z-drug: Proportion to exceed continuous prescribing threshold of 6 months, by age |
| Figure 19: Opioid: Proportion to exceed continuous prescribing threshold of 6 months, by age |

| Figure 20: GABAergic medicines: Proportion to exceed continuous prescribing threshold of 6 months, by age |
|---|
| Figure 21: Proportion of continuous prescribing periods to exceed 6 months among people prescribed DFM and living in the least (Q1) and most (Q5) deprived areas |
| Figure 22: Gabapentin: Proportion of continuous prescribing periods to exceed 6 months, by age |
| Figure 23: Pregabalin: Proportion of continuous prescribing periods to exceed 6 months, by age |
| Figure 24: Proportion of continuous prescribing periods to exceed 6 months among people prescribed gabapentin living in the least (Q1) and most (Q5) deprived areas |
| Figure 25: Proportion of continuous prescribing periods exceeding 6 months among people prescribed pregabalin living in the least (Q1) and most (Q5) deprived areas |
| Figure 26: Total counts of individual codeine and dihydrocodeine-containing medicines |
| |
| |
| Tables Table 1: Characteristics of people prescribed benzodiazepines in excess of 6 months and in excess of 12 months, 2014 |
| Table 1: Characteristics of people prescribed benzodiazepines in excess of 6 |
| Table 1: Characteristics of people prescribed benzodiazepines in excess of 6 months and in excess of 12 months, 2014 |
| Table 1: Characteristics of people prescribed benzodiazepines in excess of 6 months and in excess of 12 months, 2014 |
| Table 1: Characteristics of people prescribed benzodiazepines in excess of 6 months and in excess of 12 months, 2014 |
| Table 1: Characteristics of people prescribed benzodiazepines in excess of 6 months and in excess of 12 months, 2014 |
| Table 1: Characteristics of people prescribed benzodiazepines in excess of 6 months and in excess of 12 months, 2014 |

Executive summary

Concerns about long-term prescribing

There has been concern about levels of prescribing of medicines with the potential to create dependence (BMA, 2015). Most research on trends in the prevalence of prescribing has been based on volume of medicines prescribed. One such study identified that increases in opioid prescribing have been most pronounced for stronger opioids (Curtis et al., 2018).

It is also important, however, to use patient-level data so that the proportion of the population prescribed to, their individual characteristics, and the length of time medicines are prescribed for can be examined (McManus, 2018). While short-term use of potentially dependence forming medicines (DFM) may be clinically appropriate, long-term use - especially for the management of chronic pain - is generally not (FPM, 2019).

Commissioning of the 2017 report

This need to understand trends in long-term DFM prescribing prompted the Department of Health and Social Care (DHSC) to commission an analysis of primary care data at the patient-level. An extract from the Clinical Practice Research Datalink (CPRD; Herrett et al., 2015) was examined to produce trends in the proportion of people prescribed DFM and to calculate the length of continuous prescribing periods.

Continuous prescribing periods here refers to the length of an initial individual prescription, combined with the length of any repeat prescriptions within a particular time frame. The results were published in a report: <u>Prescribing Patterns in Dependence Forming Medicines</u> (Cartagena Farias et al., 2017).¹

Key findings from the 2017 report

The 2017 analysis showed that the proportion of patients prescribed at least one of four types of DFM increased in England from 6.3% in 2000 to 9.5% in 2015. Benzodiazepines were the only type examined to see a reduction in prescribing. The other types all at least doubled in the proportion of the population prescribed to.

The report also showed that the mean number of days that DFM were continuously prescribed for increased. In particular, the length of opioid continuous prescribing periods nearly doubled, from 64 days in 2000 to a peak of 102 days in 2013/2014. In 2015, over half of prescribing periods for GABAergic medicines exceeded 30 days, as did around a third of benzodiazepine, opioid and Z drug prescribing periods.

¹ The full report includes a research of research, detailed CPRD methodology, and can be downloaded here: http://phrc.lshtm.ac.uk/papers/PHRC_014_Final_Report.pdf

The current report builds on and extends the previous analyses

- While the previous report focused on prescribing periods exceeding 30 days, this report presents trends for three thresholds: 30 days, 6 months and 12 months. This addresses a critical gap in knowledge about the extent of very long-term DFM prescribing in England.
- In the previous report people with an epilepsy or cancer diagnosis were analysed separately. In the current analysis those who had ever been diagnosed with cancer were excluded from the sample for opioid prescribing trends only, and those who had ever been diagnosed with epilepsy were excluded for trends in GABAergic medicines only.
- The current analyses additionally examine two individual medicines: gabapentin and pregabalin. It is important to consider trends in these separately as pregabalin was not introduced to the market until 2004.
- This report also includes trends in the volume of sales of over-the-counter (OTC) medicinal products containing codeine and dihydrocodeine (2015-2018).

Aims

- Estimate the proportion of individual prescriptions for each type of DFM to exceed 30 days, 6 months and 12 months each year, 2000 to 2015.
- Estimate the proportion of continuous prescribing periods for each type of DFM to exceed 30 days, 6 months and 12 months each year, 2000 to 2015.
- Compare the demographic and area-level characteristics of people prescribed DFM short- and long-term.
- Explore time trends in sales of OTC medicine products containing codeine and dihydrocodeine, 2015 to 2018.
- Review some published estimates of the number of people in England prescribed DFM long term.

The estimates in this report draw on a relatively small data extract, the descriptive trend lines presented should not be viewed as definitive or exhaustive. This analysis is one of several studies taking place on related topics. Public Health England (PHE) have conducted a fuller review of the evidence, including detailed analyses of other data sources (Tayler et al., 2019).

Methods

The current study reanalysed the same CPRD data extract. The extract consists of a sample of patients prescribed at least one of the four types of DFM between 2000 and 2015. The CPRD contains information about individual prescriptions issued by GPs (including the specific product and number of tablets prescribed) and about the patients prescribed to (such as age, sex, and area).

The length of individual prescriptions was calculated. The length of continuous prescribing periods was estimated by combining the length of the initial prescription with that of all repeat prescriptions for a medicine of the same DFM type within a relatively short period. The estimates presented should be considered conservative. A range of approaches can be used to estimate what proportion of prescribing periods exceeded particular thresholds. The approach taken in this report was to calculate the prevalence of long-term prescribing in all prescribing periods occurring in a given year. This is ideal for understanding temporal trends, maximises the relatively small sample, and is consistent with the approach taken in the previous report.

Time trends for longer-term prescribing periods (6 months or more) are presented for 2001 to 2014. This is because prescribing at the start (2000) and end (2015) of the CPRD data extract period may not fully capture the completed period of continuous prescribing. Data on over the counter (OTC) sales of codeine and dihydrocodeine-containing medicines were acquired from IRI Group Ltd.

Definitions

The following definitions are used in this report:

- **Prescriptions**: individual prescriptions not linked at the patient level.
- Continuous prescribing periods: a series of one or more prescriptions (an initial prescription combined with all repeat prescriptions) making up a continuous episode of prescribing.


Key findings

Patterns for the proportions of prescriptions and prescribing periods exceeding 30 days were broadly consistent with the previous report, they are summarised here for completeness. The more nuanced sample inclusion criteria (relating to those with a cancer or epilepsy diagnosis) had relatively minimal effects on the results.

Trends in length of individual prescriptions


A downward trend has been evident for benzodiazepines and - since 2004 - for GABAergic medicines in the proportion of individual prescriptions to exceed 30 days. The rate has been more stable for opioids and Z drugs. As expected, individual prescriptions exceeding 6 months were too rare for trends to be reliably examined with this data extract.

Proportion of individual prescriptions to exceed 30 days, by DFM type



When the two main GABAergic medicines were examined separately, we can see that the overall fall in 2004 in the proportion of prescriptions exceeding 30 days was due to the introduction of pregabalin in that year. The proportion of pregabalin prescriptions to exceed 30 days has remained about half the rate for gabapentin. Trends for GABAergic medicines as a whole are likely to reflect changes in how much of the total prescribing was for pregabalin and how much was for gabapentin widely prescribed (Montastruc et al., 2018).

Proportion of individual prescriptions to exceed 30 days, gabapentin and pregabalin




Trends in length of continuous prescribing periods

Benzodiazepines: continuous prescribing periods


While there has been a steep fall in the proportion of benzodiazepine prescribing periods to exceed 30 days (from about half of prescribing periods in 2001 to a third in 2014), reductions in the proportions of prescribing periods to exceed 6 and 12 months have been less clear.

Benzodiazepines: Proportion of continuous prescribing periods to exceed 30 days



In 2014, 12% of continuous benzodiazepine prescribing periods exceeded 6 months. Half of these (6% of prescribing periods that year) exceeded 12 months. There are indications of a slight downward trend over time. Around 13% of prescribing periods exceeded 6 months in 2001 to 2003, while since 2007 10% to 12% were prescribed at this level.


Benzodiazepine: proportion of continuous prescribing periods to exceed 6 and 12 months



Z drugs: continuous prescribing periods


In 2014, 37% of continuous Z drug prescribing periods exceeded 30 days. The rate has had a slight downward trend over time.

Z-drugs: Proportion of continuous prescribing periods to exceed 30 days



Similar to benzodiazepines, in 2014 13% of Z drug continuous prescribing periods exceeded 6 months, and 6% exceeded a year. While the rate exceeding 30 days appears to have decreased, the rate exceeding these longer thresholds may be increasing.


Z-drugs: proportion of continuous prescribing periods to exceed 6 and 12 months




Opioids: continuous prescribing periods

Similarly, while the proportion of continuous prescribing periods to exceed 30 days reduced (from 38% in 2001 to 34% in 2014) no reduction was evident at the longer thresholds. Rather, the results were indicative of a slight upward trend over time.

Opioids: Proportion of continuous prescribing periods to exceed 30 days




Opioids: proportion of continuous prescribing periods to exceed 6 and 12 months



Gabapentin: continuous prescribing periods


Since 2010, the proportion of gabapentin prescribing periods to exceed 30 days fell from 76% to 60% in 2014. to 2014 (Figure 14).

Gabapentin: Proportion of continuous prescribing periods to exceed 30 days



The rates exceeding 6 and 12 months fluctuate in the data but are not consistent with a pattern of long-time decline.


Gabapentin: proportion of continuous prescribing periods to exceed 6 and 12 months




Pregabalin: continuous prescribing periods

While the proportion of pregabalin prescribing periods to exceed 30 days remained around 60% between 2004 and 2014, the proportion to exceed longer thresholds steadily grew over this period. In 2014, 16% of continuous pregabalin prescribing periods exceed 6 months and 8% exceeded a year.

Pregabalin: Proportion of continuous prescribing thresholds to exceed 30 days



Pregabalin: proportion of continuous prescribing periods to exceed 6 and 12 months




Among those prescribed DFM: characteristics of people prescribed DFM long-term:

Among those patients prescribed DFM, the characteristics of those prescribed to longer-term can be compared with the characteristics of those prescribed to shorter-term. It is important to bear in mind, however, that these comparisons do not take account of the characteristics associated with being prescribed DFM at all.

Older people

For most types of DFM (except opioids), older people prescribed the drug were more likely than younger people to be prescribed it for more than 6 months. For example, in 2014 18.7% of those aged 81 and over who were prescribed Z drugs were prescribed to for more than six months, compared with 5.1% of those prescribed the drug who were 18 to 40-year olds.

Z drugs: proportion of continuous prescribing periods to exceed 6 months, by age



Men

In 2014, for all types of DFM, continuous prescribing periods in excess of 12 months were more likely among men prescribed the drug than among women who were prescribed to. Differences by sex, however, were not evident at other prescribing thresholds or in earlier years.


North of England

The sample was grouped into five regions: North of England, Midlands and East of England, London, South East, and South West. People prescribed DFM and living in the North were more likely than those living in other regions to be prescribed to for periods in excess of 6 and 12 months. This was the case for all classes of DFM.

Deprived neighbourhoods

Among people prescribed DFM, those living in deprived areas were generally more likely to be prescribed to for longer. This association, for example, has remained evident and pronounced among those prescribed opioids in every year of the data extract.


Opioids: proportion of continuous prescribing periods to exceed 6 months, in areas of least (Q1) and most (Q5) deprivation



Trends sales of over the counter (OTC) medicines

There was a small reduction in the volume of OTC sales of medicines containing codeine or dihydrocodeine between 2015/2016 and 2017/2018. Apart from a slight increase in sales of ibuprofen and codeine combinations, all products showed a small fall in sales volume between 2015/6 and 2017/18.

Total counts of sales of individual codeine and dihydrocodeine-containing medicines



Estimates of the prevalence of prescribed medicine dependence

We need to be cautious about drawing conclusions about dependence on prescribed medicines from primary care data, even where the analyses take account of length of prescribing. We discuss three papers that have been used to extrapolate such a prevalence and conclude that they should not be used in this way.

Limitations

The analyses presented in this report have limitations, including that they:

- Examine trends by broad class of medicine, and do not take account of medicine strength or variations in dose.
- Are based on a relatively small sample (<50,000 patients).
- Relate to prescribing practice: these data do not indicate whether medicines were dispensed, whether they were taken, and whether they were taken by the patient they were prescribed to or by someone else.
- Cannot tell us whether long-term prescribing was clinically appropriate in individual cases, nor whether patients were dependent on prescribed medicines.
- Apply strict rules on what constitutes a continuous prescribing period; estimates should be regarded as conservative.
- Do not capture recent trends (since 2015).

Conclusions and implications

Prevalence: Individual DFM prescriptions almost never exceeded 12 months. However, once repeat prescribing is accounted for a sizeable minority of continuous prescribing periods exceed this level: about one in twenty in 2014 lasted more than a year. This ranged from about 4% of prescribing periods for opioids and gabapentin, to 6% for benzodiazepines and gabapentin, and 8% for pregabalin.

Change over time: While the prevalence of continuous prescribing over 30 days tended to show signs of decline, there is little evidence of any decline at longer-term thresholds.

Pregabalin trends: Prescribing for periods in excess of 12 months increased particularly sharply for **pregabalin** since its introduction in 2004. As pregabalin and gabapentin become controlled as class C drugs (Home Office, 2018), it will be important to monitor the impact of reclassification on prescribing practice (Torjesen, 2019). Awareness of this steep increase in long-term pregabalin prescribing is relevant across sectors, and should inform service provision in primary care, specialist services (such as pain management), and psychiatry.

Variation: Those most likely to be prescribed DFM at this level were older, living in deprived neighbourhoods, and living in the North of England. In 2014 being male emerged as associated with prescribing for periods exceeding 12 months. It is possible that changing prescribing practices are not addressing the issue of very long-term prescribing in some groups of men.

Further research

- The current analyses drew on a relatively small extract of data from the CPRD. There is a need for further research with larger CPRD extracts which explore patterns in long-term prescribing for individual medicines.
- In relation to the monitoring of opioid prescribing periods, this should include taking account of oral morphine equivalence (OME) (Curtis et al., 2019).
- Further analyses could draw on other information available in the CPRD, both
 to more fully profile the extent of DFM prescribing and the characteristics of
 who is prescribed to long term. As information on diagnoses and symptom
 recording improves, this could include exploring psychiatric comorbidities in
 this population.
- Different methods can be used to calculate the length of continuous prescribing periods and to monitor change over time in these. The analyses in this report should be replicated using other approaches.
- OTC sales data can be used to examine trends at the aggregate level. Further
 work exploring individual level purchasing and use of OTC products containing
 codeine and dihydrocodeine is needed, and could be achieved with the
 inclusion of questions on general population surveys such as the Health
 Survey for England (HSE) and the Adult Psychiatric Morbidity Survey (APMS).
- Up-to-date monitoring of duration of prescribing is essential, in particular review of current trends in the long-term prescribing of pregabalin.
- Alongside monitoring prescribing data, other types of research are also needed to understand the nature and extent of dependence on prescription medicines.

1 Introduction

1.1 Background

There has been concern about levels of prescribing (Curtis et al. 2019), particularly the risks associated with long-term prescribing (McManus, 2018), of medicines with the potential to create dependence. Dependence here refers to the need to continue taking a medicine to maintain a state of feeling normal and to avoid symptoms of withdrawal.

In response to this concern, a report was commissioned through the Public Health Research Consortium (PHRC) by the National Institute for Health Research (NIHR) Policy Research Programme: *Prescribing Patterns in Dependence Forming Medicines* (Cartagena Farias *et al.*, 2017).² The National Centre for Social Research - with clinical and methodological support from the Institute of Psychiatry, Psychology and Neuroscience and University of Bristol - undertook the descriptive study of prescribing trends. This report included a rapid review of literature and should be referred to for details of previous research on the topic. It also included analysis of an extract from the Clinical Practice Research Datalink (CPRD) and methodological details about that extract. The report described:

- Trends (2000-2015) in the proportion of people prescribed medicines known to have the potential to be dependence forming, and prescribing trends for four different types of dependence forming medicines (benzodiazepines, Z-drugs, opioids and GABAergic medicines).
- Trends in the characteristics of prescribing periods: mean number of days prescribed for and the proportion exceeding 30 days.
- Trends in the characteristics of people prescribed dependence forming medicines long term: age, sex, region, health, and area-level deprivation.

The report produced new information at the patient level necessary to understand trends in prescribing practice. It confirmed falls at the patient level in the extent and length of benzodiazepine prescribing, but also found increases in the prescribing of other types of dependence forming medicines (DFM).

The findings raised additional questions at Department of Health and Social Care (DHSC), Public Health England (PHE) and for the All Party Parliamentary Group (APPG) for prescribed drug dependence, specifically around continuous prescribing periods to exceed even longer thresholds.

1.2 New analyses in this report

The following definitions are used in this report:

Prescriptions: individual prescriptions that are not linked at the patient level.

Continuous prescribing periods: a series of one or more prescriptions (an initial prescription combined with all repeat prescriptions) making up a continuous episode of prescribing.

The current study revisits the same data extract as that used in the previous report, but builds on the analysis in several new ways:

² http://phrc.lshtm.ac.uk/papers/PHRC_014_Final_Report.pdf

Different thresholds of long-term prescribing

The previous report presented the proportion of prescriptions and the proportion of continuous prescribing periods of benzodiazepines, Z-drugs, opioids or GABAergic medicines to exceed 30 days (the recommended maximum for some drug classes). The current analyses extend this by presenting the proportion of prescriptions and continuous prescribing periods to exceed three different prescribing length thresholds: 30 days, 6 months, and 12 months.

Differently defined patient population

People with a diagnosis of epilepsy or cancer were analysed separately from other people in the previous analysis. This was done to reduce the inclusion of prescribing linked to the management of epilepsy and cancer-related pain and the use of medicines in community-based palliative care. In the current analysis, however, a more nuanced approach has been taken. Those with a cancer diagnosis were excluded from the analyses of opioid prescribing trends only (and retained in the sample for other drug classes), and those with an epilepsy diagnosis were excluded from the analysis of GABAergic medicines (and retained in the sample for the other drug classes).

Separate focus on gabapentin and pregabalin

The previous report examined prescribing for four classes of drugs: benzodiazepines, Z-drugs, opioids, and GABAergic medicines. The current analyses also examine two individual medicines included in that combined GABAergic medicine grouping - gabapentin and pregabalin – separately. It is important to examine trends in these separately as pregabalin was not introduced to the market until 2004.

Inclusion of separate analysis of over-the-counter (OTC) opioid sales

The previous report drew only on analysis of primary care prescribing data, while this report includes analysis of scale and trends in sales of OTC medicinal products containing codeine and dihydrocodeine (2015-2018).

1.3 Objectives

The main objectives were to examine the extent to which DFM are prescribed long-term and to profile the characteristics of people who are prescribed to long-term. We also include analysis of the sales of over-the-counter medicines containing (dihydro)codeine and evaluate some published estimates of the extent of long-term DFM prescribing (see Appendix B).

Objective 1

Estimate the proportion of individual prescriptions for each type of DFM to exceed 30 days, 6 months and 12 months each year between 2000 and 2015 (Chapters 3 and 4).

Objective 2

Estimate the proportion of continuous prescribing periods for each type of DFM to exceed 30 days, 6 months and 12 months each year between 2000 and 2015 (Chapters 3 and 4).

Objective 3

Describe the demographic (age, sex) and area-level (deprivation and region) characteristics of people prescribed DFMs short- and longer-term each year between 2000 and 2015 (Chapters 5 and 6).

Objective 4

Produce time trends in the volume of annual sales of OTC medicine products containing codeine and dihydrocodeine, between 2015 and 2018 (Chapter 7).

Objective 5

Extrapolate from the proportion of prescribing periods in 2014/5 to estimate the proportion of the population to be prescribed each type of DFM continuously in excess of 30 days, 6 months and 12 months (Chapter 8).

Objective 6

Review some published estimates of the number of people in England who are dependent on prescribed medicines (Appendix B).

Data tables are presented in Appendix A. Methods are described in Chapter 2, with further detail about the CPRD extract in Appendix D and the medicines included in the analysis listed in Appendix E.

2 Methods

2.1 Data extraction

The Clinical Practice Research Datalink (CPRD) contains primary care data for approximately 8% of the UK population (Herrett at el., 2015). A geographically representative random sample of 49,999 individuals living in England and prescribed any benzodiazepine, Z-drug, opioid or GABAergic medicines between 2000 and 2015 was extracted for analysis.

Further detail about the CPRD extract are provided in Appendix D and a list of all medicines included in each of DFM groups examined is provided in Appendix E. A full description of the structure, content and generalisability of the CPRD dataset can also be found in the previous report (Cartagena Farias et al., 2017), including further description of the data extract used for analysis.

2.2 Estimating length of individual prescriptions

The length of an individual prescription was defined as the number of treatment days a drug is prescribed for a specific therapy event on a specific occasion. Each consultation provided information regarding the total dose and numeric daily dose of a particular drug. To calculate the length of each individual prescription, the total quantity of drug prescribed was divided by the numeric daily dose for each given therapy (e.g. 90 tablets / 3 tablets per day = 30 days duration). Variations in dose levels were not accounted for.

2.3 Estimating length of continuous prescribing periods

The number of days for which a particular class of drug was continuously prescribed to a person for was estimated by adding the length of an initial prescription to the length of any repeat prescriptions. A repeat prescription was identified when the same drug (or another drug in the same class) was prescribed to the same person within a period. To identify a series of prescriptions that could reliably be classified as repeat prescriptions, the following rules were applied:

- 1. The number of days between each consultation is less than or equal to 35 days, or the length of the previous prescription is equal to, or less than, the number of the days between consultations (plus 5 days). This ensured that a repeat prescription was identified, as opposed to multiple prescriptions over a longer period.
- 2. In some instances, consultation identifiers varied despite being held on the same day; it was assumed that all events that occurred on the same day were part of the same consultation.
- 3. For analyses of the four major classes of drugs (benzodiazepines, Z-drugs, opioids and GABAergic medicines): if a patient switched from one type of benzodiazepine to another type of benzodiazepine, this still counted as a continuous prescribing period; the same applied to other classes of drug.
- 4. If two different drugs belonging to the same class of drug were prescribed in the same consultation (e.g. diazepam and lorazepam), the longer of the two derived

prescription lengths was used. Similarly, if two different prescription lengths were prescribed in the same consultation then the longer length was used.

5. The first consultation date when a drug was prescribed was noted as the start date of a continuous prescribing period; this ensured that repeat prescriptions following from one year to the next were counted from the original start date.

When reporting on prescribing periods that span two calendar years, duration length is based on when the prescribing period ends as opposed to curtailing periods at the end of a calendar year. As a result, periods spanning more than one calendar year are effectively reported twice, in both consecutive years. Likewise, periods spanning three years are captured in the data for each of those years. For example, if someone was prescribed opioids from May 2001 to August 2002, that would be counted as a continuous prescribing period exceeding 12 months in both 2001 and 2002. This phenomenon will affect longer prescribing periods more than shorter term prescribing. Because prevalence per year is reported, the validity of rates of longer-term prescribing is maintained as the relevant prescribing period did indeed occur in both consecutive years.

It is important to note that the CPRD data extract is cut off at the end of 2015, so it is likely that a proportion of people had not yet completed their continuous prescribing period. This is an issue when estimating the proportion of prescribing periods exceeding the longer thresholds, such as 6 or 12 months. Because of this, estimates for the final year of data (2015) have not been fully presented, and are given in grey text in the tables. Similarly, a significant proportion of people would not have their full continuous prescribing period captured at the start of the study period. For instance, we would expect low estimates of the proportion of people with a continuous prescribing period of 12 months within the first 12 months of the data as many people would have received their initial prescription before the start date of the extract. This report shows the rates for 2001 to 2014 for continuous prescribing periods exceeding 6 months and 12 months due to concern about the reliability of longer-term estimates for 2000 and 2015; rates for prescriptions, however, include estimates for 2000 as truncation should not affect this estimate.

We use the following conventions in all results tables, '-' denotes no observations (zero value), and '0' denotes non-zero values of less than 0.05% and are thus rounded to zero.

2.4 Base population

The unit of analysis in the tables varied according to the subject being examined and is referred to in the text. Analyses examining trends over time used either prescriptions or continuous prescribing periods, where appropriate, whereas analyses of the characteristics of long-term prescribing in 2014 refers to people. For the latter analyses, prescribing patterns were investigated by area level deprivation of the GP practice, region and by the gender and age of the person prescribed to. When examining the characteristics of people prescribed to for periods in excess of six or 12 months, some age bands were combined due to small base sizes.

2.5 Analysis

The mean length of prescriptions and continuous prescribing periods for each class of DFM and for two types of GABAergic medicines – pregabalin and gabapentin – were derived for each year of the data. The results presenting trends over time in prescriptions and continuous prescribing periods are descriptive, unadjusted, and not tested for statistical significance.

Further analysis estimated the proportion of prescriptions and continuous prescribing periods for each class of DFM which were in excess of 30 days, 6 months, and 12 months. These thresholds are not mutually exclusive, meaning that a continuous prescribing period in excess of 12 months was also counted as a prescribing period in excess of 30 days and in excess of 6 months.

The unit of analysis for the trends described above was prescribing events. These were also used to extrapolate a crude estimate of the proportion of the total population in 2014 to be prescribed each DFM long-term. For example, we know that 5.2% of the total CPRD population were prescribed opioids in 2014. We also know that 33.7% of continuous opioid prescribing periods in 2014 exceeded 30 days. Extrapolating from this, we can estimate that approximately 33.7% of the people prescribed opioids were prescribed long-term (33.7% of 5.2% = 1.8% of the total population). The key assumption here is that there was little association between the number of continuous prescribing periods someone experienced in a year and the length of each of those continuous prescribing periods. The mean number of prescribing periods per person prescribed to was less than 2 for all types of DFM. Because of the uncertainty and scope for systematic undercounting, we recommend that these proportions are viewed as minimum estimates and indicative only.³

2.6 Exclusion criteria

Exclusion criteria differed from those applied in the previous report. Individuals with an epilepsy diagnosis (at any point in the data extract) were excluded from the analysis of GABAergic medicines (including pregabalin- and gabapentin-specific analyses). This was because some GABAergic medicines have been approved for the long-term treatment of epilepsy. People with a cancer diagnosis (at any point in the data extract) were excluded from the analysis of opioid prescribing as the use of opioids in the treatment of severe cancer-related pain is considered standard procedure, although their suitability for treating chronic, non-cancer related pain has been questioned. These exclusion criteria were not applied to the analysis of benzodiazepines and Z-drugs.

2.7 Missing data

Overall, variation in the amount of missing data was not pronounced by gender, age or practice (see previous report). However, bias resulting from missing data could not be ruled out for other characteristics for which we could not make assessments. Therefore, missing data need to be acknowledged as a potential limitation which could have biased the results.

2.8 Over-the-counter sales

Data on the sales of over-the-counter (OTC) medicines containing codeine or dihydrocodeine between February 2015 and February 2018 were acquired from IRI Worldwide. Total counts of individual tablets were derived for each 12-month period from week ending February 28th, 2015. Counts were calculated from the sales volume multiplied by the pack size.

³ These extrapolations are approximate and should be regarded as likely underestimate.

3 Trends in DFM prescribing

This chapter presents trends in the prescribing of four classes of DFM: benzodiazepines, Z-drugs, opioids and GABAergic medicines. Trends for each of gabapentin and pregabalin (subclasses of GABAergic medicine) are presented separately in Chapter 4.

3.1 Proportion of people prescribed to

In the previous report analyses of these data showed that the overall proportion of people prescribed DFM at any point in the year increased from 6.3% in 2000 to a peak of 9.5% in 2012. The pattern varied by type of DFM. Opioids, Z-drugs and GABAergic medicines saw pronounced increases in the proportion of people for whom they were prescribed. The proportion prescribed opioids and Z-drugs doubled between 2000 and 2012. GABAergic medicine prescribing increased even more steeply, from 0.2% of people in 2000 (following approval in October 2000) to 2.1% in 2015. This context should be borne in mind when interpreting trends in the length of time medicines were prescribed for.

3.2 Mean average length of prescribing

The mean average number of days of individual prescriptions and of continuous prescribing periods are presented for each type of DFM in Appendices 1 and 2. These were produced using different exclusion criteria from the previous report (see Section 2.5). The revised mean lengths are similar to the results previously reported. GABAergic medicine prescriptions tended to be issued for longer than the other types of DFM; benzodiazepines, Z-drug and opioid prescriptions averaged a mean length of between 24-25 days in 2014 compared with 31 days for GABAergic medicines. The average length of prescriptions for GABAergic medicines remained stable between 2000 and 2015, but average prescription lengths decreased for all other drugs; decreases were particularly marked for opioids (36 days in 2000 to 24 days in 2015).

Continuous periods of DFM prescribing in 2014 averaged between 103 and 123 days. The average prescribing period for opioids increased from 64 days in 2000, to a peak of 103 days in 2014. The continuous prescribing periods for other DFMs remained more stable over time.

3.3 Prescriptions exceeding thresholds

The length of prescriptions was examined in relation to three thresholds. Rates were calculated for the proportion of prescriptions for benzodiazepines, Z-drugs, opioids and GABAergic medicines (including separately for gabapentin and for pregabalin, presented in the next chapter) to exceed:

- 30 days
- 6 months
- 12 months

Consistent with the trends presented here, the previous report (Cartagena Farias et al., 2017) also showed that the proportion of prescriptions for benzodiazepine and GABAergic medicines to exceed 30 days fluctuated but with general downward trends over time (Figure 1). For opioids, the proportion of prescriptions exceeding 30 days was around 15% from 2000-2007, and between 16-18% from 2007-2015. For Z-drugs, the proportion of prescriptions exceeding 30 days fluctuated between 10-12% between 2000 and 2010 and remained below 10% 2011-2015.

Figure 1: Proportion of prescriptions exceeding 30 days by DFM type

Source: CPRD (see Appendix 3)

The proportion of individual prescriptions to exceed 6 months and 12 months by DFM type was examined, but as expected, the proportion at each year was close to zero for these longer thresholds and no trend in prescribing was detectable.

3.4 Continuous prescribing periods exceeding thresholds


Cartagena Farias and colleagues (2017) described the proportion of continuous prescribing periods to exceed 30 days. The current re-analysis of the same data reproduces these trends using the updated set of exclusion criteria, as well as also presenting trends for the proportion of continuous prescribing periods to exceed 6 and 12 months.

Figures 2-5 below show the proportion of continuous prescribing periods for benzodiazepines, Z-drugs, opioids and GABAergic medicines to exceed 30 days, 6 months and 12 months.

3.4.1 Benzodiazepines

The proportion of continuous benzodiazepine prescribing periods to exceed 30 days fell from about half in 2001 to about a third in 2011-2014 (Figure 2). The proportion to exceed six months was around 12%-13% between 2001 and 2007, and 10%-11% between 2008 and 2014 (Figure 3). Prescribing periods exceeding 12 months followed a similar pattern: the rate was about 6% between 2001 and 2007, and about 5% between 2008 and 2014.

Figure 2: Benzodiazepines: Proportion of continuous prescribing periods exceeding thresholds



(Source: Appendix 4 to Appendix 6)

Figure 3: Benzodiazepines: Proportion of continuous prescribing periods exceeding 6 and 12 months



3.4.2 Z-drugs

Less than half of continuous prescribing periods of Z-drugs were in excess of 30 days between 2001 and 2012 (varying between 41% and 44%), and after 2012 the prescribing rate varied between 37%-40% (Figure 4). The proportion to exceed 6 months remained mostly below about 12% between 2001 and 2010, and above 12% since 2011 (Figure 5). Prescribing periods exceeding 12 months followed a similar pattern: about 4% to 5% between 2001 and 2010, since 2011 the rate was 6%.

50 45 40 35 30 Over 30 days % 25 Over 6 months 20 Over 12 months 15 10 5 0 2010 2006

Figure 4: Z-drugs: Proportion of continuous prescribing periods exceeding thresholds

(Source: Appendix 4 to Appendix 6)

Figure 5: Z-drugs: Proportion of continuous prescribing periods exceeding 6 and 12 months



3.4.3 Opioids

The proportion of continuous opioid prescribing periods to exceed 30 days diminished from 2012 onwards, following a consistent rate of around 37%-39% between 2001-2011 (Figure 6). The proportion to exceed 6 months remained around 6%-7% between 2001 and 2008, and 7%-8% between 2009 and 2014 (Figure 8). Prescribing periods exceeding 12 months followed a similar pattern: about 2%-3% between 2001 and 2008, and 3%-4% between 2009 and 2014.

Figure 6: Opioids: Proportion of continuous prescribing periods exceeding thresholds

(Source: Appendix 4 to Appendix 6)





Figure 7: Opioids: Proportion of continuous prescribing periods exceeding 6 and 12 months

3.4.4 GABAergic medicines


The proportion of continuous GABAergic medicine prescribing periods to exceed 30 days increased from 58% in 2001 to 71% in 2005 (Figure 8); this covers the period in which pregabalin was first introduced. Prescribing rates diminished from 69% in 2010 to 58% in 2014. The proportion to exceed 6 months remained around 10%-12% between 2001 and 2005, and 13%-14% between 2006 and 2014 (Figure 9). Prescribing periods exceeding 12 months diminished slightly from 5% to 3% between 2001 and 2005, and then increased back up to 5% in 2014.

Figure 8: GABAergic medicines: Proportion of continuous prescribing periods exceeding thresholds



(Source: Appendix 4 to Appendix 6)

Figure 9: GABAergic medicines: Proportion of continuous prescribing periods exceeding 6 and 12 months



4 Trends in gabapentin and pregabalin prescribing

As well as producing trends for the combined category of GABAergic medicines, we also examined two of the main individual GABAergic medicines separately: gabapentin and pregabalin.

4.1 Items prescribed in England 2008-2018

Figure 10 draws on recently published NHS Digital data showing the volume of prescriptions dispensed in England between 2008 and 2018 (NHSD 2019). These figures include prescribing for all indications, while our analysis of GABAergic medicines excluded prescribing to those with an epilepsy diagnosis. However, the NHS Digital figures do provide useful context on the increasing volume of overall GABAergic prescribing and demonstrate that most prescribing is for gabapentin or pregabalin.





Figure 10: Number of GABAergic medicine items dispensed in 2008-2018

4.2 Average length of prescriptions and continuous prescribing periods

The average number of days individual gabapentin prescription were issued for remained stable over time at around 36 days (Figure 11). Average pregabalin prescriptions increased in length from 2004 (when pregabalin was first approved for use in the European Union) to 2008; from 2008 onwards, the number of days covered by individual pregabalin prescriptions remained stable at a mean of around 35 days.


Figure 11: Average number of days of prescriptions by GABAergic medicine type



Source: CPRD (see Appendix 7)

The average number of days that gabapentin was continuously prescribed for remained relatively stable, but with some fluctuations (Figure 12). The average number of days that pregabalin was prescribed for increased steeply over this time, almost doubling between 2004 (67 days) and 2014 (127 days).

Figure 12: Average number of days of continuous prescribing periods by GABAergic medicine type




Source: CPRD (see Appendix 8)

4.3 Prescriptions exceeding thresholds

Figure 13 shows the proportions of prescriptions exceeding 30 days for people prescribed gabapentin and pregabalin. The proportion of gabapentin prescriptions to exceed 30 days increased over time until around 2011, at which time the proportion declined from 65% to 52% in 2014. The proportion of pregabalin prescriptions to exceed 30 days has also broadly increased over time, from about 13% when the drug was first introduced to about 30% in 2013 and 2014.

Figure 13: Proportion of prescriptions exceeding 30 days by GABAergic medicine type



Source: CPRD (see Appendix 9)


4.4 Continuous prescribing periods exceeding thresholds

The figures below show the proportion of continuous prescribing periods for gabapentin and pregabalin to exceed 30 days, 6 months and 12 months.

4.4.1 Gabapentin

The proportion of continuous prescribing periods exceeding 30 days remained stable up to 2010, when rates fell sharply for the next 4 years. Rates of prescribing for thresholds exceeding 6 months and 12 months remained relatively stable from 2001 to 2014 (Figure 14).

Figure 14: Gabapentin: Proportion of continuous prescribing periods exceeding thresholds




(Source: CPRD Appendix 10 to Appendix 12)

4.4.2 Pregabalin


The proportion of continuous prescribing periods exceeding 30 days fluctuated from 2004 to 2014, but there was no particular upward or downward trend over this time (Figure 15). Conversely, rates of prescribing exceeding 6 months and 12 months steadily increased over time (Figure 16)

Figure 15: Pregabalin: Proportion of continuous prescribing periods exceeding thresholds



Source: CPRD (see Appendix 10 to Appendix 12)

Figure 16: Pregabalin: Proportion of continuous prescribing periods exceeding 6 and 12 months



Source: CPRD (see Appendix 10 to Appendix 12)

5 Characteristics associated with DFM prescribing in excess of 6 months


5.1 Variation in continuous prescribing by agegroup

Among patients prescribed DFM, the characteristics of those prescribed to continuously for six months or more could be compared with the characteristics of those prescribed to for less than six months. It is important to bear in mind, however, that these comparisons do not take account of the characteristics associated with being prescribed DFM at all. Too few patients in the sample were prescribed to for more than 12 months for these results to be robust. They are therefore not discussed in the text but are presented in the tables in Appendix A for reference.

5.1.1 Benzodiazepines

Older people prescribed benzodiazepines were more likely to be prescribed to in excess of 6 months than younger people prescribed the drug. This association with age has remained consistent over time.

Figure 17: Benzodiazepines: Proportion to exceed continuous prescribing threshold of 6 months, by age




Base: People prescribed benzodiazepines each year. Source: CPRD (see Appendix 14)

5.1.2 Z-drugs

Similar to benzodiazepine prescribing, older people prescribed Z-drugs were more likely than younger people prescribed the drug to be prescribed for periods exceeding 6 months (and 12 months, see Appendix 18). This pattern has remained consistent over time.

Figure 18: Z-drug: Proportion to exceed continuous prescribing threshold of 6 months, by age




Base: People prescribed Z-drugs each year. Source: CPRD (see Appendix 17)

5.1.3 Opioids

Among people prescribed opioids, the association between age and being prescribed to longer-term (6+ months) was less clear cut. While younger people prescribed the drug were generally less likely to be prescribed it longer term, the rates for other age groups were similar.

Figure 19: Opioid: Proportion to exceed continuous prescribing threshold of 6 months, by age




Base: People prescribed opioids each year. Source: CPRD (see Appendix 20)

5.1.4 GABAergic medicines

Age related differences in prescribing of GABAergic medicines in excess of 6 months have changed over the study period. The pattern of prescribing is variable from 2001 to 2007, but an age association emerged from 2008 with longer-term prescribing becoming more common in older than younger people.

Figure 20: GABAergic medicines: Proportion to exceed continuous prescribing threshold of 6 months, by age



Base: People prescribed GABAergic medicines each year. Source: CPRD (see Appendix 23)

5.2 Variation in continuous prescribing by gender


The previous report found little in terms of an association with gender for any DFM prescribing in excess of 30 days. The results from this report largely replicate these findings for prescribing exceeding 6 months and 12 months. Notable exceptions to this overall pattern include evidence of gender differences in prescribing exceeding 6 months up to 2008/2009 for benzodiazepines and GABAergic medicines, and men were more likely to be prescribed to in 2014 for periods exceeding 12 months across all DFMs (see Appendix 25 to Appendix 27).

5.3 Variation in continuous prescribing by area level deprivation


Variations in continuous prescribing by area level deprivation was examined using measures of IMD at the lower level super output area (LSOA) of the patient's home address⁴. Figure 21 shows that people prescribed DFM who lived in the most deprived areas were more likely to be prescribed benzodiazepines, Z-drugs and opioids for 6 months or more than those in the least deprived areas; the variation in long term prescribing periods exceeding 6 months for GABAergic medicines by deprivation level was more mixed. Similarly, people prescribed DFM who lived in the most deprived areas were more likely to be prescribed benzodiazepines, Z-drugs and opioids for 12 months or more compared to those in the least deprived areas, while the pattern for GABAergic medicines by area deprivation level was again more mixed.

Figure 21: Proportion of continuous prescribing periods to exceed 6 months among people prescribed DFM and living in the least (Q1) and most (Q5) deprived areas







⁴ Practice level IMD was available in CPRD but we chose to use patient level IMD as we are interested in patient characteristics as opposed to those of the GP practice.



Opioids



GABAergic medicines



Source: CPRD (see Appendix 29, Appendix 32, Appendix 35, Appendix 38) Based on people living in the fifth of areas with the most deprivation (Q5) and the fifth of areas with the least deprivation (Q1) according to the 2007 Index of Multiple Deprivation

5.4 Characteristics associated with long-term prescribing in 2014

CPRD data for 2014 were examined to compare the circumstances of people prescribed DFMs short term (that is, for continuous periods of 30 days or less) and long-term (more than 6 and 12 months). See Table 1 to Table 4. Comparisons by characteristics were made separately for each threshold.

5.4.1 Gender variations in long-term prescribing

Among people prescribed DFM, men were more likely than women to be prescribed to for more than 12 months; this was the case for every type of DFM. For instance, among women prescribed benzodiazepines, 5.1% were prescribed to for periods exceeding 12 months, compared to 7.3% of men; gender differences were not evident for relatively shorter long-term prescribing periods of over 6 months.

5.4.2 Age-group variations in long-term prescribing

A continuous measure of age was used to examine whether there were age-related variations in long-term prescribing in excess of 6 months and 12 months, so no reference category was assigned. Older people were more likely than younger people to be prescribed for longer than 6 months.⁵; this was the case for all types of DFM apart from opioids. Apart from Z-drugs, older people were no more likely than younger people to be prescribed DFMs for longer than 12 months.

5.4.3 Area level deprivation variations in long-term prescribing

Comparisons in the rates of long-term prescribing among people living in the most (V) compared to least (I) deprived areas was examined. People prescribed DFM and living in the most deprived areas were more likely to prescribed to for periods in excess of 6 months. For example, among people prescribed benzodiazepines and living in the least deprived neighbourhoods, 8.8% were prescribed to for longer than 6 months, while among those living in the most deprived neighbourhoods 19.4% were prescribed to at this level. For GABAergic medicines, this was evident only in prescribing in excess of 12 months.

5.4.4 Regional variations in long-term prescribing

The ten available regions in the CPRD were grouped in to five larger regions, North of England, Midlands and East of England, London, South East and South West, in order to maximise power. North of England was chosen as the reference category as it was the most populated region in this sample.

People prescribed DFM and living in the North of England were the most likely to be prescribed to long-term for periods in excess of 6 and 12 months. For instance, 15.5% of those living in North of England were prescribed to for longer than 6 months

⁵ Mean age (years) for short-term periods lasting under 30 days: 59.3 (benzodiazepines), 58.9 (Z-drugs), 60.6 (opioids) and 58.2 (GABAergic medicines).

compared to 11.7% living in the South East. This was the case for all classes of DFM, although the pattern was less pronounced for GABAergic medicines.

Please refer to Appendix 40 to 44 for related analyses. These present the characteristics associated with being prescribed each type of DFM for up to 3 months, between 3 to 6 months, between 6 to 12 months and for more than 12 months.

Table 1: Characteristics of people prescribed benzodiazepines in excess of 6 months and in excess of 12 months, 2014

| Benzodiazepines | | | | |
|----------------------|-----------|---------|--|--|
| | Long-term | | | |
| | Over 6 | Over 12 | | |
| | months | months | | |
| Mean age (years) | 64.0** | 62.3 | | |
| IMD quintile (%) | | | | |
| I (least deprived) § | 8.8 | 3.3 | | |
| II | 9.6 | 3.5 | | |
| III | 12.0* | 4.9 | | |
| IV | 11.5* | 7.5*** | | |
| V (most deprived) | 19.4*** | 10.6*** | | |
| Gender (%) | | | | |
| Male § | 12.9 | 7.3 | | |
| Female | 11.7 | 5.1*** | | |
| Region (%) | | | | |
| North of England § | 15.5 | 9.7 | | |
| Midlands and East of | 11.3*** | 4.8*** | | |
| England | | | | |
| London | 9.9*** | 7.3 | | |
| South East | 11.7** | 3.7*** | | |
| South West | 10.1*** | 5.0*** | | |

^{*} P \leq 0.05, ** P \leq 0.01, *** P \leq 0.001, § - reference category Source: CPRD

Table 2: Characteristics of people prescribed Z-drugs in excess of 6 months and in excess of 12 months, 2014

| Z-drugs | | | |
|----------------------|-----------|---------|--|
| | Long-term | | |
| | Over 6 | Over 12 | |
| | months | months | |
| Mean age (years) | 67.9*** | 67.5*** | |
| IMD quintile (%) | | | |
| I (least deprived) § | 13.0 | 6.2 | |
| II | 13.1 | 6.0 | |
| III | 9.6* | 5.4 | |
| IV | 11.8 | 4.8 | |
| V (most deprived | 17.6* | 8.0 | |
| Gender (%) | | | |
| Male § | 13.9 | 7.7 | |
| Female | 12.2 | 5.1** | |
| Region (%) | | | |
| North of England § | 16.7 | 9.5 | |
| Midlands and East of | | | |
| England | 11.4*** | 4.7*** | |
| London | 13.5 | 5.6* | |
| South East | 11.6*** | 5.4*** | |
| South West | 11.7* | 4.2*** | |

^{*} P \leq 0.05, ** P \leq 0.01, *** P \leq 0.001, § - reference category Source: CPRD

Table 3: Characteristics of people prescribed opioids in excess of 6 months and in excess of 12 months, 2014

| Opioids | | | |
|----------------------|-----------|---------|--|
| | Long-term | | |
| | Over 6 | Over 12 | |
| | months | months | |
| Mean age (years) | 63.3 | 63.1 | |
| IMD quintile (%) | | | |
| I (least deprived) § | 4.6 | 0.9 | |
| II | 7.4*** | 4.6*** | |
| III | 7.1*** | 3.7*** | |
| IV | 8.1*** | 4.2*** | |
| V (most deprived | 8.3*** | 3.8*** | |
| Gender (%) | | | |
| Male § | 7.3 | 4.1 | |
| Female | 7.3 | 3.3** | |
| Region (%) | | | |
| North of England § | 9.2 | 4.7 | |
| Midlands and East of | | | |
| England | 6.5*** | 3.6* | |
| London | 5.3*** | 2.2*** | |
| South East | 6.9*** | 3.3*** | |
| South West | 5.3*** | 2.0*** | |

^{*} P \leq 0.05, ** P \leq 0.01, *** P \leq 0.001, § - reference category *Source: CPRD*

Table 4: Characteristics of people prescribed GABAergic medicines in excess of 6 months and in excess of 12 months, 2014

| GABAergic medicines | | | |
|----------------------|-----------|---------|--|
| | Long-term | | |
| | Over 6 | Over 12 | |
| | months | months | |
| Mean age (years) | 63.2*** | 61.1 | |
| IMD quintile (%) | | | |
| I (least deprived) § | 14.3 | 3.9 | |
| II | 12.0 | 5.6* | |
| III | 13.0 | 3.3 | |
| IV | 16.2 | 6.0* | |
| V (most deprived | 14.5 | 6.0** | |
| Gender (%) | | | |
| Male § | 14.3 | 6.9 | |
| Female | 13.8 | 3.9*** | |
| Region (%) | | | |
| North of England § | 15.2 | 4.8 | |
| Midlands and East of | | | |
| England | 15.6 | 7.4*** | |
| London | 12.1 | 2.2* | |
| South East | 13.6 | 4.4 | |
| South West | 9.0*** | 4.8 | |

^{*} P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, § - reference category Source: CPRD

6 Characteristics associated with gabapentin and pregabalin prescribing in excess of 6 months


6.1 Variation in continuous prescribing by agegroup

The previous chapter examined the characteristics of people prescribed benzodiazepines, opioids, Z-drugs and GABAergic medicines long-term; the focus in this chapter is on the characteristics of people prescribed gabapentin and pregabalin. One of the reasons why these drugs are looked at separately is because pregabalin only entered the market in 2004. Again, due to the small number of people in the sample prescribed to for more than 12 months, the focus here is on those prescribed to for more than 6 months. Even at this threshold, numbers are relatively low, and results should be treated with caution. The data tables for 30 days, 6 months and 12 months can be found in Appendix A.

6.1.1 Gabapentin

From 2007 older people prescribed gabapentin were more likely than younger people prescribed the drug to receive it long-term (Figure 22, and Appendix 45). Prior to 2007, associations with age were unclear.

Figure 22: Gabapentin: Proportion of continuous prescribing periods to exceed 6 months, by age




Source: CPRD (see Appendix 48)

6.1.2 Pregabalin

The youngest age group of 18-40 had the lowest rate of pregabalin prescribed for longer than 6 months from 2007 onwards (apart from a spike in prescriptions at 2012); there were no age-related differences in long-term prescribing prior to 2007.

Figure 23: Pregabalin: Proportion of continuous prescribing periods to exceed 6 months, by age



Source: CPRD (see Appendix 51)

6.2 Variation in continuous prescribing by gender

The results did not show any clear gender differences over time. However, overall, women prescribed gabapentin and pregabalin for periods exceeding 6 months tended to have a lower prescription rate for some periods of time. For example, rates between 2001-2006 were lower for women prescribed gabapentin, and rates between 2006-2010 were lower for women prescribed pregabalin.

6.3 Variation in continuous prescribing by area level deprivation

Rates of continuous gabapentin prescribing exceeding 6 months varied over time among people living in the most compared to least deprived areas (Figure 24); a similar trend is observed for prescribing periods exceeding 12 months.

Overall, people prescribed pregabalin living in the most compared to least deprived areas had a higher rate for continuous prescribing periods exceeding 6 months (Figure 25); however, rates of prescribing for people living in the least deprived areas spiked in 2008-2009 and were higher than those in the most deprived areas. There was no clear trend for prescribing periods exceeding 12 months, as rates between the most and least deprived areas varies over time.

Figure 24: Proportion of continuous prescribing periods to exceed 6 months among people prescribed gabapentin living in the least (Q1) and most (Q5) deprived areas





Figure 25: Proportion of continuous prescribing periods exceeding 6 months among people prescribed pregabalin living in the least (Q1) and most (Q5) deprived areas



Source: CPRD (see Appendix 57 and Appendix 60)

6.4 Characteristics associated with long-term prescribing in 2014

CPRD data for 2014 were examined to compare the circumstances of people prescribed gabapentin and pregabalin short- and long-term (see Tables 9 to 12 below).

6.4.1 Gender variations in long-term prescribing

There were no gender differences in prescribing rates for long-term prescriptions of gabapentin, but women receiving pregabalin were less likely to be prescribed to for more than both 6 months and 12 months. For instance, 4.9% of women were prescribed to for longer than 12 months compared to 13% of men.

6.4.2 Age-group variations in long-term prescribing

Long-term prescribing was more common in older than younger people prescribed gabapentin (for periods exceeding 6 months only), but there were no age-related differences in long-term pregabalin prescribing⁶.

6.4.3 Area level deprivation variations in long-term prescribing

People living in the most compared to least deprived areas were no more likely to be prescribed gabapentin or pregabalin long-term.

6.4.4 Regional variations in long-term prescribing

Overall, there were no regional differences in long-term prescribing (over 6 and 12 months) of gabapentin and pregabalin relative to the North of England (Tables 9 and 10). The one exception was a lower prescribing rate of pregabalin for continuous prescribing periods exceeding 12 months in the South East (4%) compared to the North of England (11.7%).

Please refer to Appendices 62 and 63 for related analyses. These present the characteristics associated with being prescribed gabapentin and pregabalin for up to 3 months, between 3 to 6 months, between 6 to 12 months and for more than 12 months.

44

⁶ Mean age (years) for short-term periods lasting under 30 days: 61.0 (gabapentin) and 55.0 (pregabalin).

Table 5: Characteristics of people prescribed gabapentin in excess of 6 months and in excess of 12 months, 2014

| Gabapentin | | | |
|----------------------|-----------|---------|--|
| | Long-term | | |
| | Over 6 | Over 12 | |
| | months | months | |
| Mean age (years) | 64.0** | 64.1 | |
| IMD quintile (%) | | | |
| I (least deprived) § | 15.2 | 4.4 | |
| II | 12.4 | 4.6 | |
| III | 9.6 | 1.5 | |
| IV | 14.6 | 7.3 | |
| V (most deprived | 10.9 | 3.4 | |
| Gender (%) | | | |
| Male § | 10.3 | 4.5 | |
| Female | 13.5 | 4.1 | |
| Region (%) | | | |
| North of England § | 12.5 | 4.1 | |
| Midlands and East of | | | |
| England | 15.7 | 5.2 | |
| London | 8.1 | 0 | |
| South East | 12.7 | 5.1 | |
| South West | 5.4 | 3.2 | |

^{*} P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, § - reference category

Source: CPRD

Table 6: Characteristics of people prescribed pregabalin in excess of 6 months and in excess of 12 months, 2014

| Pregabalin | | | |
|----------------------|-----------|---------|--|
| | Long-term | | |
| | Over 6 | Over 12 | |
| | months | months | |
| Mean age (years) | 60.7 | 59.4 | |
| IMD quintile (%) | | | |
| I (least deprived) § | 10.7 | 3.3 | |
| II | 15.1 | 8.2 | |
| III | 16.1 | 7.1 | |
| IV | 22.2* | 11.1* | |
| V (most deprived | 14.6 | 7.9 | |
| Gender (%) | | | |
| Male § | 19.9 | 13.0 | |
| Female | 13.9* | 4.9*** | |
| Region (%) | | | |
| North of England § | 19.4 | 11.7 | |
| Midlands and East of | | | |
| England | 15.6 | 6.5 | |
| London | 14.0 | 10.5 | |
| South East | 14.3 | 4.0** | |
| South West | 14.8 | 11.5 | |

^{*} P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, § - reference category

Source: CPRD

7 Time trends in sales of codeine- and dihydrocodeine-containing medications

A data extract of over the counter (OTC) sales of codeine and dihydrocodeine-containing medicines was acquired from IRI Group Ltd, a market research company which provides market intelligence data.⁷ Total counts of individual tablets were derived for three consecutive 12-month periods, ending in the final week of February 2016, 2017 and 2018.




Figure 26: Total counts of sales in individual codeine and dihydrocodeine-containing medicines

See Appendix 64 for precise counts.

There was a small decrease in total OTC sales of medicines containing codeine or dihydrocodeine between 2015/2016 and 2017/2018, broadly consistent across categories. Apart from a small increase over time in sales of ibuprofen and codeine combinations (1% increase), all types of medicine showed small decreases in sales volume between 2015/6 and 2017/18. The largest drop in sales was in aspirin and codeine products (15% reduction).

These trends are broadly consistent with those reported for all analgesics in *The Pharmaceutical Journal* using sales data supplied by IRI Ltd (Connelly, 2017), as well as by Nielson Scantrack (Connelly, 2018). The more recent article described sales of adult oral analgesics as 'stagnated'.

⁷ Data provided by IRI Group Ltd. Data are for sales of OTC products (including own-label and generic) from all multiples and independent pharmacies and most supermarkets and grocery outlets in Great Britain. These included: Asda, Iceland, Morrisons, Sainsburys (inc. local), Tesco (inc. Express, Metro), Co-operative, Waitrose, Boots, Superdrug, Lloyds, Wilkinson, and all independent chemists. Independent grocers were generally not included in the sales data. This means that trends over time are likely to be reliable, but that the total volume is an underestimate.

8 Conclusions

Prevalence

Individual DFM prescriptions almost never exceeded 12 months. However, once repeat prescribing was accounted for a sizeable minority of prescribing periods exceed this level: with about one in twenty lasting more than a year. This ranged from about 4% of prescribing periods for opioids and gabapentin, to 6% for benzodiazepines and gabapentin, and 8% for pregabalin.

Change over time

While the prevalence of continuous prescribing over 30 days tended to show signs of decline, there is little evidence of decline at the longer-term thresholds. Prescribing for periods in excess of 12 months increased particularly sharply for pregabalin since its introduction in 2004.

Pregabalin trends

As pregabalin and gabapentin become controlled as class C drugs (Home Office, 2018), it will be important to monitor the impact of reclassification on prescribing practice (Torjesen, 2019). Awareness of this steep increase in long-term pregabalin prescribing is relevant across sectors, and should inform service provision in primary care, specialist services (such as pain management), and psychiatry. Reclassification has made it illegal to supply pregabalin and gabapentin through repeat dispensing and restricts prescription lengths to 28 days. In an analysis of the impact of tramadol reclassification, which occurred in 2014 due to concerns about potential misuse, overprescribing, and increasing numbers of associated deaths, prescriptions and related deaths fell (Chen et al., 2018).

Variation

Among people prescribed DFM, those most likely to end up being prescribed to for longer were older, living in deprived neighbourhoods, and living in the North of England. In 2014, being male emerged as associated with being prescribing DFM for periods exceeding 12 months. It is possible that changing prescribing practices are not addressing the issue of very long-term prescribing in some groups of men.

Further research

The current analyses drew on a relatively small extract of data from the CPRD. There is a need for further research with larger extracts which explore patterns in long-term prescribing for individual medicines. In relation to the monitoring of opioid prescribing periods, this should include taking account of oral morphine equivalence (OME) (Curtis et al., 2019). The current research should be viewed as an initial and descriptive visualisation of patterns, future work should extend this with more detailed analyses.

9 References

BMA (2015) Prescribed drugs associated with dependence and withdrawal – building a consensus for action.

http://bmaopac.hosted.exlibrisgroup.com/exlibris/aleph/a23_1/apache_media/H6IB5G1BL8SX1KJ7VY4MCRCXVG6EV7.pdf

British Pain Society. Opioids for persistent pain: summary of guidance on good practice from the British Pain Society. *Br J Pain* 2012; **6:** 9–10.

Cartagena Farias J, Porter L, McManus S, Strang J, Hickman M, Reed K, Smith N. (2017) *Prescribing Patterns in Dependence Forming Medicines*. London. Available at: http://phrc.lshtm.ac.uk/project_2011-2019_014.html.

Chen TC, Chen LC, Knaggs RDA. A 15-year overview of increasing tramadol utilisation and associated mortality and the impact of tramadol classification in the United Kingdom. Pharmacoepidemiol Drug Saf2018;27:487-94. doi:10.1002/pds.4320 pmid:28944519

Connelly D. A breakdown of the over-the-counter medicines market in Britain in 2016. *The Pharmaceutical Journal*, April 2017, Vol 298, No 7900, online | DOI: 10.1211/PJ.2017.20202662.

Connelly D. A breakdown of the over-the-counter medicines market in Britain in 2017. *The Pharmaceutical Journal*, May 2018, Vol 300, No 7913,

Curtis H, Croker R, Walker AJ, Richards GC, Quinlan J, Goldacre B. Opioid Prescribing Trends and geographical variation in England, 1998-2018 a retrospective database study. *Lancet Psychiatry*. 2019, 6: 2; 140-150.

Faculty of Pain Medicine (2019) Opioids Aware: A resource for patients and healthcare professionals to support prescribing of opioid medicines for pain. https://www.rcoa.ac.uk/faculty-of-pain-medicine/opioids-aware

Herrett E, Gallagher AM, Bhaskaran K, Forbes H, Mathur R, van Staa T, Smeeth L. Data Resource Profile: Clinical Practice Research Datalink (CPRD). *International Journal of Epidemiology*, Volume 44, Issue 3, 1 June 2015, Pages 827–836.

Home Office. Atkins V. Pregabalin and gabapentin to be controlled as class C drugs. 15 Oct 2018. www.gov.uk/government/news/pregabalin-and-gabapentin-to-be-controlled-as-class-c-drugs.

McManus S. Opioids: long-term prescribing trends. *Lancet Psychiatry*. 2019, 6: 2; 140-150doi.or/10.1016/S2215-0366(18)30507-8

Montastruc F, Loo SY, Renoux C. Trends in First Gabapentin and Pregabalin Prescriptions in Primary Care in the United Kingdom, 1993-2017 *JAMA*. 2018;320(20):2149-2151. doi:10.1001/jama.2018.12358

NHS Digital (2019). *Prescription cost analysis-England 2018*. https://digital.nhs.uk/data-and-information/publications/statistical/prescription-cost-analysis/2018. (Accessed May 9th, 2019)

Reed K, Bond A, Witton J, Cornish R, Hickman M, Strang J. (2011) *The changing use of prescribed benzodiazepines and z-drugs and of over-the-counter codeine-containing products in England*. Kings: London.

Taylor S, Annand F, Burkinshaw P, Greaves F, Kelleher M, Knight J, Perkins C, Tran A, White M, Marsden J. *An evidence review of dependence and withdrawal associated with some prescribed medicines*. 2019. Public Health England, London.

Torjesen I. Pregabalin and gabapentin: what impact will reclassification have on doctors and patients? *BMJ* 2019; 364: I1107.

Appendix A. Data tables

Appendix 1 Average number of days of prescriptions by DFM type

| Year | Benzodiazepines | Z-drugs | Opioids | GABAergic medicines |
|-------------------|-----------------|---------|---------|------------------------|
| 2000 ⁸ | 31.0 | 27.9 | 35.8 | 30.9 |
| 2001 | 30.9 | 26.8 | 30.2 | 32.3 |
| 2002 | 30.8 | 26.7 | 26.2 | 32.0 |
| 2003 | 30.0 | 26.4 | 22.9 | 31.9 |
| 2004 | 29.0 | 26.6 | 23.2 | 32.6 |
| 2005 | 28.3 | 26.6 | 23.0 | 31.5 |
| 2006 | 27.1 | 26.6 | 22.9 | 31.1 |
| 2007 | 26.5 | 26.2 | 22.6 | 31.5 |
| 2008 | 25.6 | 26.7 | 22.8 | 31.4 |
| 2009 | 26.2 | 26.3 | 23.3 | 31.7 |
| 2010 | 25.4 | 25.7 | 23.7 | 31.2 |
| 2011 | 25.5 | 25.2 | 23.8 | 31.1 |
| 2012 | 24.8 | 24.3 | 23.8 | 30.9 |
| 2013 | 23.9 | 23.7 | 24.0 | 31.3 |
| 2014 | 24.7 | 24.2 | 24.2 | 30.6 |
| 2015 | 25.5 | 24.6 | 23.9 | 30.7 |

Appendix 2 Average number of days of continuous prescribing periods by DFM type

| Year | Benzodiazepines | Z-drugs | Opioids | GABAergic medicines |
|------|-----------------|---------|---------|------------------------|
| 2000 | 95.0 | 78.2 | 64.1 | 80.2 |
| 2001 | 106.4 | 94.9 | 72.1 | 112.0 |
| 2002 | 115.9 | 94.6 | 73.9 | 100.5 |
| 2003 | 117.8 | 91.4 | 77.5 | 99.6 |
| 2004 | 113.5 | 87.3 | 76.5 | 98.8 |
| 2005 | 117.2 | 107.9 | 81.7 | 105.2 |
| 2006 | 117.5 | 100.3 | 83.0 | 116.8 |
| 2007 | 113.7 | 104.1 | 80.9 | 111.7 |
| 2008 | 118.3 | 104.6 | 84.8 | 108.8 |
| 2009 | 120.7 | 103.2 | 90.3 | 113.5 |
| 2010 | 121.9 | 103.4 | 92.0 | 121.1 |
| 2011 | 112.0 | 109.9 | 99.4 | 115.3 |
| 2012 | 123.2 | 113.3 | 96.0 | 119.4 |
| 2013 | 110.0 | 106.6 | 102.6 | 122.8 |
| 2014 | 117.0 | 107.7 | 103.2 | 123.1 |
| 2015 | 108.9 | 112.8 | 93.8 | 113.5 |

⁸ In the tables data for 2000 and 2015 are presented but should be treated with caution due to the impact of truncation near the start and end of the data extract. To emphasis this, these figures are presented in grey.

51

Appendix 3 Proportion of prescriptions exceeding 30 days by DFM type.

| % Long-term prescriptions (over 30 days) | | | | | |
|--|-----------------|---------|---------|-----------|--|
| | | | | GABAergic | |
| Year | Benzodiazepines | Z-drugs | Opioids | medicines | |
| 2000 | 21.7 | 12.9 | 15.7 | 24.7 | |
| 2001 | 21.3 | 10.0 | 15.4 | 29.4 | |
| 2002 | 21.0 | 11.8 | 16.0 | 31.2 | |
| 2003 | 20.7 | 11.6 | 15.0 | 36.9 | |
| 2004 | 19.8 | 11.0 | 15.4 | 38.0 | |
| 2005 | 18.5 | 11.5 | 15.3 | 32.9 | |
| 2006 | 16.7 | 11.6 | 15.7 | 30.8 | |
| 2007 | 14.5 | 10.5 | 15.1 | 32.5 | |
| 2008 | 12.0 | 10.7 | 15.3 | 31.8 | |
| 2009 | 12.9 | 12.1 | 15.8 | 32.4 | |
| 2010 | 12.4 | 11.8 | 16.1 | 31.5 | |
| 2011 | 12.2 | 9.5 | 16.2 | 31.9 | |
| 2012 | 11.5 | 8.9 | 16.7 | 29.0 | |
| 2013 | 11.0 | 8.6 | 18.2 | 26.7 | |
| 2014 | 12.1 | 9.2 | 17.5 | 23.6 | |
| 2015 | 11.7 | 9.9 | 15.7 | 25.0 | |

Appendix 4 Proportion of continuous prescribing periods exceeding 30 days by DFM type

| | % Long-term prescribing (over 30 days) | | | | |
|------|--|---------|---------|------------------------|--|
| Year | Benzodiazepines | Z-drugs | Opioids | GABAergic medicines | |
| 2000 | 50.7 | 43.8 | 37.1 | 55.8 | |
| 2001 | 49.1 | 42.7 | 38.5 | 58.0 | |
| 2002 | 46.0 | 44.0 | 37.9 | 62.2 | |
| 2003 | 45.6 | 42.2 | 37.3 | 69.3 | |
| 2004 | 44.0 | 43.7 | 38.8 | 67.3 | |
| 2005 | 43.5 | 42.9 | 37.8 | 70.7 | |
| 2006 | 42.6 | 42.4 | 37.3 | 65.5 | |
| 2007 | 39.6 | 40.6 | 36.8 | 66.7 | |
| 2008 | 37.9 | 41.6 | 37.0 | 67.4 | |
| 2009 | 35.5 | 42.8 | 37.3 | 67.8 | |
| 2010 | 34.3 | 40.7 | 38.1 | 69.3 | |
| 2011 | 33.0 | 41.8 | 36.7 | 64.8 | |
| 2012 | 34.2 | 43.5 | 35.9 | 61.6 | |
| 2013 | 32.3 | 40.2 | 34.2 | 58.7 | |
| 2014 | 32.7 | 37.0 | 33.7 | 58.0 | |
| 2015 | 31.8 | 37.1 | 32.3 | 56.3 | |

Appendix 5 Proportion of continuous prescribing periods exceeding 6 months by DFM type

| | % Long-term prescribing (over 6 months) | | | | |
|------|---|---------|---------|-----------|--|
| | | | | GABAergic | |
| Year | Benzodiazepines | Z-drugs | Opioids | medicines | |
| 2000 | 12.3 | 7.9 | 5.7 | 9.1 | |
| 2001 | 12.6 | 12.3 | 7.2 | 10.4 | |
| 2002 | 13.2 | 10.8 | 5.7 | 12.3 | |
| 2003 | 13.2 | 9.8 | 6.8 | 12.5 | |
| 2004 | 11.2 | 11.2 | 6.9 | 11.6 | |
| 2005 | 12.8 | 11.9 | 7.4 | 11.3 | |
| 2006 | 12.7 | 11.7 | 7.2 | 13.5 | |
| 2007 | 11.7 | 12.6 | 6.8 | 13.0 | |
| 2008 | 11.0 | 11.1 | 7.1 | 13.4 | |
| 2009 | 10.6 | 11.9 | 8.2 | 13.3 | |
| 2010 | 11.2 | 10.7 | 7.5 | 14.5 | |
| 2011 | 10.8 | 12.5 | 7.5 | 13.7 | |
| 2012 | 11.5 | 13.3 | 7.5 | 13.6 | |
| 2013 | 10.0 | 12.1 | 7.4 | 14.3 | |
| 2014 | 12.1 | 12.8 | 7.3 | 14.0 | |
| 2015 | 9.0 | 12.0 | 6.6 | 11.9 | |

Appendix 6 Proportion of continuous prescribing periods exceeding 12 months by DFM type

| | % Long-term prescribing (over 12 months) | | | | |
|------|--|---------|---------|------------------------|--|
| Year | Benzodiazepines | Z-drugs | Opioids | GABAergic medicines | |
| 2000 | 3.8 | 2.8 | 1.8 | 1.7 | |
| 2001 | 5.4 | 5.1 | 2.5 | 5.1 | |
| 2002 | 6.4 | 4.7 | 2.5 | 3.4 | |
| 2003 | 6.1 | 4.7 | 3.0 | 4.0 | |
| 2004 | 5.3 | 4.0 | 2.4 | 3.0 | |
| 2005 | 5.8 | 5.4 | 2.8 | 3.1 | |
| 2006 | 5.7 | 5.1 | 2.9 | 5.0 | |
| 2007 | 5.5 | 5.1 | 2.6 | 3.7 | |
| 2008 | 5.2 | 4.7 | 2.7 | 4.3 | |
| 2009 | 5.1 | 5.5 | 3.3 | 4.6 | |
| 2010 | 5.5 | 4.3 | 3.0 | 5.7 | |
| 2011 | 5.0 | 5.9 | 3.1 | 5.0 | |
| 2012 | 5.2 | 6.2 | 3.4 | 4.4 | |
| 2013 | 4.5 | 6.0 | 3.3 | 5.7 | |
| 2014 | 5.9 | 6.0 | 3.6 | 5.0 | |
| 2015 | 4.4 | 5.5 | 1.8 | 3.6 | |

Appendix 7 Average number of days of prescriptions by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 31.8 | - |
| 2001 | 35.4 | - |
| 2002 | 37.0 | - |
| 2003 | 35.1 | - |
| 2004 | 35.3 | 26.6 |
| 2005 | 36.3 | 31.1 |
| 2006 | 36.5 | 31.6 |
| 2007 | 35.6 | 32.9 |
| 2008 | 35.7 | 34.9 |
| 2009 | 36.7 | 34.3 |
| 2010 | 37.0 | 34.9 |
| 2011 | 36.4 | 34.5 |
| 2012 | 36.5 | 34.7 |
| 2013 | 36.3 | 35.1 |
| 2014 | 36.5 | 35.3 |
| 2015 | 35.1 | 36.3 |

Appendix 8 Average number of days of continuous prescribing periods by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 75.3 | - |
| 2001 | 91.4 | - |
| 2002 | 93.1 | - |
| 2003 | 95.3 | - |
| 2004 | 84.7 | 64.2 |
| 2005 | 103.4 | 66.8 |
| 2006 | 98.2 | 90.2 |
| 2007 | 95.2 | 91.4 |
| 2008 | 100.8 | 110.3 |
| 2009 | 99.4 | 110.5 |
| 2010 | 104.5 | 104.4 |
| 2011 | 95.2 | 108.6 |
| 2012 | 96.9 | 107.6 |
| 2013 | 100.2 | 114.2 |
| 2014 | 102.4 | 126.6 |
| 2015 | 94.0 | 116.3 |

Appendix 9 Proportion of prescriptions exceeding 30 days by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 29.2 | - |
| 2001 | 49.2 | - |
| 2002 | 51.4 | - |
| 2003 | 56.6 | - |
| 2004 | 55.9 | 13.3 |
| 2005 | 57.7 | 15.6 |
| 2006 | 57.1 | 15.6 |
| 2007 | 58.4 | 21.6 |
| 2008 | 59.8 | 28.8 |
| 2009 | 64.4 | 25.4 |
| 2010 | 64.0 | 25.4 |
| 2011 | 65.1 | 25.6 |
| 2012 | 58.6 | 27.5 |
| 2013 | 54.8 | 29.6 |
| 2014 | 52.2 | 29.6 |
| 2015 | 49.7 | 30.1 |

Appendix 10 Proportion of continuous prescribing periods exceeding 30 days by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 55.5 | - |
| 2001 | 68.5 | - |
| 2002 | 68.2 | - |
| 2003 | 72.6 | - |
| 2004 | 71.5 | 60.0 |
| 2005 | 75.5 | 56.7 |
| 2006 | 73.4 | 54.6 |
| 2007 | 72.7 | 59.5 |
| 2008 | 74.0 | 62.6 |
| 2009 | 74.8 | 63.9 |
| 2010 | 75.7 | 60.8 |
| 2011 | 71.3 | 58.5 |
| 2012 | 68.7 | 57.3 |
| 2013 | 62.2 | 57.6 |
| 2014 | 59.8 | 62.7 |
| 2015 | 58.3 | 61.8 |

Appendix 11 Proportion of continuous prescribing periods exceeding 6 months by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 9.2 | - |
| 2001 | 9.6 | - |
| 2002 | 8.0 | - |
| 2003 | 12.3 | - |
| 2004 | 8.7 | 6.7 |
| 2005 | 11.9 | 3.0 |
| 2006 | 11.5 | 11.4 |
| 2007 | 10.5 | 11.3 |
| 2008 | 11.3 | 12.2 |
| 2009 | 12.7 | 13.2 |
| 2010 | 11.7 | 12.0 |
| 2011 | 9.1 | 14.1 |
| 2012 | 11.3 | 13.7 |
| 2013 | 11.5 | 15.9 |
| 2014 | 12.3 | 16.0 |
| 2015 | 9.2 | 13.8 |

Appendix 12 Proportion of continuous prescribing periods exceeding 12 months by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 0.0 | - |
| 2001 | 4.1 | - |
| 2002 | 2.6 | - |
| 2003 | 4.0 | - |
| 2004 | 2.6 | 0.0 |
| 2005 | 4.3 | 0.8 |
| 2006 | 3.1 | 4.0 |
| 2007 | 3.1 | 2.8 |
| 2008 | 4.0 | 3.8 |
| 2009 | 3.8 | 4.5 |
| 2010 | 3.9 | 4.7 |
| 2011 | 3.1 | 5.7 |
| 2012 | 3.4 | 4.5 |
| 2013 | 4.0 | 6.1 |
| 2014 | 4.2 | 7.7 |
| 2015 | 1.7 | 5.2 |

Appendix 13 Benzodiazepines: Proportion of continuous prescribing periods exceeding 30 days, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 35.9 | 43.8 | 52.3 | 62.1 |
| 2001 | 34.1 | 42.5 | 50.7 | 60.7 |
| 2002 | 26.5 | 37.9 | 49.5 | 58.1 |
| 2003 | 27.5 | 38.7 | 46.8 | 59.9 |
| 2004 | 28.5 | 35.9 | 45.8 | 56.6 |
| 2005 | 30.4 | 35.9 | 45.2 | 56.9 |
| 2006 | 26.6 | 36.5 | 44.9 | 55.5 |
| 2007 | 26.6 | 34.7 | 41.7 | 51.2 |
| 2008 | 28.8 | 31.9 | 39.5 | 48.7 |
| 2009 | 23.4 | 30.2 | 36.9 | 47.0 |
| 2010 | 23.3 | 31.1 | 35.8 | 41.7 |
| 2011 | 16.0 | 28.5 | 35.7 | 43.1 |
| 2012 | 20.8 | 29.8 | 37.4 | 44.0 |
| 2013 | 16.4 | 26.9 | 37.8 | 42.0 |
| 2014 | 22.7 | 31.4 | 35.6 | 37.0 |
| 2015 | 19.2 | 31.0 | 33.1 | 40.9 |

Appendix 14 Benzodiazepines: Proportion of continuous prescribing periods exceeding 6 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 6.5 | 8.7 | 13.3 | 17.4 |
| 2001 | 8.9 | 11.1 | 12.4 | 16.4 |
| 2002 | 7.2 | 10.2 | 12.5 | 21.1 |
| 2003 | 6.5 | 9.6 | 13.4 | 20.2 |
| 2004 | 5.5 | 9.5 | 11.5 | 15.0 |
| 2005 | 9.1 | 11.4 | 13.1 | 16.2 |
| 2006 | 5.9 | 11.3 | 13.3 | 17.1 |
| 2007 | 8.9 | 10.1 | 11.6 | 16.0 |
| 2008 | 5.7 | 8.1 | 11.7 | 16.9 |
| 2009 | 5.4 | 8.9 | 10.5 | 15.9 |
| 2010 | 9.5 | 10.3 | 11.0 | 13.5 |
| 2011 | 4.9 | 9.9 | 10.4 | 15.6 |
| 2012 | 7.7 | 8.4 | 12.0 | 17.5 |
| 2013 | 2.3 | 10.0 | 10.0 | 16.0 |
| 2014 | 10.8 | 10.7 | 12.5 | 14.7 |
| 2015 | 4.7 | 8.6 | 8.7 | 13.6 |

Appendix 15 Benzodiazepines: Proportion of continuous prescribing periods exceeding 12 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 2.2 | 3.2 | 4.4 | 4.1 |
| 2001 | 3.4 | 4.7 | 5.6 | 6.7 |
| 2002 | 4.0 | 4.2 | 6.7 | 9.7 |
| 2003 | 3.9 | 4.0 | 5.5 | 10.7 |
| 2004 | 3.2 | 5.1 | 5.5 | 6.1 |
| 2005 | 4.0 | 4.8 | 6.0 | 7.8 |
| 2006 | 2.6 | 6.3 | 5.9 | 6.6 |
| 2007 | 2.5 | 4.7 | 5.5 | 8.4 |
| 2008 | 3.6 | 4.1 | 5.3 | 7.6 |
| 2009 | 3.2 | 4.0 | 4.6 | 8.4 |
| 2010 | 4.8 | 5.6 | 6.1 | 4.5 |
| 2011 | 2.2 | 5.2 | 4.4 | 7.2 |
| 2012 | 3.3 | 3.7 | 5.9 | 7.1 |
| 2013 | 0.3 | 3.9 | 5.4 | 7.0 |
| 2014 | 6.3 | 6.4 | 5.9 | 4.6 |
| 2015 | 0.8 | 4.8 | 5.4 | 4.7 |

Appendix 16 Z-drugs: Proportion of continuous prescribing periods exceeding 30 days, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 30.8 | 43.7 | 44.4 | 61.7 |
| 2001 | 34.0 | 37.1 | 45.0 | 60.8 |
| 2002 | 30.5 | 40.2 | 47.9 | 58.5 |
| 2003 | 35.2 | 35.9 | 49.2 | 50.6 |
| 2004 | 33.4 | 36.9 | 51.8 | 52.5 |
| 2005 | 28.4 | 36.4 | 49.3 | 54.5 |
| 2006 | 31.5 | 35.8 | 46.0 | 61.1 |
| 2007 | 30.5 | 37.1 | 45.6 | 47.8 |
| 2008 | 30.5 | 36.3 | 45.7 | 57.7 |
| 2009 | 29.6 | 36.9 | 46.9 | 59.6 |
| 2010 | 31.0 | 34.3 | 43.1 | 61.7 |
| 2011 | 27.9 | 37.2 | 50.2 | 47.4 |
| 2012 | 30.0 | 37.6 | 48.5 | 56.0 |
| 2013 | 27.8 | 32.7 | 46.2 | 53.5 |
| 2014 | 21.8 | 32.7 | 41.8 | 46.0 |
| 2015 | 21.2 | 33.7 | 41.4 | 45.8 |

Appendix 17 Z-drugs: Proportion of continuous prescribing periods exceeding 6 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 5.3 | 6.3 | 7.9 | 14.7 |
| 2001 | 5.2 | 10.8 | 13.6 | 21.4 |
| 2002 | 6.6 | 10.9 | 10.7 | 15.4 |
| 2003 | 8.4 | 8.5 | 9.5 | 15.7 |
| 2004 | 5.8 | 9.8 | 13.4 | 15.3 |
| 2005 | 8.7 | 9.8 | 12.8 | 16.9 |
| 2006 | 5.2 | 8.7 | 12.9 | 22.6 |
| 2007 | 9.5 | 11.6 | 13.2 | 16.9 |
| 2008 | 6.7 | 9.2 | 11.6 | 19.0 |
| 2009 | 5.5 | 7.9 | 14.6 | 21.3 |
| 2010 | 6.0 | 8.7 | 10.7 | 21.0 |
| 2011 | 4.5 | 11.2 | 13.8 | 20.4 |
| 2012 | 8.1 | 10.8 | 13.7 | 22.2 |
| 2013 | 7.0 | 11.6 | 11.4 | 19.8 |
| 2014 | 5.1 | 10.5 | 14.7 | 18.7 |
| 2015 | 6.5 | 11.1 | 11.4 | 18.7 |

Appendix 18 Z-drugs: Proportion of continuous prescribing periods exceeding 12 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 2.2 | 2.1 | 3.2 | 4.3 |
| 2001 | 1.8 | 4.2 | 6.0 | 8.8 |
| 2002 | 3.5 | 5.3 | 4.0 | 6.6 |
| 2003 | 4.4 | 4.0 | 4.8 | 6.1 |
| 2004 | 3.3 | 3.8 | 3.9 | 5.5 |
| 2005 | 4.6 | 4.7 | 5.4 | 7.9 |
| 2006 | 3.5 | 3.2 | 6.2 | 8.9 |
| 2007 | 3.6 | 5.9 | 4.9 | 5.3 |
| 2008 | 2.3 | 4.1 | 5.1 | 8.0 |
| 2009 | 2.4 | 3.9 | 6.3 | 10.4 |
| 2010 | 2.6 | 4.9 | 3.6 | 6.7 |
| 2011 | 1.7 | 5.9 | 6.3 | 9.1 |
| 2012 | 5.1 | 3.9 | 6.1 | 12.2 |
| 2013 | 6.1 | 5.7 | 4.5 | 9.6 |
| 2014 | 3.0 | 5.0 | 6.8 | 8.2 |
| 2015 | 3.7 | 4.6 | 5.2 | 9.1 |

Appendix 19 Opioids: Proportion of continuous prescribing periods exceeding 30 days, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 28.5 | 36.1 | 39.6 | 45.3 |
| 2001 | 28.4 | 36.3 | 43.3 | 42.8 |
| 2002 | 29.4 | 37.8 | 42.6 | 35.8 |
| 2003 | 25.6 | 36.3 | 40.7 | 44.7 |
| 2004 | 27.8 | 36.2 | 44.0 | 41.8 |
| 2005 | 26.9 | 36.7 | 41.1 | 42.9 |
| 2006 | 30.6 | 35.9 | 38.3 | 44.6 |
| 2007 | 28.2 | 36.0 | 39.4 | 39.2 |
| 2008 | 26.8 | 38.1 | 39.9 | 36.0 |
| 2009 | 24.6 | 37.0 | 40.4 | 39.7 |
| 2010 | 26.8 | 37.1 | 40.6 | 41.9 |
| 2011 | 28.6 | 36.7 | 38.2 | 39.1 |
| 2012 | 28.9 | 34.9 | 37.0 | 41.3 |
| 2013 | 28.1 | 34.6 | 36.8 | 31.7 |
| 2014 | 25.1 | 35.1 | 36.0 | 31.5 |
| 2015 | 23.4 | 35.2 | 33.1 | 29.7 |

Appendix 20 Opioids: Proportion of continuous prescribing periods exceeding 6 months, by age

| | | | <u> </u> | |
|------|-------|-------|----------|-----|
| Year | 18-40 | 41-60 | 61-80 | 81+ |
| 2000 | 5.4 | 5.4 | 5.8 | 6.7 |
| 2001 | 5.9 | 6.9 | 7.5 | 8.6 |
| 2002 | 4.5 | 7.1 | 5.6 | 4.8 |
| 2003 | 6.2 | 7.2 | 6.7 | 7.5 |
| 2004 | 4.8 | 6.4 | 8.2 | 6.5 |
| 2005 | 6.9 | 7.3 | 7.7 | 7.4 |
| 2006 | 6.4 | 7.5 | 6.6 | 9.2 |
| 2007 | 5.1 | 6.8 | 7.0 | 8.3 |
| 2008 | 5.0 | 7.0 | 7.2 | 8.7 |
| 2009 | 5.9 | 7.4 | 9.5 | 8.4 |
| 2010 | 6.0 | 7.3 | 7.7 | 8.5 |
| 2011 | 5.6 | 8.3 | 7.8 | 6.5 |
| 2012 | 5.0 | 6.4 | 8.4 | 9.6 |
| 2013 | 6.7 | 8.0 | 7.8 | 6.1 |
| 2014 | 6.4 | 7.5 | 7.9 | 6.2 |
| 2015 | 4.7 | 6.5 | 7.3 | 6.1 |

Appendix 21 Opioids: Proportion of continuous prescribing periods exceeding 12 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|-----|
| 2000 | 1.9 | 1.6 | 1.9 | 1.8 |
| 2001 | 2.1 | 2.5 | 2.8 | 2.4 |
| 2002 | 2.5 | 3.1 | 2.5 | 1.2 |
| 2003 | 2.8 | 3.6 | 2.5 | 3.1 |
| 2004 | 1.8 | 2.4 | 2.6 | 2.5 |
| 2005 | 2.9 | 2.7 | 3.0 | 2.6 |
| 2006 | 2.7 | 3.0 | 2.7 | 3.9 |
| 2007 | 2.8 | 3.2 | 2.1 | 2.8 |
| 2008 | 3.0 | 2.2 | 2.9 | 3.2 |
| 2009 | 2.4 | 3.5 | 3.6 | 2.6 |
| 2010 | 3.1 | 3.0 | 2.8 | 3.3 |
| 2011 | 2.4 | 3.6 | 3.2 | 2.6 |
| 2012 | 2.8 | 2.9 | 3.8 | 3.9 |
| 2013 | 3.3 | 4.2 | 2.9 | 2.3 |
| 2014 | 3.8 | 3.6 | 4.0 | 2.7 |
| 2015 | 2.1 | 1.9 | 1.7 | 2.0 |

Appendix 22 GABAergic medicines: Proportion of continuous prescribing periods exceeding 30 days, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 38.1 | 62.1 | 60.1 | 76.9 |
| 2001 | 57.5 | 66.8 | 57.8 | 53.6 |
| 2002 | 52.0 | 69.2 | 74.8 | 44.2 |
| 2003 | 63.8 | 77.9 | 73.2 | 65.7 |
| 2004 | 61.8 | 68.8 | 70.3 | 71.0 |
| 2005 | 69.8 | 70.5 | 75.6 | 62.5 |
| 2006 | 62.6 | 67.5 | 64.7 | 64.6 |
| 2007 | 68.5 | 68.6 | 68.2 | 60.5 |
| 2008 | 66.0 | 64.0 | 74.3 | 63.2 |
| 2009 | 64.8 | 65.3 | 74.7 | 61.1 |
| 2010 | 59.1 | 68.2 | 74.9 | 69.2 |
| 2011 | 56.1 | 66.8 | 66.9 | 64.8 |
| 2012 | 50.8 | 62.6 | 65.0 | 61.2 |
| 2013 | 50.3 | 58.7 | 60.7 | 63.9 |
| 2014 | 44.2 | 58.2 | 61.6 | 63.9 |
| 2015 | 56.9 | 51.8 | 61.1 | 58.1 |

Appendix 23 GABAergic medicines: Proportion of continuous prescribing periods exceeding 6 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 5.8 | 13.0 | 6.2 | 0.0 |
| 2001 | 9.7 | 10.8 | 10.4 | 8.9 |
| 2002 | 9.7 | 16.1 | 9.5 | 7.2 |
| 2003 | 9.4 | 11.3 | 14.9 | 14.3 |
| 2004 | 18.2 | 8.1 | 12.0 | 11.2 |
| 2005 | 19.0 | 9.4 | 10.9 | 7.1 |
| 2006 | 20.0 | 11.7 | 12.2 | 14.1 |
| 2007 | 13.7 | 16.1 | 10.6 | 11.3 |
| 2008 | 12.0 | 13.9 | 12.7 | 15.0 |
| 2009 | 7.7 | 14.0 | 13.1 | 21.3 |
| 2010 | 10.8 | 15.6 | 14.3 | 18.6 |
| 2011 | 8.5 | 12.6 | 16.4 | 15.9 |
| 2012 | 12.0 | 11.3 | 15.8 | 15.8 |
| 2013 | 10.6 | 11.4 | 16.5 | 19.5 |
| 2014 | 9.5 | 12.9 | 14.4 | 21.0 |
| 2015 | 15.4 | 11.0 | 11.7 | 12.3 |

Appendix 24 GABAergic medicines: Proportion of continuous prescribing periods exceeding 12 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|-----|
| 2000 | 0.0 | 1.9 | 0.0 | 0.0 |
| 2001 | 5.8 | 5.3 | 5.8 | 0.0 |
| 2002 | 1.6 | 6.0 | 1.0 | 3.6 |
| 2003 | 0.0 | 4.9 | 4.0 | 5.7 |
| 2004 | 7.1 | 0.2 | 4.5 | 2.2 |
| 2005 | 8.5 | 2.5 | 2.3 | 0.4 |
| 2006 | 5.4 | 4.0 | 4.9 | 7.2 |
| 2007 | 5.5 | 3.6 | 3.3 | 3.9 |
| 2008 | 1.8 | 5.2 | 4.0 | 6.7 |
| 2009 | 3.3 | 4.5 | 4.1 | 9.9 |
| 2010 | 3.3 | 6.9 | 5.0 | 7.8 |
| 2011 | 1.8 | 5.1 | 5.9 | 7.1 |
| 2012 | 3.9 | 4.0 | 4.6 | 5.7 |
| 2013 | 5.2 | 4.4 | 5.9 | 9.3 |
| 2014 | 5.0 | 4.3 | 5.5 | 5.9 |
| 2015 | 2.6 | 3.6 | 4.3 | 2.4 |

Appendix 25 Proportion of continuous prescribing periods exceeding 30 days, by DFM type and gender

| | Benzodiazepines Z-drugs | | Opi | Opioids | | GABAergic medicines | | |
|-------|-------------------------|--------|------|---------|------|---------------------|------|--------|
| Years | Male | Female | Male | Female | Male | Female | Male | Female |
| 2000 | 48.9 | 51.5 | 42.7 | 44.4 | 36.2 | 37.7 | 54.6 | 56.8 |
| 2001 | 47.3 | 50.0 | 41.0 | 43.5 | 38.5 | 38.6 | 60.7 | 55.7 |
| 2002 | 44.1 | 47.0 | 45.9 | 43.0 | 38.0 | 37.8 | 65.2 | 60.3 |
| 2003 | 44.9 | 45.9 | 44.5 | 41.0 | 36.7 | 37.7 | 73.5 | 66.9 |
| 2004 | 41.5 | 45.2 | 46.4 | 42.2 | 36.0 | 40.4 | 73.5 | 63.2 |
| 2005 | 41.8 | 44.3 | 45.5 | 41.5 | 36.2 | 38.8 | 75.6 | 67.5 |
| 2006 | 41.6 | 43.1 | 43.9 | 41.5 | 35.3 | 38.5 | 72.5 | 61.2 |
| 2007 | 39.4 | 39.7 | 42.1 | 39.8 | 37.3 | 36.5 | 68.4 | 65.7 |
| 2008 | 38.3 | 37.7 | 40.1 | 42.4 | 35.1 | 38.1 | 72.5 | 64.6 |
| 2009 | 34.5 | 35.9 | 43.1 | 42.5 | 36.4 | 37.8 | 68.7 | 67.3 |
| 2010 | 35.8 | 33.5 | 40.6 | 40.8 | 38.5 | 37.8 | 69.4 | 69.2 |
| 2011 | 33.5 | 32.7 | 41.8 | 41.8 | 36.1 | 37.1 | 67.0 | 63.2 |
| 2012 | 33.6 | 34.6 | 42.3 | 44.1 | 35.3 | 36.2 | 63.4 | 60.3 |
| 2013 | 34.1 | 31.3 | 40.2 | 40.2 | 34.2 | 34.2 | 59.1 | 58.5 |
| 2014 | 36.0 | 31.0 | 36.9 | 37.0 | 31.9 | 34.7 | 56.8 | 58.7 |
| 2015 | 32.9 | 31.2 | 36.4 | 37.4 | 31.5 | 32.7 | 55.9 | 56.5 |

Appendix 26 Proportion of continuous prescribing periods exceeding 6 months, by DFM type and gender

| | Benzodi | azepines | Z-dru | ıgs | Opio | ids | GABAergic | medicines |
|-------|---------|----------|-------|--------|------|--------|-----------|-----------|
| Years | Male | Female | Male | Female | Male | Female | Male | Female |
| 2000 | 10.5 | 13.1 | 7.5 | 8.0 | 6.6 | 5.2 | 8.5 | 9.6 |
| 2001 | 10.4 | 13.7 | 12.8 | 12.1 | 6.5 | 7.6 | 12.5 | 8.5 |
| 2002 | 11.9 | 13.8 | 11.5 | 10.4 | 5.4 | 5.9 | 15.8 | 10.0 |
| 2003 | 10.6 | 14.4 | 10.3 | 9.5 | 7.2 | 6.6 | 17.1 | 9.9 |
| 2004 | 9.6 | 12.0 | 11.3 | 11.2 | 6.2 | 7.2 | 16.8 | 8.2 |
| 2005 | 11.7 | 13.3 | 11.7 | 12.0 | 7.5 | 7.3 | 14.5 | 9.2 |
| 2006 | 11.6 | 13.3 | 10.7 | 12.3 | 6.6 | 7.5 | 15.8 | 12.0 |
| 2007 | 11.3 | 12.0 | 13.5 | 12.1 | 6.6 | 7.0 | 13.6 | 12.7 |
| 2008 | 11.0 | 11.0 | 10.1 | 11.5 | 6.4 | 7.4 | 16.2 | 11.9 |
| 2009 | 10.5 | 10.7 | 10.8 | 12.5 | 7.6 | 8.6 | 12.5 | 13.8 |
| 2010 | 11.6 | 11.0 | 10.4 | 10.8 | 6.9 | 7.8 | 12.7 | 15.7 |
| 2011 | 10.0 | 11.2 | 11.2 | 13.1 | 7.9 | 7.1 | 13.5 | 13.9 |
| 2012 | 10.8 | 11.9 | 13.3 | 13.3 | 6.6 | 8.1 | 13.3 | 13.7 |
| 2013 | 11.1 | 9.4 | 11.5 | 12.5 | 7.4 | 7.4 | 13.9 | 14.6 |
| 2014 | 12.9 | 11.7 | 13.9 | 12.2 | 7.3 | 7.3 | 14.3 | 13.8 |
| 2015 | 8.2 | 9.4 | 9.0 | 13.6 | 6.3 | 6.7 | 13.0 | 11.3 |

Appendix 27 Proportion of continuous prescribing periods exceeding 12 months, by DFM type and gender

| | Benzodiazepines | | Z-d | Z-drugs | | Opioids | | GABAergic medicines | |
|-------|-----------------|--------|------|---------|------|---------|------|---------------------|--|
| Years | Male | Female | Male | Female | Male | Female | Male | Female | |
| 2000 | 4.0 | 3.7 | 3.3 | 2.6 | 1.8 | 1.8 | 2.4 | 1.0 | |
| 2001 | 4.6 | 5.8 | 5.0 | 5.1 | 2.5 | 2.6 | 5.6 | 4.7 | |
| 2002 | 5.0 | 7.1 | 5.6 | 4.3 | 2.1 | 2.7 | 5.4 | 2.1 | |
| 2003 | 4.8 | 6.7 | 4.7 | 4.6 | 3.3 | 2.8 | 3.9 | 4.1 | |
| 2004 | 4.4 | 5.7 | 4.5 | 3.7 | 2.2 | 2.5 | 4.4 | 2.1 | |
| 2005 | 5.8 | 5.8 | 5.3 | 5.5 | 2.8 | 2.8 | 4.1 | 2.4 | |
| 2006 | 5.7 | 5.7 | 4.7 | 5.4 | 2.5 | 3.2 | 5.8 | 4.6 | |
| 2007 | 5.3 | 5.6 | 6.2 | 4.5 | 2.6 | 2.7 | 5.0 | 3.0 | |
| 2008 | 4.1 | 5.8 | 3.4 | 5.4 | 2.2 | 3.0 | 4.9 | 4.0 | |
| 2009 | 5.3 | 5.0 | 5.6 | 5.4 | 3.3 | 3.2 | 4.2 | 4.8 | |
| 2010 | 6.6 | 4.9 | 3.9 | 4.5 | 2.8 | 3.1 | 5.6 | 5.7 | |
| 2011 | 4.2 | 5.4 | 5.1 | 6.3 | 3.3 | 2.9 | 4.6 | 5.3 | |
| 2012 | 5.5 | 5.1 | 6.3 | 6.2 | 3.3 | 3.4 | 4.0 | 4.6 | |
| 2013 | 4.2 | 4.7 | 6.2 | 5.8 | 3.5 | 3.2 | 5.6 | 5.7 | |
| 2014 | 7.3 | 5.1 | 7.7 | 5.1 | 4.2 | 3.3 | 6.9 | 3.9 | |
| 2015 | 3.8 | 4.7 | 4.9 | 5.8 | 1.3 | 2.1 | 4.2 | 3.2 | |

Appendix 28 Proportion of continuous benzodiazepine prescribing periods exceeding 30 days, by IMD quintile

| xy = 0 | TTTD qui | | | | | | | | | | |
|--------|---|------|------|------|-----------|--|--|--|--|--|--|
| | Proportion of benzodiazepine prescribing | | | | | | | | | | |
| | Index of multiple deprivation (IMD) quintiles | | | | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | | | | |
| 2000 | 47.4 | 49.2 | 52.4 | 51.0 | 53.7 | | | | | | |
| 2001 | 44.9 | 47.1 | 50.0 | 52.2 | 51.6 | | | | | | |
| 2002 | 43.0 | 39.6 | 51.4 | 47.9 | 49.7 | | | | | | |
| 2003 | 44.4 | 41.3 | 48.6 | 46.8 | 47.8 | | | | | | |
| 2004 | 38.3 | 40.8 | 44.8 | 47.0 | 50.6 | | | | | | |
| 2005 | 35.8 | 42.8 | 44.0 | 47.0 | 48.0 | | | | | | |
| 2006 | 43.9 | 40.8 | 43.5 | 41.8 | 43.9 | | | | | | |
| 2007 | 35.9 | 36.0 | 44.8 | 35.9 | 47.7 | | | | | | |
| 2008 | 33.2 | 36.4 | 40.0 | 37.7 | 43.4 | | | | | | |
| 2009 | 32.3 | 36.0 | 36.3 | 35.1 | 38.3 | | | | | | |
| 2010 | 33.0 | 31.8 | 35.1 | 34.3 | 38.9 | | | | | | |
| 2011 | 29.4 | 31.4 | 36.9 | 35.0 | 31.9 | | | | | | |
| 2012 | 31.7 | 30.2 | 40.2 | 37.7 | 31.2 | | | | | | |
| 2013 | 29.4 | 31.3 | 34.9 | 33.9 | 31.3 | | | | | | |
| 2014 | 28.1 | 27.3 | 36.0 | 33.4 | 39.5 | | | | | | |
| 2015 | 29.8 | 31.1 | 31.9 | 32.4 | 34.4 | | | | | | |

Appendix 29 Proportion of continuous benzodiazepine prescribing periods exceeding 6 months, by IMD quintile

| | Proportion of benzodiazepine prescribing | | | | | | | | |
|---|--|------|------|------|-----------|--|--|--|--|
| Index of multiple deprivation (IMD) quintiles | | | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | | |
| 2000 | 9.3 | 11.2 | 12.1 | 14.9 | 14.2 | | | | |
| 2001 | 9.5 | 11.3 | 11.8 | 13.9 | 16.9 | | | | |
| 2002 | 9.6 | 12.5 | 16.4 | 12.0 | 16.3 | | | | |
| 2003 | 11.9 | 11 | 12.9 | 13.7 | 17.2 | | | | |
| 2004 | 9.5 | 8.5 | 12.3 | 12.3 | 14.8 | | | | |
| 2005 | 8.9 | 10.7 | 14.5 | 13.7 | 17.3 | | | | |
| 2006 | 12.9 | 10.8 | 12.0 | 12.7 | 16.4 | | | | |
| 2007 | 8.3 | 9.7 | 13.6 | 10.8 | 17.4 | | | | |
| 2008 | 7.1 | 10.7 | 11.8 | 11.6 | 14.2 | | | | |
| 2009 | 9.5 | 11.0 | 11.1 | 11.2 | 10.3 | | | | |
| 2010 | 8.7 | 10.3 | 11.3 | 9.9 | 17.5 | | | | |
| 2011 | 10.5 | 10.7 | 8.7 | 13.3 | 10.1 | | | | |
| 2012 | 9.7 | 11.3 | 13.9 | 10.4 | 12.7 | | | | |
| 2013 | 7.3 | 9.7 | 10.0 | 10.8 | 12.5 | | | | |
| 2014 | 8.8 | 9.6 | 12.0 | 11.5 | 19.4 | | | | |
| 2015 | 7.9 | 6.6 | 7.2 | 12.9 | 11.5 | | | | |

Appendix 30 Proportion of continuous benzodiazepine prescribing periods exceeding 12 months, by IMD quintile

| Drangetion of honzadiazonina proceeding | | | | | | | | |
|---|-----------|-----|-----|-----|-----------|--|--|--|
| Proportion of benzodiazepine prescribing | | | | | | | | |
| Index of multiple deprivation (IMD) quintiles | | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | |
| 2000 | 3.0 | 3.5 | 4.1 | 3.5 | 5.2 | | | |
| 2001 | 3.0 | 5.5 | 6.0 | 6.1 | 6.5 | | | |
| 2002 | 4.1 | 5.8 | 8.0 | 6.9 | 7.4 | | | |
| 2003 | 5.8 | 4.8 | 5.9 | 6.5 | 7.8 | | | |
| 2004 | 4.1 | 3.4 | 5.6 | 5.9 | 8.6 | | | |
| 2005 | 3.1 | 3.6 | 7.7 | 6.1 | 9.7 | | | |
| 2006 | 4.9 | 4.0 | 6.4 | 6.3 | 7.9 | | | |
| 2007 | 3.4 | 4.2 | 6.7 | 5.2 | 8.8 | | | |
| 2008 | 2.9 | 5.2 | 5.8 | 4.5 | 8.2 | | | |
| 2009 | 4.2 | 5.3 | 6.0 | 5.1 | 5.0 | | | |
| 2010 | 3.3 | 4.0 | 6.4 | 5.7 | 9.0 | | | |
| 2011 | 4.3 | 4.9 | 3.1 | 5.8 | 7.3 | | | |
| 2012 | 2.7 | 6.5 | 5.4 | 4.5 | 6.9 | | | |
| 2013 | 4.1 | 3.4 | 4.5 | 5.3 | 5.6 | | | |
| 2014 | 3.3 | 3.5 | 4.9 | 7.5 | 10.6 | | | |
| 2015 | 3.7 | 3.4 | 2.9 | 6.2 | 6.3 | | | |

Appendix 31 Proportion of continuous Z-drugs prescribing periods exceeding 30 days, by IMD quintile

| | Proportion of Z-drug prescribing | | | | | | |
|------|---|------|------|------|--------------------|--|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | | |
| | | | | | | | |
| Year | Q1 (least deprived) | Q2 | Q3 | Q4 | Q5 (most deprived) | | |
| 2000 | 50.1 | 42.1 | 41.9 | 40.5 | 45 | | |
| 2001 | 39.3 | 49.3 | 44.5 | 40.9 | 39.9 | | |
| 2002 | 41.7 | 48.6 | 43.4 | 44.6 | 40.9 | | |
| 2003 | 38.8 | 48.1 | 44.6 | 34.8 | 43.3 | | |
| 2004 | 42.4 | 43.8 | 43.8 | 42.6 | 47.5 | | |
| 2005 | 42.2 | 41.2 | 47.1 | 48.6 | 34.1 | | |
| 2006 | 41.7 | 43.9 | 43.8 | 41.0 | 40.7 | | |
| 2007 | 34.8 | 44.1 | 34.6 | 45.1 | 44.3 | | |
| 2008 | 44.7 | 41.9 | 42.8 | 37.5 | 41.1 | | |
| 2009 | 43.1 | 40.3 | 44.8 | 40.9 | 46.3 | | |
| 2010 | 37.7 | 41.0 | 39.4 | 45.0 | 41.7 | | |
| 2011 | 42.0 | 42.1 | 36.9 | 43.8 | 44.6 | | |
| 2012 | 43.7 | 42.8 | 40.0 | 47.7 | 43.6 | | |
| 2013 | 37.5 | 48.1 | 39.3 | 35.9 | 37.2 | | |
| 2014 | 42.2 | 37.5 | 36.8 | 29.5 | 38.2 | | |
| 2015 | 40.3 | 39 | 37.5 | 33.9 | 32.7 | | |

Appendix 32 Proportion of continuous Z-drugs prescribing periods exceeding 6 months, by IMD quintile

| Proportion of Z-drug prescribing | | | | | | | | |
|----------------------------------|---|------|------|------|-----------|--|--|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | | | |
| | | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | |
| 2000 | 6.3 | 8.8 | 9.1 | 7.1 | 7.8 | | | |
| 2001 | 9.3 | 17.9 | 15.3 | 8.5 | 10.8 | | | |
| 2002 | 11.3 | 11.2 | 12.8 | 9.3 | 9.4 | | | |
| 2003 | 8.5 | 7.7 | 10.6 | 8.5 | 14.4 | | | |
| 2004 | 9.4 | 9.6 | 10.3 | 10.8 | 17.3 | | | |
| 2005 | 8.6 | 12.5 | 11.7 | 14.1 | 12.7 | | | |
| 2006 | 9.3 | 11.7 | 15.8 | 9.9 | 11.7 | | | |
| 2007 | 12.3 | 12.2 | 12.3 | 12.2 | 14.7 | | | |
| 2008 | 10.4 | 9.7 | 11.8 | 10.9 | 13.3 | | | |
| 2009 | 11.9 | 11.8 | 11.4 | 11.1 | 13.8 | | | |
| 2010 | 9.4 | 10.7 | 9.4 | 12.7 | 11.8 | | | |
| 2011 | 12.3 | 9.2 | 10.2 | 15.5 | 18.0 | | | |
| 2012 | 14.1 | 10.9 | 13.3 | 14.2 | 15.1 | | | |
| 2013 | 9.0 | 12.4 | 10.8 | 12.4 | 17.3 | | | |
| 2014 | 13.0 | 13.1 | 9.6 | 11.8 | 17.5 | | | |
| 2015 | 11.4 | 11.5 | 14.2 | 14.0 | 8.4 | | | |

Appendix 33 Proportion of continuous Z-drugs prescribing periods exceeding 12 months, by IMD quintile

| | Proportion of Z-drug prescribing | | | | | | |
|------|---|-----|-----|-----|-----------|--|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | | |
| | | | | | | | |
| | Q1 (least | | | | Q5 (most | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | |
| 2000 | 2.4 | 4.0 | 1.6 | 3.2 | 2.8 | | |
| 2001 | 4.8 | 7.5 | 5.2 | 3.0 | 4.7 | | |
| 2002 | 3.9 | 5.1 | 5.0 | 5.6 | 3.9 | | |
| 2003 | 3.3 | 3.2 | 6.2 | 4.2 | 6.6 | | |
| 2004 | 4.0 | 3.6 | 2.9 | 4.0 | 6.0 | | |
| 2005 | 4.2 | 6.6 | 5.1 | 5.8 | 5.3 | | |
| 2006 | 4.0 | 3.6 | 8.9 | 4.7 | 4.4 | | |
| 2007 | 3.7 | 5.3 | 3.9 | 5.1 | 8.1 | | |
| 2008 | 4.7 | 3.5 | 6.2 | 3.2 | 6.7 | | |
| 2009 | 5.7 | 5.3 | 6.3 | 3.8 | 6.5 | | |
| 2010 | 3.4 | 3.2 | 3.4 | 6.6 | 5.7 | | |
| 2011 | 5.0 | 3.6 | 5.8 | 7.2 | 10.1 | | |
| 2012 | 5.9 | 5.0 | 7.1 | 6.4 | 7.2 | | |
| 2013 | 4.0 | 5.8 | 4.5 | 5.1 | 11.6 | | |
| 2014 | 6.2 | 6.0 | 5.4 | 4.7 | 7.9 | | |
| 2015 | 3.3 | 4.3 | 8.4 | 8.1 | 3.8 | | |

Appendix 34 Proportion of continuous opioids prescribing periods exceeding 30 days, by IMD quintile

| | Proportion of opioid prescribing | | | | | | | |
|------|---|------|-------|------|-----------|--|--|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | |
| 2000 | 41.9 | 37.0 | 41.92 | 32.4 | 34.6 | | | |
| 2001 | 38.7 | 41.1 | 42.8 | 34.6 | 36.3 | | | |
| 2002 | 35.6 | 40.4 | 41.1 | 34.5 | 37.6 | | | |
| 2003 | 34.7 | 37.4 | 41.5 | 34.0 | 38.5 | | | |
| 2004 | 35.4 | 39.3 | 40.3 | 41.6 | 35.6 | | | |
| 2005 | 34.5 | 37.9 | 38.8 | 39.9 | 36.5 | | | |
| 2006 | 35.3 | 37.7 | 42.1 | 35.5 | 35.9 | | | |
| 2007 | 37.8 | 35.4 | 38.8 | 36.6 | 35.9 | | | |
| 2008 | 37.1 | 39.1 | 40.2 | 35.3 | 34.0 | | | |
| 2009 | 37.5 | 39.4 | 39.4 | 36.0 | 34.2 | | | |
| 2010 | 37.8 | 38.8 | 39.5 | 35.9 | 38.2 | | | |
| 2011 | 37.0 | 38.4 | 34.8 | 36.4 | 36.7 | | | |
| 2012 | 33.3 | 36.9 | 35.2 | 36.7 | 36.4 | | | |
| 2013 | 30.6 | 35.6 | 33.5 | 34.5 | 35.4 | | | |
| 2014 | 33.6 | 34.1 | 31.1 | 36.3 | 32.7 | | | |
| 2015 | 33.8 | 35.0 | 28.6 | 33.3 | 31.0 | | | |

Appendix 35 Proportion of continuous opioids prescribing periods exceeding 6 months, by IMD quintile

| Proportion of Opioid prescribing | | | | | | | | |
|----------------------------------|---|------|-----|-----|-----------|--|--|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | |
| 2000 | 6.6 | 5.2 | 7.7 | 4.8 | 4.7 | | | |
| 2001 | 5.9 | 8.4 | 8.1 | 5.7 | 7.5 | | | |
| 2002 | 4.3 | 6.6 | 4.2 | 5.9 | 6.8 | | | |
| 2003 | 6.6 | 5.8 | 7.4 | 6.5 | 7.8 | | | |
| 2004 | 5.4 | 7.6 | 6.2 | 5.9 | 8.6 | | | |
| 2005 | 5.5 | 7.4 | 7.6 | 6.5 | 9.4 | | | |
| 2006 | 6.6 | 6.6 | 8.3 | 7.0 | 7.3 | | | |
| 2007 | 6.8 | 6.5 | 7.3 | 6.3 | 7.3 | | | |
| 2008 | 5.0 | 7.6 | 7.1 | 6.6 | 8.3 | | | |
| 2009 | 6.5 | 10.2 | 9.5 | 6.9 | 7.7 | | | |
| 2010 | 6.2 | 7.7 | 7.9 | 7.4 | 7.9 | | | |
| 2011 | 6.4 | 8.0 | 6.2 | 7.7 | 8.7 | | | |
| 2012 | 6.2 | 7.7 | 7.4 | 7.5 | 8.3 | | | |
| 2013 | 5.0 | 7.6 | 6.7 | 8.2 | 8.5 | | | |
| 2014 | 4.6 | 7.4 | 7.1 | 8.1 | 8.3 | | | |
| 2015 | 6.5 | 8.4 | 5.8 | 5.6 | 6.8 | | | |

Appendix 36 Proportion of continuous opioids prescribing periods exceeding 12 months, by IMD quintile

| | Proportion of opioid prescribing | | | | | | |
|---|----------------------------------|-----|-----|-----|-----------|--|--|
| Index of multiple deprivation (IMD) quintiles | | | | | | | |
| | Q1 (least | | | | Q5 (most | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | |
| 2000 | 2.9 | 2.0 | 2.2 | 1.4 | 1.1 | | |
| 2001 | 1.5 | 3.4 | 3.1 | 2.5 | 1.8 | | |
| 2002 | 1.4 | 3.2 | 2.0 | 2.5 | 2.7 | | |
| 2003 | 3.4 | 1.3 | 2.7 | 3.1 | 4.6 | | |
| 2004 | 1.4 | 2.5 | 3.0 | 1.5 | 3.5 | | |
| 2005 | 2.3 | 1.8 | 3.4 | 3.0 | 3.3 | | |
| 2006 | 3.2 | 3.1 | 2.9 | 2.8 | 2.9 | | |
| 2007 | 2.5 | 2.6 | 2.4 | 2.7 | 2.9 | | |
| 2008 | 1.5 | 2.8 | 2.9 | 2.4 | 3.8 | | |
| 2009 | 1.9 | 3.4 | 4.4 | 2.9 | 3.6 | | |
| 2010 | 1.4 | 2.8 | 3.3 | 3.4 | 3.6 | | |
| 2011 | 1.7 | 3.5 | 2.0 | 3.0 | 4.8 | | |
| 2012 | 2.5 | 3.1 | 3.1 | 3.9 | 3.9 | | |
| 2013 | 2.1 | 4.2 | 2.2 | 3.8 | 3.5 | | |
| 2014 | 0.8 | 4.6 | 3.6 | 4.2 | 3.9 | | |
| 2015 | 1.4 | 2.6 | 1.5 | 1.5 | 2.1 | | |

Appendix 37 Proportion of continuous GABAergic medicines prescribing periods exceeding 30 days, by IMD quintile

| Р | roportion of | GABAergi | c medicines | s prescribin | g | | | |
|------|---|----------|-------------|--------------|-----------|--|--|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | |
| 2000 | 39.2 | 31.2 | 72.2 | 64.1 | 65.6 | | | |
| 2001 | 56.2 | 47.7 | 63.3 | 65.7 | 57.0 | | | |
| 2002 | 58.4 | 66.0 | 63.0 | 58.0 | 66.5 | | | |
| 2003 | 66.8 | 74.1 | 70.7 | 66.7 | 68.3 | | | |
| 2004 | 71.3 | 69.8 | 68.8 | 67.0 | 60.6 | | | |
| 2005 | 71.6 | 66.5 | 80.3 | 68.1 | 68.4 | | | |
| 2006 | 65.5 | 62.4 | 67.3 | 65.8 | 66.8 | | | |
| 2007 | 68.5 | 63.8 | 62.5 | 63.0 | 77.5 | | | |
| 2008 | 65.1 | 68.5 | 68.8 | 64.9 | 69.8 | | | |
| 2009 | 72.9 | 66.1 | 70.5 | 68.2 | 62.8 | | | |
| 2010 | 70.5 | 71.3 | 69.7 | 72.0 | 63.1 | | | |
| 2011 | 62.9 | 67.3 | 64.6 | 69.3 | 59.1 | | | |
| 2012 | 61.9 | 62.6 | 59.5 | 64.8 | 58.8 | | | |
| 2013 | 58.2 | 58.1 | 63.4 | 59.9 | 53.3 | | | |
| 2014 | 56.5 | 54.6 | 62.8 | 59.6 | 56.3 | | | |
| 2015 | 54.1 | 56.9 | 56.0 | 57.4 | 56.7 | | | |

Appendix 38 Proportion of continuous GABAergic medicines prescribing periods exceeding 6 months, by IMD quintile

| Proportion of GABAergic medicines prescribing | | | | | | | | |
|---|---|------|------|------|-----------|--|--|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | |
| 2000 | 16.7 | 7.9 | 8.2 | 7.8 | 7.8 | | | |
| 2001 | 15.4 | 8.8 | 8.4 | 12.5 | 7.9 | | | |
| 2002 | 12.2 | 19.6 | 6.8 | 10.4 | 12.9 | | | |
| 2003 | 13.9 | 16.1 | 16.5 | 10.3 | 5.5 | | | |
| 2004 | 12.5 | 11.3 | 13.2 | 15.0 | 6.1 | | | |
| 2005 | 14.1 | 7.0 | 10.2 | 15.4 | 10.0 | | | |
| 2006 | 12.2 | 10.9 | 11.9 | 12.6 | 19.7 | | | |
| 2007 | 17.0 | 8.4 | 11.1 | 14.6 | 13.7 | | | |
| 2008 | 14.8 | 10.6 | 16.5 | 13.7 | 12.3 | | | |
| 2009 | 15.4 | 14.4 | 15.3 | 13.1 | 9.4 | | | |
| 2010 | 15.4 | 18.3 | 14.2 | 15.5 | 9.7 | | | |
| 2011 | 11.6 | 15.0 | 14.3 | 14.5 | 13.1 | | | |
| 2012 | 13.3 | 11.9 | 13.6 | 15.3 | 13.6 | | | |
| 2013 | 11.9 | 16.7 | 12 | 17.5 | 12.3 | | | |
| 2014 | 14.5 | 11.9 | 12.8 | 16.1 | 14.9 | | | |
| 2015 | 8.5 | 13.8 | 12.6 | 10.3 | 14.5 | | | |

Appendix 39 Proportion of continuous GABAergic medicines prescribing periods exceeding 12 months, by IMD quintile

| Pı | roportion of | GABAergi | c medicine | s prescribir | ng | | | |
|------|---|----------|------------|--------------|-----------|--|--|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | | | |
| | Q1 (least | | | | Q5 (most | | | |
| Year | deprived) | Q2 | Q3 | Q4 | deprived) | | | |
| 2000 | 0.0 | 2.0 | 3.0 | 4.0 | 7.8 | | | |
| 2001 | 6.2 | 3.5 | 4.2 | 8.9 | 2.3 | | | |
| 2002 | 1.7 | 6.4 | 3.2 | 1.3 | 4.7 | | | |
| 2003 | 5.4 | 3.6 | 5.3 | 1.9 | 4.9 | | | |
| 2004 | 0.5 | 2.4 | 4.4 | 3.7 | 3.5 | | | |
| 2005 | 2.2 | 0.7 | 4.4 | 5.6 | 2.6 | | | |
| 2006 | 6.3 | 4.3 | 4.3 | 4.3 | 5.8 | | | |
| 2007 | 5.5 | 2.2 | 3.1 | 4.2 | 2.9 | | | |
| 2008 | 4.9 | 3.2 | 5.4 | 4.1 | 4.3 | | | |
| 2009 | 3.5 | 5.5 | 4.5 | 5.8 | 3.5 | | | |
| 2010 | 6.7 | 7.0 | 4.2 | 5.7 | 5.0 | | | |
| 2011 | 2.9 | 7.0 | 5.7 | 4.8 | 4.7 | | | |
| 2012 | 4.6 | 3.6 | 2.2 | 4.9 | 6.4 | | | |
| 2013 | 5.7 | 7.0 | 4.4 | 7.1 | 3.8 | | | |
| 2014 | 3.8 | 5.7 | 3.2 | 5.9 | 6.2 | | | |
| 2015 | 1.7 | 4.7 | 3.1 | 2.7 | 5.8 | | | |

Appendix 40 Characteristics of people prescribed benzodiazepines in 2014, by continuous prescribing period length

| Benzodiazepines | | | | | | | |
|----------------------|-------------------|------------------|-------------------|-------------------|--|--|--|
| | Up to 3 months | 3 to 6 months | 6 to 12 months | Over 12 months | | | |
| Mean age (years) | 61.0*** | 63.8* | 65.6*** | 62.3 | | | |
| IMD quintile (%) | | | | | | | |
| I (least deprived) § | 57.5 | 6.0 | 6.0 | 3.3 | | | |
| | 56.3 | 6.0 | 6.8 | 3.5 | | | |
| III | 43.6*** | 9.3** | 8.3* | 4.9 | | | |
| IV | 49.7*** | 11.7*** | 4.5 | 7.5*** | | | |
| V (most deprived | 43.0*** | 12.0*** | 9.4** | 10.6*** | | | |
| Gender (%) | | | | | | | |
| Male § | 51.6 | 10.0 | 6.1 | 7.3 | | | |
| Female | 49.1 | 8.4* | 7.5 | 5.1*** | | | |
| Region (%) | | | | | | | |
| North of England § | 46.6 | 10.1 | 6.7 | 9.7 | | | |
| Midlands and East of | | | | | | | |
| England | 54.5*** | 10.2 | 6.9 | 4.8*** | | | |
| London | 51.9* | 3.7*** | 3.4** | 7.3 | | | |
| South East | 53.3*** | 9.6 | 9.2** | 3.7*** | | | |
| South West | 35.6*** | 6.9* | 5.1 | 5.0*** | | | |

^{*} P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, § - reference category

Appendix 41 Characteristics of people prescribed Z-drugs in 2014, by continuous prescribing period length

| Z-drugs | | | | | | |
|----------------------|-------------------|------------------|-------------------|-------------------|--|--|
| | Up to 3 months | 3 to 6 months | 6 to 12 months | Over 12 months | | |
| Mean age (years) | 60.4*** | 65.0*** | 68.1*** | 67.5*** | | |
| IMD quintile (%) | | | | | | |
| I (least deprived) § | 61.7 | 11.8 | 7.2 | 6.2 | | |
| II | 67.1* | 9.1 | 7.6 | 6.0 | | |
| III | 64.1 | 9.1 | 5.0* | 5.4 | | |
| IV | 67.8** | 6.3*** | 7.8 | 4.8 | | |
| V (most deprived | 52.7*** | 3.6*** | 11.3** | 8.0 | | |
| Gender (%) | | | | | | |
| Male § | 27.5 | 16.5 | 14.8 | 7.7 | | |
| Female | 26.9** | 17.4* | 15.3 | 5.1** | | |
| Region (%) | | | | | | |
| North of England § | 61.6 | 7.0 | 9.1 | 9.5 | | |
| Midlands and East of | | | | | | |
| England | 68.8*** | 7.3 | 7.2 | 4.7*** | | |
| London | 71.6*** | 6.6 | 7.9 | 5.6* | | |
| South East | 62.7 | 11.2*** | 7.0 | 5.4*** | | |
| South West | 51.8*** | 5.2 | 7.5 | 4.2*** | | |

^{*} $P \le 0.05$, ** $P \le 0.01$, *** $P \le 0.001$, § - reference category

Appendix 42 Characteristics of people prescribed opioids in 2014, by continuous prescribing period length

| Opioids | | | | | | |
|----------------------|-------------------|------------------|-------------------|-------------------|--|--|
| | Up to 3 months | 3 to 6 months | 6 to 12 months | Over 12 months | | |
| Mean age (years) | 61.6*** | 63.3 | 63.8 | 63.1 | | |
| IMD quintile (%) | | | | | | |
| I (least deprived) § | 52.1 | 10.4 | 4.2 | 0.9 | | |
| II | 45.4*** | 10.6 | 3.5 | 4.6*** | | |
| III | 51.8 | 8.1** | 3.8 | 3.6*** | | |
| IV | 55.0* | 9.6 | 4.4 | 4.*** | | |
| V (most deprived | 50.5 | 7.3*** | 5.1 | 3.9*** | | |
| Gender (%) | | | | | | |
| Male § | 52.3 | 8.7 | 3.6 | 4.2 | | |
| Female | 49.7*** | 9.4 | 4.6** | 3.3** | | |
| Region (%) | | | | | | |
| North of England § | 53.6 | 8.8 | 5.3 | 4.7 | | |
| Midlands and East of | | | | | | |
| England | 52.9 | 9.9 | 3.1*** | 3.6* | | |
| London | 52.4 | 10.1 | 3.8* | 2.2*** | | |
| South East | 46.0*** | 8.9 | 4.0** | 3.3*** | | |
| South West | 49.9** | 8.3 | 3.8* | 2.0*** | | |

^{*} P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, § - reference category

Appendix 43 Characteristics of people prescribed GABAergic medicines in 2014, by continuous prescribing period length

| GABAergic medicines | | | | | |
|----------------------|-------------------|------------------|-------------------|-------------------|--|
| | Up to 3 months | 3 to 6 months | 6 to 12 months | Over 12 months | |
| Mean age (years) | 59.7 | 60.1 | 63.7*** | 61.1 | |
| IMD quintile (%) | | | | | |
| I (least deprived) § | 41.0 | 10.7 | 13.2 | 3.8 | |
| II | 38.1 | 17.7*** | 7.3*** | 5.6* | |
| III | 44.0 | 21.1*** | 12.5 | 3.2 | |
| IV | 41.5 | 19.1*** | 11.6 | 5.9* | |
| V (most deprived | 47.5*** | 12.0 | 10.0** | 6.2** | |
| Gender (%) | | | | | |
| Male § | 43.5 | 15.1 | 8.4 | 6.9 | |
| Female | 42.2 | 16.8 | 12.2*** | 3.9*** | |
| Region (%) | | | | | |
| North of England § | 44.9 | 14.2 | 12.2 | 4.8 | |
| Midlands and East of | | | | | |
| England | 43.1 | 16.0 | 9.6* | 7.4*** | |
| London | 43.2 | 13.7 | 11.7 | 2.2* | |
| South East | 41.3* | 16.8* | 11.0 | 4.4 | |
| South West | 37.4*** | 23.9*** | 7.3*** | 4.8 | |

^{*} P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, § - reference category

Appendix 44 Proportion of continuous prescribing periods exceeding 30 days by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 55.5 | - |
| 2001 | 68.5 | - |
| 2002 | 68.2 | - |
| 2003 | 72.6 | - |
| 2004 | 71.5 | 60.0 |
| 2005 | 75.5 | 56.7 |
| 2006 | 73.4 | 54.6 |
| 2007 | 72.7 | 59.5 |
| 2008 | 74.0 | 62.6 |
| 2009 | 74.8 | 63.9 |
| 2010 | 75.7 | 60.8 |
| 2011 | 71.3 | 58.5 |
| 2012 | 68.7 | 57.3 |
| 2013 | 62.2 | 57.6 |
| 2014 | 59.8 | 62.7 |
| 2015 | 58.3 | 61.8 |

Appendix 45 Proportion of continuous prescribing periods exceeding 6 months by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 9.2 | - |
| 2001 | 9.6 | - |
| 2002 | 8.0 | - |
| 2003 | 12.3 | - |
| 2004 | 8.7 | 6.7 |
| 2005 | 11.9 | 3.0 |
| 2006 | 11.5 | 11.4 |
| 2007 | 10.5 | 11.3 |
| 2008 | 11.3 | 12.2 |
| 2009 | 12.7 | 13.2 |
| 2010 | 11.7 | 12.0 |
| 2011 | 9.1 | 14.1 |
| 2012 | 11.3 | 13.7 |
| 2013 | 11.5 | 15.9 |
| 2014 | 12.3 | 16.0 |
| 2015 | 9.2 | 13.8 |

Appendix 46 Proportion of continuous prescribing periods exceeding 12 months by GABAergic medicine type

| Year | Gabapentin | Pregabalin |
|------|------------|------------|
| 2000 | 0.0 | - |
| 2001 | 4.1 | - |
| 2002 | 2.6 | - |
| 2003 | 4.0 | - |
| 2004 | 2.6 | 0.0 |
| 2005 | 4.3 | 0.8 |
| 2006 | 3.1 | 4.0 |
| 2007 | 3.1 | 2.8 |
| 2008 | 4.0 | 3.8 |
| 2009 | 3.8 | 4.5 |
| 2010 | 3.9 | 4.7 |
| 2011 | 3.1 | 5.7 |
| 2012 | 3.4 | 4.5 |
| 2013 | 4.0 | 6.1 |
| 2014 | 4.2 | 7.7 |
| 2015 | 1.7 | 5.2 |

Appendix 47 Gabapentin: Proportion of continuous prescribing periods exceeding 30 days, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 32.0 | 67.5 | 54.3 | 75.0 |
| 2001 | 61.8 | 78.3 | 61.3 | 70.6 |
| 2002 | 57.4 | 74.7 | 76.1 | 41.9 |
| 2003 | 54.1 | 78.0 | 75.8 | 68.4 |
| 2004 | 62.2 | 73.9 | 72.4 | 75.5 |
| 2005 | 66.2 | 76.4 | 78.5 | 73.2 |
| 2006 | 64.0 | 77.8 | 73.5 | 74.6 |
| 2007 | 67.5 | 75.8 | 71.0 | 73.5 |
| 2008 | 76.4 | 71.9 | 78.1 | 63.0 |
| 2009 | 70.0 | 71.7 | 80.1 | 69.5 |
| 2010 | 66.9 | 76.6 | 79.2 | 71.6 |
| 2011 | 66.7 | 71.6 | 73.5 | 67.3 |
| 2012 | 60.1 | 70.2 | 71.3 | 63.1 |
| 2013 | 58.5 | 64.6 | 61.0 | 62.6 |
| 2014 | 48.9 | 63.5 | 62.9 | 53.2 |
| 2015 | 53.2 | 55.4 | 63.0 | 58.4 |

Appendix 48 Gabapentin: Proportion of continuous prescribing periods exceeding 6 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2000 | 4.0 | 17.5 | 6.5 | 0.0 |
| 2001 | 2.9 | 10.1 | 10.7 | 11.8 |
| 2002 | 5.9 | 8.4 | 9.4 | 6.5 |
| 2003 | 9.8 | 10.1 | 15.8 | 10.5 |
| 2004 | 8.5 | 4.9 | 11.4 | 13.2 |
| 2005 | 11.7 | 9.3 | 14.6 | 10.7 |
| 2006 | 17.4 | 8.0 | 11.5 | 15.3 |
| 2007 | 9.1 | 11.0 | 8.8 | 15.7 |
| 2008 | 5.5 | 11.4 | 10.8 | 18.0 |
| 2009 | 9.0 | 12.2 | 11.8 | 23.2 |
| 2010 | 5.2 | 11.9 | 12.4 | 17.4 |
| 2011 | 7.4 | 6.0 | 11.1 | 14.5 |
| 2012 | 9.1 | 8.1 | 14.1 | 14.9 |
| 2013 | 9.4 | 10.8 | 11.0 | 17.4 |
| 2014 | 6.8 | 11.7 | 13.5 | 16.9 |
| 2015 | 12.1 | 8.0 | 8.5 | 12.9 |

Appendix 49 Gabapentin: Proportion of continuous prescribing periods exceeding 12 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|-----|
| 2000 | 0.0 | 0.0 | 0.0 | 0.0 |
| 2001 | 2.9 | 4.3 | 4.0 | 0.0 |
| 2002 | 2.9 | 4.2 | 0.9 | 3.2 |
| 2003 | 3.3 | 3.1 | 4.2 | 7.9 |
| 2004 | 1.2 | 0.5 | 5.4 | 1.9 |
| 2005 | 7.8 | 2.7 | 5.3 | 1.8 |
| 2006 | 3.5 | 2.2 | 2.3 | 8.5 |
| 2007 | 2.6 | 2.9 | 2.8 | 4.8 |
| 2008 | 1.8 | 4.7 | 3.6 | 6.0 |
| 2009 | 2.0 | 3.3 | 3.9 | 7.3 |
| 2010 | 2.6 | 5.6 | 2.6 | 5.5 |
| 2011 | 0.0 | 3.3 | 2.5 | 9.1 |
| 2012 | 3.5 | 2.9 | 3.4 | 5.0 |
| 2013 | 1.9 | 3.3 | 4.7 | 6.1 |
| 2014 | 1.5 | 4.2 | 4.3 | 7.3 |
| 2015 | 0.8 | 1.9 | 2.0 | 1.0 |

Appendix 50 Pregabalin: Proportion of continuous prescribing periods exceeding 30 days, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2004 | 100.0 | 33.3 | 83.3 | 50.0 |
| 2005 | 47.1 | 71.4 | 52.6 | 27.3 |
| 2006 | 62.2 | 61.0 | 46.0 | 44.4 |
| 2007 | 55.0 | 57.4 | 68.5 | 37.0 |
| 2008 | 60.0 | 54.2 | 72.7 | 74.3 |
| 2009 | 67.9 | 58.1 | 70.4 | 57.1 |
| 2010 | 58.6 | 59.2 | 63.8 | 62.2 |
| 2011 | 56.4 | 58.2 | 60.2 | 59.0 |
| 2012 | 47.1 | 56.1 | 62.4 | 64.7 |
| 2013 | 47.1 | 55.6 | 63.6 | 60.9 |
| 2014 | 48.1 | 62.2 | 66.7 | 75.0 |
| 2015 | 61.6 | 61.7 | 63.0 | 65.5 |

Appendix 51 Pregabalin: Proportion of continuous prescribing periods exceeding 6 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2004 | 0.0 | 0.0 | 16.7 | 0.0 |
| 2005 | 5.9 | 4.1 | 1.8 | 0.0 |
| 2006 | 13.5 | 10.2 | 13.0 | 5.6 |
| 2007 | 13.3 | 9.2 | 12.9 | 11.1 |
| 2008 | 10.0 | 10.7 | 16.5 | 8.6 |
| 2009 | 4.9 | 14.7 | 14.8 | 14.3 |
| 2010 | 5.4 | 14.3 | 12.1 | 16.2 |
| 2011 | 9.8 | 13.0 | 17.2 | 16.4 |
| 2012 | 15.9 | 9.8 | 16.4 | 16.2 |
| 2013 | 8.6 | 17.3 | 17.8 | 16.1 |
| 2014 | 13.7 | 14.0 | 18.2 | 19.7 |
| 2015 | 13.4 | 14.4 | 14.0 | 13.8 |

Appendix 52 Pregabalin: Proportion of continuous prescribing periods exceeding 12 months, by age

| Year | 18-40 | 41-60 | 61-80 | 81+ |
|------|-------|-------|-------|------|
| 2004 | 0.0 | 0.0 | 0.0 | 0.0 |
| 2005 | 0.0 | 0.0 | 1.8 | 0.0 |
| 2006 | 8.1 | 2.5 | 5.0 | 0.0 |
| 2007 | 5.0 | 1.4 | 4.0 | 0.0 |
| 2008 | 0.0 | 3.4 | 5.8 | 5.7 |
| 2009 | 1.2 | 4.1 | 5.6 | 7.1 |
| 2010 | 0.0 | 7.2 | 4.3 | 5.4 |
| 2011 | 3.8 | 4.5 | 7.3 | 9.8 |
| 2012 | 5.8 | 1.8 | 5.2 | 11.8 |
| 2013 | 5.0 | 7.8 | 4.4 | 8.0 |
| 2014 | 8.4 | 6.5 | 8.3 | 9.2 |
| 2015 | 2.7 | 6.6 | 4.3 | 8.6 |

Appendix 53 Proportion of continuous prescribing periods exceeding 30 days, by GABAergic medicine type and gender

| | Gaba | pentin | Prega | abalin |
|-------|------|--------|-------|--------|
| Years | Male | Female | Male | Female |
| 2000 | 52.6 | 58.1 | - | - |
| 2001 | 68.2 | 68.8 | - | - |
| 2002 | 70.1 | 67.0 | - | - |
| 2003 | 78.6 | 69.7 | - | - |
| 2004 | 74.3 | 69.8 | 75.0 | 54.5 |
| 2005 | 79.7 | 72.9 | 46.9 | 62.4 |
| 2006 | 79.8 | 69.2 | 55.8 | 54.1 |
| 2007 | 77.8 | 69.6 | 65.5 | 56.7 |
| 2008 | 77.0 | 72.3 | 65.7 | 61.0 |
| 2009 | 73.7 | 75.4 | 66.5 | 62.7 |
| 2010 | 79.1 | 73.3 | 59.0 | 61.8 |
| 2011 | 75.3 | 68.3 | 57.4 | 59.1 |
| 2012 | 70.6 | 67.4 | 55.5 | 58.5 |
| 2013 | 65.5 | 60.1 | 50.3 | 61.6 |
| 2014 | 59.9 | 59.8 | 60.2 | 64.0 |
| 2015 | 59.9 | 57.4 | 62.2 | 61.5 |

Appendix 54 Proportion of continuous prescribing periods exceeding 6 months, by GABAergic medicine type and gender

| | Gabar | pentin | Prega | abalin |
|-------|-------|--------|-------|--------|
| Years | Male | Female | Male | Female |
| 2000 | 3.5 | 14.5 | - | - |
| 2001 | 12.9 | 7.1 | - | - |
| 2002 | 11.1 | 6.2 | - | - |
| 2003 | 17.1 | 9.9 | - | - |
| 2004 | 10.7 | 7.5 | 0.0 | 9.1 |
| 2005 | 13.0 | 11.2 | 2.0 | 3.5 |
| 2006 | 11.9 | 11.2 | 14.3 | 10.2 |
| 2007 | 10.4 | 10.6 | 13.3 | 10.4 |
| 2008 | 12.6 | 10.5 | 16.4 | 10.0 |
| 2009 | 11.9 | 13.2 | 16.5 | 11.6 |
| 2010 | 11.3 | 11.9 | 12.1 | 11.8 |
| 2011 | 9.6 | 8.7 | 13.1 | 14.6 |
| 2012 | 11.0 | 11.5 | 12.0 | 14.8 |
| 2013 | 10.1 | 12.4 | 15.0 | 16.4 |
| 2014 | 10.3 | 13.5 | 19.9 | 13.9 |
| 2015 | 9.9 | 8.7 | 14.2 | 13.6 |

Appendix 55 Proportion of continuous prescribing periods exceeding 12 months, by GABAergic medicine type and gender

| | Gabar | pentin | Pregabalin | | |
|-------|-------|--------|------------|--------|--|
| Years | Male | Female | Male | Female | |
| 2000 | 0.0 | 0.0 | - | - | |
| 2001 | 5.9 | 2.7 | - | - | |
| 2002 | 5.1 | 1.0 | - | - | |
| 2003 | 5.7 | 3.2 | - | - | |
| 2004 | 2.1 | 2.8 | 0.0 | 0.0 | |
| 2005 | 5.2 | 3.7 | 0.0 | 1.2 | |
| 2006 | 5.1 | 1.8 | 2.6 | 4.6 | |
| 2007 | 3.1 | 3.0 | 3.5 | 2.5 | |
| 2008 | 4.4 | 3.8 | 6.0 | 2.7 | |
| 2009 | 3.1 | 4.2 | 6.6 | 3.4 | |
| 2010 | 4.4 | 3.5 | 5.1 | 4.5 | |
| 2011 | 2.8 | 3.4 | 3.9 | 6.8 | |
| 2012 | 2.9 | 3.7 | 3.5 | 5.1 | |
| 2013 | 3.6 | 4.3 | 5.8 | 6.3 | |
| 2014 | 4.5 | 4.1 | 13.0 | 4.9 | |
| 2015 | 2.4 | 1.3 | 6.9 | 4.2 | |

Appendix 56 Proportion of continuous gabapentin prescribing periods exceeding 30 days, by IMD quintile

| Proportion of gabapentin prescribing | | | | | | |
|--------------------------------------|---|------|------|------|------|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | |
| Year | Q1 | Q2 | Q3 | Q4 | Q5 | |
| 2000 | 47.4 | 42.9 | 61.3 | 61.5 | 59.1 | |
| 2001 | 72.0 | 61.4 | 71.4 | 77.1 | 60.5 | |
| 2002 | 64.9 | 71.4 | 70.8 | 63.4 | 70.9 | |
| 2003 | 68.8 | 80.2 | 73.5 | 67.3 | 73.7 | |
| 2004 | 76.3 | 72.1 | 76.1 | 70.7 | 64.6 | |
| 2005 | 67.7 | 80.6 | 81.2 | 72.9 | 73.9 | |
| 2006 | 75.7 | 66.0 | 78.3 | 78.7 | 70.2 | |
| 2007 | 69.6 | 71.0 | 69.9 | 75.5 | 76.4 | |
| 2008 | 70.0 | 73.3 | 78.8 | 73.0 | 75.1 | |
| 2009 | 78.4 | 74.5 | 74.9 | 77.2 | 69.6 | |
| 2010 | 81.6 | 78.7 | 76.7 | 80.8 | 64.5 | |
| 2011 | 74.8 | 71.8 | 72.9 | 77.3 | 60.3 | |
| 2012 | 71.1 | 67.5 | 69.5 | 70.6 | 66.1 | |
| 2013 | 70.9 | 60.8 | 68.3 | 61.4 | 52.6 | |
| 2014 | 63.3 | 54.6 | 59.3 | 66.1 | 57.5 | |
| 2015 | 61.0 | 58.2 | 57.0 | 58.4 | 57.5 | |

Appendix 57 Proportion of continuous gabapentin prescribing periods exceeding 6 months, by IMD quintile

| | Proportion of gabapentin prescribing | | | | | |
|------|---|------|------|------|------|--|
| | Index of multiple deprivation (IMD) quintiles | | | | | |
| Year | Q1 | Q2 | Q3 | Q4 | Q5 | |
| 2000 | 15.8 | 9.5 | 9.7 | 11.5 | 0.0 | |
| 2001 | 4.0 | 9.1 | 11.9 | 12.5 | 7.9 | |
| 2002 | 5.3 | 14.3 | 4.6 | 8.5 | 7.3 | |
| 2003 | 8.8 | 15.6 | 16.2 | 12.5 | 7.9 | |
| 2004 | 12.5 | 7.7 | 8.7 | 12.9 | 2.7 | |
| 2005 | 11.1 | 10.5 | 13.7 | 12.8 | 11.2 | |
| 2006 | 11.2 | 8.8 | 7.8 | 14.0 | 15.3 | |
| 2007 | 15.2 | 7.1 | 6.5 | 13.8 | 10.4 | |
| 2008 | 10.8 | 6.8 | 14.4 | 13.3 | 10.4 | |
| 2009 | 16.3 | 13.7 | 13.8 | 12.5 | 8.8 | |
| 2010 | 11.8 | 15.3 | 11.9 | 13.5 | 7.0 | |
| 2011 | 7.7 | 11.1 | 8.6 | 12.4 | 5.0 | |
| 2012 | 10.0 | 11.7 | 11.9 | 12.1 | 10.9 | |
| 2013 | 9.7 | 11.7 | 9.8 | 12.3 | 13.3 | |
| 2014 | 15.2 | 12.4 | 9.5 | 14.6 | 10.9 | |
| 2015 | 7.6 | 10.3 | 7.3 | 9.0 | 11.3 | |

Appendix 58 Proportion of continuous gabapentin prescribing periods exceeding 12 months, by IMD quintile

| Proportion of gabapentin prescribing | | | | | |
|--------------------------------------|-------|-------------|------------|------------|----------|
| | Index | of multiple | deprivatio | n (IMD) qu | iintiles |
| Year | Q1 | Q2 | Q3 | Q4 | Q5 |
| 2000 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 2001 | 0.0 | 2.3 | 7.1 | 6.3 | 2.6 |
| 2002 | 0.0 | 4.8 | 3.1 | 1.4 | 3.6 |
| 2003 | 2.5 | 4.2 | 5.9 | 2.9 | 5.3 |
| 2004 | 5.0 | 1.0 | 3.3 | 4.3 | 0.0 |
| 2005 | 2.0 | 3.2 | 4.3 | 6.0 | 5.2 |
| 2006 | 3.7 | 2.7 | 1.7 | 5.1 | 2.3 |
| 2007 | 6.3 | 0.6 | 1.3 | 5.3 | 2.1 |
| 2008 | 5.0 | 3.1 | 4.4 | 3.5 | 4.5 |
| 2009 | 3.9 | 3.1 | 4.2 | 4.0 | 3.7 |
| 2010 | 4.6 | 5.0 | 2.5 | 3.9 | 3.9 |
| 2011 | 2.6 | 3.7 | 2.7 | 4.4 | 1.7 |
| 2012 | 3.3 | 3.3 | 2.5 | 3.8 | 3.9 |
| 2013 | 2.9 | 4.6 | 3.1 | 4.7 | 4.4 |
| 2014 | 4.4 | 4.6 | 1.5 | 7.3 | 3.6 |
| 2015 | 0.6 | 0.0 | 0.0 | 2.7 | 4.2 |

Appendix 59 Proportion of continuous pregabalin prescribing periods exceeding 30 days, by IMD quintile

| Proportion of pregabalin prescribing | | | | | |
|--------------------------------------|-------|-------------|--------------|--------------|--------|
| | Index | of multiple | e deprivatio | n (IMD) quir | ntiles |
| Year | Q1 | Q2 | Q3 | Q4 | Q5 |
| 2004 | 33.3 | 100.0 | 100.0 | 100.0 | 33.3 |
| 2005 | 57.7 | 50.0 | 66.7 | 66.7 | 41.7 |
| 2006 | 54.3 | 61.7 | 56.6 | 47.3 | 51.7 |
| 2007 | 57.4 | 65.8 | 58.5 | 48.9 | 69.0 |
| 2008 | 75.5 | 60.4 | 59.4 | 62.5 | 59.8 |
| 2009 | 72.9 | 66.9 | 57.0 | 61.0 | 62.2 |
| 2010 | 61.8 | 62.0 | 60.2 | 58.2 | 61.6 |
| 2011 | 55.2 | 65.1 | 57.4 | 59.5 | 55.8 |
| 2012 | 59.0 | 66.5 | 53.2 | 55.5 | 53.7 |
| 2013 | 59.5 | 55.5 | 57.2 | 59.1 | 56.4 |
| 2014 | 59.8 | 61.6 | 67.1 | 66.7 | 57.6 |
| 2015 | 60.9 | 66.4 | 65.4 | 57.7 | 59.8 |

Appendix 60 Proportion of continuous pregabalin prescribing periods exceeding 6 months, by IMD quintile

| Proportion of pregabalin prescribing | | | | | |
|--------------------------------------|-------|-------------|------------|--------------|--------|
| | Index | of multiple | deprivatio | n (IMD) quir | ntiles |
| Year | Q1 | Q2 | Q3 | Q4 | Q5 |
| 2004 | 0.0 | 0.0 | 100.0 | 0.0 | 0.0 |
| 2005 | 3.8 | 0.0 | 0.0 | 4.8 | 8.3 |
| 2006 | 8.7 | 8.3 | 20.8 | 7.3 | 10.3 |
| 2007 | 11.1 | 6.8 | 16.9 | 8.0 | 14.1 |
| 2008 | 18.9 | 12.1 | 5.8 | 12.5 | 13.0 |
| 2009 | 18.8 | 15.3 | 9.7 | 12.7 | 10.4 |
| 2010 | 8.8 | 13.4 | 15.0 | 12.1 | 11.0 |
| 2011 | 12.7 | 11.6 | 16.2 | 12.0 | 18.2 |
| 2012 | 11.5 | 14.6 | 10.9 | 15.9 | 14.7 |
| 2013 | 13.5 | 16.8 | 13.9 | 18.2 | 16.8 |
| 2014 | 10.7 | 15.1 | 16.1 | 22.2 | 14.5 |
| 2015 | 12.2 | 19.8 | 14.2 | 8.0 | 16.4 |

Appendix 61 Proportion of continuous pregabalin prescribing periods exceeding 12 months, by IMD quintile

| Proportion of pregabalin prescribing | | | | | |
|--------------------------------------|-------|-------------|------------|-------------|--------|
| | Index | of multiple | deprivatio | n (IMD) qui | ntiles |
| Year | Q1 | Q2 | Q3 | Q4 | Q5 |
| 2004 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 2005 | 0.0 | 0.0 | 0.0 | 4.8 | 0.0 |
| 2006 | 4.3 | 3.3 | 5.7 | 1.8 | 3.4 |
| 2007 | 0.0 | 1.4 | 4.6 | 0.0 | 7.0 |
| 2008 | 1.9 | 5.5 | 2.9 | 3.4 | 4.3 |
| 2009 | 4.2 | 5.1 | 4.3 | 6.8 | 2.2 |
| 2010 | 4.4 | 4.2 | 4.4 | 6.4 | 4.1 |
| 2011 | 6.0 | 6.2 | 5.9 | 4.0 | 7.1 |
| 2012 | 3.6 | 5.7 | 3.8 | 3.8 | 5.3 |
| 2013 | 4.1 | 6.5 | 5.6 | 6.6 | 8.1 |
| 2014 | 3.3 | 8.2 | 7.1 | 11.1 | 7.9 |
| 2015 | 5.2 | 7.6 | 4.7 | 1.8 | 7.4 |

Appendix 62 Characteristics of people prescribed gabapentin in 2014, by continuous prescribing period length

| Gabapentin | | | | | |
|----------------------|-------------------|------------------|-------------------|-------------------|--|
| | Up to 3 months | 3 to 6 months | 6 to 12 months | Over 12 months | |
| Mean age (years) | 59.9 | 59.5 | 62.7 | 64.1 | |
| IMD quintile (%) | | | | | |
| I (least deprived) § | 44.9 | 8.2 | 12.0 | 4.4 | |
| II | 39.5 | 15.6* | 8.7 | 4.6 | |
| III | 48.7 | 17.6* | 8.5 | 1.5 | |
| IV | 51.0 | 15.6* | 8.3 | 7.3 | |
| V (most deprived | 50.6 | 10.9 | 9.3 | 3.6 | |
| Gender (%) | | | | | |
| Male § | 48.8 | 15.6 | 6.3 | 4.5 | |
| Female | 46.0 | 12.7 | 11.0* | 4.1 | |
| Region (%) | | | | | |
| North of England § | 48.1 | 12.8 | 10.2 | 4.1 | |
| Midlands and East of | | | | | |
| England | 45.2 | 17.0 | 10.9 | 5.2 | |
| London | 46.0 | 10.8 | 9.5 | - | |
| South East | 46.7 | 10.5 | 8.3 | 5.1 | |
| South West | 49.5 | 21.5* | 4.3 | 3.2 | |

^{*} P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, § - reference category

Appendix 63 Characteristics of people prescribed pregabalin in 2014, by continuous prescribing period length

| Pregabalin | | | | | |
|----------------------|-------------------|------------------|-------------------|----------------|--|
| | Up to 3 months | 3 to 6 months | 6 to 12 months | Over 12 months | |
| Mean age (years) | 56.6* | 59.8 | 62.4* | 59.4 | |
| IMD quintile (%) | | | | | |
| I (least deprived) § | 59.0 | 18.0 | 9.8 | 3.3 | |
| II | 48.6 | 21.9 | 8.2 | 8.2 | |
| III | 51.0 | 25.8 | 11.0 | 7.1 | |
| IV | 43.2** | 20.4 | 13.0 | 11.1* | |
| V (most deprived | 57.0 | 18.2 | 7.3 | 7.9 | |
| Gender (%) | | | | | |
| Male § | 52.6 | 17.2 | 7.7 | 13.0 | |
| Female | 50.9 | 22.9 | 11.0 | 4.9*** | |
| Region (%) | | | | | |
| North of England § | 48.5 | 19.4 | 8.7 | 11.7 | |
| Midlands and East of | | | | | |
| England | 54.6 | 20.8 | 10.4 | 6.5 | |
| London | 52.6 | 19.3 | 3.5 | 10.5 | |
| South East | 54.4 | 21.3 | 12.5 | 4.0** | |
| South West | 39.3 | 26.2 | 6.6 | 11.5 | |

^{*} P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, § - reference category

Source: CPRD

Appendix 64 Total counts of sales of individual codeine and dihydrocodeine-containing medicines

| | 3/2015 to 2/2016 | 3/2016 to 2/2017 | 3/2017 to 2/2018 |
|-----------------------------------|---------------------|---------------------|---------------------|
| Paracetamol & codeine | 7,713,634 | 7,133,170 | 7,269,506 |
| Paracetamol & codeine & caffeine | 3,848,998 | 3,753,986 | 3,322,000 |
| Ibuprofen & codeine | 2,223,748 | 2,160,476 | 2,254,624 |
| Paracetamol & dihydrocodeine | 1,129,152 | 1,119,408 | 986,080 |
| Buclizine & paracetamol & codeine | 291,948 | 278,772 | 271,008 |
| Aspirin & codeine | 115,872 | 109,184 | 98,368 |
| Total | 15,323,352 | 14,554,996 | 14,201,586 |

Appendix B. Evaluation of published estimates of dependence on prescribed medicine

What proportion of adults in England are dependent on medicines they are being prescribed?

Information needed

There is concern about the extent of dependence on medicines that people are prescribed. To understand the scale of the problem, an estimate is needed of the proportion of adults in England who are dependent on (in terms of showing signs of addiction to) the medicines they are prescribed. With a reliable estimate of the proportion, an estimate of the number of people can also be extrapolated.

To produce such an estimate, a large sample representative of (and therefore generalisable to) the wider national population is needed. Information about what potentially DFMs each individual in the sample was taking would need to be combined at the patient level with information about whether they have any indicators of dependence.

The focus here is on four classes of medicine that have been identified as potentially dependence forming: opioids, benzodiazepines, Z-drugs, and GABAergic medicines. Ideally, prescribing that was clinically appropriate despite dependence risk would be excluded. For example, the risk of dependence may be secondary when managing pain in palliative care, or implicit when treating a person for established dependence with maintenance therapy. The focus here is also on medicines prescribed through NHS primary care, and not on those obtained either illicitly or through private practice. Other research has addressed the abuse and misuse of prescription medicines, for example through diversion. Those findings are only relevant here where they relate to individuals taking the medicines that have been prescribed to them. Given enormous international variation in health care systems, prescribing practices, and the epidemiology of prescription drug dependence, understanding of the context in England needs to be based on data from England.

Information available

From a research (and clinical) perspective, identifying dependence is complex. The subjective experience of signs and symptoms of dependence could be captured using a screening tool (where people are able and willing to report those signs openly). However, many of the available screening tools focus on a specific class of medicine, on drug misuse rather than dependence on a medication that is being taken as prescribed, or they take a long time, and so are costly, to administer.⁹

⁹ See this paper for discussion of some available tools: Wickersham JA, Azar MM, Cannon CM et al. Validation of a Brief Measure of Opioid Dependence: ⁶The Rapid Opioid Dependence Screen (RODS) *J Correct Health Care*. 2015 21(1): 12-26.

Length of prescribing (at various thresholds, e.g. four, six, and 12 months) has been used in research as an indicator of dependence. However, not everyone prescribed to long-term is dependent, and some prescribed to shorter-term will be dependent. The Royal College of Anaesthetists produced a list of eighteen 'indicators that suggest the possibility of dependence should be explored in those on a long term opioid prescription'. ¹⁰ Indications that someone being prescribed opioids long-term might be dependent include: presence of psychiatric illness; concern expressed by family members, carers, or pharmacists; and resistance to attending appointments to review opioid prescribing or addiction assessment. What the list demonstrates is that length of prescribing alone may not be enough, and identification is not straightforward.

The validity of using length of prescribing as a single proxy for dependence on prescribed medicines is not considered here. Instead, we briefly consider the statistical robustness of the following three studies as sources for estimating the prevalence of long-term prescribing in the adult population in England. These are reviewed because they are commonly quoted sources for estimates of there being 1-1.5m people dependent, cited by the media and others to demonstrate the scale of the need for action. The first two papers focus on long-term prescribing (exceeding four and six months) of benzodiazepines around 18 years ago, the third on long-term term prescribing (exceeding 12 months) of benzodiazepines and Z-drugs in 2014/5.

Heather et al.: Benzodiazepine prescribing in 1997-1999 exceeding six months

A research study carried out in seven GP practices in Newcastle and North Tyneside in 1997-1999 identified 1,297 patients who had been prescribed benzodiazepines for longer than six months. These figures were subsequently scaled up (not by the study authors) to a projected 1.5 million people in England being prescribed benzodiazepines for periods exceeding six months.

Representativeness of the sample: The papers present findings from a well-run evaluation. The sample was designed to evaluate an intervention and the selection criteria were not intended to produce a sample representative of the wider population. In personal communication one of the authors said: 'There is no way our 2004 paper could be used to estimate prevalence [of long-term benzodiazepine prescribing]. Apart from geographical restrictions, there were several kinds of restrictions on entry to the study.'

Exclusions: Most of the 1,297 patients identified in the trial were not considered suitable for an intervention aimed at reducing their long-term benzodiazepine prescribing. This was for a range of reasons, including because they were taking benzodiazepines for a serious acute illness (such as cancer) or for a medical

https://www.rcoa.ac.uk/system/files/FPM-OA-identification-treatment.pdf

¹¹ Heather N, Bowie A, Ashton H, et al. Randomised controlled trial of two brief interventions against long-term benzodiazepine use: outcome of intervention. *Addiction Research and Theory* 2004; 12(2): 141–154. http://www.benzo.org.uk/amisc/rndoi.pdf

¹² Godfrey C, Heather N, Bowie A, et al. Randomised controlled trial of two brief interventions against long-term benzodiazepine use: Cost-effectiveness. *Addiction Research and Theory.* 2008; 16(4): 309–317.

condition (such as epilepsy). These conditions have been applied as exclusion criteria in subsequent research on long-term prescribing of potentially DFMs, and further reinforce the conclusion that the rates generated in this paper are not comparable with rates generated in the latter studies.

Recency: While the papers were published in 2004 and 2008, it is important to note that they are based on data collected in 1997 to 1999, and so relate to benzodiazepine prescribing practice in those GP surgeries two decades ago. The proportion of people to be prescribed benzodiazepines has fallen in recent years and estimates based on data collected at that time are unlikely to reflect current prescribing levels. ¹³

BBC Panorama poll: Benzodiazepine prescribing in 2000-2001 exceeding four months

In 2001, Panorama commissioned a poll of about two thousand adults which found that 3% of the sample had been prescribed a benzodiazepine for longer than four months. ¹⁴ Based on the 2001 adult population, this was extrapolated to a projected 1.5 million people in England being prescribed benzodiazepines long-term (for four months or more).

Assessing methods: Attempts have been made to obtain information about the sampling or survey methods, both with the polling agency (NOP) and with Panorama. However, no details could be located. We are therefore unable to confirm whether this poll was robust.

Outcome definition: It is also not clear what questions survey participants were asked, and therefore precisely what was being measured. For example, whether the 3% of participants were being prescribed a benzodiazepine for a period exceeding four months at the time of the interview, or whether these patients had been prescribed at that level in the previous12 months, or at some point in their life. Analysis of CPRD data¹⁵ has found that about 3.5% of adults had been prescribed benzodiazepines at some point in 2000, but that most prescribing periods were around 30 days or less. Irrespective of the measure used on the Panorama poll, this suggests that the rate generated by the poll may have been an overestimate. The poll also likely relied on people being aware of what medicines they were being prescribed, other research has found that self-reports may be inaccurate in relation to prescribing.¹⁶

Recency: As mentioned above, other work has since found that the proportion of the population to be prescribed this class of medicine to have fallen since 2000/2001. This indicates that results from 2000-2001 are unlikely to reflect recent prescribing patterns.

93

¹³ Cartagena Farias J, Porter L, McManus S, Strang J, Hickman M, Reed K, Smith N. (2017) *Prescribing patterns in dependence forming medicines*. London: NatCen.

¹⁴ http://news.bbc.co.uk/1/hi/programmes/panorama/tranquillisers/1325909.stm

¹⁵ Cartagena Farias J, Porter L, McManus S, Strang J, Hickman M, Reed K, Smith N. (2017) *Prescribing patterns in dependence forming medicines*. London: NatCen.

https://www.sciencedaily.com/releases/2009/12/091210000845.htm

Davies et al.: Benzodiazepine and Z-drugs prescribing exceeding 12 months

Analysis of data on patients registered at a number of GP surgeries in Bradford in 2014-2015 found that the proportion of 16 to 80 year olds taking benzodiazepines and/or Z-drugs very long-term (for more than one year) was 0.69%.¹⁷ This figure was then applied by the authors to the registered NHS patient population that age in England. The projection was that there were between 265,000 and 295,000 patients age 16-80 taking benzodiazepines and/or Z-drugs long-term in England. The study found that 35% of all patients prescribed benzodiazepines or Z-drugs had been prescribed them very long term (for more than 12 months).

Assessing methods: While the results of this survey are published as a paper in the *British Journal of General Practice*, relatively limited methodological detail was provided. For example, it is not stated how many GP surgeries were visited, and whether any surgeries were selected but did not take part.

Exclusions are well defined, appropriate, and lean towards the rate of long-term prescribing generated being a conservative estimate. The results were filtered to exclude children and those aged 80 years and over, those in receipt of palliative care, those with a diagnosis of epilepsy, those with severe and enduring mental health issues, and those 'with illness at the time of the survey' (although how this latter exclusion was applied is not defined).

Representativeness of the sample: The key issue is with the appropriateness of extrapolating to a national prevalence based on an unadjusted sample of patients living in and around one city. Considerable variation in prescribing is evident between clinical commissioning groups within a region, as well as between regions. ¹⁸ In order to justify scaling estimates from one area to a national prevalence, the characteristics of that sample would need to be profiled so that they could be compared with the national profile. This is particularly relevant for those characteristics known to be associated with prescribing propensity, such as age, sex, ethnicity and area level deprivation. Weighting factors could then be applied to adjust the profile of the patient sample to better approximate that of the national population. Without knowing how representative the patient sample is of the national population, and how representative the GP practice sample is of GP practices nationally, it is not possible to establish how robust an estimate this is of long-term benzodiazepine and Z-drug prescribing nationally.

¹⁷ Davies J, Rae TC, Montagu L (2017) Long-term benzodiazepine and Z-drugs use in England: a survey of general research. British Journal of General Practice 67(662): e:609-e613.

http://bjgp.org/content/67/662/e609

¹⁸ OpenPrescribing.net allows for variation in volume of items prescribed per 1,000 registered patients to be charted and finds significant variation between practices, CCGs, and regions (Curtis et al. 2019).

Appendix C. Estimating the minimum proportion of the population prescribed DFM long-term in 2014

Estimates of the proportion of people in primary care to have been prescribed dependence forming medicines long-term can be extrapolated from the prescribing data. It should be noted that the estimates that we could derive from the available data were crude, based on several assumptions, and likely to significantly underestimate rates. The figures presented should therefore be considered the minimum. The methods used to generate these estimates, and their limitations, are set out in Chapter 2.

The proportion of people in primary care in 2014 who had been prescribed DFM long-term varied by type of drug. Opioids had the highest prevalence of the population being prescribed in excess of 30 days: at least 1.8%. 1.1% of adults were prescribed GABAergic medicines in excess of 30 days, 0.9% were prescribed benzodiazepines and 0.7% were prescribed Z-drugs at this level.

Very broadly, at least one person in 500 in 2014 had been continuously prescribed an opioid for over a year; with a similar minimum proportion prescribed benzodiazepines for over a year. The rates for Z-drugs and GABAergic medicines appeared to be lower, at a minimum of at least one person in a 1000.

It should be noted that there is a lot of uncertainty in these estimates and they should be regarded as minimum estimates.

Table 7: Minimum proportion of the population exceeding DFM continuous prescribing periods in 2014, by length (%)

| | >30 days | >6 months | >12 months |
|---------------------|----------|-----------|------------|
| Benzodiazepines | 0.9 | 0.3 | 0.2 |
| Z-drugs | 0.7 | 0.2 | 0.1 |
| Opioids | 1.8 | 0.4 | 0.2 |
| GABAergic medicines | 1.1 | 0.3 | 0.1 |

Appendix D. Clinical Practice Research Datalink

A full consideration of the CPRD data extract analysed can be found in the previous report. Some of that information is provided below for reference.

CPRD data collection and structure

The Clinical Practice Research Datalink (CPRD) GOLD¹⁹ is an ongoing primary care database of anonymised medical records. There are 701 contributing GP practices in the UK which includes 15.8 million patients that meet the quality criteria. From those, 4.4 million patients are active (currently registered and/or alive)(CPRD, July 2016), representing approximately 6.9% of the UK population. Data collection begins when a patient is registered with a contributing practice. Thereafter the information recorded for each patient consultation or contact with the practice is uploaded to the CPRD on a monthly basis.

The CPRD primary care database is a rich source of health data for research. It includes data on:

- Demographics (gender and age)
- Strategic Health Authority
- Medical symptoms and diagnoses
- Tests and vaccinations
- Therapies
- Health-related behaviours
- Referrals to secondary care.
- Drug prescriptions.


Prescriptions are recorded by the GP using the GEMscript product code system (which includes the generic name of the product) with a therapy record containing the British National Formulary (BNF) code, daily dose, quantity prescribed, numbers of packs, and issue sequence (if it is a repeat prescription). Additionally, the CPRD also allows for the study of geographical differences using the ten former Strategic Health Authorities in England (SHA).

In terms of the data structure, the CPRD is a longitudinal dataset that combines location information for each GP practice with the consultation record for each registered patient. Each consultation in the CPRD is associated with a unique patient identifier and a date. A patient can have several consultations over time, with each consultation containing all the clinical and associated therapies and referrals information connected to it. However, it was not possible to directly associate diagnoses with treatments. For example, in a single consultation, a patient may be diagnosed with anxiety and flu and be prescribed paracetamol, ibuprofen, gabapentin

¹⁹ The main primary care database held by CPRD is known as GOLD (formerly GPRD). GOLD contains the anonymised, longitudinal medical records of patients registered with contributing primary care practices across the UK.

and diazepam with no systematic way to link each drug to each diagnosis. This represents a significant challenge to the management of the data within the CPRD.

Data structure of the CPRD



Arrows indicate that several diagnoses and several treatments may be present for each clinical encounter, but a definitive link between a specific diagnosis and a specific treatment is not available in the dataset.

CPRD representativeness

For the analysis, it was important to establish how representative the CPRD information was in order to know with what confidence results could be generalised to the wider patient population. In terms of representativeness of patients within the CPRD, a study concluded that patients are broadly representative of the UK general population in terms of age, sex and ethnicity when compared with the 2011 UK Census and are comparable to the Health Survey for England (HSE) for body mass index (BMI) distribution (Herrett et al., 2015).

In terms of representation of practices, historically the CPRD has gathered data from practices using the Vision IT system.²⁰ Over the last two years there has been a significant fall in the number of practices using Vision, and even a complete drop off in practices in certain geographical areas. This is attributable to some Clinical Commissioning Groups (CCGs) requiring different software to be used. Therefore, the

²⁰ Vision is an IT package designed to provide NHS patients with joined up services. It makes relevant information available to the whole healthcare team, intended to facilitate collaboration and reduce duplication of effort.

CPRD may not be representative of all practices in the UK based on geography and size.

Profile of the CPRD extract

The CPRD extract contains 49,999 patients who have been prescribed at least one of the four types of DFMs of interest at some point between 2000 and 2015. The extract is a geographically representative random sample of patients in England on the CPRD, with the proportion of patients in each area determined by the proportion of the overall population living there at the 2011 Census. For example, from the total number of patients requested in the extract, 10% are located in the South West, 10% are in Yorkshire and 15% are in London. See Table A.8.

Table A.8: Geographical distribution of patients

| Region | % | N |
|-----------------|-----|--------|
| North East | 5 | 2,449 |
| North West | 13 | 6,651 |
| Yorkshire | 10 | 4,983 |
| East Midlands | 9 | 4,276 |
| West Midlands | 11 | 5,284 |
| East of England | 11 | 5,515 |
| London | 15 | 7,709 |
| South East | 16 | 8,144 |
| South West | 10 | 4,988 |
| Total | 100 | 49,999 |

As well as providing a geographically representative sample for national analysis, the number of patients across each of the nine regions listed above is also sufficient for regional analysis. It should be noted that the nature of data collection from a self-selecting sample of GP practices does not allow for the production of probability weights which could account for non-response bias.

The 49,999 patients have an associated 13,868,419 consultations distributed between 2000 and 2015. This is an average of 23 consultations per patient per year. Of those patients, 59.5% are women. The patients span a wide age range, including some children and some adults in their nineties and over. Median age in 2000 was 50 years old, increasing to 52 in 2007, and 56 years in 2015. It is important to note that this increase in patients' age over time. Some patients are included in the extract from 2000 and continue to be prescribed over time until 2015, thereby contributing their own ageing effect to the age distribution of the sample. Conversely, patients who were first prescribed in later years contribute fewer years to the age distribution.

-

²¹ Mean age in 2015 was 56.19 and the standard deviation was 18.71.

The CPRD does not contain individual level information on socioeconomic characteristics. However, it does contain area level deprivation based on the location of the GP practice which acts as a proxy measure of socioeconomic circumstances of the patient. Overall, the extract contained roughly similar proportions of patients at each decile of deprivation.

Appendix E. Medicines included in the analysis

| Drug name | Product name | BNF code | BNF header |
|--------------------------------|--|--|---|
| Benzodiazepines | | | |
| Temazepam | Temazepam 10mg tablets | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Nitrazepam | Nitrazepam 5mg tablets | 4010100 | Hypnotics |
| Diazepam | Diazepam 2mg tablets | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 5mg tablets | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Lorazepam | Lorazepam 4mg/1ml solution for injection ampoules | 04010201/04080200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Benzodiazepines (peri-operative) |
| Temazepam | Temazepam 10mg/5ml oral solution sugar free | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Clonazepam | Clonazepam 1mg/1ml solution for injection ampoules and diluent | 4080200 | Drugs Used In Status Epilepticus |
| Temazepam | Temazepam 10mg capsules | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Lorazepam | Lorazepam 1mg tablets | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 10mg tablets | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg capsules | 04010201/04105000/10020200 | Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants |
| Clonazepam | Clonazepam 500microgram tablets | 4080100 | Control Of Epilepsy |
| Temazepam | Temazepam 20mg tablets | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Clonazepam | Clonazepam 2mg tablets | 4080100 | Control Of Epilepsy |
| Diazepam | Diazepam 5mg RecTubes (Wockhardt UK) | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Diazepam | Diazepam 10mg RecTubes (Wockhardt UK) | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Lorazepam | Lorazepam 2.5mg tablets | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 5mg capsules | 04010201/04105000/10020200 | Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants |
| Diazepam | Diazepam 2mg/5ml oral solution | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Diazepam | Valium 2mg Tablet (Roche Products) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Temazepam | Temazepam 20mg capsules | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Triazolam | Triazolam 250microgram Tablet | 4010100 | Hypnotics |
| Nitrazepam | Nitrazepam 5mg Capsule | 4010100 | Hypnotics |
| Flurazepam hydrochloride | Dalmane 30mg capsules (Meda Pharmaceuticals) | 4010201 | Benzodiazepines |
| Clobazam | Clobazam 10mg capsules | 04010200/04080100 | Anxiolytics/Control Of Epilepsy |
| Clobazam | Clobazam 10mg tablets | 4080100 | Control Of Epilepsy |
| Chlordiazepoxide Hydrochloride | Chlordiazepoxide 10mg tablets | 4010200 | Anxiolytics |
| Diazepam | Diazepam 5mg | 04010100/04010200 | Hypnotics/Anxiolytics |

4010100 Lormetazepam Lormetazepam 1mg tablets Hypnotics Lormetazepam Lormetazepam 1mg Capsule 4010100 Hypnotics Nitrazepam Mogadon 5mg Capsule (Roche Products) 4010100 Hypnotics 4010100 Nitrazepam Nitrazepam 10mg Tablet Hypnotics Lormetazepam 500microgram tablets 4010100 Lormetazepam Hypnotics 04010100/04010200 Hypnotics/Anxiolytics Diazepam 2mg capsules Diazepam Flurazepam hydrochloride Flurazepam 15mg capsules 4010201 Benzodiazepines 4010201 Flurazepam hydrochloride Dalmane 15mg capsules (Meda Pharmaceuticals) Renzodiazenines Tensium 10mg tablets (DDSA Pharmaceuticals) 04010201/04080200/04080300/10020200/15010401 Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Oxazepam Oxazepam 30mg Capsule 4010200 Anxiolytics Oxazepam Oxazepam 15mg tablets 4010201 Benzodiazepines 04010201/04080200/10020200/15010401/54140000 Diazepam 10mg/2.5ml rectal solution tube Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning Diazepam 04010100/04010200 Hypnotics/Anxiolytics Valium 5mg Tablet (Roche Products) Diazepam Stesolid 5mg rectal tube (Actavis UK) 04010201/04080200/10020200/15010401/54140000 Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning Diazepam 4080100 Clonazepam Clonazepam 2.5mg/ml drops sugar free Control Of Epilepsy 4010200 Anxiolytics Chlordiazepoxide Hydrochloride Chlordiazepoxide 5mg tablets 4010201 Benzodiazepines Oxazepam 10mg tablets Oxazepam Loprazolam mesilate Loprazolam 1mg tablets 4010100 Hypnotics 04010201/04105000/10020200 Chlordiazepoxide hydrochloride Chlordiazepoxide 10mg tablets Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants 04010200/04080200 Anxiolytics/Drugs Used In Status Epilepticus Diazepam Stesolid 10mg Rectal tubes (Dumex) 04010201/04080200/10020200/15010401/54140000 Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning Diazepam Diazepam 2.5mg/1.25ml rectal solution tube 04010201/04105000/10020200 Chlordiazepoxide hydrochloride Chlordiazepoxide 5mg tablets Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants 4010200 Anxiolytics Chlordiazepoxide 25mg tablets Chlordiazepoxide Hydrochloride Diazepam 5mg/2.5ml rectal solution tube 04010201/04080200/10020200/15010401/54140000 Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning Diazepam 04010201/04080200/15010401 Lorazepam Ativan 4mg/1ml solution for injection ampoules (Pfizer) Benzodiazepines/Drugs Used In Status Epilepticus/Benzodiazepines (peri-operative) 4010201 Flurazepam hydrochloride Benzodiazenines Flurazepam 30mg capsules 04010100/15010401 Hypnotics/Benzodiazepines (peri-operative) Temazepam Temazepam planpak Capsule (Manufacturer unknown) 04010100/15010401 Hypnotics/Benzodiazepines (peri-operative) Temazepam Temazepam 15mg capsules 4010100 Triazolam Triazolam 125microgram Tablet Hypnotics 4010100 Nitrazepam Mogadon 5mg Tablet (ICN Pharmaceuticals France S.A.) Hypnotics 4010100 Nitrazepam Nitrazepam 2.5mg/5ml oral suspension Hypnotics Clonazepam Rivotril 2mg tablets (Roche Products) 4080100 Control Of Epilepsy 04010201/04080200/10020200/15010401/54140000 Diazepam Diazepam 10mg suppositories Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning 04010200/04080200 Diazepam Diazepam 5mg suppository Anxiolytics/Drugs Used In Status Epilepticus Diazepam Stesolid 10mg rectal tube (Actavis UK) 04010201/04080200/10020200/15010401/54140000 Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning Clobazam Frisium 10mg Capsule (Aventis Pharma) 04010200/04080100 Anxiolytics/Control Of Epilepsy

Chlordiazepoxide Hydrochloride Chlordiazepoxide 25mg tablets Anxiolytics Oxazepam Oxazepam 30mg Tablet 4010200 Anxiolytics Temazepam Temazepam gelthix 10mg Capsule (Pharmacia) 04010100/15010401 Hypnotics/Benzodiazepines (peri-operative) 4010200 Chlordiazepoxide hydrochloride Librium 10mg Capsule (ICN Pharmaceuticals France S.A.) Anxiolytics Diazepam 04010100/04010200 Hypnotics/Anxiolytics Diazepam 1mg/5ml suspension Chlordiazepoxide hydrochloride Librium 5mg Tablet (ICN Pharmaceuticals France S.A.) 4010200 Anxiolytics 04010201/04080200/04080300/10020200/15010401 Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Diazepam 5mg/5ml oral solution 04010100/04010200 Diazepam Diazepam 10mg capsules Hypnotics/Anxiolytics Diazepam 2.5mg/5ml oral suspension 04010201/04080200/10020200/15010401/54140000 Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning Diazepam Alprazolam Alprazolam 250microgram tablets 4010201 Benzodiazepines Bromazepam Lexotan 3mg Tablet (Roche Products) 4010200 Anxiolytics 4010100 Nitrazepam Nitrazepam 5mg/5ml oral suspension Hypnotics 4010201 Benzodiazepines Diazepam 2mg/5ml oral solution sugar free Diazepam Valium 10mg Tablet (Roche Products) 04010100/04010200 Hypnotics/Anxiolytics Diazepam 04010100/15010401 Temazepam Temazepam 30mg capsules Hypnotics/Benzodiazepines (peri-operative) 4080100 Control Of Epilepsy Clonazepam Rivotril 500microgram tablets (Roche Products) 4010200 Anxiolytics Medazepam Nobrium 10mg Capsule (Roche Products) Medazepam Medazepam 10mg Capsule 4010200 Anxiolytics 4010200 Medazepam Nobrium 5mg Capsule (Roche Products) Anxiolytics 4010201 Xanax 250microgram tablets (Pfizer) Benzodiazepines Alprazolam Ativan 1mg Tablet (Wyeth Pharmaceuticals) 04010200/15010401 Anxiolytics/Benzodiazepines (peri-operative) Lorazepam Alprazolam Alprazolam 500microgram tablets 4010201 Benzodiazepines 4010200 Anxiolytics Lexotan 1.5mg Tablet (Roche Products) Bromazepam 04010201/04080200/04080300/10020200/15010401 Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam 10mg/2ml solution for injection ampoules Diazepam Ketazolam Anxon 30mg Capsule (Beecham Research Laboratories) 4010200 04010201/04080200/10020200/15010401/54140000 Diazepam Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning Diazepam 10mg/2ml emulsion for injection ampoules 04010100/15010401 Hypnotics/Benzodiazepines (peri-operative) Temazepam Normison 10mg Capsule (Wyeth Pharmaceuticals) 4010200 Anxiolytics Ketazolam Anxon 15mg Capsule (Beecham Research Laboratories) 04010100/15010401 Hypnotics/Benzodiazepines (peri-operative) Temazepam Temazepam gelthix 20mg Capsule (Pharmacia) 4010200 Anxiolytics Chlordiazepoxide hydrochloride Librium 5mg Capsule (ICN Pharmaceuticals France S.A.) 4010201 Alprazolam Xanax 500microgram tablets (Pfizer) 04010201/04080200/10020200/15010401/54140000 Diazepam Diazepam 10mg/5ml oral suspension Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning 4080100 Clonazepam Clonazepam 2mg/5ml oral solution sugar free Control Of Epilepsy 04010201/15010401 Lorazepam Lorazepam 500micrograms/5ml oral suspension Benzodiazepines/Benzodiazepines (peri-operative) Diazenam Diazemuls 10mg/2ml emulsion for injection ampoules (Actavis UK) 04010201/04080200/10020200/15010401/54140000 Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning Clobazam Frisium 10mg tablets (Sanofi) 4080100 Control Of Epilepsy

| Clonazepam | Clonazepam 500micrograms/5ml oral suspension | 4080100 | Control Of Epilepsy |
|--------------------------------|---|--|---|
| Temazepam | Temazepam 10mg/5ml Oral solution (Generics (UK)) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Nitrazepam | Nitrados 5mg Tablet (Rorer Pharmaceuticals) | 4010100 | Hypnotics |
| Chlordiazepoxide Hydrochloride | Librium 100mg Injection (Roche Products) | 4010200 | Anxiolytics |
| Diazepam | Valium 5mg/ml Injection (Roche Products) | 04010200/04080200 | Anxiolytics/Drugs Used In Status Epilepticus |
| Diazepam | Diazepam rectubes 20mg Rectal tubes (C P Pharmaceuticals) | 04010200/04080200 | Anxiolytics/Drugs Used In Status Epilepticus |
| Clonazepam | Rivotril 2.5mg/1ml drops (Imported (Switzerland)) | 4080100 | Control Of Epilepsy |
| Clobazam | Clobazam 5mg/5ml oral suspension | 4080100 | Control Of Epilepsy |
| Chlordiazepoxide hydrochloride | Librium 10mg Tablet (ICN Pharmaceuticals France S.A.) | 4010200 | Anxiolytics |
| Clonazepam | Clonazepam 500micrograms/5ml solution sugar free | 4080100 | Control Of Epilepsy |
| Lorazepam | Ativan 2.5mg Tablet (Wyeth Pharmaceuticals) | 04010200/15010401 | Anxiolytics/Benzodiazepines (peri-operative) |
| Chlordiazepoxide hydrochloride | Librium 5mg capsules (Meda Pharmaceuticals) | 04010201/04105000/10020200 | Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants |
| Lormetazepam | Noctamid 1mg Tablet (Schering Health Care) | 4010100 | Hypnotics |
| Diazepam | Diazepam 2.5mg RecTubes (Wockhardt UK) | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Diazepam | Valium 5mg Capsule (Roche Products) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Nitrazepam | Mogadon 5mg tablets (Meda Pharmaceuticals) | 4010100 | Hypnotics |
| Bromazepam | Bromazepam 3mg tablets | 4010200 | Anxiolytics |
| Diazepam | Valium 2mg Capsule (Roche Products) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Temazepam | Temazepam 10mg gel-fill capsules | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Valium 10mg Suppository (Roche Products) | 04010200/04080200 | Anxiolytics/Drugs Used In Status Epilepticus |
| | CHLORDIAZEPOXIDE | 0 | Unknown |
| Temazepam | Temazepam gelthix 30mg Capsule (Pharmacia) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Clobazam | Clobazam 10mg/5ml oral suspension | 4080100 | Control Of Epilepsy |
| Diazepam | Diazepam 2mg/5ml oral solution sugar free (Actavis UK) | 4010201 | Benzodiazepines |
| Amitriptyline | Amitriptyline 12.5mg / Chlordiazepoxide 5mg capsules | 4030100 | Tricyclic And Related Antidepressant Drugs |
| Lormetazepam | Loramet 1mg Capsule (Wyeth Pharmaceuticals) | 4010100 | Hypnotics |
| Temazepam | Normison 20mg Capsule (Wyeth Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Medazepam | Medazepam 5mg Capsule | 4010200 | Anxiolytics |
| | CHLORDIAZEPOXIDE | 0 | Unknown |
| Bromazepam | Bromazepam 1.5mg tablets | 4010200 | Anxiolytics |
| Temazepam | Temazepam gelthix 15mg Capsule (Pharmacia) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Loprazolam mesilate | Dormonoct 1mg Tablet (Hoechst Marion Roussel) | 4010100 | Hypnotics |
| Ketazolam | Ketazolam 15mg Capsule | 4010200 | Anxiolytics |
| Diazepam | Diazepam 20mg rectal tubes | 04010200/04080200 | Anxiolytics/Drugs Used In Status Epilepticus |
| Nitrazepam | Somnite 2.5mg/5ml oral suspension (Norgine Pharmaceuticals) | 4010100 | Hypnotics |
| Flurazepam hydrochloride | Paxane 30mg Capsule (M A Steinhard) | 4010100 | Hypnotics |
| | | | |

| Chlordiazepoxide hydrochloride | Librium 10mg capsules (Meda Pharmaceuticals) | 04010201/04105000/10020200 | Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants |
|--------------------------------|---|--|---|
| Chlordiazepoxide Hydrochloride | Chlordiazepozide 100mg injection | 4010200 | Anxiolytics |
| Oxazepam | Oxanid 10mg Tablet (M A Steinhard) | 4010200 | Anxiolytics |
| Prazepam | Centrax 10mg Tablet (Parke-davis Research Laboratories) | 4010200 | Anxiolytics |
| Clonazepam | Rivotril 1mg/1ml concentrate for solution for injection ampoules (Roche Products) | 4080200 | Drugs Used In Status Epilepticus |
| Temazepam | Temazepam 10mg/5ml oral solution sugar free (Rosemont Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Nitrazepam | Surem 5mg Capsule (Galen) | 4010100 | Hypnotics |
| Chlordiazepoxide hydrochloride | Tropium 5mg capsules (Dr Reddy's Laboratories (UK)) | 04010201/10020200 | Benzodiazepines/Skeletal Muscle Relaxants |
| | CHLORDIAZEPOXIDE | 0 | Unknown |
| Diazepam | Diazepam 2mg Tablet (Crosspharma) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Ketazolam | Ketazolam 30mg Capsule | 4010200 | Anxiolytics |
| Diazepam | Valium 5mg Suppository (Roche Products) | 04010200/04080200 | Anxiolytics/Drugs Used In Status Epilepticus |
| Diazepam | Evacalm 5mg Tablet (Unimed Pharmaceuticals) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 5mg Tablet (DDSA Pharmaceuticals) | 4010200 | Anxiolytics |
| Temazepam | Euhypnos forte 20mg Capsule (Pharmacia) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 2mg tablets (Ranbaxy (UK)) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Chlordiazepoxide hydrochloride | Tropium 5mg tablets (Dr Reddy's Laboratories (UK)) | 04010201/10020200 | Benzodiazepines/Skeletal Muscle Relaxants |
| Diazepam | Valium 2mg/5ml Oral solution (Roche Products) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Temazepam | Temazepam 20mg Tablet (Wyeth Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Temazepam | Temazepam 20mg tablets (IVAX Pharmaceuticals UK) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Valclair 10mg suppositories (Durbin Plc) | 0 | Unknown |
| Chlordiazepoxide Hydrochloride | Librium 25mg Tablet (ICN Pharmaceuticals France S.A.) | 4010200 | Anxiolytics |
| Diazepam | Diazepam 5mg tablets (Actavis UK) | 04010201/04080200/04080300/10020200/15010401 | $Benzo diaze pines/Drugs\ Used\ In\ Status\ Epilepticus/Febrile\ Convulsions/Skeletal\ Muscle\ Relaxants/Benzo diaze pines\ (peri-operative)$ |
| Temazepam | Temazepam 20mg gel-fill capsules | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 10mg/2ml solution for injection ampoules (Wockhardt UK) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Clonazepam | Clonazepam 250micrograms/5ml oral solution | 4080100 | Control Of Epilepsy |
| Temazepam | Temazepam 10mg/5ml oral solution sugar free (A A H Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 2mg/5ml oral solution (Sandoz) | 04010201/04080200/10020200/15010401/54140000 | $Benzo diazepines/Drugs\ Used\ In\ Status\ Epilepticus/Skeletal\ Muscle\ Relaxants/Benzo diazepines\ (peri-operative)/Therapy\ Of\ Convulsions\ Due\ To\ Poisoning\ Due\ Due\ Due\ Due\ Due\ Due\ Due\ Due$ |
| Diazepam | Solis 5mg Capsule (Galen) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Lorazepam | Lorazepam 1mg tablets (Teva UK) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Temazepam | Temazepam 10mg Tablet (Wyeth Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 2mg tablets (A A H Pharmaceuticals) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 10mg/2ml solution for injection ampoules (Hameln Pharmaceuticals) | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Temazepam | Temazepam 10mg tablets (IVAX Pharmaceuticals UK) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 5mg/2.5ml rectal solution tube (Sandoz) | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Diazepam | Diazepam 5mg/5ml oral solution (Sandoz) | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| | | | |

| Lormetazepam | Lormetazepam 500microgram tablets (A A H Pharmaceuticals) | 4010100 | Hypnotics |
|--------------------------------|--|--|---|
| Diazepam | Diazepam 10mg tablets (Mylan) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Temazepam | Temazepam 10mg tablets (A A H Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 2mg tablets (Teva UK) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 2mg tablets (Mylan) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 10mg tablets (Ranbaxy (UK)) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Lormetazepam | Lormetazepam 500microgram tablets (Genus Pharmaceuticals) | 4010100 | Hypnotics |
| Temazepam | Temazepam 10mg tablets (Teva UK) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Nitrazepam | Nitrazepam 5mg tablets (A A H Pharmaceuticals) | 4010100 | Hypnotics |
| Diazepam | Diazepam 5mg tablets (Teva UK) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Clonazepam | Clonazepam 2mg/5ml oral solution sugar free (Rosemont Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| Temazepam | Temazepam 10mg tablets (Mylan) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Lormetazepam | Lormetazepam 1mg tablets (Mylan) | 4010100 | Hypnotics |
| Diazepam | Diazepam 2mg tablets (Actavis UK) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Lormetazepam | Lormetazepam 1mg Tablet (Wyeth Pharmaceuticals) | 4010100 | Hypnotics |
| Nitrazepam | Nitrazepam 5mg tablets (Wockhardt UK) | 4010100 | Hypnotics |
| Diazepam | Diazepam 2mg Tablet (Regent Laboratories) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Temazepam | Temazepam 20mg tablets (Mylan) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 10mg/2.5ml rectal solution tube (Sandoz) | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Diazepam | Diazepam 5mg tablets (Mylan) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 5mg tablets (A A H Pharmaceuticals) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Lormetazepam | Lormetazepam 500microgram tablets (Mylan) | 4010100 | Hypnotics |
| Diazepam | Diazepam 2mg tablets (IVAX Pharmaceuticals UK) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 5mg Tablet (Crosspharma) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Nitrazepam | Nitrazepam 5mg tablets (Teva UK) | 4010100 | Hypnotics |
| Lormetazepam | Lormetazepam 1mg tablets (Genus Pharmaceuticals) | 4010100 | Hypnotics |
| Nitrazepam | Nitrazepam 5mg Tablet (DDSA Pharmaceuticals) | 4010100 | Hypnotics |
| Nitrazepam | Nitrazepam 5mg tablets (Mylan) | 4010100 | Hypnotics |
| Diazepam | Diazepam 10mg tablets (Actavis UK) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 2mg Tablet (Berk Pharmaceuticals) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Diazepam | Diazepam 5mg Tablet (Berk Pharmaceuticals) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 5mg Capsule (DDSA Pharmaceuticals) | 4010200 | Anxiolytics |
| Nitrazepam | Nitrazepam 5mg Tablet (Berk Pharmaceuticals) | 4010100 | Hypnotics |
| Lorazepam | Lorazepam 2.5mg tablets (Teva UK) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 5mg capsules (A A H Pharmaceuticals) | 04010201/04105000/10020200 | Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants |
| Lorazepam | Lorazepam 1mg tablets (Mylan) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| | | | |

| Diazepam | Atensine 10mg Tablet (Rorer Pharmaceuticals) | 04010100/04010200 | Hypnotics/Anxiolytics |
|--------------------------------|--|--|---|
| Temazepam | Temazepam 20mg Tablet (Pharmacia) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Oxazepam | Oxazepam 10mg tablets (A A H Pharmaceuticals) | 4010201 | Benzodiazepines |
| Temazepam | Euhypnos 10mg/5ml Oral solution (Pharmacia) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Nitrazepam | Remnos 10mg Tablet (DDSA Pharmaceuticals) | 4010100 | Hypnotics |
| Lorazepam | Lorazepam 1mg/5ml oral suspension | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Clobazam | Clobazam 2.5mg capsules | 4080100 | Control Of Epilepsy |
| Lorazepam | Lorazepam 1mg/5ml oral solution | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 5mg Rectal tubes (Hillcross Pharmaceuticals) | 04010200/04080200 | Anxiolytics/Drugs Used In Status Epilepticus |
| Temazepam | Temazepam 10mg Capsule (Berk Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Temazepam | Temazepam 20mg Capsule (Berk Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Lorazepam | Lorazepam 1mg tablets (Genus Pharmaceuticals) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Chlordiazepoxide Hydrochloride | Chlordiazepoxide 25mg Tablet (DDSA Pharmaceuticals) | 4010200 | Anxiolytics |
| Clobazam | Clobazam 25mg/5ml oral solution | 4080100 | Control Of Epilepsy |
| Nitrazepam | Nitrazepam 5mg tablets (Actavis UK) | 4010100 | Hypnotics |
| Lorazepam | Lorazepam 1mg tablets (Arrow Generics) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Oxazepam | Oxazepam 15mg tablets (A A H Pharmaceuticals) | 4010201 | Benzodiazepines |
| Temazepam | Temazepam 10mg Tablet (IVAX Pharmaceuticals UK) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Oxazepam | Oxazepam 10mg tablets (Actavis UK) | 4010201 | Benzodiazepines |
| Oxazepam | Oxazepam 10mg Tablet (IVAX Pharmaceuticals UK) | 4010200 | Anxiolytics |
| Oxazepam | Oxazepam 10mg tablets (Genus Pharmaceuticals) | 4010201 | Benzodiazepines |
| Temazepam | Temazepam 10mg Tablet (Pharmacia) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg Capsule (Approved Prescription Services) | 4010200 | Anxiolytics |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg Capsule (IVAX Pharmaceuticals UK) | 4010200 | Anxiolytics |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 5mg capsules (Actavis UK) | 04010201/10020200 | Benzodiazepines/Skeletal Muscle Relaxants |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg capsules (Actavis UK) | 04010201/10020200 | Benzodiazepines/Skeletal Muscle Relaxants |
| Loprazolam mesilate | Loprazolam 1mg tablets (Zentiva) | 4010100 | Hypnotics |
| Oxazepam | Oxazepam 15mg Tablet (IVAX Pharmaceuticals UK) | 4010200 | Anxiolytics |
| Oxazepam | Oxazepam 15mg tablets (Actavis UK) | 4010201 | Benzodiazepines |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 5mg Capsule (Approved Prescription Services) | 4010200 | Anxiolytics |
| Diazepam | Diazepam 10mg tablets (Teva UK) | 04010201/04080200/04080300/10020200/15010401 | $Benzo diaze pines/Drugs\ Used\ In\ Status\ Epilepticus/Febrile\ Convulsions/Skeletal\ Muscle\ Relaxants/Benzo diaze pines\ (peri-operative)$ |
| Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg Capsule (DDSA Pharmaceuticals) | 4010200 | Anxiolytics |
| Diazepam | Diazepam 10mg tablets (A A H Pharmaceuticals) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Temazepam | Temazepam 20mg Capsule (Hillcross Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 10mg Suppository (Sinclair IS Pharma Plc) | 04010200/04080200 | Anxiolytics/Drugs Used In Status Epilepticus |
| Temazepam | Temazepam 20mg tablets (A A H Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| 1 | | | |

| ĺ | Temazepam | Temazepam 20mg tablets (A A H Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
|---|--------------------------------|--|--|---|
| | Temazepam | Temazepam 10mg Capsule (Hillcross Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| | Temazepam | Temazepam 10mg Capsule (Hillcross Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| | Triazolam | Triazolam 0.25mg Tablet (Berk Pharmaceuticals) | 4010100 | Hypnotics |
| | Triazolam | Triazolam 0.25mg Tablet (Berk Pharmaceuticals) | 4010100 | Hypnotics |
| | Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg Tablet (DDSA Pharmaceuticals) | 4010200 | Anxiolytics |
| | Diazepam | Diazepam 5mg/5ml oral solution (A A H Pharmaceuticals) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| | Lorazepam | Lorazepam 2.5mg tablets (Genus Pharmaceuticals) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| | Chlordiazepoxide hydrochloride | Chlordiazepoxide 5mg tablets (A A H Pharmaceuticals) | 04010201/10020200 | Benzodiazepines/Skeletal Muscle Relaxants |
| | Midazolam | Midazolam Oral solution | 04080200/15010401 | Drugs Used In Status Epilepticus/Benzodiazepines (peri-operative) |
| | Chlordiazepoxide hydrochloride | Tropium 10mg tablets (Dr Reddy's Laboratories (UK)) | 04010201/10020200 | Benzodiazepines/Skeletal Muscle Relaxants |
| | Clonazepam | Clonazepam 500microgram/5ml Oral solution (Rosemont Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| | Diazepam | Diazepam 2mg Tablet (M & A Pharmachem) | 04010100/04010200 | Hypnotics/Anxiolytics |
| | Diazepam | Diazepam 5mg tablets (Ranbaxy (UK)) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| | Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg capsules (A A H Pharmaceuticals) | 04010201/04105000/10020200 | Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants |
| | Diazepam | Diazepam 5mg tablets (Sandoz) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| | Temazepam | Temazepam 10mg tablets (Actavis UK) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| | Temazepam | Temazepam 10mg tablets (Genus Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| | Diazepam | Diazepam 2mg tablets (Almus Pharmaceuticals) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| | Midazolam | Midazolam 2.5mg/ml oral solution | 15010401 | Benzodiazepines (peri-operative) |
| | Lorazepam | Lorazepam 1mg tablets (Sandoz) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| | Temazepam | Temazepam 20mg tablets (Genus Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| | Clobazam | Clobazam 10mg/5ml oral solution | 4080100 | Control Of Epilepsy |
| | Clobazam | Clobazam 5mg/5ml oral solution | 4080100 | Control Of Epilepsy |
| | Lorazepam | Lorazepam 500micrograms/5ml oral solution | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| | Triazolam | Triazolam (roi) 125microgram Tablet | 4010100 | Hypnotics |
| | Diazepam | Diazepam 10mg tablets (IVAX Pharmaceuticals UK) | 04010201/04080200/04080300/10020200/15010401 | $Benzo diaze pines/Drugs\ Used\ In\ Status\ Epilepticus/Febrile\ Convulsions/Skeletal\ Muscle\ Relaxants/Benzo diaze pines\ (peri-operative)$ |
| | Temazepam | Temazepam 20mg tablets (Teva UK) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| | Oxazepam | Oxazepam 15mg tablets (Genus Pharmaceuticals) | 4010201 | Benzodiazepines |
| | Nitrazepam | Nitrazepam 5mg tablets (Ranbaxy (UK)) | 4010100 | Hypnotics |
| | Temazepam | Temazepam 20mg tablets (Actavis UK) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| | Diazepam | Diazepam 5mg tablets (IVAX Pharmaceuticals UK) | 04010201/04080200/04080300/10020200/15010401 | $Benzo diaze pines/Drugs\ Used\ In\ Status\ Epilepticus/Febrile\ Convulsions/Skeletal\ Muscle\ Relaxants/Benzo diaze pines\ (peri-operative)$ |
| | Clonazepam | Clonazepam 500micrograms/5ml oral solution | 4080100 | Control Of Epilepsy |
| | Clobazam | Clobazam 25mg/5ml oral suspension | 4080100 | Control Of Epilepsy |
| | Lormetazepam | Lormetazepam 1mg/5ml oral suspension | 4010100 | Hypnotics |
| | Clonazepam | Clonazepam 500micrograms/5ml oral solution sugar free | 4080100 | Control Of Epilepsy |
| | | | | |

| Midazolam | Midazolam 10mg/1ml oromucosal solution pre-filled oral syringes | 0 | Unknown |
|------------|--|--|---|
| Midazolam | Midazolam 5mg/0.5ml oromucosal solution pre-filled syringes | 4080200 | Drugs Used In Status Epilepticus |
| Temazepam | Temazepam 10mg tablets (Sandoz) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Clonazepam | Clonazepam 2mg/5ml oral solution sugar free | 4080100 | Control Of Epilepsy |
| Clobazam | Clobazam 4mg/5ml oral solution | 4080100 | Control Of Epilepsy |
| Diazepam | Diazepam 10mg/5ml oral solution | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Clobazam | Clobazam 2.5mg/5ml oral solution | 4080100 | Control Of Epilepsy |
| Midazolam | Epistatus 10mg/1ml oromucosal solution (Special Products) | 4080200 | Drugs Used In Status Epilepticus |
| Diazepam | Diazepam 10mg/2ml solution for injection ampoules (A A H Pharmaceuticals) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Midazolam | Epistatus 7.5mg/0.75ml oromucosal solution pre-filled oral syringes (Special Products) | 4080200 | Drugs Used In Status Epilepticus |
| | Diazepam 2mg/5ml oral suspension | 4010201 | Benzodiazepines |
| Clobazam | Clobazam 4mg/5ml oral suspension | 4080100 | Control Of Epilepsy |
| Clobazam | Clobazam 2.5mg/5ml oral suspension | 4080100 | Control Of Epilepsy |
| Clobazam | Clobazam 100mg/5ml oral suspension | 4080100 | Control Of Epilepsy |
| Midazolam | Epistatus 10mg/1ml oromucosal solution pre-filled oral syringes (Special Products) | 4080200 | Drugs Used In Status Epilepticus |
| Clonazepam | Clonazepam 500microgram tablets (Phoenix Healthcare Distribution) | 4080100 | Control Of Epilepsy |
| Diazepam | Diazepam 2mg/5ml oral solution sugar free (A A H Pharmaceuticals) | 4010201 | Benzodiazepines |
| Diazepam | Diazepam 2.5mg/5ml oral solution | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Clonazepam | Clonazepam 2mg tablets (Sigma Pharmaceuticals Plc) | 4080100 | Control Of Epilepsy |
| Midazolam | Midazolam 7.5mg/0.75ml oromucosal solution pre-filled syringes | 4080200 | Drugs Used In Status Epilepticus |
| Diazepam | Diazepam 10mg Tablet (M & A Pharmachem) | 04010100/04010200 | Hypnotics/Anxiolytics |
| Midazolam | Epistatus 5mg/0.5ml oromucosal solution pre-filled oral syringes (Special Products | 4080200 | Drugs Used In Status Epilepticus |
| Clobazam | Tapclob 5mg/5ml oral suspension (Martindale Pharmaceuticals) | 04010201/04080100 | Benzodiazepines/Control Of Epilepsy |
| Triazolam | Triazolam 0.125mg Tablet (Berk Pharmaceuticals) | 4010100 | Hypnotics |
| Clobazam | Clobazam 10mg/5ml oral suspension sugar free | 04010201/04080100 | Benzodiazepines/Control Of Epilepsy |
| Bromazepam | Bromazepam 3mg tablets | 4010000 | Hypnotics And Anxiolytics |
| Temazepam | Temazepam 10mg/5ml oral solution sugar free (Focus Pharmaceuticals) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Midazolam | Midazolam 10mg/1ml oromucosal solution pre-filled oral syringes | 15010401 | Benzodiazepines (peri-operative) |
| Diazepam | Diazepam 2mg tablets (Wockhardt UK) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Lorazepam | Lorazepam 5mg/5ml oral solution | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Clobazam | Tapclob 10mg/5ml oral suspension (Martindale Pharmaceuticals) | 04010201/04080100 | Benzodiazepines/Control Of Epilepsy |
| Temazepam | Temazepam 10mg tablets (F.Maltby & Sons) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Temazepam | Temazepam 10mg tablets (Ethigen) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Lorazepam | Lorazepam 2.5mg tablets (Sandoz) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Clobazam | Frisium 10mg tablets (Waymade Healthcare Plc) | 4080100 | Control Of Epilepsy |
| Clonazepam | Clonazepam 250micrograms/5ml oral suspension | 4080100 | Control Of Epilepsy |
| | | | |

| Consignan Cuesargam 12 Zimmergrams/mile and supervision (see A Parameterically) Consequence Consequence (see A Parameterically) Consequence Consequence (see A Parameterically) Consequence Consequence Consequence Consequence (see A Parameterically) Consequence Consequence Consequence (see A Parameterically) Consequence Consequence Consequence (see A Parameterically) Consequence Consequence Consequence Consequence (see Consequence Conse | | | | |
|--|--------------------------------|---|--|---|
| Consisteran Consisteran Standard Consisteran Standard Consisteran | Diazepam | Diazepam 5mg tablets (Waymade Healthcare Plc) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Colorate (Colorate Colorate Co | Diazepam | Diazepam 5mg tablets (Sovereign Medical) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Disaspam Disparent famility from the statistication (Michaelbustines) (Michaelbustin | Clonazepam | Clonazepam 312.5micrograms/5ml oral suspension | 4080100 | Control Of Epilepsy |
| Disageam (Disageam (Figlian) oil siludion (Mi Distribution) (Verbalter) sensor (Verbalter | Clonazepam | Clonazepam 500microgram tablets (A A H Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| Consequent Sommitted Sommi | Diazepam | Diazepam 10mg/5ml oral solution (AM Distributions (Yorkshire)) | 04010201/04080200/10020200/15010401/54140000 | Benzodiazepines/Drugs Used In Status Epilepticus/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative)/Therapy Of Convulsions Due To Poisoning |
| Disaspam Disaspam Sing ballets (CR Pharmaceuticals) Clobazam (Clobazam Sing Part and suspension (Special Order) Cloracepam (Clobazam Sing Part and Suspension (Spe | Diazepam | Diazepam 5mg/5ml oral solution (AM Distributions (Yorkshire)) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Coloxizam Coloxizam Freignian of suspension (Special Circler) 6000000000000000000000000000000000000 | Clonazepam | Clonazepam 500microgram tablets (Almus Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| Temazapam Temazapam Mini galacitic (Sandors) designostrosassis (Sandors) designostrosas (Sandors) designostrosassis (Sandors) designostrosassis (Sandors) designostrosassis (Sandors) designostrosassis (Sandors) designostrosassis (Sandors) designostrosassis (Sandors) designostrosas (Sandors) designostrosassis (Sandors) designostrosas (Sandors) designostrosassis (Sandors) designostrosas (Sandors) designos | Diazepam | Diazepam 5mg tablets (DE Pharmaceuticals) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diszepam Diazepam Sing ballets (Arrow Generics) Diszepam Diszepam Sing ballets (Arrow Generics) Diszepam Sing b | Clobazam | Clobazam 5mg/5ml oral suspension (Special Order) | 4080100 | Control Of Epilepsy |
| Lorazepam Lorazepam Ing tablets (A H Pharmacouticals) Clonazepam Consequent Soluminary Control of Episery Lorazepam Consequent Soluminary Control of Episery Lorazepam Consequent Soluminary Control of Episery Lorazepam Diazepam Soluminary Control of Episery Diazepam Diazepam Soluminary Control of Episery Lorazepam Diazepam Diazepam Soluminary Control of Episery Lorazepam Diazepam Diazepam Soluminary Control of Episery Lorazepam Control of Episery Lorazepam Control of Episery Lorazepam Lorazepam Soluminary Control of Episery Lorazepam Lorazepam Soluminary C | Temazepam | Temazepam 20mg tablets (Sandoz) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Clonazepam (Conazepam Somirorgama tabletis (Activas UK) 460100 Control Of Epispay Nitrazepam Zmg/Smil oral suspension 480100 Activas UK) 460100 Centrol Of Epispay Diazepam (Diazepam Smg/Smil oral suspension 480100 Activas UK) 460100 Activa | Diazepam | Diazepam 5mg tablets (Arrow Generics) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Nitrazepam Zong Solitor (Sapper Control Suspension Autous (Sapper Control Suspension Suspension Suspension Suspension Suspension Suspension Su | Lorazepam | Lorazepam 1mg tablets (A A H Pharmaceuticals) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Lorazepam (Lorazepam Smg/Sml oral suspension (Lorazepam Smg/Sml oral suspension) (Lorazepam Smg/Sml oral suspension) (Lorazepam Ding tablets (Almus Pharmaceuticals) (Lorazepam Ding tablets (Almus Pharmaceuticals) (Lorazepam Ding tablets (A Al Pharmaceuticals) (Lorazepam Ding tablets (A Al Pharmaceuticals) (Lorazepam Ding tablets (A Al Pharmaceuticals) (Lorazepam Ding tablets (Sovereign Medical) (Lorazepam Ding Tablets (Maymade Healthcare Distribution) (Lorazepam Dinazepam Dinazep | Clonazepam | Clonazepam 500microgram tablets (Actavis UK) | 4080100 | Control Of Epilepsy |
| Diazepam Diazepam Mographical | | Nitrazepam 2mg/5ml oral suspension | 4010100 | Hypnotics |
| Chlordizepoxide hydrochloride Chlordizepoxide Jong tablets (Sovereign Medical) Diazepam 2mg tablets (Legap) Diazepam 2mg tablets (Legap) Diazepam 2mg tablets (Waymade Healthcare Pic) Diazepam Diazepam 2mg tablets (Waymade Healthcare Pic) Diazepam Diazepam 2mg tablets (Waymade Healthcare Pic) Diazepam Noctamid 0.5mg Tablet (Shering Health Care) Diazepam Diazepam 2mg tablets (Sovereign Medical) Diazepam 2mg tablets (Maymade Healthcare Pic) Diazepam Noctamid 0.5mg Tablet (Shering Health Care) Diazepam Diazepam 2mg tablets (Phoenix Healthcare Distribution) Diazepam Diazepam 2mg/5ml oral suspension Diazepam 2mg/5ml oral suspension Diazepam 2mg/5ml oral suspension Diazepam 2mg/5ml oral | Lorazepam | Lorazepam 5mg/5ml oral suspension | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Diazepam Diazepam mg tablets (Sovereign Medical) Temazepam (Temazepam 20mg Tablet (Lagap) Diazepam (Diazepam 20mg Tablet (Suyamade Healthcare Pic) Diazepam (Diazepam 20mg Tablet (Suyamade Healthcare Pic) Lormetazepam (Notamid 0.5mg Tablet (Schering Health Care) Diazepam (Temazepam 30mg gel-filir capsules Diazepam 20mg Tablet (Schering Health Care) Diazepam (Temazepam 30mg gel-filir capsules Diazepam 20mg tablets (Phoenix Healthcare Distribution) Diazepam (Diazepam 2mg tablets (Phoenix Healthcare Distribution) Diazepam (Diazepam 2mg tablets (Phoenix Healthcare Distribution) Diazepam (Diazepam 2mg tablets (Phoenix Healthcare Distribution) Diazepam (Diazepam 30mg gel-filir capsules) Diazep | Diazepam | Diazepam 10mg tablets (Almus Pharmaceuticals) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Temazepam Temazepam Zomg Tablet (Lagap) Clobazam Clobazam Clobazam Clobazam - 7.5mg/5ml oral suspension Diazepam Diazepam Zomg tablets (Waymade Healthcare Pic) Lormetazepam Noctamid 0.5mg Tablet (Schering Health Care) Mortanid 1.5mg Tablets (Phoenix Healthcare Distribution) Mortanid 1.5mg Tablets (Phoenix Healthcare Distribution) Mortanid 0.5mg Tablet (Schering Healthcare Distribution) Mortanid 1.5mg Tablets (Phoenix Healthcare Distribution) Mortanid 1.5mg Tablets (Phoenix Healthcare Distribution) Mortanid 1.5mg Tablets (Phoenix Healthcare Distribution) Mortanid 2.5mg Tablets (Manusche Relaxant/Berozodiazepines (Deri-operative) Mortanid 2.5mg Tablets (Manusche Relaxant/Berozodiazepines (Der | Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg tablets (A A H Pharmaceuticals) | 04010201/04105000/10020200 | Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants |
| Clobazam Clobazam 7.5mg/5ml oral suspension 440000 Control Of Epilepsy Diazepam 2mg tablets (Waymade Healthcare PIc) 440000 Hypnotics Temazepam Noctamid 0.5mg Tablet (Schering Health Care) 4400000 Hypnotics Temazepam 2mg tablets (Phoenix Healthcare Distribution) 64000001/100000001/100000001/100000001/1000000 | Diazepam | Diazepam 2mg tablets (Sovereign Medical) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam 2mg tablets (Waymade Healthcare Pic) 0401000/150010001 Hypnotics Lormetazepam Noctamid 0.5mg Tablet (Schering Health Care) 4010100 Hypnotics Temazepam Opazepam 2mg tablets (Phoenix Healthcare Distribution) 0401000/15001001 Hypnotics/Benzodiazepines (peri-operative) Diazepam 2mg tablets (Phoenix Healthcare Distribution) 0401000/15001001 Hypnotics/Benzodiazepines (peri-operative) Diazepam Diazepam 2mg tablets (Phoenix Healthcare Distribution) 0401000/150010001 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Diazepam 2mg tablets (Alliance Healthcare (Distribution)) 04010001/04000000/100000000/150000001 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Chordiazepoxide hydrochloride Chlordiazepoxide 20mg tablets (Phoenix Healthcare Distribution) 0401000/14000000/15000000000000000000000 | Temazepam | Temazepam 20mg Tablet (Lagap) | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Lormetazepam Noctamid 0.5mg Tablet (Schering Health Care) 4011100 Hypnotics Temazepam 30mg gel-fill capsules 04011001/15011401 Hypnotics/Benzodiazepines (peri-operative) Diazepam Diazepam 2mg tablets (Phoenix Healthcare Distribution) 04010021/040800000/04080000/15010401 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Diazepam 2mg tablets (Alliance Healthcare (Distribution)) 04010021/04080000/04080000/150104001 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Chlordiazepoxide hydrochloride Chlordiazepoxide 10mg tablets (Phoenix Healthcare Distribution) 0401002/04080000/15002000 Benzodiazepines/Drugs Used in Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants Clonazepam Clonazepam 5mg/5ml oral suspension 400100 Control Of Epilepsy Lorazepam 1 torazepam 10mg/5ml oral suspension 0401000 Hypnotics Lorazepam 10mg/5ml oral suspension 0401001/5ml and 100100 Benzodiazepines/Benzodiazepines (peri-operative) Clobazam 5mg/5ml oral suspension 0401001/5ml Benzodiazepines/Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/5ml oral suspension 0401001/5ml Benzodiazepines/Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/5ml oral suspension sugar free (A A H Pharmaceuticals) 0401000 Benzodiazepines/Benzodiazepines/Control Of Epilepsy Clobazam 5mg/5ml oral suspension sugar free (A A H Pharmaceuticals) 0401000 Control Of Epilepsy Clobazam 5mg/5ml oral solution 0401000 Neuropathic Pain/Control Of Epilepsy | Clobazam | Clobazam 7.5mg/5ml oral suspension | 4080100 | Control Of Epilepsy |
| Temazepam Temazepam and gel-fill capsules (Prio-perative) Diazepam 2mg tablets (Phoenix Healthcare Distribution) o4010001/04080200/04080000/10020000/15010401 Benzodiazepines (peri-operative) Diazepam 2mg tablets (Alliance Healthcare (Distribution) o4010001/04080200/04080000/10020000/15010401 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Chlordiazepoxide hydrochloride Chlordiazepoxide 10mg tablets (Phoenix Healthcare Distribution) o4010001/0100000/10020000 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Clonazepam Clonazepam Smg/Sml oral suspension 400100 Control Of Epilepsy Lorazepam Img tablets (Genesis Pharmaceuticals) 0401000 Control Of Epilepsy Nitrazepam Nitrazepam 10mg/Sml oral suspension 401000 Hypnotics Lorazepam Lorazepam 2mg/Sml oral suspension 0401000 Hypnotics Lorazepam Clobazam 5mg/Sml oral suspension 0401001/15010401 Benzodiazepines/Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/Sml oral suspension 0401001/15010401 Benzodiazepines/Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/Sml oral suspension 0401001/15010401 Benzodiazepines/Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/Sml oral suspension sugar free (A A H Pharmaceuticals) 0401000/140800100 Benzodiazepines/Benzodiazepines/Benzodiazepines (peri-operative) Clonazepam 10mg/Sml oral solution 04010001/15010401 Benzodiazepines/Benz | Diazepam | Diazepam 2mg tablets (Waymade Healthcare Plc) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Diazepam Diazepam Zmg tablets (Phoenix Healthcare Distribution) 04010001/04080000/10002000/15010401 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Zmg tablets (Alliance Healthcare (Distribution)) 04010001/04080000/100020000 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Chlordiazepoxide 4 10mg tablets (Phoenix Healthcare Distribution) 04010001/04105000/10002000 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Zmg tablets (Phoenix Healthcare Distribution) 04010001/04105000/10002000 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Zmg tablets (Phoenix Healthcare Distribution) 04010001/04105000/10002000 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Zmg tablets (Phoenix Healthcare Distribution) 04010001/0410000 Benzodiazepines/Drugs Used in Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Diazepam Zmg Tablets (Phoenix Healthcare Distribution) 04010000/04000000 Denzepam Zmg Tablets (Phoenix Healthcare Distribution) 04010001/15010001 Benzodiazepines/Drugs (Peri-operative) Denzepam Zmg Tablets (Phoenix Healthcare Distribution) 04010001/1501001 Benzodiazepines/Drugs (Peri-operative) Denzepam Zmg Tablets (Phoenix Healthcare Distribution) 04010001/1501001 Benzodiazepines/Drugs (Peri-operative) Denzepam Zmg Tablets (Phoenix Healthcare Distribution) 04010001/1501001 Benzodiazepines/Drugs (Peri-operative) Denzepam Zmg Tablets (Phoenix Healthcare Distribution) 04010001/1501001 Benzodiazepines/Drugs (Peri-operative) Denzepam Zmg Tablets (Phoenix Healthcare Distribution) 04010001/1501001 Benzodiazepines/Drugs (Peri-operative) Denzepam Zmg Tablets (P | Lormetazepam | Noctamid 0.5mg Tablet (Schering Health Care) | 4010100 | Hypnotics |
| Diazepam Diazepam 2mg tablets (Alliance Healthcare (Distribution)) o4010201/04080200/04080200/100202000 Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) Chlordiazepoxide hydrochloride Chlordiazepoxide 10mg tablets (Phoenix Healthcare Distribution) o4010201/04105000/10020200 Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants Clonazepam Clonazepam 5mg/5ml oral suspension 4080100 Control Of Epilepsy Lorazepam Ing tablets (Genesis Pharmaceuticals) 04010201/15010401 Benzodiazepines/Benzodiazepines (peri-operative) Nitrazepam Ingrapam 10mg/5ml oral suspension 40010201/5010401 Benzodiazepines/Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/5ml oral suspension sugar free (A A H Pharmaceuticals) 04010201/5010401 Benzodiazepines/Benzodiazepines (peri-operative) Clonazepam Clonazepam 10mg/5ml oral suspension sugar free (A A H Pharmaceuticals) 04010201/04080100 Benzodiazepines/Control Of Epilepsy Clonazepam 10mg/5ml oral solution 4001000 Control Of Epilepsy GABAergics Gabapentin Gabapentin 100mg capsules 04010000 Neuropathic Pain/Control Of Epilepsy | Temazepam | Temazepam 30mg gel-fill capsules | 04010100/15010401 | Hypnotics/Benzodiazepines (peri-operative) |
| Chlordiazepoxide hydrochloride Chlordiazepoxide 10mg tablets (Phoenix Healthcare Distribution) Clonazepam Clonazepam Smg/5ml oral suspension Lorazepam Lorazepam Ing tablets (Genesis Pharmaceuticals) Nitrazepam Nitrazepam Ingram 1 suspension Lorazepam Nitrazepam Ingram 1 suspension Lorazepam Lorazepam Ingram 1 suspension Adolto Pharmaceuticals Adolto Pharmac | Diazepam | Diazepam 2mg tablets (Phoenix Healthcare Distribution) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Clonazepam Clonazepam 5mg/5ml oral suspension 4080100 Control Of Epilepsy Lorazepam Ing tablets (Genesis Pharmaceuticals) 04010201/15010401 Benzodiazepines (peri-operative) Nitrazepam Nitrazepam 10mg/5ml oral suspension 401000 Hypnotics Lorazepam Lorazepam 2mg/5ml oral suspension 04010201/15010401 Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/5ml oral suspension 04010201/15010401 Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/5ml oral suspension sugar free (A A H Pharmaceuticals) 04010201/04080100 Benzodiazepines (peri-operative) Clonazepam Clonazepam 10mg/5ml oral solution 4080100 Control Of Epilepsy GABAergics Gabapentin Gabapentin 100mg capsules 04070300/04080100 Neuropathic Pain/Control Of Epilepsy | Diazepam | Diazepam 2mg tablets (Alliance Healthcare (Distribution)) | 04010201/04080200/04080300/10020200/15010401 | Benzodiazepines/Drugs Used In Status Epilepticus/Febrile Convulsions/Skeletal Muscle Relaxants/Benzodiazepines (peri-operative) |
| Lorazepam Lorazepam 1mg tablets (Genesis Pharmaceuticals) Nitrazepam Nitrazepam 10mg/5ml oral suspension Autorous Lorazepam 2mg/5ml oral suspension Lorazepam Lorazepam 2mg/5ml oral suspension Clobazam Clobazam 5mg/5ml oral suspension sugar free (A A H Pharmaceuticals) Clobazepam 10mg/5ml oral solution Autorous 408010 Control Of Epilepsy GABAergics Gabapentin Gabapentin 100mg capsules Neuropathic Pain/Control Of Epilepsy | Chlordiazepoxide hydrochloride | Chlordiazepoxide 10mg tablets (Phoenix Healthcare Distribution) | 04010201/04105000/10020200 | Benzodiazepines/Drugs Used In Substance Dependence - Alcohol Dependence/Skeletal Muscle Relaxants |
| Nitrazepam Nitrazepam 10mg/5ml oral suspension 401000 Hypnotics Lorazepam Lorazepam 2mg/5ml oral suspension 04010201/15010401 Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/5ml oral suspension sugar free (A A H Pharmaceuticals) 04010201/04080100 Benzodiazepines/Control Of Epilepsy Clonazepam Clonazepam 10mg/5ml oral solution 4080100 Control Of Epilepsy GABAergics Gabapentin Gabapentin 100mg capsules 04070300/04080100 Neuropathic Pain/Control Of Epilepsy | Clonazepam | Clonazepam 5mg/5ml oral suspension | 4080100 | Control Of Epilepsy |
| Lorazepam Lorazepam 2mg/Sml oral suspension 04010201/15010401 Benzodiazepines (peri-operative) Clobazam Clobazam 5mg/Sml oral suspension sugar free (A A H Pharmaceuticals) 04010201/04080100 Benzodiazepines/Control Of Epilepsy Clonazepam Clonazepam 10mg/Sml oral solution 4080100 Control Of Epilepsy GABAergics Gabapentin Gabapentin 100mg capsules 04010201/04080100 Neuropathic Pain/Control Of Epilepsy | Lorazepam | Lorazepam 1mg tablets (Genesis Pharmaceuticals) | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| Clobazam Clobazam 5mg/5ml oral suspension sugar free (A A H Pharmaceuticals) Clonazepam 10mg/5ml oral solution Clonazepam 10mg/5ml oral solution GABAergics Gabapentin Gabapentin 100mg capsules O4070300/04080100 Neuropathic Pain/Control Of Epilepsy | Nitrazepam | Nitrazepam 10mg/5ml oral suspension | 4010100 | Hypnotics |
| Clonazepam 10mg/5ml oral solution 4080100 Control Of Epilepsy GABAergics Gabapentin Gabapentin 100mg capsules 04070300/04080100 Neuropathic Pain/Control Of Epilepsy | Lorazepam | Lorazepam 2mg/5ml oral suspension | 04010201/15010401 | Benzodiazepines/Benzodiazepines (peri-operative) |
| GABAergics Gabapentin Gabapentin 100mg capsules O4070300/04080100 Neuropathic Pain/Control Of Epillepsy | Clobazam | Clobazam 5mg/5ml oral suspension sugar free (A A H Pharmaceuticals) | 04010201/04080100 | Benzodiazepines/Control Of Epilepsy |
| Gabapentin Gabapentin 100mg capsules 04070300/04080100 Neuropathic Pain/Control Of Epilepsy | Clonazepam | Clonazepam 10mg/5ml oral solution | 4080100 | Control Of Epilepsy |
| Gabapentin Gabapentin 100mg capsules 04070300/04080100 Neuropathic Pain/Control Of Epilepsy | GABAergics | | | |
| Prographic Prographic 2 Fing capsulor 04030000/4070300/04080100 Antidepressant Drugs/Neuronathic Pain/Control Of Enilepsy | | Gabapentin 100mg capsules | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| riegauaiii zniig capsules | Pregabalin | Pregabalin 25mg capsules | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |

04030000/04070300/04080100

Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy

Pregabalin

Pregabalin 75mg capsules

| Gabapentin | Gabapentin 300mg capsules | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
|-------------|---|-------------------------------------|--|
| Lamotrigine | Lamotrigine 25mg tablets | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 50mg tablets | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 5mg dispersible tablets sugar free | 4080100 | Control Of Epilepsy |
| Vigabatrin | Vigabatrin 500mg tablets | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 200mg tablets | 4080100 | Control Of Epilepsy |
| Vigabatrin | Vigabatrin 500mg oral powder sachets sugar free | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamictal 25mg tablets (GlaxoSmithKline UK) | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 400mg capsules | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamotrigine 100mg tablets | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 600mg tablets | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamotrigine 25mg dispersible tablets sugar free | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamictal 100mg dispersible tablets (GlaxoSmithKline UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 100mg dispersible tablets sugar free | 4080100 | Control Of Epilepsy |
| Gabapentin | Neurontin 300mg capsules (Pfizer) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamotrigine 2mg dispersible tablets sugar free | 4080100 | Control Of Epilepsy |
| Pregabalin | Lyrica 75mg capsules (Pfizer) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Pregabalin | Pregabalin 150mg capsules | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| Pregabalin | Pregabalin 50mg capsules | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| Pregabalin | Pregabalin 200mg capsules | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| Pregabalin | Pregabalin 100mg capsules | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| Pregabalin | Pregabalin 300mg capsules | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamictal 50mg tablets (GlaxoSmithKline UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamictal 100mg tablets (GlaxoSmithKline UK) | 4080100 | Control Of Epilepsy |
| Pregabalin | Lyrica 100mg capsules (Pfizer) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Pregabalin | Lyrica 50mg capsules (Pfizer) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Pregabalin | Lyrica 200mg capsules (Pfizer) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Gabapentin | Gabapentin 800mg tablets | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Neurontin 100mg capsules (Pfizer) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamictal 200mg tablets (GlaxoSmithKline UK) | 4080100 | Control Of Epilepsy |
| Pregabalin | Lyrica 300mg capsules (Pfizer) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Vigabatrin | Sabril 500mg tablets (Sanofi) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamictal 25mg dispersible tablets (GlaxoSmithKline UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamictal 5mg dispersible tablets (GlaxoSmithKline UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 50mg dispersible tablets sugar free | 4080100 | Control Of Epilepsy |
| Gabapentin | Neurontin 400mg capsules (Pfizer) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | | | |

| Pregabalin | Lyrica 150mg capsules (Pfizer) | 04030000/04070300/04080100/06010500 | $Antidepressant\ Drugs/Neuropathic\ Pain/Control\ Of\ Epilepsy/Treatment\ Of\ Diabetic\ Nephropathy\ And\ Neuropathy$ |
|-------------|---|-------------------------------------|---|
| Pregabalin | Lyrica 25mg capsules (Pfizer) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Vigabatrin | Vigabatrin 125mg capsules | 4080100 | Control Of Epilepsy |
| Gabapentin | Neurontin 600mg tablets (Pfizer) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Vigabatrin | Sabril 500mg oral powder sachets (Sanofi) | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 250mg/5ml oral solution | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | TIAGABINE | 0 | Unknown |
| Lamotrigine | Lamictal 2mg dispersible tablets (GlaxoSmithKline UK) | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 400mg/5ml oral suspension | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Gabapentin 300mg capsules (Teva UK) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Neurontin 800mg tablets (Pfizer) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamotrigine 25mg tablets (Actavis UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 100mg tablets (Actavis UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 25mg tablets (Mylan) | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 100mg capsules (Teva UK) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Gabapentin 100mg capsules (A A H Pharmaceuticals) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Gabapentin 300mg capsules (A A H Pharmaceuticals) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Gabapentin 600mg tablets (Teva UK) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamotrigine 25mg dispersible tablets sugar free (Arrow Generics) | 4080100 | Control Of Epilepsy |
| Pregabalin | Pregabalin 225mg capsules | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| Pregabalin | Lyrica 225mg capsules (Pfizer) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Lamotrigine | Lamotrigine 50mg tablets (Actavis UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 25mg tablets (Teva UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 100mg tablets (Teva UK) | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 250mg/5ml oral suspension | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Gabapentin 600mg tablets (A A H Pharmaceuticals) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Gabapentin 250mg/5ml Oral solution (Boots Pharmaceuticals) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamotrigine 50mg Suppository | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 50mg tablets (Teva UK) | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 100mg capsules (Sandoz) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Lamotrigine | Lamotrigine 50mg tablets (IVAX Pharmaceuticals UK) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 25mg tablets (Fannin UK) | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 300mg capsules (Mylan) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Gabapentin | Gabapentin 300mg capsules (Sandoz) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin Soonig capsules (Sandoz) | | |
| Lamotrigine | Lamotrigine 25mg dispersible tablets sugar free (A A H Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| | | 4080100 4080100 | |

| I | Pregabalin | Lyrica 150mg capsules (Lexon (UK)) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
|---|-------------|---|-------------------------------------|--|
| | Lamotrigine | Lamotrigine 25mg tablets (Accord Healthcare) | 4080100 | Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 25mg tablets (A A H Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| | Gabapentin | Gabapentin 300mg capsules (Ranbaxy (UK)) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | | Pregabalin 20mg/ml oral solution sugar free | 04030000/04070300/04080100/06010500 | $Antidepressant\ Drugs/Neuropathic\ Pain/Control\ Of\ Epilepsy/Treatment\ Of\ Diabetic\ Nephropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ Neuropathy\$ |
| | Lamotrigine | Lamictal 25mg tablets (DE Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| | Lamotrigine | Lamictal 25mg tablets (Lexon (UK)) | 4080100 | Control Of Epilepsy |
| | | Lyrica 20mg/ml oral solution (Pfizer) | 04030000/04070300/04080100/06010500 | $Antidepressant\ Drugs/Neuropathic\ Pain/Control\ Of\ Epilepsy/Treatment\ Of\ Diabetic\ Nephropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ Neuropathy\$ |
| | Pregabalin | Lyrica 50mg capsules (Waymade Healthcare Plc) | 04030000/04070300/04080100/06010500 | $Antidepressant\ Drugs/Neuropathic\ Pain/Control\ Of\ Epilepsy/Treatment\ Of\ Diabetic\ Nephropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ And\ Neuropathy\ Neuropathy\$ |
| | Gabapentin | Gabapentin 300mg capsules (Actavis UK) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin | Gabapentin 300mg/5ml oral solution | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 5mg dispersible tablets sugar free (Actavis UK) | 4080100 | Control Of Epilepsy |
| | | Gabapentin 50mg/ml oral solution sugar free | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin | Gabapentin 300mg capsules (Almus Pharmaceuticals) | 0 | Unknown |
| | Gabapentin | Gabapentin 300mg capsules (Bristol Laboratories) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 25mg tablets (Bristol Laboratories) | 4080100 | Control Of Epilepsy |
| | | Pregabalin 150mg/5ml oral solution | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 25mg dispersible tablets sugar free (Tillomed Laboratories) | 4080100 | Control Of Epilepsy |
| | Gabapentin | Gabapentin 100mg capsules (Mylan) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin | Neurontin 300mg capsules (Dowelhurst) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin | Gabapentin 300mg capsules (Arrow Generics) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 50mg tablets (Alliance Healthcare (Distribution)) | 4080100 | Control Of Epilepsy |
| | | Gabapentin 600mg/5ml oral solution | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin | Gabapentin 100mg capsules (Actavis UK) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin | Gabapentin 400mg capsules (Almus Pharmaceuticals) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin | Gabapentin 300mg/5ml oral suspension | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Gabapentin | Gabapentin 100mg capsules (Zentiva) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 100mg tablets (A A H Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 100mg tablets (Alliance Healthcare (Distribution)) | 4080100 | Control Of Epilepsy |
| | Vigabatrin | Vigabatrin 500mg/5ml oral solution | 4080100 | Control Of Epilepsy |
| | Gabapentin | Gabapentin 600mg tablets (Zentiva) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | | Gabapentin 400mg/5ml oral solution | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 100mg dispersible tablets sugar free (Teva UK) | 4080100 | Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 25mg dispersible tablets sugar free (Teva UK) | 4080100 | Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 50mg tablets (A A H Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| | Lamotrigine | Lamotrigine 25mg tablets (Alliance Healthcare (Distribution)) | 4080100 | Control Of Epilepsy |
| | | | | |

| Lamotrigine | Lamotrigine 100mg dispersible tablets sugar free (A A H Pharmaceuticals) | 4080100 | Control Of Epilepsy | |
|-------------|--|----------------------------|---|--|
| Lamotrigine | Lamotrigine 200mg tablets (A A H Pharmaceuticals) | 4080100 | Control Of Epilepsy | |
| Gabapentin | Gabapentin 300mg capsules (Creo Pharma) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy | |
| Lamotrigine | Lamictal 25mg dispersible tablets (Lexon (UK)) | 4080100 | Control Of Epilepsy | |
| | Pregabalin 75mg/5ml oral solution | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy | |
| Lamotrigine | Lamotrigine 200mg tablets (Waymade Healthcare Plc) | 4080100 | Control Of Epilepsy | |
| Lamotrigine | Lamotrigine 50mg/5ml oral suspension | 4080100 | Control Of Epilepsy | |
| Gabapentin | Gabapentin 300mg capsules (Waymade Healthcare Plc) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy | |
| Lamotrigine | Lamotrigine 50mg tablets (Accord Healthcare) | 4080100 | Control Of Epilepsy | |
| Lamotrigine | Lamotrigine 100mg tablets (Accord Healthcare) | 4080100 | Control Of Epilepsy | |
| Lamotrigine | Lamotrigine 200mg tablets (Accord Healthcare) | 4080100 | Control Of Epilepsy | |
| Lamotrigine | Lamotrigine 50mg tablets (Sigma Pharmaceuticals Plc) | 4080100 | Control Of Epilepsy | |
| Pregabalin | Rewisca 75mg capsules (Consilient Health) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Rewisca 50mg capsules (Consilient Health) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Rewisca 300mg capsules (Consilient Health) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Rewisca 150mg capsules (Consilient Health) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Rewisca 100mg capsules (Consilient Health) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Rewisca 200mg capsules (Consilient Health) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Rewisca 25mg capsules (Consilient Health) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Rewisca 225mg capsules (Consilient Health) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Gabapentin | Gabapentin 300mg capsules (Zentiva) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy | |
| Pregabalin | Pregabalin 50mg capsules (A A H Pharmaceuticals) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Alzain 25mg capsules (Dr Reddy's Laboratories (UK)) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Alzain 100mg capsules (Dr Reddy's Laboratories (UK)) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Lamotrigine | Lamotrigine 25mg dispersible tablets sugar free (Actavis UK) | 4080100 | Control Of Epilepsy | |
| Pregabalin | Pregabalin 100mg capsules (A A H Pharmaceuticals) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Alzain 300mg capsules (Dr Reddy's Laboratories (UK)) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Alzain 225mg capsules (Dr Reddy's Laboratories (UK)) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Alzain 150mg capsules (Dr Reddy's Laboratories (UK)) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Alzain 75mg capsules (Dr Reddy's Laboratories (UK)) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Alzain 50mg capsules (Dr Reddy's Laboratories (UK)) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Pregabalin | Alzain 200mg capsules (Dr Reddy's Laboratories (UK)) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy | |
| Gabapentin | Gabapentin 600mg tablets (DE Pharmaceuticals) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy | |
| Pregabalin | Pregabalin 150mg/5ml oral suspension | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy | |
| Gabapentin | Gabapentin 300mg capsules (Alliance Healthcare (Distribution)) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy | |
| Gabapentin | Gabapentin 50mg/ml oral solution sugar free (Alliance Healthcare (Distribution)) | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy | |
| | | | | |

| Pregabalin | Lyrica 75mg capsules (Waymade Healthcare Plc) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
|-----------------------------|--|-------------------------------------|--|
| Pregabalin | Lyrica 150mg capsules (Sigma Pharmaceuticals Plc) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Lamotrigine | Lamotrigine 100mg tablets (Kent Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| Gabapentin | Gabapentin 100mg/5ml oral solution | 04070300/04080100 | Neuropathic Pain/Control Of Epilepsy |
| Pregabalin | Pregabalin 75mg capsules (Teva UK) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy |
| Pregabalin | Pregabalin 25mg capsules (Teva UK) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy |
| Pregabalin | Lecaent 50mg capsules (Actavis UK) | 04030000/04080100 | Antidepressant Drugs/Control Of Epilepsy |
| Pregabalin | Pregabalin 50mg capsules (Accord Healthcare) | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| Pregabalin | Pregabalin 50mg capsules (Sandoz) | 04030000/04070300/04080100 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy |
| Pregabalin | Lyrica 75mg capsules (Stephar (U.K.)) | 04030000/04070300/04080100/06010500 | Antidepressant Drugs/Neuropathic Pain/Control Of Epilepsy/Treatment Of Diabetic Nephropathy And Neuropathy |
| Lamotrigine | Lamictal 50mg tablets (DE Pharmaceuticals) | 4080100 | Control Of Epilepsy |
| Lamotrigine | Lamotrigine 5mg/5ml oral solution | 4080100 | Control Of Epilepsy |
| Opioids | | | |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules | 4070200 | Opioid Analgesics |
| Pentazocine Lactate | Fortral 30mg/ml Injection (Sterwin Medicines) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml linctus | 3090100 | Cough Suppressants |
| Codeine phosphate | Codeine 30mg tablets | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Zydol 50mg capsules (Grunenthal) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 10mg/5ml oral solution | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 25mg/5ml oral solution | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 25mg tablet | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 10mg/ml injection | 4070200 | Opioid Analgesics |
| Buprenorphine Hydrochloride | Buprenorphine HCl 300micrograms injection | 4070200 | Opioid Analgesics |
| Pentazocine hydrochloride | Pentazocine 50mg capsules | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulfate 100mg/50ml solution for infusion vials | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Buprenorphine hydrochloride | Buprenorphine 200microgram sublingual tablets sugar free | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Pethidine Hydrochloride | Pethidine 50mg/ml injection | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine 15mg Suppository | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Morphine sulfate | MST Continus 10mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg tablets | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Morphine sulfate | MST Continus Suspension 20mg granules sachets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Fentanyl 50microgram/ml Injection | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| Fentanyl | Fentanyl 25micrograms/hour transdermal patches | 4070200 | Opioid Analgesics |
| | | | |

| Codeine Phosphate | Codeine 30mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
|-----------------------------|--|-------------------|--|
| Morphine sulfate | Morphine sulfate 10mg/5ml oral solution unit dose vials sugar free | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine 30mg Suppository | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride/ | Tramacet 37.5mg/325mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg modified-release capsules | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate 1mg/ml Injection | 4070200 | Opioid Analgesics |
| Fentanyl | Durogesic 25micrograms transdermal patches (Janssen-Cilag) | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 50micrograms/hour transdermal patches | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 50mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Morphine sulfate | Oramorph 10mg/5ml oral solution (Boehringer Ingelheim) | 4070200 | Opioid Analgesics |
| | Migraleve tablets (McNeil Products) | 04070200/04070401 | Opioid Analgesics/Treatment Of Acute Migraine |
| Buclizine hydrochloride/ | Migraleve Pink tablets (McNeil Products) | 04070200/04070401 | Opioid Analgesics/Treatment Of Acute Migraine |
| Codeine phosphate | Codeine 15mg/5ml linctus sugar free | 3090100 | Cough Suppressants |
| Dihydrocodeine tartrate | Dihydrocodeine 60mg modified-release tablets | 4070200 | Opioid Analgesics |
| Pentazocine hydrochloride | Pentazocine 25mg tablets | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 50mg tablets | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 3mg/5ml linctus paediatric sugar free | 3090100 | Cough Suppressants |
| Methadone hydrochloride | Methadone 1mg/ml oral solution | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine sulfate | MST Continus 30mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 50mg/ml injection | 4070200 | Opioid Analgesics |
| Morphine sulfate | Oramorph sr 10mg Tablet (Boehringer Ingelheim) | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Buprenorphine 400microgram sublingual tablets sugar free | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Diamorphine Hydrochloride | Diamorphine 10mg Injection (Manufacturer unknown) | 4070200 | Opioid Analgesics |
| Meptazinol hydrochloride | Meptazinol 200mg tablets | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg soluble tablets sugar free | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Temgesic 200microgram sublingual tablets (RB Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Zydol SR 100mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 50mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Df118 40mg Tablet (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Sevredol 10mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dextromoramide tartrate | Dextromoramide 5mg tablets | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 100mg modified-release capsules | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 100mg modified-release tablets | 4070200 | Opioid Analgesics |
| Dextromoramide tartrate | Palfium 5mg tablets (Roche Products) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 10mg tablets | 4070200 | Opioid Analgesics |
| Morphine sulfate | MST Continus 5mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| į. | | | |

| Codeine phosphate | Galcodine 3mg/5ml Oral solution (Thornton & Ross) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics | l |
|-----------------------------|--|----------------------------|---|---|
| Morphine sulfate | MST Continus Suspension 60mg granules sachets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Morphine sulfate | MST Continus 60mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Fentanyl | Fentanyl 100micrograms/hour transdermal patches | 4070200 | Opioid Analgesics | İ |
| Morphine sulfate | Oramorph 10mg/5ml oral solution unit dose vials (Boehringer Ingelheim) | 4070200 | Opioid Analgesics | İ |
| Codeine Phosphate | Codeine phosphate 15mg/5ml diabetic oral solution | 03090100/04070200 | Cough Suppressants/Opioid Analgesics | İ |
| Dihydrocodeine tartrate | Dihydrocodeine 40mg tablets | 4070200 | Opioid Analgesics | İ |
| Tramadol hydrochloride | Tramadol 150mg modified-release capsules | 4070200 | Opioid Analgesics | İ |
| Tramadol Hydrochloride | Tramadol 24 Modified-release tablet | 4070200 | Opioid Analgesics | İ |
| Tramadol Hydrochloride | Tramadol 24 Modified-release tablet | 4070200 | Opioid Analgesics | İ |
| Fentanyl | Durogesic 50micrograms transdermal patches (Janssen-Cilag) | 4070200 | Opioid Analgesics | İ |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 5mg powder for injection solution | 4070200 | Opioid Analgesics | İ |
| Hydromorphone hydrochloride | Hydromorphone 2.6mg capsules | 4070200 | Opioid Analgesics | İ |
| Hydromorphone hydrochloride | Hydromorphone 1.3mg capsules | 4070200 | Opioid Analgesics | İ |
| Tramadol hydrochloride | Zydol SR 200mg tablets (Grunenthal) | 4070200 | Opioid Analgesics | İ |
| Methadone hydrochloride | Methadone 2mg/5ml linctus | 3090100 | Cough Suppressants | İ |
| Tramadol Hydrochloride | Tramadol 12 Modified-release tablet | 4070200 | Opioid Analgesics | İ |
| Methadone hydrochloride | Physeptone 5mg tablets (Martindale Pharmaceuticals) | 03090100/04070200/04105200 | Cough Suppressants/Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence | İ |
| Morphine sulfate | Sevredol 10mg/5ml oral solution (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Morphine Sulphate | Morphine sulphate 12 20mg Modified-release capsule | 4070200 | Opioid Analgesics | İ |
| Codeine phosphate | Galcodine 15mg/5ml linctus (Thornton & Ross) | 3090100 | Cough Suppressants | İ |
| Oxycodone hydrochloride | Oxycodone 10mg capsules | 4070200 | Opioid Analgesics | İ |
| Oxycodone hydrochloride | OxyContin 10mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Fentanyl citrate | Fentanyl 400microgram lozenges | 4070200 | Opioid Analgesics | İ |
| Morphine Sulphate | Morphine sulphate 12 50mg Modified-release capsule | 4070200 | Opioid Analgesics | İ |
| Fentanyl | Durogesic 75micrograms transdermal patches (Janssen-Cilag) | 4070200 | Opioid Analgesics | İ |
| Morphine hydrochloride | Morphine hydrochloride 15mg suppositories | 4070200 | Opioid Analgesics | İ |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 30mg powder for injection solution | 4070200 | Opioid Analgesics | İ |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 10mg powder for injection solution | 4070200 | Opioid Analgesics | İ |
| Morphine sulfate | Morphine 10mg modified-release tablets | 4070200 | Opioid Analgesics | İ |
| Fentanyl citrate | Fentanyl 600microgram lozenges | 4070200 | Opioid Analgesics | İ |
| Fentanyl citrate | Fentanyl 800microgram lozenges | 4070200 | Opioid Analgesics | İ |
| Morphine sulfate | MST Continus 15mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Morphine tartrate/Cyclizine | Cyclimorph 10 solution for injection 1ml ampoules (AMCo) | 4070200 | Opioid Analgesics | ١ |
| Morphine sulfate | Morphine sulfate 10mg/5ml oral solution | 4070200 | Opioid Analgesics | İ |
| Oxycodone hydrochloride | Oxycodone 10mg modified-release tablets | 4070200 | Opioid Analgesics | İ |
| | | | | |

| Buprenorphine | Transtec 35micrograms/hour transdermal patches (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
|---|---|---|---|
| Morphine sulfate | MST Continus 100mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 10mg modified-release capsules | 4070200 | Opioid Analgesics |
| Buprenorphine | Transtec 52.5micrograms/hour transdermal patches (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Buprenorphine 8mg sublingual tablets sugar free | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Zydol SR 150mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Buprenorphine | Transtec 70micrograms/hour transdermal patches (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Subutex 8mg sublingual tablets (RB Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Tramadol 200mg modified-release capsules | 4070200 | Opioid Analgesics |
| Morphine sulfate | Sevredol 20mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 120mg modified-release tablets | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 20mg/ml oral solution sugar free | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 75micrograms/hour transdermal patches | 4070200 | Opioid Analgesics |
| Morphine sulfate | Sevredol 50mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone Hydrochloride | Oxycodone hydrochloride 10mg/ml injection | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 5mg tablets | 03090100/04070200/04105200 | Cough Suppressants/Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 500mg powder for injection solution | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 100mg powder for injection solution | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Buprenorphine 2mg sublingual tablets sugar free | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | | | |
| Oxycodone hydrochloride | OxyNorm 5mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol | OxyNorm 5mg capsules (Napp Pharmaceuticals) Tramadol 37.5mg / Paracetamol 325mg tablets | 4070200 4070200 | Opioid Analgesics Opioid Analgesics |
| | | | |
| Tramadol | Tramadol 37.5mg / Paracetamol 325mg tablets | 4070200 | Opioid Analgesics |
| Tramadol hydrochlorida/Paracetamol Oxycodone hydrochloride | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets | 4070200 4070200 | Opioid Analgesics Opioid Analgesics |
| Tramadol hydrochlorida/Paracetamol Oxycodone hydrochloride Oxycodone hydrochloride | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free | 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol hudsochloride/Parasatamol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets | 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol hudsochloride/Parasotamol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free | 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol Nudrochloride / Daracetamol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets | 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol Dycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets Oxycodone 5mg capsules | 4070200 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol bydrochloride/Dayscotamol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride Oxycodone hydrochloride Buprenorphine | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets Oxycodone 5mg capsules Buprenorphine 35micrograms/hour transdermal patches | 4070200 4070200 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol bydrochloride/Paracatanol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride Oxycodone hydrochloride Buprenorphine Morphine Sulphate | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets Oxycodone 5mg capsules Buprenorphine 35micrograms/hour transdermal patches Morphine sulphate 10mg/ml Injection | 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol budrochloride/Paracatanol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride Oxycodone hydrochloride Buprenorphine Morphine Sulphate Buprenorphine | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets Oxycodone 5mg capsules Buprenorphine 35micrograms/hour transdermal patches Morphine sulphate 10mg/ml Injection Buprenorphine 52.5micrograms/hour transdermal patches | 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol hydrochloride/Darscatamol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride Oxycodone hydrochloride Buprenorphine Morphine Sulphate Buprenorphine Oxycodone hydrochloride | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets Oxycodone 5mg capsules Buprenorphine 35micrograms/hour transdermal patches Morphine sulphate 10mg/ml Injection Buprenorphine 52.5micrograms/hour transdermal patches Oxycodone 80mg modified-release tablets | 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol hydrochloride/Parsoctamol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride Oxycodone hydrochloride Buprenorphine Morphine Sulphate Buprenorphine Oxycodone hydrochloride Fentanyl | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets Oxycodone 5mg capsules Buprenorphine 35micrograms/hour transdermal patches Morphine sulphate 10mg/ml Injection Buprenorphine 52.5micrograms/hour transdermal patches Oxycodone 80mg modified-release tablets Durogesic DTrans 25micrograms transdermal patches (Janssen-Cilag) | 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol hydrochloride/Parsoctamol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride Oxycodone hydrochloride Buprenorphine Morphine Sulphate Buprenorphine Oxycodone hydrochloride Fentanyl Fentanyl | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets Oxycodone 5mg capsules Buprenorphine 35micrograms/hour transdermal patches Morphine sulphate 10mg/ml Injection Buprenorphine 52.5micrograms/hour transdermal patches Oxycodone 80mg modified-release tablets Durogesic DTrans 25micrograms transdermal patches (Janssen-Cilag) Durogesic DTrans 50micrograms transdermal patches (Janssen-Cilag) | 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |
| Tramadol hydrochloride/Parsostamol Oxycodone hydrochloride Oxycodone hydrochloride Oxycodone hydrochloride Morphine sulfate Oxycodone hydrochloride Oxycodone hydrochloride Buprenorphine Morphine Sulphate Buprenorphine Oxycodone hydrochloride Fentanyl Fentanyl Diamorphine Hydrochloride | Tramadol 37.5mg / Paracetamol 325mg tablets Oxycodone 20mg modified-release tablets Oxycodone 5mg/5ml oral solution sugar free Oxycodone 40mg modified-release tablets Morphine 20mg modified-release granules sachets sugar free Oxycodone 5mg modified-release tablets Oxycodone 5mg capsules Buprenorphine 35micrograms/hour transdermal patches Morphine sulphate 10mg/ml Injection Buprenorphine 52.5micrograms/hour transdermal patches Oxycodone 80mg modified-release tablets Oxycodone 80mg modified-release tablets Durogesic DTrans 25micrograms transdermal patches (Janssen-Cilag) Durogesic DTrans 50micrograms transdermal patches (Janssen-Cilag) Diamorphine 3mg/5ml oral solution | 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 4070200 | Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics Opioid Analgesics |

| Morphine Sulphate | Morphine sulphate 12 30mg Modified-release capsule | 4070200 | Opioid Analgesics | l |
|-----------------------------|---|-------------------|--|---|
| Buprenorphine | Buprenorphine 10micrograms/hour transdermal patches | 4070200 | Opioid Analgesics | |
| Buprenorphine | Buprenorphine 20micrograms/hour transdermal patches | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | Oxycodone 20mg capsules | 4070200 | Opioid Analgesics | |
| Buprenorphine | Buprenorphine 5micrograms/hour transdermal patches | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 20mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 20mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Fentanyl | Durogesic DTrans 75micrograms transdermal patches (Janssen-Cilag) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 10mg/ml concentrate oral solution (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Buprenorphine Hydrochloride | Temgesic 0.3mg/ml Injection (Reckitt Benckiser Healthcare (UK)) | 4070200 | Opioid Analgesics | |
| Dihydrocodeine tartrate | Df118 10mg/5ml Oral solution (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine sulfate 15mg suppositories | 4070200 | Opioid Analgesics | |
| Buprenorphine | BuTrans 5micrograms/hour transdermal patches (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Morphine Hydrochloride | Morphine hcl Oral solution (Thornton and Ross) | 4070200 | Opioid Analgesics | |
| Dextromoramide Tartrate | Palfium 5mg/ml Injection (Roche Products) | 4070200 | Opioid Analgesics | |
| Diamorphine Hydrochloride | Diamorphine 100mg Injection (Manufacturer unknown) | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine 30mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Dextromoramide tartrate | Dextromoramide 10mg tablets | 0 | Unknown | |
| Diamorphine Hydrochloride | Diamorphine 5mg Injection (Manufacturer unknown) | 4070200 | Opioid Analgesics | |
| Buprenorphine hydrochloride | Temgesic 400microgram sublingual tablets (RB Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence | |
| Morphine sulfate | MST Continus 200mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Diamorphine Hydrochloride | Diamorphine 30mg Injection (Manufacturer unknown) | 4070200 | Opioid Analgesics | |
| Morphine Sulphate | Morphine sulphate 30mg/ml Injection | 4070200 | Opioid Analgesics | |
| Pentazocine hydrochloride | Fortral 25mg tablets (Zentiva) | 4070200 | Opioid Analgesics | |
| Tramadol Hydrochloride | Tramadol 12 Modified-release tablet | 4070200 | Opioid Analgesics | |
| Meptazinol hydrochloride | Meptid 200mg Tablet (Shire Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Dihydrocodeine tartrate | DHC Continus 60mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Diamorphine hydrochloride | Diamorphine 5mg/5ml Oral solution (Manufacturer unknown) | 4070200 | Opioid Analgesics | |
| Morphine hydrochloride | Morphine hydrochloride 30mg suppositories | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine 60mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Diamorphine hydrochloride | Diamorphine 10mg Tablet (Manufacturer unknown) | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Oramorph 20mg/ml concentrated oral solution (Boehringer Ingelheim) | 4070200 | Opioid Analgesics | |
| Cyclizine hydrochloride/ | Diconal tablets (Amdipharm Plc) | 4070200 | Opioid Analgesics | |
| Morphine tartrate/Cyclizine | Cyclimorph 15 solution for injection 1ml ampoules (AMCo) | 4070200 | Opioid Analgesics | |
| Diamorphine Hydrochloride | Diamorphine 500mg Injection (Manufacturer unknown) | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine 20mg tablets | 4070200 | Opioid Analgesics | 1 |
| | | | | |

| Morphine sulfate | Morphine 100mg modified-release tablets | 4070200 | Opioid Analgesics |
|-----------------------------|--|-------------------|--|
| Dihydrocodeine tartrate | DHC Continus 90mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | DHC Continus 120mg tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 90mg modified-release tablets | 4070200 | Opioid Analgesics |
| Hydromorphone hydrochloride | Hydromorphone 4mg modified-release capsules | 4070200 | Opioid Analgesics |
| Hydromorphone hydrochloride | Palladone 2.6mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Hydromorphone hydrochloride | Palladone SR 4mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Hydromorphone hydrochloride | Palladone SR 2mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | MXL 30mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | MXL 60mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | MXL 120mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | MXL 90mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zamadol SR 50mg capsules (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zamadol SR 100mg capsules (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate 24 60mg Modified-release capsule | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 15mg modified-release tablets | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 5mg modified-release tablets | 4070200 | Opioid Analgesics |
| Hydromorphone hydrochloride | Palladone 1.3mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 100mg modified-release granules sachets sugar free | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 1mg/ml oral solution sugar free | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Tramadol 100mg effervescent powder sachets sugar free | 0 | Unknown |
| Oxycodone hydrochloride | OxyNorm liquid 5mg/5ml oral solution (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | OxyContin 40mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone Hydrochloride | Oxynorm 10mg/ml Solution for injection (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 10mg Tablet (Aurum Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate 12 60mg Modified-release capsule | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | OxyNorm 10mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | OxyContin 80mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Subutex 2mg sublingual tablets (RB Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Buprenorphine | BuTrans 10micrograms/hour transdermal patches (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | MXL 150mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Df118 50mg/ml Injection (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dextromoramide tartrate | Palfium 10mg tablets (Roche Products) | 0 | Unknown |
| Pentazocine Lactate | Pentazocine 30mg/ml injection | 4070200 | Opioid Analgesics |
| Morphine Anhydrous | Morphine 10mg/ml Tincture | 4070200 | Opioid Analgesics |
| Morphine sulfate | Sevredol 10mg Suppository (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| 1 | | | |

| Pentazocine hydrochloride | Fortral 50mg Capsule (Sanofi-Synthelabo) | 4070200 | Opioid Analgesics | ĺ |
|------------------------------|---|-------------------|--|---|
| Oxycodone hydrochloride | Oxycodone hydrochloride 30mg suppositories | 0 | Unknown | |
| Fentanyl | Durogesic 100micrograms transdermal patches (Janssen-Cilag) | 4070200 | Opioid Analgesics | l |
| Meptazinol hydrochloride | Meptid 100mg/1ml solution for injection ampoules (Almirall) | 4070200 | Opioid Analgesics | l |
| Tramadol hydrochloride | Zydol 50mg soluble tablets (Grunenthal) | 4070200 | Opioid Analgesics | l |
| Papaveretum | Papaveretum 15.4mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics | l |
| Tramadol hydrochloride | Zydol 100mg/2ml solution for injection ampoules (Grunenthal) | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Oramorph 30mg/5ml oral solution unit dose vials (Boehringer Ingelheim) | 4070200 | Opioid Analgesics | l |
| Oxycodone hydrochloride | Oxycodone 10mg/ml oral solution sugar free | 4070200 | Opioid Analgesics | l |
| Tramadol hydrochloride | Tramadol 100mg/2ml solution for injection ampoules | 4070200 | Opioid Analgesics | |
| Tramadol hydrochloride | Tramadol 75mg modified-release tablets | 0 | Unknown | |
| Tramadol hydrochloride | Tramadol 50mg effervescent powder sachets sugar free | 0 | Unknown | l |
| Buprenorphine | Buprenorphine 70micrograms/hour transdermal patches | 4070200 | Opioid Analgesics | |
| Morphine Sulphate | Morphine sulphate 24 30mg Modified-release capsule | 4070200 | Opioid Analgesics | |
| Methadone hydrochloride | Methadone 10mg/ml oral solution sugar free | 4105200 | Drugs Used In Substance Dependence - Opioid Dependence | |
| Tramadol hydrochloride | Tramadol 50mg orodispersible tablets sugar free | 4070200 | Opioid Analgesics | l |
| Tramadol hydrochloride | Tramadol 300mg modified-release tablets | 4070200 | Opioid Analgesics | l |
| Tramadol hydrochloride | Tramadol 400mg modified-release tablets | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Morcap SR 20mg capsules (Hospira UK) | 0 | Unknown | |
| Meptazinol hydrochloride | Meptazinol 100mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Morphine 200mg modified-release tablets | 4070200 | Opioid Analgesics | l |
| Fentanyl citrate | Fentanyl 200microgram lozenges | 4070200 | Opioid Analgesics | l |
| Fentanyl | Durogesic DTrans 12micrograms transdermal patches (Janssen-Cilag) | 4070200 | Opioid Analgesics | l |
| Cyclizine hydrochloride/ | Dipipanone 10mg / Cyclizine 30mg tablets | 4070200 | Opioid Analgesics | l |
| Dextropropoxyphene Napsylate | Dextropropoxyphene 60mg capsules | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine sulfate 15mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Morphine 60mg modified-release granules sachets sugar free | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Morphine sulfate 10mg suppositories | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | MST Continus Suspension 100mg granules sachets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Codeine Phosphate/Ibuprofen | Ibuprofen and codeine 200mg + 12.5mg Tablet | 04070200/10010100 | Opioid Analgesics/Non-steroidal Anti-inflammatory Drugs | l |
| Morphine sulfate | Oramorph 100mg/5ml oral solution unit dose vials (Boehringer Ingelheim) | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | MST Continus Suspension 30mg granules sachets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Buprenorphine hydrochloride | Subutex 0.4mg sublingual tablets (RB Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence | l |
| Fentanyl citrate | Actiq 200microgram lozenges with integral oromucosal applicator (Teva UK) | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Zomorph 10mg modified-release capsules (ProStrakan) | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Zomorph 30mg modified-release capsules (ProStrakan) | 4070200 | Opioid Analgesics | l |
| | | | | |

| Morphine sulfate | Morphine sulfate 20mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics | ĺ |
|-----------------------------|---|----------------------------|--|---|
| Morphine sulfate | Morphine sulfate 30mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine sulfate 60mg/2ml solution for injection ampoules | 4070200 | Opioid Analgesics | l |
| Buprenorphine | BuTrans 20micrograms/hour transdermal patches (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | l |
| Diamorphine hydrochloride | Diamorphine 15mg/5ml Oral solution (Manufacturer unknown) | 4070200 | Opioid Analgesics | l |
| Morphine Sulphate | Morcap sr 20mg Modified-release capsule (Faulding Pharmaceuticals (Dbl)) | 4070200 | Opioid Analgesics | |
| Tramadol hydrochloride | Zamadol 50mg capsules (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics | l |
| Morphine Sulphate | Morphine sulphate 24 200mg Modified-release capsule | 4070200 | Opioid Analgesics | l |
| Morphine Sulphate | Morphine sulphate 100mg Modified-release capsule | 4070200 | Opioid Analgesics | l |
| Morphine Sulphate | Morphine sulphate 12 100mg Modified-release capsule | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Zomorph 100mg modified-release capsules (ProStrakan) | 4070200 | Opioid Analgesics | l |
| Methadone Hydrochloride | Methadone 10mg/ml Injection | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence | l |
| Morphine sulfate | Morphine sulfate 30mg/5ml oral solution unit dose vials sugar free | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Morphine 30mg modified-release granules sachets sugar free | 4070200 | Opioid Analgesics | l |
| Phenazocine Hydrobromide | Phenazocine hydrobromide 5mg tablet | 4070200 | Opioid Analgesics | l |
| Tramadol hydrochloride | Tramake 50mg capsules (Galen) | 4070200 | Opioid Analgesics | l |
| Fentanyl | Durogesic DTrans 100micrograms transdermal patches (Janssen-Cilag) | 4070200 | Opioid Analgesics | l |
| Fentanyl citrate | Fentanyl 1.2mg lozenges | 4070200 | Opioid Analgesics | l |
| Diamorphine Hydrochloride | Diamorphine 10mg/5ml oral solution | 4070200 | Opioid Analgesics | l |
| Papaveretum | Papaveretum 7.7mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics | |
| Codeine phosphate | Codeine phosphate powder | 50030000 | Extemporaneous Item | l |
| Morphine Sulphate | Morphine sulphate 24 90mg Modified-release capsule | 4070200 | Opioid Analgesics | |
| Hydromorphone hydrochloride | Hydromorphone 2mg modified-release capsules | 4070200 | Opioid Analgesics | |
| Diamorphine hydrochloride | Diamorphine hydrochloride powder | 4070200 | Opioid Analgesics | |
| Hydromorphone hydrochloride | Hydromorphone 8mg modified-release capsules | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine 50mg tablets | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Zomorph 200mg modified-release capsules (ProStrakan) | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Zomorph 60mg modified-release capsules (ProStrakan) | 4070200 | Opioid Analgesics | l |
| Codeine phosphate | Codeine 6.75mg/5ml oral solution | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | |
| Morphine Sulphate | Morphine sulphate 10mg/ml Injection | 4070200 | Opioid Analgesics | l |
| Tramadol hydrochloride | Zydol XL 300mg tablets (Grunenthal) | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Oramorph sr 30mg Tablet (Boehringer Ingelheim) | 4070200 | Opioid Analgesics | l |
| Morphine tartrate/Cyclizine | Morphine tartrate 10mg/1ml / Cyclizine tartrate 50mg/1ml solution for injection | 4070200 | Opioid Analgesics | l |
| Tramadol hydrochloride | Zydol XL 200mg tablets (Grunenthal) | 4070200 | Opioid Analgesics | l |
| Fentanyl | Tilofyl 75micrograms/hour transdermal patches (Tillomed Laboratories) | 4070200 | Opioid Analgesics | l |
| Nalbuphine Hydrochloride | Nubain 10mg/ml Injection (Bristol-Myers Squibb Pharmaceuticals) | 4070200 | Opioid Analgesics | l |
| | | | | |

| Pethidine hydrochloride | Pethidine 100mg/2ml solution for injection ampoules | 4070200 | Opioid Analgesics |
|---|---|-------------------|--|
| Phenazocine Hydrobromide | Narphen 5mg Tablet (Smith & Nephew Healthcare) | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 50mg Tablet (Roche Products) | 4070200 | Opioid Analgesics |
| Morphine tartrate/ Cyclizine | Morphine tartrate 15mg/1ml / Cyclizine tartrate 50mg/1ml solution for injection | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 50mg/1ml solution for injection ampoules | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Dextromoramide Tartrate | Dextromoramide 5mg/ml injection | 4070200 | Opioid Analgesics |
| Pentazocine lactate | Pentazocine 50mg suppositories | 0 | Unknown |
| Morphine sulfate | Oramorph sr 60mg Tablet (Boehringer Ingelheim) | 4070200 | Opioid Analgesics |
| Methadone Hydrochloride | Physeptone 10mg/ml Injection (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine sulfate | MXL 200mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Sevredol 20mg/ml concentrated oral solution (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate 12 200mg Modified-release capsule | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Actiq 400microgram lozenges with integral oromucosal applicator (Teva UK) | 4070200 | Opioid Analgesics |
| Morphine anhydrous/ | J Collis Browne's mixture (Thornton & Ross) | 01020200/04070200 | Other Antispasmodics/Opioid Analgesics |
| Pethidine hydrochloride/ | Pamergan P100 solution for injection 2ml ampoules (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | SRM-RHOTARD 10mg Modified-release tablet (Pharmacia) | 4070200 | Opioid Analgesics |
| Morphine sulfate | SRM-RHOTARD 30mg Modified-release tablet (Pharmacia) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Oramorph sr 100mg Tablet (Boehringer Ingelheim) | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 10mg tablets | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morcap SR 100mg capsules (Hospira UK) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphgesic SR 10mg tablets (AMCo) | 4070200 | Opioid Analgesics |
| Dextropropoxyphene napsylate | Dextropropoxyphene 60mg capsules | 0 | Unknown |
| Morphine Sulphate | Morcap sr 50mg Modified-release capsule (Faulding Pharmaceuticals (Dbl)) | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride/ | Pethidine 100mg/2ml / Promethazine 50mg/2ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Nalbuphine Hydrochloride | Nalbuphine hc 10mg/ml Injection | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 100mg/5ml oral solution unit dose vials sugar free | 4070200 | Opioid Analgesics |
| Codeine Phosphate/Papaverine | Omnopon 10mg Tablet (Roche Products) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphgesic SR 30mg tablets (AMCo) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphgesic SR 60mg tablets (AMCo) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphgesic SR 100mg tablets (AMCo) | 4070200 | Opioid Analgesics |
| Codeine Phosphate/ Papaverine | Papaveretum 10mg tablet | 4070200 | Opioid Analgesics |
| Hydrochloride/Morphine Hydromorphone hydrochloride | Palladone SR 16mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Hydromorphone hydrochloride | Hydromorphone 16mg modified-release capsules | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Dromadol SR 100mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/10ml solution for injection pre-filled syringes | 4070200 | Opioid Analgesics |
| Levorphanol Tartrate | Dromoran roche 1.5mg Tablet (Roche Products) | 4070200 | Opioid Analgesics |

| Tramadol hydrochloride | Zamadol Melt 50mg tablets (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
|-----------------------------|---|-------------------|--|
| Morphine sulfate | Morphine sulfate 30mg suppositories | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 20mg suppositories | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramake Insts 50mg sachets (Galen) | 0 | Unknown |
| Hydromorphone hydrochloride | Palladone SR 8mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Hydromorphone hydrochloride | Palladone SR 24mg capsules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zydol XL 400mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Physeptone 1mg/ml mixture sugar free (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Dromadol SR 200mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zamadol SR 200mg capsules (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Cocaine/Diamorphine | Diamorphine hydrochloride and cocaine oral solution | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zydol XL 150mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Rhotard Morphine SR 10mg tablets (Sovereign Medical) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Rhotard Morphine SR 30mg tablets (Sovereign Medical) | 4070200 | Opioid Analgesics |
| Fentanyl | Tilofyl 50micrograms/hour transdermal patches (Tillomed Laboratories) | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate 24 120mg Modified-release capsule | 4070200 | Opioid Analgesics |
| Morphine sulfate | Filnarine SR 30mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Morphine sulfate | MST Continus Suspension 200mg granules sachets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 20mg/ml oral solution sugar free | 4105200 | Drugs Used In Substance Dependence - Opioid Dependence |
| Dextromoramide Tartrate | Palfium 10mg Suppository (Roche Products) | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 50mg/5ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Dromadol SR 150mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Alfentanil hydrochloride | Alfentanil 5mg/1ml solution for injection ampoules | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Codeine phosphate | Codeine 15mg/5ml linctus (Thornton & Ross) | 3090100 | Cough Suppressants |
| Tramadol hydrochloride | Zamadol SR 150mg capsules (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diagesil 10mg Injection (Berk Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml linctus (A A H Pharmaceuticals) | 3090100 | Cough Suppressants |
| Codeine phosphate | Codeine 15mg/5ml Oral solution (Approved Prescription Services) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Methadone hydrochloride | Methadose 10mg/ml oral solution concentrate (Rosemont Pharmaceuticals) | 4105200 | Drugs Used In Substance Dependence - Opioid Dependence |
| Methadone hydrochloride | Methadose 20mg/ml oral solution concentrate (Rosemont Pharmaceuticals) | 4105200 | Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Tramake Insts 100mg sachets (Galen) | 0 | Unknown |
| Methadone hydrochloride | Methodex 1mg/ml Mixture (Link Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine Sulphate | Morcap sr 100mg Modified-release capsule (Faulding Pharmaceuticals (Dbl)) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 50mg/2ml solution for injection ampoules | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Diamorphine Hydrochloride | Diaphine 100mg Injection (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Hydromorphone hydrochloride | Hydromorphone 24mg modified-release capsules | 4070200 | Opioid Analgesics |
| 1 | | | |

| Alfentanil Hydrochloride | Alfentanil 500micrograms/ml injection | 15010403 | Opioid Analgesics (in Anaesthesia) |
|------------------------------|--|-------------------|--|
| Morphine Sulphate | Morphine sulphate 20mg/ml Injection | 4070200 | Opioid Analgesics |
| Pethidine | Pethidine with levallorphan tartrate injection | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Fentanyl 1.6mg lozenges | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Sublimaze 50microgram/ml Injection (Janssen-Cilag) | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| Fentanyl citrate | Actiq 600microgram lozenges with integral oromucosal applicator (Teva UK) | 4070200 | Opioid Analgesics |
| Morphine Anhydrous | Nepenthe 8.4mg/ml Injection (Celltech Pharma Europe) | 4070200 | Opioid Analgesics |
| Codeine Phosphate | Codeine phosphate with other ingredient pastilles | 3090100 | Cough Suppressants |
| Dextromoramide Tartrate | Palfium 10mg/ml Injection (Roche Products) | 4070200 | Opioid Analgesics |
| Pentazocine lactate | Fortral 50mg suppositories (Sterwin Medicines) | 0 | Unknown |
| Dextropropoxyphene napsylate | Doloxene 60mg Capsule (Eli Lilly and Company) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Actiq 1.6mg lozenges with integral oromucosal applicator (Teva UK) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Physeptone 1mg/ml mixture (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine sulfate | Filnarine SR 10mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Filnarine SR 100mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Dromadol XL 300mg tablets (IVAX Pharmaceuticals UK) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Actiq 800microgram lozenges with integral oromucosal applicator (Teva UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zamadol 24hr 200mg modified-release tablets (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Filnarine SR 60mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| | OXYCODONE | 0 | Unknown |
| Tramadol hydrochloride | Zamadol 24hr 150mg modified-release tablets (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate 24 150mg Modified-release capsule | 4070200 | Opioid Analgesics |
| Fentanyl | Tilofyl 100micrograms/hour transdermal patches (Tillomed Laboratories) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Metharose 1mg/ml oral solution sugar free (Rosemont Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine Sulphate | Moraxen 35mg Rectal tampon (Schwarz Pharma) | 4070200 | Opioid Analgesics |
| Alfentanil Hydrochloride | Rapifen 100microgram/ml Paediatric injection (Janssen-Cilag) | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Diamorphine Hydrochloride | Diamorphine hydrochloride bpc 1973 3mg/5ml oral solution | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zamadol 24hr 300mg modified-release tablets (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dextromoramide Tartrate | Dextromoramide 10mg suppository | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diaphine 10mg Injection (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 200mg modified-release granules sachets sugar free | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Physeptone 10mg/1ml solution for injection ampoules (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Dromadol SR 75mg tablets (IVAX Pharmaceuticals UK) | 0 | Unknown |
| Pethidine Hydrochloride | Pethidine 50mg/ml Injection (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 5mg/5ml oral solution | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Actiq 1.2mg lozenges with integral oromucosal applicator (Teva UK) | 4070200 | Opioid Analgesics |
| | | | |

| Methadone hydrochloride | Methadone 2mg/5ml Oral solution (Martindale Pharmaceuticals) | 3090100 | Cough Suppressants |
|---------------------------|---|----------------------------|--|
| Tramadol hydrochloride | Tramadol 50mg capsules (IVAX Pharmaceuticals UK) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate powder | 04070200/50030000 | Opioid Analgesics/Extemporaneous Item |
| Methadone hydrochloride | Methadone 2mg/5ml linctus (Thornton & Ross) | 3090100 | Cough Suppressants |
| Morphine Sulphate | Morphine sulphate rapiject 1mg/ml Injection (International Medication Systems (UK)) | 4070200 | Opioid Analgesics |
| Morphine Anhydrous | Morphine 8.4mg/ml elixir | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine powder | 50000000 | Ingredients |
| Diamorphine Hydrochloride | Diaphine 30mg Injection (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 1mg/ml oral solution sugar free (Rosemont Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Levorphanol Tartrate | Levorphanol 1.5mg Tablet | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diagesil 500mg Injection (Berk Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diamorphine 10mg/5ml Oral solution (Manufacturer unknown) | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 3mg/5ml oral solution | 4070200 | Opioid Analgesics |
| Fentanyl | Tilofyl 25micrograms/hour transdermal patches (Tillomed Laboratories) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Dromadol XL 200mg tablets (IVAX Pharmaceuticals UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Dromadol XL 150mg tablets (IVAX Pharmaceuticals UK) | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 50mg/ml Injection (Auden McKenzie (Pharma Division)) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (Wockhardt UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Pentazocine lactate | Pentazocine 30mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Pentazocine lactate | Pentazocine 60mg/2ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulfate 30mg/30ml solution for infusion vials | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (Actavis UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Dromadol XL 400mg tablets (IVAX Pharmaceuticals UK) | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 100mg/10ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 50mg Tablet (Roche Products) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (IVAX Pharmaceuticals UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 15mg/5ml oral solution | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg Capsule (Generics (UK)) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methex 1mg/ml Mixture (Generics (UK)) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Alfentanil hydrochloride | Rapifen 5mg/ml Concentrate for solution for infusion (Janssen-Cilag) | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Oxycodone hydrochloride | OxyNorm 10mg/1ml solution for injection ampoules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 3mg/5ml linctus paediatric (Thornton & Ross) | 0 | Unknown |
| Tramadol hydrochloride | Zamadol 24hr 400mg modified-release tablets (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine anhydrous | Opium tincture | 0 | Unknown |
| Diamorphine Hydrochloride | Diagesil 5mg Injection (Berk Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 100mg/2ml Injection (Roche Products) | 4070200 | Opioid Analgesics |
| | | | |

| Diamorphine hydrochloride | Diamorphine 5mg powder for solution for injection ampoules (Novartis) | 4070200 | Opioid Analgesics |
|-----------------------------|--|----------------------------|--|
| Methadone hydrochloride | Methadone 10mg/1ml solution for injection ampoules | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Methadone Hydrochloride | Methadone 35mg/ml Injection | 4105200 | Drugs Used In Substance Dependence - Opioid Dependence |
| Codeine phosphate | Codeine 15mg/5ml linctus sugar free (A A H Pharmaceuticals) | 3090100 | Cough Suppressants |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Wockhardt UK) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 1mg/ml oral solution (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Pethidine Hydrochloride/ | Promethazine hydrochloride 50mg with pethidine 100mg/2ml injection | 03040102/04070200 | Sedating Antihistamines/Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (IVAX Pharmaceuticals UK) | 4070200 | Opioid Analgesics |
| Tramadol Hydrochloride | Tramadol sr 150mg Modified-release tablet (Winthrop Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Ethylmorphine hydrochloride | Collins 1.5mg/5ml elixir (Collins Elixir Co) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine Oral solution (Berk Pharmaceuticals) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml linctus sugar free (Pinewood Healthcare) | 3090100 | Cough Suppressants |
| Codeine phosphate | Codeine 15mg/5ml Oral solution (Celltech Pharma Europe) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml linctus sugar free (Thornton & Ross) | 3090100 | Cough Suppressants |
| Codeine phosphate | Codeine 15mg/5ml Oral solution (William Ransom) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml Oral solution (Rusco) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Tramadol sr 100mg Modified-release tablet (Winthrop Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol Hydrochloride | Tramadol sr 200mg Modified-release tablet (Winthrop Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (A A H Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine Oral solution (Co-operative) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (A A H Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Actavis UK) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml Oral solution (Nucare Plc) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (Wockhardt UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/5ml oral solution (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 10mg Injection (Hillcross Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (Actavis UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Teva UK) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Genus Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Mylan) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg Tablet (Berk Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate 30mg/ml Injection (Celltech Pharma Europe) | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 10mg powder for solution for injection ampoules (Novartis) | 4070200 | Opioid Analgesics |
| | | | |

| Diamorphine hydrochloride | Diamorphine 30mg powder for solution for injection ampoules (Novartis) | 4070200 | Opioid Analgesics |
|-----------------------------|--|----------------------------|--|
| Codeine phosphate | Codeine 30mg tablets (Kent Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (PLIVA Pharma) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | OxyNorm 20mg/2ml solution for injection ampoules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Oxycodone 20mg/2ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulfate 50mg/50ml solution for infusion vials | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Temgesic 300micrograms/1ml solution for injection ampoules (RB Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Fentanyl citrate | Fentanyl 100micrograms/2ml solution for injection ampoules | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| Oxycodone hydrochloride | Oxycodone 10mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Tramadol Hydrochloride | Tramadol 24 Modified-release tablet | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramquel SR 100mg capsules (Beechmere Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Alfentanil hydrochloride | Alfentanil 1mg/2ml solution for injection ampoules | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Methadone hydrochloride | Methadone 20mg/2ml solution for injection ampoules | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Tradorec XL 200mg tablets (Paladin Labs Europe) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tradorec XL 100mg tablets (Paladin Labs Europe) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 25mg/5ml oral solution (Thornton & Ross) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Larapam SR 100mg tablets (Sandoz) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Fentanyl 500micrograms/10ml solution for injection ampoules | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| Fentanyl | Matrifen 25micrograms/hour transdermal patches (Teva UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tradorec XL 300mg tablets (Paladin Labs Europe) | 4070200 | Opioid Analgesics |
| Fentanyl | Matrifen 100micrograms/hour transdermal patches (Teva UK) | 4070200 | Opioid Analgesics |
| Alfentanil hydrochloride | Alfentanil 5mg/10ml solution for injection ampoules | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Fentanyl | Matrifen 50micrograms/hour transdermal patches (Teva UK) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 50mg/5ml solution for injection ampoules | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Tramadol hydrochloride | Mabron 200mg modified-release tablets (Morningside Healthcare) | 4070200 | Opioid Analgesics |
| Tramadol Hydrochloride | Tramadol 50mg modified-release tablets | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zydol SR 50mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramquel SR 50mg capsules (Beechmere Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 5mg/ml oral solution | 0 | Unknown |
| Tramadol hydrochloride | Tramadol 150mg modified-release tablets | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 200mg modified-release tablets | 4070200 | Opioid Analgesics |
| Alfentanil hydrochloride | Alfentanil 0.1% nasal spray | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Physeptone 50mg/1ml solution for injection ampoules (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Methadone hydrochloride | Physeptone 50mg/1ml solution for injection ampoules (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Methadone hydrochloride | Methadone 35mg/3.5ml solution for injection ampoules | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Pethidine hydrochloride | Pethidine 50mg/1ml solution for injection ampoules (AMCo) | 4070200 | Opioid Analgesics |
| 1 | | | |

| Fentanyl | Fentalis Reservoir 100micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
|--|--|-------------------|--|
| Fentanyl | Fentalis Reservoir 25micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Mabron 100mg modified-release tablets (Morningside Healthcare) | 4070200 | Opioid Analgesics |
| Tramadol Hydrochloride | Tramadol (roi) Tablet | 4070200 | Opioid Analgesics |
| Fentanyl | Fentalis Reservoir 50micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| Fentanyl | Matrifen 12micrograms/hour transdermal patches (Teva UK) | 4070200 | Opioid Analgesics |
| Fentanyl | Mezolar Matrix 12micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| Fentanyl | Osmach 50micrograms/hour transdermal patches (Teva UK) | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 40micrograms/dose transdermal system | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 50mg capsules | 4070200 | Opioid Analgesics |
| Fentanyl | Mezolar Matrix 25micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| Pentazocine Lactate | Pentazocine 30mg/ml Injection (Sterwin Medicines) | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 25mg Tablet (Roche Products) | 4070200 | Opioid Analgesics |
| Remifentanyl hydrochloride | Remifentanil 1mg powder for solution for injection vials | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Tramadol hydrochloride | Larapam SR 200mg tablets (Sandoz) | 4070200 | Opioid Analgesics |
| Buprenorphine Hydrochloride/ | Naloxone hc 2mg + 8mg Tablet | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Fentanyl | Mezolar Matrix 75micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| Fentanyl | Matrifen 75micrograms/hour transdermal patches (Teva UK) | 4070200 | Opioid Analgesics |
| Fentanyl | Fentalis Reservoir 75micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Alfentanil hydrochloride | Rapifen 1mg/2ml solution for injection ampoules (Janssen-Cilag) | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Tramadol hydrochloride | Tramadol 50mg Capsule (Tillomed Laboratories) | 4070200 | Opioid Analgesics |
| Fentanyl | Mezolar Matrix 100micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zamadol 100mg/2ml solution for injection ampoules (Meda Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramquel SR 200mg capsules (Beechmere Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | DF 118 Forte 40mg tablets (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Galcodine 3mg/5ml linctus paediatric (Thornton & Ross) | 3090100 | Cough Suppressants |
| Fentanyl | Osmach 100micrograms/hour transdermal patches (Ratiopharm UK) | 4070200 | Opioid Analgesics |
| Fentanyl | Mezolar Matrix 50micrograms/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| Fentanyl | Osmach 25micrograms/hour transdermal patches (Teva UK) | 4070200 | Opioid Analgesics |
| Dextromoramide tartrate | Palfium 5mg Tablet (IDIS World Medicines) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Eptadone 1mg/ml oral solution (Dee Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Ibuprofen/Codeine phosphate | Solpadeine Migraine Ibuprofen & Codeine tablets (Omega Pharma) | 04070200/10010100 | Opioid Analgesics/Non-steroidal Anti-inflammatory Drugs |
| Fentanyl citrate | Fentanyl 100microgram sublingual tablets sugar free | 4070200 | Opioid Analgesics |
| Naloxone Hydrochloride/ oxycodone Hydrochloride | Oxycodone 10mg / Naloxone 5mg modified-release tablets | 4070200 | Opioid Analgesics |

| Naloxone Hydrochloride/ oxycodone Hydrochloride | Targinact 10mg/5mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
|--|---|-------------------|--|
| Naloxone hydrochloride/ Oxycodone hydrochloride | Targinact 20mg/10mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Naloxone hydrochloride /Oxycodone hydrochloride | Oxycodone 20mg / Naloxone 10mg modified-release tablets | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Marol 100mg modified-release tablets (Morningside Healthcare) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Abstral 100microgram sublingual tablets (ProStrakan) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Zentiva) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Fentanyl 200microgram sublingual tablets sugar free | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Marol 200mg modified-release tablets (Morningside Healthcare) | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Fentanyl 100microgram buccal tablets sugar free | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Effentora 100microgram buccal tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Marol 150mg modified-release tablets (Morningside Healthcare) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Abstral 200microgram sublingual tablets (ProStrakan) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Nobligan retard 100mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Abstral 800microgram sublingual tablets (ProStrakan) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Maxitram SR 200mg capsules (Chiesi) | 4070200 | Opioid Analgesics |
| Meptazinol hydrochloride | Meptid 200mg tablets (Almirall) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Effentora 200microgram buccal tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Abstral 400microgram sublingual tablets (ProStrakan) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Fentanyl 200microgram buccal tablets sugar free | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramulief SR 100mg tablets (AMCo) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramulief SR 200mg tablets (AMCo) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramulief SR 150mg tablets (AMCo) | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Fentanyl 400microgram sublingual tablets sugar free | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Effentora 400microgram buccal tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 10mg/5ml oral solution (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Niche Generics) | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Buprenorphine 2mg sublingual tablets sugar free (Teva UK) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Buprenorphine hydrochloride | Buprenorphine 8mg sublingual tablets sugar free (Teva UK) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Pethidine hydrochloride | Pethidine 50mg tablets (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Maxitram SR 100mg capsules (Chiesi) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Maxitram SR 50mg capsules (Chiesi) | 4070200 | Opioid Analgesics |
| Alfentanil Hydrochloride | Rapifen 500microgram/ml Injection (Janssen-Cilag) | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Fentanyl citrate | Effentora 600microgram buccal tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Buprenorphine 300micrograms/1ml solution for injection ampoules | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |

| Fentanyl citrate | Abstral 600microgram sublingual tablets (ProStrakan) | 4070200 | Opioid Analgesics |
|---------------------------|--|----------------------------|--|
| Morphine sulfate | Morphine 30mg modified-release capsules | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Fentanyl 400microgram buccal tablets sugar free | 4070200 | Opioid Analgesics |
| Naloxone Hydrochloride/ | Oxycodone 5mg / Naloxone 2.5mg modified-release tablets | 4070200 | Opioid Analgesics |
| Naloxone Hydrochloride/ | Targinact 5mg/2.5mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone Hydrochloride | Oxycodone 50mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Almus Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | OxyNorm 50mg/1ml solution for injection ampoules (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride/ | Oxycodone 40mg / Naloxone 20mg modified-release tablets | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramquel SR 150mg capsules (Beechmere Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Maxitram SR 150mg capsules (Chiesi) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Larapam SR 150mg tablets (Sandoz) | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Fentanyl 300microgram sublingual tablets sugar free | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Effentora 800microgram buccal tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride/ | Targinact 40mg/20mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Fentanyl 50micrograms/dose nasal spray | 4070200 | Opioid Analgesics |
| Fentanyl | Osmach 75micrograms/hour transdermal patches (Teva UK) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml Oral solution (Thornton & Ross) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Fentanyl Citrate | Abstral 300microgram sublingual tablets (ProStrakan) | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Fentanyl 600microgram sublingual tablets sugar free | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg tablets (Wockhardt UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine Oral solution (Family Health) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (Teva UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 100mg/2ml Injection (C P Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (Teva UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Methadone hydrochloride | Methadone 1mg/ml oral solution (Rosemont Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine hydrochloride | Morphine hcl 15mg Suppository (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulphate 10mg Suppository (Aurum Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulphate 15mg Suppository (Celltech Pharma Europe) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 1mg/ml Mixture (Macarthy Medical) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Methadone hydrochloride | Methadone 1mg/ml Mixture (Macarthy Medical) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Diamorphine hydrochloride | Diamorphine 30mg powder for solution for injection ampoules (Wockhardt UK) | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 30mg powder for solution for injection ampoules (Wockhardt UK) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/1ml solution for injection ampoules (Martindale) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 100mg modified-release tablets (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Fentanyl | Victanyl 100micrograms/hour transdermal patches (Actavis UK) | 4070200 | Opioid Analgesics |
| · | | | |

| Buprenorphine Hydrochloride/ | Naloxone hc 500micrograms + 2mg Tablet | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
|------------------------------|--|----------------------------|--|
| Pentazocine lactate | Fortral 30mg/1ml solution for injection ampoules (Zentiva) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Df118 30mg Tablet (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride/ | Tramadol 37.5mg / Paracetamol 325mg effervescent tablets sugar free | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride/ | Tramacet 37.5mg/325mg effervescent tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate 10mg Modified-release capsule | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Fentanyl 800microgram sublingual tablets sugar free | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Fentanyl 600microgram buccal tablets sugar free | 4070200 | Opioid Analgesics |
| Fentanyl | Victanyl 75micrograms/hour transdermal patches (Actavis UK) | 4070200 | Opioid Analgesics |
| Fentanyl | Osmanil 75micrograms/hour transdermal patches (Zentiva) | 4070200 | Opioid Analgesics |
| Fentanyl | Osmanil 25micrograms/hour transdermal patches (Zentiva) | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 50mg/ml intramuscular injection (Roche Products) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml Oral solution (Actavis UK) | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 150mg modified-release tablets (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 10mg powder for solution for injection ampoules (Wockhardt UK) | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Instanyl 100micrograms/dose nasal spray (Takeda UK) | 4070200 | Opioid Analgesics |
| Fentanyl | Osmanil 50micrograms/hour transdermal patches (Zentiva) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol sr 50mg Capsule (Hillcross Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Methadone Hydrochloride | Methadone Oral solution | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/1ml suspension for injection vials | 4070200 | Opioid Analgesics |
| Alfentanil hydrochloride | Rapifen Intensive Care 5mg/1ml solution for injection ampoules (Janssen-Cilag) | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Codeine phosphate | Codeine phosphate Oral solution (William Ransom) | 01040200/04070200 | Antimotility Drugs/Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Zentiva) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (Ranbaxy (UK)) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Fentanyl Citrate | Instanyl 200micrograms/dose nasal spray (Takeda UK) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 100mg modified-release capsules | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 200mg modified-release capsules | 4070200 | Opioid Analgesics |
| Alfentanil hydrochloride | Rapifen 5mg/10ml solution for injection ampoules (Janssen-Cilag) | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Methadone hydrochloride | Eptadone 5mg/ml oral solution (Dee Pharmaceuticals) | 0 | Unknown |
| Fentanyl citrate | Sublimaze 500micrograms/10ml solution for injection ampoules (Janssen-Cilag) | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| Fentanyl citrate | Sublimaze 500micrograms/10ml solution for injection ampoules (Janssen-Cilag) | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| Tramadol hydrochloride | Mabron 150mg modified-release tablets (Morningside Healthcare) | 4070200 | Opioid Analgesics |
| Fentanyl | Osmanil 12micrograms/hour transdermal patches (Zentiva) | 4070200 | Opioid Analgesics |
| Fentanyl | Victanyl 50micrograms/hour transdermal patches (Actavis UK) | 4070200 | Opioid Analgesics |
| Pentazocine lactate | Fortral 60mg/2ml solution for injection ampoules (Zentiva) | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Fentanyl 200micrograms/dose nasal spray | 4070200 | Opioid Analgesics |
| | | | |

| Pethidine Hydrochloride | Pethidine injection | 4070200 | Opioid Analgesics | Ī |
|--------------------------|--|-------------------|--|---|
| Fentanyl Citrate | PecFent 100micrograms/dose nasal spray (ProStrakan) | 4070200 | Opioid Analgesics | |
| Fentanyl | Osmanil 100micrograms/hour transdermal patches (Zentiva) | 4070200 | Opioid Analgesics | |
| Fentanyl | Victanyl 25micrograms/hour transdermal patches (Actavis UK) | 4070200 | Opioid Analgesics | |
| Fentanyl citrate | Instanyl 50micrograms/dose nasal spray (Takeda UK) | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine 60mg modified-release capsules | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 30mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 60mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 15mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | Oxycodone 15mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Tapentadol hydrochloride | Tapentadol 200mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Tapentadol hydrochloride | Tapentadol 50mg tablets | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | Oxycodone 30mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 120mg modified-release tablets (Napp Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Fentanyl Citrate | Fentanyl 100micrograms/dose nasal spray | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | Oxycodone 60mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Tapentadol hydrochloride | Palexia 50mg tablets (Grunenthal) | 4070200 | Opioid Analgesics | |
| Tapentadol hydrochloride | Palexia SR 50mg tablets (Grunenthal) | 4070200 | Opioid Analgesics | |
| Tapentadol Hydrochloride | Tapentadol 100mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Tapentadol Hydrochloride | Tapentadol 150mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Tapentadol Hydrochloride | Palexia SR 100mg tablets (Grunenthal) | 4070200 | Opioid Analgesics | |
| Tapentadol Hydrochloride | Tapentadol 50mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Tapentadol hydrochloride | Palexia 75mg tablets (Grunenthal) | 4070200 | Opioid Analgesics | |
| Tapentadol Hydrochloride | Palexia SR 150mg tablets (Grunenthal) | 4070200 | Opioid Analgesics | |
| Oxycodone Hydrochloride | Oxycodone 120mg modified-release tablets | 4070200 | Opioid Analgesics | |
| Tramadol hydrochloride | Tramadol 200mg modified-release capsules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Fentanyl citrate | PecFent 400micrograms/dose nasal spray (ProStrakan) | 4070200 | Opioid Analgesics | |
| Tapentadol hydrochloride | Tapentadol 75mg tablets | 4070200 | Opioid Analgesics | |
| Fentanyl citrate | Fentanyl 800microgram buccal tablets sugar free | 4070200 | Opioid Analgesics | |
| Fentanyl | Fencino 12micrograms/hour transdermal patches (DB Ashbourne) | 4070200 | Opioid Analgesics | |
| Fentanyl | Fencino 50micrograms/hour transdermal patches (DB Ashbourne) | 4070200 | Opioid Analgesics | |
| Methadone hydrochloride | Physeptone 20mg/2ml solution for injection ampoules (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence | |
| Tramadol Hydrochloride | Tramadol 100mg/ml oral drops | 4070200 | Opioid Analgesics | |
| Tramadol hydrochloride | Zeridame SR 150mg tablets (Actavis UK) | 4070200 | Opioid Analgesics | |
| Fentanyl | Fencino 75micrograms/hour transdermal patches (DB Ashbourne) | 4070200 | Opioid Analgesics | |
| Fentanyl | Fencino 100micrograms/hour transdermal patches (DB Ashbourne) | 4070200 | Opioid Analgesics | |
| | | | | 1 |

| Tapentadol hydrochloride | Palexia SR 200mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
|---------------------------|---|----------------------------|--|
| Fentanyl | Fencino 25micrograms/hour transdermal patches (DB Ashbourne) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg tablets (Ranbaxy (UK)) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Papaveretum | Omnopon 15.4mg/ml Injection (Roche Products) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Filnarine SR 200mg tablets (Teva UK) | 4070200 | Opioid Analgesics |
| Tapentadol Hydrochloride | Tapentadol 250mg modified-release tablets | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 75micrograms/hr Transdermal patch (Sandoz) | 4070200 | Opioid Analgesics |
| Tapentadol hydrochloride | Palexia SR 250mg tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/10ml solution for injection Minijet pre-filled syringes (UCB) | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 10mg Injection (Approved Prescription Services) | 4070200 | Opioid Analgesics |
| Diamorphine Hydrochloride | Diamorphine hydrochloride 30mg Injection (Hillcross Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 1mg/ml oral solution sugar free (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine sulfate | Morphine 90mg modified-release capsules | 4070200 | Opioid Analgesics |
| Fentanyl Citrate | Fentanyl 400micrograms/dose nasal spray | 4070200 | Opioid Analgesics |
| Tramadol Hydrochloride | Tramadol (roi) Tablet | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 150mg modified-release capsules | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine phosphate 15mg Tablet (Wockhardt UK) | 01040200/04070200 | Antimotility Drugs/Opioid Analgesics |
| Morphine sulfate | Morphine 120mg modified-release capsules | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine phosphate 30mg Tablet (Wockhardt UK) | 01040200/04070200 | Antimotility Drugs/Opioid Analgesics |
| Morphine sulfate | M-eslon 10mg Capsule (Trinity Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (Ranbaxy (UK)) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Codeine Phosphate | Codeine phosphate oral solution | 01040200/03090100 | Antimotility Drugs/Cough Suppressants |
| Tramadol hydrochloride | Tramadol 200mg modified-release tablets (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 100mg/2ml solution for injection ampoules (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Almus Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (Almus Pharmaceuticals) | 0 | Unknown |
| Pethidine hydrochloride | Pethidine 100mg/2ml solution for injection ampoules (Actavis UK) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg tablets (Actavis UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 5mg powder for solution for injection vials (Teva UK) | 4070200 | Opioid Analgesics |
| Fentanyl citrate | Fentanyl 100micrograms/2ml solution for injection ampoules (Martindale Pharma) | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| Diamorphine hydrochloride | Diamorphine 5mg powder for solution for injection ampoules | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 5mg powder for solution for injection vials | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 10mg powder for solution for injection ampoules | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 30mg powder for solution for injection ampoules | 4070200 | Opioid Analgesics |
| Fentanyl | Durogesic DTrans 50micrograms transdermal patches (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| | | | |

| Diamorphine hydrochloride | Diamorphine 10mg powder for solution for injection vials | 4070200 | Opioid Analgesics | |
|-----------------------------|---|----------------------------|--|--|
| Diamorphine | Diamorphine 30mg powder for solution for injection vials | 4070200 | Opioid Analgesics | |
| Diamorphine hydrochloride | Diamorphine 100mg powder for solution for injection ampoules | 4070200 | Opioid Analgesics | |
| Diamorphine | Diamorphine 100mg powder for solution for injection vials | 4070200 | Opioid Analgesics | |
| Codeine phosphate | Codeine 3mg/5ml linctus paediatric | 0 | Unknown | |
| Tramadol hydrochloride | Marol 150mg modified-release tablets (Teva UK) | 4070200 | Opioid Analgesics | |
| Tramadol hydrochloride | Marol 100mg modified-release tablets (Teva UK) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 5mg modified-release tablets (DE Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 20mg modified-release tablets (Lexon (UK)) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 20mg/2ml solution for injection ampoules (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 5mg capsules (Lexon (UK)) | 4070200 | Opioid Analgesics | |
| Morphine sulfate | Morphine sulfate 10mg/1ml solution for injection ampoules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 5mg capsules (DE Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Buprenorphine hydrochloride | Subutex 2mg sublingual tablets (Lexon (UK)) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence | |
| Codeine phosphate | Codeine phosphate 15mg Tablet (Celltech Pharma Europe) | 01040200/04070200 | Antimotility Drugs/Opioid Analgesics | |
| Codeine phosphate | Codeine 15mg tablets (Alliance Healthcare (Distribution)) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | |
| Morphine sulfate | Morphine sulfate 10mg/1ml solution for injection ampoules (UCB Pharma) | 4070200 | Opioid Analgesics | |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Bristol Laboratories) | 4070200 | Opioid Analgesics | |
| Codeine phosphate | Codeine 60mg tablets (A A H Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | |
| Fentanyl | Fentanyl 12micrograms/hour transdermal patches (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 10mg/1ml solution for injection ampoules (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 10mg modified-release tablets (Mawdsley-Brooks & Company) | 4070200 | Opioid Analgesics | |
| Tramadol hydrochloride | Marol 200mg modified-release tablets (Teva UK) | 4070200 | Opioid Analgesics | |
| Fentanyl | Durogesic DTrans 12micrograms transdermal patches (Mawdsley-Brooks & Company) | 4070200 | Opioid Analgesics | |
| Tramadol hydrochloride | Tramadol 100mg modified-release capsules (Alliance Healthcare (Distribution)) | 4070200 | Opioid Analgesics | |
| Fentanyl | Fentanyl 25micrograms/hour transdermal patches (Phoenix Healthcare Distribution) | 4070200 | Opioid Analgesics | |
| Codeine phosphate | Codeine 30mg tablets (Alliance Healthcare (Distribution)) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 20mg capsules (DE Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 10mg/1ml solution for injection ampoules (Mawdsley-Brooks & Company) | 4070200 | Opioid Analgesics | |
| Codeine phosphate | Codeine 30mg tablets (Bristol Laboratories) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | |
| Oxycodone hydrochloride | OxyNorm 10mg capsules (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | OxyContin 80mg modified-release tablets (Mawdsley-Brooks & Company) | 4070200 | Opioid Analgesics | |
| Codeine phosphate | Codeine 15mg tablets (Bristol Laboratories) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | |
| Fentanyl citrate | Fentanyl 100micrograms/2ml solution for injection ampoules (AMCo) | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) | |
| Oxycodone hydrochloride | Longtec 5mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| Oxycodone hydrochloride | Longtec 10mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics | |
| | | | | |

| Oxycodone hydrochloride | Longtec 40mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics | l |
|-----------------------------|---|----------------------------|---|---|
| Pethidine hydrochloride | Pethidine 100mg/10ml solution for injection ampoules (Alliance Healthcare | 4070200 | Opioid Analgesics | İ |
| Tramadol hydrochloride | Tramadol 50mg capsules (Bristol Laboratories) | 4070200 | Opioid Analgesics | İ |
| Oxycodone hydrochloride | OxyNorm 10mg capsules (DE Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Tramadol hydrochloride | Tramadol 50mg capsules (Accord Healthcare) | 4070200 | Opioid Analgesics | İ |
| Oxycodone hydrochloride | OxyContin 10mg modified-release tablets (Lexon (UK)) | 4070200 | Opioid Analgesics | İ |
| Codeine phosphate | Codeine 15mg tablets (Kent Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | İ |
| Codeine phosphate | Codeine 25mg/5ml oral solution (A A H Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | İ |
| Codeine phosphate | Codeine 15mg tablets (Sigma Pharmaceuticals Plc) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | İ |
| Tramadol hydrochloride | Tramadol 100mg modified-release capsules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Hydromorphone hydrochloride | Hydromorphone 10mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics | İ |
| Morphine sulfate | Morphine sulfate 5mg/5ml solution for injection ampoules | 4070200 | Opioid Analgesics | İ |
| Oxycodone hydrochloride | OxyContin 10mg modified-release tablets (DE Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Oxycodone hydrochloride | Longtec 20mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Diamorphine hydrochloride | Diamorphine 10mg powder for solution for injection vials (Teva UK) | 4070200 | Opioid Analgesics | İ |
| Morphine hydrochloride | Morphine hydrochloride 10mg/5ml oral solution (Special Order) | 4070200 | Opioid Analgesics | İ |
| Diamorphine hydrochloride | Diamorphine 10mg powder for solution for injection ampoules (Actavis UK) | 4070200 | Opioid Analgesics | İ |
| Codeine phosphate | Codeine 60mg tablets (Alliance Healthcare (Distribution)) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics | İ |
| Ibuprofen/Codeine phosphate | Ibuprofen and codeine 200mg+12.8mg Tablet (Almus Pharmaceuticals) | 04070200/10010100 | Opioid Analgesics/Non-steroidal Anti-inflammatory Drugs | İ |
| Morphine sulfate | Morphine 10mg modified-release tablets (Sigma Pharmaceuticals Plc) | 4070200 | Opioid Analgesics | İ |
| Pethidine Hydrochloride | Pethidine oral liquid | 4070200 | Opioid Analgesics | İ |
| Morphine sulfate | Morphine sulfate 10mg/1ml solution for injection ampoules (Hameln | 4070200 | Opioid Analgesics | İ |
| Pethidine hydrochloride | Pharmacouticals) Pethidine 50mg/1ml solution for injection ampoules (A A H Pharmacouticals) | 4070200 | Opioid Analgesics | İ |
| Codeine phosphate | Codeine phosphate 60mg Tablet (Wockhardt UK) | 01040200/04070200 | Antimotility Drugs/Opioid Analgesics | İ |
| Morphine Sulphate | Morphine sulphate Capsule | 4070200 | Opioid Analgesics | İ |
| Tramadol hydrochloride | Tramadol 50mg modified-release capsules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Pethidine Hydrochloride | Pethidine capsule | 4070200 | Opioid Analgesics | İ |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Kent Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Morphine sulfate | Morphine sulfate 30mg suppositories (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Morphine Sulphate | Morphine sulphate Oral solution | 4070200 | Opioid Analgesics | İ |
| Oxycodone hydrochloride | Longtec 80mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics | İ |
| Pethidine hydrochloride | Pethidine 50mg tablets (Actavis UK) | 4070200 | Opioid Analgesics | İ |
| Buprenorphine | Transtec 52.5micrograms/hour transdermal patches (DE Pharmaceuticals) | 4070200 | Opioid Analgesics | ١ |
| Fentanyl | Fentanyl 50micrograms/hour transdermal patches (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics | l |
| Morphine sulfate | Morcap SR 50mg capsules (Hospira UK) | 0 | Unknown | ١ |
| Morphine sulfate | Morphine 20mg modified-release capsules | 0 | Unknown | İ |
| | | | | |

| Diamorphine hydrochloride | Diamorphine 500mg powder for solution for injection ampoules | 4070200 | Opioid Analgesics |
|-----------------------------|---|----------------------------|--|
| Nalbuphine hydrochloride | Nubain 10mg/1ml solution for injection ampoules (Bristol-Myers Squibb | 0 | Unknown |
| Codeine phosphate | Codeine 15mg tablets (Phoenix Healthcare Distribution) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 50mg/50ml solution for infusion vials (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Dihydrocodeine Tartrate | Dihydrocodeine 10mg tablets | 03090100/04070200 | Cough Suppressants/Opioid Analgesics |
| Diamorphine | Diamorphine 500mg powder for solution for injection vials | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 100micrograms/hour transdermal patches (Phoenix Healthcare Distribution) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 1mg/ml oral solution sugar free (Thornton & Ross) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine sulfate | Morphine sulphate 15mg/ml Injection (Celltech Pharma Europe) | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 50mg/ml Injection (Roche Products) | 4070200 | Opioid Analgesics |
| Pethidine Hydrochloride | Pethidine 10mg/ml Injection (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 50mg Capsule (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg/1ml solution for injection ampoules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine Sulphate | Morphine sulphate Injection | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/2ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zamadol SR 200mg capsules (Lexon (UK)) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 50mg/50ml solution for infusion vials (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (Phoenix Healthcare Distribution) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Alfentanil Hydrochloride | Alfentanil 5mg/5ml buccal spray | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Oxylan 10mg modified-release tablets (Chanelle Medical UK) | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 25micrograms/hour transdermal patches (Sigma Pharmaceuticals Plc) | 4070200 | Opioid Analgesics |
| Buprenorphine | BuTrans 5micrograms/hour transdermal patches (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/5ml oral solution (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (Almus Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Pethidine hydrochloride | Pethidine 50mg tablets (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Oxylan 10mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Oxylan 20mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg tablets (Teva UK) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Buprenorphine hydrochloride | Prefibin 8mg sublingual tablets (Sandoz) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Codeine phosphate | Codeine 30mg tablets (Waymade Healthcare Plc) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 15mg/1ml solution for injection ampoules (UCB Pharma) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 15mg/1ml solution for injection ampoules (Wockhardt UK) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 10mg/5ml oral solution | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Oxycodone hydrochloride | Oxycodone 5mg/5ml oral solution | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Lynlor 20mg capsules (Actavis UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Zeridame SR 100mg tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| | | | |

| 1 | Codeine phosphate | Codeine 15mg/5ml linctus sugar free (DE Pharmaceuticals) | 3090100 | Cough Suppressants |
|---|-----------------------------|--|-------------------|--|
| | Pethidine hydrochloride | Pethidine 50mg/5ml solution for injection ampoules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Oxycodone hydrochloride | Lynlor 5mg capsules (Actavis UK) | 4070200 | Opioid Analgesics |
| | Buprenorphine hydrochloride | Buprenorphine 2mg sublingual tablets sugar free (A A H Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Diamorphine hydrochloride | Diamorphine 10mg tablets (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Morphine sulfate | Morphine sulfate 10mg suppositories (Martindale Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Tramadol hydrochloride | Tramadol 50mg modified-release capsules (DE Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Oxycodone hydrochloride | Oxylan 20mg modified-release tablets (Chanelle Medical UK) | 4070200 | Opioid Analgesics |
| | Diamorphine hydrochloride | Diamorphine 3mg/5ml oral solution | 4070200 | Opioid Analgesics |
| | Morphine sulfate | Morphine sulfate 1mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| | Pethidine hydrochloride | Pethidine 50mg tablets (Alliance Healthcare (Distribution)) | 4070200 | Opioid Analgesics |
| | Buprenorphine | BuTrans 10micrograms/hour transdermal patches (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| | | Generic Migraleve tablets | 04070200/04070401 | Opioid Analgesics/Treatment Of Acute Migraine |
| | Methadone sulfate | Morphine sulfate 5mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| | Oxycodone hydrochloride | Lynlor 10mg capsules (Actavis UK) | 4070200 | Opioid Analgesics |
| | Morphine hydrochloride | Morphine hydrochloride 10mg/5ml oral solution | 4070200 | Opioid Analgesics |
| | Codeine phosphate | Codeine 15mg/5ml linctus sugar free (Kent Pharmaceuticals) | 3090100 | Cough Suppressants |
| | Fentanyl citrate | Fentanyl 400microgram buccal films sugar free | 4070200 | Opioid Analgesics |
| | | BuTrans 20micrograms/hour transdermal patches (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| | Methadone | Methadone 100mg capsules | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Buprenorphine | Hapoctasin 70micrograms/hour transdermal patches (Actavis UK) | 4070200 | Opioid Analgesics |
| | Fentanyl | Fentanyl 200microgram buccal films sugar free | 4070200 | Opioid Analgesics |
| | Buprenorphine | Hapoctasin 52.5micrograms/hour transdermal patches (Actavis UK) | 4070200 | Opioid Analgesics |
| | Fentanyl | Fentanyl 37.5microgram/hour transdermal patches | 4070200 | Opioid Analgesics |
| | Fentanyl | Mezolar Matrix 37.5microgram/hour transdermal patches (Sandoz) | 4070200 | Opioid Analgesics |
| | Morphine sulfate | Morphine sulfate 10mg/5ml oral solution (Alliance Healthcare (Distribution)) | 4070200 | Opioid Analgesics |
| | Buprenorphine | Transtec 35micrograms/hour transdermal patches (Mawdsley-Brooks & Company) | 4070200 | Opioid Analgesics |
| | Fentanyl citrate | Fentanyl 100micrograms/2ml solution for injection ampoules (A A H Pharmaceuticals) | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| | Oxycodone hydrochloride | Shortec 10mg capsules (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Buprenorphine hydrochloride | Buprenorphine 2mg sublingual tablets sugar free (Actavis UK) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| | Dihydrocodeine tartrate | Dihydrocodeine 30mg tablets (Ranbaxy (UK)) | 4070200 | Opioid Analgesics |
| | Buclizine hydrochloride/ | Generic Migraleve Pink tablets | 04070200/04070401 | Opioid Analgesics/Treatment Of Acute Migraine |
| | Buprenorphine hydrochloride | Tephine 200microgram sublingual tablets (Sandoz) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Alfentanil hydrochloride | Alfentanil 500micrograms/1ml solution for injection ampoules | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| | Morphine sulfate | Morphine sulfate 5mg/10ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| | | | | |

| Tramadol hydrochloride | Tramadol 50mg modified-release capsules (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
|-----------------------------|---|----------------------------|---|
| Oxycodone hydrochloride | Shortec 5mg capsules (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Shortec 20mg capsules (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Buprenorphine | Hapoctasin 35micrograms/hour transdermal patches (Actavis UK) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Oxylan 40mg modified-release tablets (Chanelle Medical UK) | 4070200 | Opioid Analgesics |
| Remifentanyl hydrochloride | Remifentanil 5mg powder for solution for injection vials | 15010403 | Opioid Analgesics (in Anaesthesia) |
| Fentanyl | Fentanyl 25micrograms/hour transdermal patches (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Codeine 30mg suppositories | 4070200 | Opioid Analgesics |
| | Morphine 0.1% in Intrasite gel | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 500micrograms/5ml oral solution | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (Waymade Healthcare Plc) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 5mg powder for solution for injection ampoules (Actavis UK) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tilodol SR 200mg tablets (Sandoz) | 4070200 | Opioid Analgesics |
| | Tapentadol 20mg/ml oral solution sugar free | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 25micrograms/hour transdermal patches (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| Buprenorphine | Transtec 35micrograms/hour transdermal patches (Sigma Pharmaceuticals Plc) | 4070200 | Opioid Analgesics |
| Methadone hydrochloride | Methadone 5mg capsules | 03090100/04070200/04105200 | Cough Suppressants/Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine sulfate | Morphine sulfate 5mg/5ml oral solution | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 15mg tablets (DE Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Codeine phosphate | Codeine 15mg/5ml linctus (Waymade Healthcare Plc) | 3090100 | Cough Suppressants |
| | Opiodur 12micrograms/hour transdermal patches (Pfizer) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg tablets (Waymade Healthcare Plc) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Buprenorphine hydrochloride | Tephine 400microgram sublingual tablets (Sandoz) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Fentanyl | Fentanyl 12micrograms/hour transdermal patches (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 50mg/50ml solution for infusion vials (Alliance Healthcare | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Phoenix Healthcare Distribution) | 4070200 | Opioid Analgesics |
| Fentanyl | Mylafent 75micrograms/hour transdermal patches (Mylan) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 30mg/1ml solution for injection ampoules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine 30mg modified-release tablets (Sigma Pharmaceuticals Plc) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 10mg/1ml solution for injection ampoules (Wockhardt UK) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 50mg/50ml solution for infusion vials (Torbay Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Morningside Healthcare) | 4070200 | Opioid Analgesics |
| | Recivit 267microgram sublingual tablets (Grunenthal) | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 100mg/50ml solution for infusion vials (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tapentadol hydrochloride | Palexia 20mg/ml oral solution (Grunenthal) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Sigma Pharmaceuticals Plc) | 4070200 | Opioid Analgesics |
| | | | |

| ĺ | Oxycodone hydrochloride | Reltebon 40mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
|---|-----------------------------|--|----------------------------|--|
| | Oxycodone hydrochloride | Reltebon 20mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| | Morphine sulfate | Morphine sulfate 20mg/2ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| | Oxycodone hydrochloride | Reltebon 10mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| | Oxycodone hydrochloride | Reltebon 5mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| | Morphine sulfate | Morphine sulfate 2.5mg/5ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| | Codeine phosphate | Codeine 30mg tablets (DE Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| | Oxycodone hydrochloride | Reltebon 80mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| | Buprenorphine hydrochloride | Buprenorphine 200microgram sublingual tablets sugar free (A A H Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Morphine hydrochloride | Morphine hydrochloride 1mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| | Methadone hydrochloride | Methadone hydrochloride powder | 50030000 | Extemporaneous Item |
| | Buprenorphine hydrochloride | Buprenorphine 4mg sublingual tablets sugar free | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Tramadol | Tramacet 37.5mg/325mg tablets (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| | Buprenorphine hydrochloride | Buprenorphine 6mg sublingual tablets sugar free | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Buprenorphine hydrochloride | Buprenorphine 8mg sublingual tablets sugar free (Zentiva) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Tramadol hydrochloride | Tramadol 100mg modified-release capsules (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| | Fentanyl | Fentanyl 12micrograms/hour transdermal patches (Phoenix Healthcare Distribution) | 4070200 | Opioid Analgesics |
| | Pethidine hydrochloride | Pethidine 50mg/1ml solution for injection ampoules (Alliance Healthcare | 4070200 | Opioid Analgesics |
| | Oxycodone hydrochloride | OxyContin 40mg modified-release tablets (DE Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Oxycodone hydrochloride | Reltebon 30mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| | Fentanyl | Fentanyl 133microgram sublingual tablets sugar free | 4070200 | Opioid Analgesics |
| | Fentanyl citrate | Fentanyl 2.5mg/50ml solution for infusion vials | 04070200/15010403 | Opioid Analgesics/Opioid Analgesics (in Anaesthesia) |
| | Morphine sulfate | Morphine sulfate 10mg/1ml solution for injection ampoules (DE Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Codeine phosphate | Codeine 1mg suppositories | 4070200 | Opioid Analgesics |
| | Morphine sulfate | Morphine sulfate 10mg/5ml oral solution (Actavis UK) | 4070200 | Opioid Analgesics |
| | Codeine phosphate | Codeine phosphate powder (Martindale Pharmaceuticals) | 50030000 | Extemporaneous Item |
| | Buprenorphine hydrochloride | Subutex 2mg sublingual tablets (DE Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Oxycodone hydrochloride | Reltebon 60mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| | Tramadol hydrochloride | Tramadol 50mg modified-release capsules (J M McGill) | 4070200 | Opioid Analgesics |
| | Dihydrocodeine tartrate | Dihydrocodeine 10mg/5ml oral solution (Waymade Healthcare Plc) | 4070200 | Opioid Analgesics |
| | Codeine phosphate | Codeine 15mg tablets (Crescent Pharma) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| | Oxycodone hydrochloride | Oxylan 5mg modified-release tablets (Chanelle Medical UK) | 4070200 | Opioid Analgesics |
| | Buprenorphine hydrochloride | Buprenorphine 400microgram sublingual tablets sugar free (Teva UK) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Oxycodone hydrochloride | Reltebon 15mg modified-release tablets (Actavis UK) | 4070200 | Opioid Analgesics |
| | Oxycodone hydrochloride | Longtec 30mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics |
| | Morphine sulfate | Morphine sulfate 2mg/5ml oral solution | 4070200 | Opioid Analgesics |
| | | | | |

| Oxycodone hydrochloride | Longtec 15mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics |
|-----------------------------|---|----------------------------|--|
| Tramadol | Tramadol 37.5mg / Paracetamol 325mg tablets (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 100mg modified-release capsules (Ennogen Healthcare) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Longtec 60mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 100mg modified-release capsules (Icarus Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg/1ml solution for injection ampoules (Alliance Healthcare (Distribution)) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (Sigma Pharmaceuticals Plc) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 50mg/1ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Morphine sulfate | Morphine sulfate 50mg/5ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| | Longtec 120mg modified-release tablets (Qdem Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Buprenorphine hydrochloride | Gabup 8mg sublingual tablets (Martindale Pharmaceuticals) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| Morphine sulfate | Morphine sulfate 10mg/10ml solution for injection ampoules | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Maneo 100mg modified-release tablets (Mylan) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Oxycodone 5mg/5ml oral solution sugar free (Wockhardt UK) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (Crescent Pharma) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Buprenorphine hydrochloride | Buprenorphine 2mg sublingual tablets sugar free (Sigma Pharmaceuticals Plc) | 04070200/04105200 | Opioid Analgesics/Drugs Used In Substance Dependence - Opioid Dependence |
| | Mylafent 12micrograms/hour transdermal patches (Mylan) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 30mg tablets (Mawdsley-Brooks & Company) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg capsules (Kent Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Codeine phosphate | Codeine 60mg tablets (DE Pharmaceuticals) | 01040200/03090100/04070200 | Antimotility Drugs/Cough Suppressants/Opioid Analgesics |
| Fentanyl | Mylafent 50micrograms/hour transdermal patches (Mylan) | 4070200 | Opioid Analgesics |
| Diamorphine hydrochloride | Diamorphine 5mg powder for solution for injection ampoules (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Abtard 30mg modified-release tablets (DB Ashbourne) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Abtard 20mg modified-release tablets (DB Ashbourne) | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 100micrograms/hour transdermal patches (A A H Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Fentanyl | Fentanyl 267microgram sublingual tablets sugar free | 4070200 | Opioid Analgesics |
| Dihydrocodeine tartrate | Dihydrocodeine 30mg/5ml oral solution | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Abtard 40mg modified-release tablets (DB Ashbourne) | 4070200 | Opioid Analgesics |
| Oxycodone hydrochloride | Abtard 80mg modified-release tablets (DB Ashbourne) | 4070200 | Opioid Analgesics |
| Tramadol hydrochloride | Tramadol 50mg modified-release capsules (Cubic Pharmaceuticals) | 4070200 | Opioid Analgesics |
| Z-drugs | | | |
| Zopiclone | Zopiclone 7.5mg tablets | 4010100 | Hypnotics |
| Zopiclone | Zopiclone 3.75mg tablets | 4010100 | Hypnotics |
| Zolpidem tartrate | Zolpidem 5mg tablets | 4010100 | Hypnotics |
| Zolpidem tartrate | Stilnoct 5mg tablets (Sanofi) | 4010100 | Hypnotics |
| Zopiclone | Zimovane 7.5mg tablets (Sanofi) | 4010100 | Hypnotics |
| 1 | | | |

| Zolpidem tartrate | Stilnoct 10mg tablets (Sanofi) | 4010100 | Hypnotics | 1 |
|-------------------|--|---------|-----------|---|
| Zopiclone | Zimovane LS 3.75mg tablets (Sanofi) | 4010100 | Hypnotics | |
| Zopiclone | Zileze 3.75 tablets (Opus Pharmaceuticals) | 4010100 | Hypnotics | |
| Zaleplon | Sonata 10mg capsules (Meda Pharmaceuticals) | 4010100 | Hypnotics | |
| Zaleplon | Zaleplon 10mg capsules | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 10mg tablets | 4010100 | Hypnotics | |
| Zaleplon | Zaleplon 5mg capsules | 4010100 | Hypnotics | |
| Zaleplon | Sonata 5mg capsules (Meda Pharmaceuticals) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg/5ml oral suspension | 4010100 | Hypnotics | |
| Zopiclone | Zileze 7.5 tablets (Opus Pharmaceuticals) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 7.5mg tablets (Actavis UK) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg tablets (Actavis UK) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 5mg Tablet (Winthrop Pharmaceuticals) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg tablets (IVAX Pharmaceuticals UK) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg tablets (Mylan) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 10mg tablets (A A H Pharmaceuticals) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 5mg tablets (A A H Pharmaceuticals) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 7.5mg tablets (IVAX Pharmaceuticals UK) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 7.5mg tablets (Mylan) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 10mg tablets (Mylan) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 7.5mg tablets (PLIVA Pharma) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg tablets (A A H Pharmaceuticals) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg tablets (Teva UK) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 7.5mg tablets (Teva UK) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 7.5mg tablets (Kent Pharmaceuticals) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg tablets (Kent Pharmaceuticals) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 10mg tablets (IVAX Pharmaceuticals UK) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 5mg tablets (Teva UK) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 5mg tablets (IVAX Pharmaceuticals UK) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 10mg Tablet (Winthrop Pharmaceuticals) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 7.5mg tablets (A A H Pharmaceuticals) | 4010100 | Hypnotics | |
| Zolpidem tartrate | Zolpidem 10mg tablets (Teva UK) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 7.5mg tablets (Sandoz) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg/5ml oral solution | 4010100 | Hypnotics | |
| Zopiclone | Zimovane 7.5mg tablets (Lexon (UK)) | 4010100 | Hypnotics | |
| Zopiclone | Zopiclone 3.75mg tablets (Almus Pharmaceuticals) | 4010100 | Hypnotics | |
| | | | | |

| Zopiclone | Zopiclone 7.5mg/5ml oral suspension | 4010100 | Hypnotics |
|-------------------|---|---------|-----------|
| Zopiclone | Zopiclone 7.5mg/5ml oral solution | 4010100 | Hypnotics |
| Zopiclone | Zopiclone 7.5mg tablets (Sigma Pharmaceuticals Plc) | 4010100 | Hypnotics |
| Zolpidem tartrate | Zolpidem 10mg tablets (Zentiva) | 4010100 | Hypnotics |
| Zopiclone | Zopiclone 7.5mg tablets (Phoenix Healthcare Distribution) | 4010100 | Hypnotics |