



This is a repository copy of *Progress and challenges in engineering cyanobacteria as chassis for light-driven biotechnology.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/155392/>

Version: Published Version

Article:

Hitchcock, A. orcid.org/0000-0001-6572-434X, Hunter, C.N. orcid.org/0000-0003-2533-9783 and Canniffe, D.P. (2019) Progress and challenges in engineering cyanobacteria as chassis for light-driven biotechnology. *Microbial Biotechnology*. ISSN 1751-7915

<https://doi.org/10.1111/1751-7915.13526>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Special issue article

Progress and challenges in engineering cyanobacteria as chassis for light-driven biotechnology

Andrew Hitchcock^{1,*}  C. Neil Hunter¹  and Daniel P. Canniffe² 

¹Department of Molecular Biology and Biotechnology, University of Sheffield, Sheffield, UK.

²Institute of Integrative Biology, University of Liverpool, Liverpool, UK.

Summary

Cyanobacteria are prokaryotic phototrophs that, in addition to being excellent model organisms for studying photosynthesis, have tremendous potential for light-driven synthetic biology and biotechnology. These versatile and resilient microorganisms harness the energy of sunlight to oxidise water, generating chemical energy (ATP) and reductant (NADPH) that can be used to drive sustainable synthesis of high-value natural products in genetically modified strains. In this commentary article for the Synthetic Microbiology Caucus we discuss the great progress that has been made in engineering cyanobacterial hosts as microbial cell factories for solar-powered biosynthesis. We focus on some of the main areas where the synthetic biology and metabolic engineering tools in cyanobacteria are not as advanced as those in more widely used heterotrophic chassis, and go on to highlight key improvements that we feel are required to unlock the full power of cyanobacteria for future green biotechnology.

Introduction

Cyanobacteria are a diverse group of prokaryotic microorganisms that inhabit a range of environmental

Received 3 July, 2019; revised 29 November, 2019; accepted 1 December, 2019.

*For correspondence. E-mail a.hitchcock@sheffield.ac.uk; Tel. (+44) 114 222 4240; Fax (+44) 114 222 2711.

Microbial Biotechnology (2019) 0(0), 1–5
doi:10.1111/1751-7915.13526

Funding Information

The authors acknowledge award number BB/M000265/1 from the Biotechnology and Biological Sciences Research Council (BBSRC UK). AH further acknowledges support from a Royal Society University Research Fellowship (award number UR/R1191548).

niches. They perform oxygenic photosynthesis like plants and algae, utilizing solar energy and water to generate chemical energy (ATP) and reducing power (NADPH) to fix atmospheric carbon dioxide (CO₂) to make carbohydrates (Fig. 1). Oxygen is a 'waste' product of this process and ancient cyanobacteria oxygenated the Earth's atmosphere, permitting the evolution of complex, multicellular life forms. Cyanobacteria are excellent model organisms for fundamental photosynthesis research because of their relation to eukaryotic plastids, amenability to genetic engineering and relatively rapid generation time. They are also promising chassis for light-powered biotechnological applications, requiring only sunlight, water, CO₂ and trace inorganic minerals for growth, and tolerating variations in light, pH, salinity and temperature. Much progress has been made towards developing cyanobacteria as green microbial cell factories for the sustainable production of valuable natural products, chemicals and biofuels (see reviews by Oliver *et al.*, 2016; Khan *et al.*, 2019; Khan and Fu, 2019; Lin and Pakrasi, 2019); however, while the 'synthetic biology toolkit' for cyanobacterial metabolic engineering is advancing rapidly, it still lags behind those of model heterotrophic microorganisms (e.g. *Escherichia coli*) and yeast (*Saccharomyces cerevisiae*). Here, we consider the current situation with genetic engineering of cyanobacteria and highlight some key limitations that must be overcome in order to realize the full biotechnological potential of these versatile microorganisms.

Modifying cyanobacterial genomes is simple but slow

For an organism to be a suitable chassis for metabolic engineering an annotated genome sequence should be available, and it must be amenable to genetic manipulation to allow efficient and rapid generation of knockout and 'knock-in' strains. Many genetically tractable cyanobacteria meet these criteria (e.g. *Synechocystis* sp. PCC 6803, *Synechococcus* sp. PCC 7002, *Synechococcus elongatus* PCC 7942 and *Anabaena* sp. PCC 7120). Fast-growing (e.g. *S. elongatus* UTEX 2973) and 'non-model' species are also promising candidates for industrial applications (Yu *et al.*, 2015; Gale *et al.*, 2019; Mukherjee *et al.*, 2019).




Fig. 1. Engineering cyanobacterial chassis for light-driven product synthesis.

A. Schematic overview of a cyanobacterial 'chassis'. The photosynthetic light reactions generate ATP and NADPH using solar energy and water. The ATP and NADPH are consumed by the Calvin–Benson–Bassham (CBB) cycle to convert atmospheric CO₂ into carbohydrates. B. Genetic engineering approaches can be used to engineer cyanobacteria so that photosynthetically generated ATP and reductant can be used for light-driven product synthesis.

C. We suggest three overlapping and complementary areas for improvement that we believe will benefit the cyanobacterial synthetic biology community. (1) First, reducing polyploidy and/or improving CRISPR-based genome editing would increase the speed, ease and efficiency of generating genetically modified strains. (2) New replicative shuttle plasmids for extra-chromosomal expression of foreign genes and pathways are needed to enhance the 'design-build-test-learn' cycle that lies at the heart of the synthetic biology process. (3) An improved 'synthetic biology toolkit' that can be used in a combinatorial fashion to allow precise temporal and orthogonal gene expression/protein production to improve product yields and minimize instability of engineered strains.

Suicide constructs for making targeted genomic modifications by double homologous recombination can be transferred into cyanobacteria by natural transformation, conjugation or electroporation. A number of 'neutral' chromosomal sites are available for integrating heterologous DNA, avoiding polar effects and undesirable changes in cell metabolism and physiology (Ng *et al.*, 2015; Pinto *et al.*, 2015). Some cyanobacteria also contain endogenous plasmids that can be used for expression of foreign genes (Xu *et al.*, 2011).

Marked genomic modifications are common in cyanobacteria as they allow positive selection and maintenance of engineered strains; however, only a limited number of antibiotic markers are available, restricting the number of sequential genetic manipulations that can be made. Furthermore, the metabolic load of multiple resistances can negatively affect cell performance, and antibiotic use is economically and ecologically undesirable in

scaled-up 'real world' applications. Strategies utilizing alternative phosphorus (phosphite) and nitrogen (melamine) sources for selection circumvent the antibiotic issue (Polyviou *et al.*, 2015; Selão *et al.*, 2019), but the problem of marker recycling remains. Marker-less editing is therefore desirable and two-stage selection/counter-selection (Lea-Smith *et al.*, 2016) and site-specific recombinase (Tan *et al.*, 2013) systems can be used in cyanobacteria. However, because many cyanobacteria are relatively slow growing (compared with *E. coli*) and display polyploidy (multiple genome copies per cell), time-consuming segregation to generate homozygous strains in which every copy of the genome has been modified slows down genetic engineering. This is doubly laborious for generating marker-less strains, which require segregation after each transformation/recombination event. A better understanding of polyploidy may permit generation of organisms with only one genome copy

per cell, which would speed up further genome modification but may come with an undesirable cost to cell fitness. The recently announced CyanoSource barcoded mutant library for the model species *Synechocystis* sp. PCC 6803 (Gale *et al.*, 2019) will be a valuable resource of knockout strains for the community.

The potential of CRISPR genome editing in cyanobacteria

A better solution to the segregation time and multi-loci/marker rescue problems is the use of clustered regularly interspaced short palindromic repeats (CRISPR) genome editing, which has been employed successfully in several cyanobacterial species (see Behler *et al.*, 2018 for a detailed review). CRISPR genome editing can target multiple genes in parallel, is precise, marker-less and scar-free, and can edit every copy of the chromosome with just one round of cell viability-based selection. To date, the CRISPR-machinery is introduced into cyanobacteria on an exogenous replicative plasmid (see below) that must be cured from the modified strain, slowing down the overall process. There are also issues with toxicity and 'off-target' effects of the CRISPR nuclease and the efficiency of editing. These difficulties must be resolved for the true power of CRISPR genome editing in cyanobacteria to be unlocked, but it looks set to be an invaluable method for strain editing in the future.

The need for replicative plasmids for propagation of exogenous DNA

While genome editing is essential in some circumstances, genes can also be introduced to cyanobacteria on self-replicating plasmids, as is common with other synthetic biology chassis. Replicative plasmids overcome the time-constraint of strain segregation and the difficulty of integrating larger inserts into the genome, and gene expression is often higher from plasmids than from the bacterial chromosome (Chen *et al.*, 2016). As discussed above, CRISPR- and recombinase-based genome engineering also require expression of plasmid-borne genes (Tan *et al.*, 2013; Wendt *et al.*, 2016). However, the use of plasmids in cyanobacterial genetic engineering is somewhat limited; only pRSF1010 derivatives or fusions of native cyanobacterial and *E. coli* plasmids have been widely used in cyanobacteria due to stability issues with other platforms (see Xia *et al.*, 2019 for an overview of replicative plasmids in cyanobacteria). A better understanding of cyanobacterial plasmid replication and the development of exogenous self-replicating vectors based on alternatives to the RSF1010 backbone would be of great benefit to the cyanobacterial metabolic engineering community. Shuttle vectors that are

maintained in both *E. coli*/ yeast and the cyanobacterial host are particularly useful as they facilitate high throughput assembly of constructs using modular DNA assembly methods in the former prior to transferring into the latter, speeding up the 'design-build-test-learn' optimization cycle of synthetic biology.

Genetic instability necessitates regulation of heterologous gene expression/protein production

Achieving specific and tuneable regulation of introduced genes and pathways is another significant hurdle in cyanobacterial metabolic engineering, and it can avoid uncontrolled expression of heterologous genes, and the accompanying negative effects on the cell (Jones, 2014). As cyanobacteria display adaptive evolution in response to stress, tight control of gene expression can prevent selection of faster-growing loss of function suppressor mutations.

Control can be exerted at the transcriptional, translational and post-translational levels, but well-characterized regulatory elements used in other organisms do not tend to work as predicted when ported into cyanobacteria (Sengupta *et al.*, 2018). There has therefore been an enormous effort to develop and characterize genetic parts for cyanobacteria that can be used with standardized assembly methods such as the SyneBrick vectors (Kim *et al.*, 2017) and the CyanoGate modular cloning (MoClo) system (Vasudevan *et al.*, 2019). For example, promoters responding to isopropyl β -D-1-thiogalactopyranoside (Huang *et al.*, 2010), anhydrotetracycline (Huang and Lindblad, 2013), sugars (Kelly *et al.*, 2018), light (Abe *et al.*, 2014), metals (Englund *et al.*, 2016), temperature (Mermet-Bouvier and Chauvat, 1994) and metabolic/environmental signals (Immethun *et al.*, 2017) have been investigated, but in general their use is affected by poor dynamic range, leakiness, toxicity or photo-lability of the inducer, and/or incompatibility with cyanobacterial growth conditions (Camsund and Lindblad, 2014). Native and synthetic promoters of different strengths can be used when constitutive gene expression is permitted/desired (Markley *et al.*, 2015; Ruffing *et al.*, 2016), although there may be unpredictable diurnal regulation as expression is often only tested under constant illumination.

Regulation of cyanobacterial transcription can also be achieved with riboswitches (Nakahira *et al.*, 2013; Ma *et al.*, 2014) or CRISPR interference (CRISPRi), which allows simultaneous repression of multiple genes (multiplexing) (Gordon *et al.*, 2016; Yao *et al.*, 2016). Other synthetic RNA-based systems such as antisense repression and trans-activating sRNAs are also effective in cyanobacteria (Ueno *et al.*, 2018). Transcriptional terminator libraries to prevent readthrough to downstream genes have been developed (Lui and Pakrasi, 2018),

while at the post-transcriptional level, ribosome binding site libraries for varying the strength of translation initiation (Wang *et al.*, 2018; Lui and Pakrasi, 2018) and degradation tags to control protein levels (Huang *et al.*, 2010; Landry *et al.*, 2013), have been described in cyanobacteria. A big challenge for the future is to combine this ever-expanding range of genetic parts to achieve tight, dynamic, specific and orthogonal regulation of multiple components, allowing stable pathway maintenance and predictable performance under conditions relevant to scaled commercial applications.

Concluding remarks: a bright, green future for cyanobacterial biotechnology

Great progress has been made in engineering cyanobacterial chassis as microbial cell factories for renewable and light-driven synthesis of biochemicals and biofuels, but further improvements are now required to develop these processes into economically viable options for bioproduction. We suggest three main areas for improvement, summarized below and in Fig. 1. (i) Increase the speed and efficiency of marker-less genome modification, for example by optimization of CRISPR-based technologies. (ii) Development of new replicative shuttle plasmids for expression of heterologous genes and operons. (iii) Achieving tighter regulatory control of introduced pathways to increase long-term strain stability. By combining such advances in the constantly developing cyanobacterial synthetic biology toolkit with knowledge from 'omics' and genome-scale metabolic flux analyses/models (Hagerman and Hess, 2018; Hendry *et al.*, 2019), and new developments in the field of protein and pathway engineering, significant optimization of strain performance and product yield should be achievable. We are confident of a bright, green future for cyanobacterial biotechnology.

Acknowledgements

The authors acknowledge award number BB/M000265/1 from the Biotechnology and Biological Sciences Research Council (BBSRC UK). AH further acknowledges support from a Royal Society University Research Fellowship (award number UR/R011915/48).

Conflict of interest

None declared.

References

- Abe, K., Miyake, K., Nakamura, N., Kojima, K., Ferri, S., Ikebukuro, K., and Sode, K. (2014) Engineering of a green-light inducible gene expression system in *Synechocystis* sp. PCC6803. *Microb Biotechnol* **7**: 177–183.
- Behler, J., Vijay, D., Hess, W.R., and Akhtar, M.K. (2018) CRISPR-based technologies for metabolic engineering in cyanobacteria. *Trends Biotechnol* **36**: 996–1010.
- Camsund, D., and Lindblad, P. (2014) Engineered transcriptional systems for cyanobacterial biotechnology. *Front Bioeng Biotechnol* **2**: 40.
- Chen, Y., Taton, A., Go, M., London, R.E., Pieper, L.M., Golden, S.S., and Golden, J.W. (2016) Self-replicating shuttle vectors based on pANS, a small endogenous plasmid of the unicellular cyanobacterium *Synechococcus elongatus* PCC 7942. *Microbiology* **162**: 2029–2041.
- Englund, E., Liang, F., and Lindberg, P. (2016) Evaluation of promoters and ribosome binding sites for biotechnological applications in the unicellular cyanobacterium *Synechocystis* sp. PCC 6803. *Sci Rep* **6**: 36640.
- Gale, G.A.R., Schiavon Osorio, A.A., Mills, L.A., Wang, B., Lea-Smith, D.J., and McCormick, A.J. (2019) Emerging species and genome editing tools: future prospects in cyanobacterial synthetic biology. *Microorganisms* **7**: E409.
- Gordon, G.C., Korosh, T.C., Cameron, J.C., Markley, A.L., Begemann, M.B., and Pfeleger, B.F. (2016) CRISPR interference as a titratable, trans-acting regulatory tool for metabolic engineering in the cyanobacterium *Synechococcus* sp. strain PCC 7002. *Metab Eng* **38**: 170–179.
- Hagerman, M., and Hess, W.R. (2018) Systems and synthetic biology for the biotechnological application of cyanobacteria. *Curr Opin Biotechnol* **49**: 94–99.
- Hendry, J.I., Bandyopadhyay, A., Srinivasan, S., Pakrasi, H.B., and Maranas, C.D. (2019) Metabolic model guided strain design of cyanobacteria. *Curr Opin Biotechnol* **64**: 17–23.
- Huang, H.H., and Lindblad, P. (2013) Wide-dynamic-range promoters engineered for cyanobacteria. *J Biol Eng* **7**: 10.
- Huang, H.H., Camsund, D., Lindblad, P., and Heidorn, T. (2010) Design and characterization of molecular tools for a Synthetic Biology approach towards developing cyanobacterial biotechnology. *Nucleic Acids Res* **38**: 2577–2593.
- Immethun, C.M., DeLorenzo, D.M., Focht, C.M., Gupta, D., Johnson, C.B., and Moon, T.S. (2017) Physical, chemical, and metabolic state sensors expand the synthetic biology toolbox for *Synechocystis* sp. PCC 6803. *Biotechnol Bioeng* **114**: 1561–1569.
- Jones, P.R. (2014) Genetic instability in cyanobacteria - an elephant in the room? *Front Bioeng Biotechnol* **2**: 12.
- Kelly, C.L., Taylor, G.M., Hitchcock, A., Torres-Méndez, A., and Heap, J.T. (2018) A rhamnose-inducible system for precise and temporal control of gene expression in cyanobacteria. *ACS Synth Biol* **7**: 1056–1066.
- Khan, S., and Fu, P. (2019) Biotechnological perspectives on algae: a viable option for next generation biofuels. *Curr Opin Biotechnol* **62**: 146–152.
- Khan, A.Z., Bilal, M., Mehmood, S., Sharma, A., and Iqbal, H.M.N. (2019) State-of-the-art genetic modalities to engineer cyanobacteria for sustainable biosynthesis of biofuel and fine-chemicals to meet bio-economy challenges. *Life* **9**: E54.
- Kim, W.J., Lee, S.M., Um, Y., Sim, S.J., and Woo, H.M. (2017) Development of synebrick vectors as a synthetic biology platform for gene expression in *Synechococcus elongatus* PCC 7942. *Front Plant Sci* **8**: 293.

- Landry, B.P., Stöckel, J., and Pakrasi, H.B. (2013) Use of degradation tags to control protein levels in the Cyanobacterium *Synechocystis* sp. Strain PCC 6803. *Appl Environ Microbiol* **79**: 2833–2835.
- Lea-Smith, D.J., Vasudevan, R., and Howe, C.J. (2016) Generation of marked and markerless mutants in model cyanobacterial species. *J Vis Exp* **111**: 54001.
- Lin, P.C., and Pakrasi, H.B. (2019) Engineering cyanobacteria for production of terpenoids. *Planta* **249**: 145–154.
- Lui, D., and Pakrasi, H.B. (2018) Exploring native genetic elements as plug-in tools for synthetic biology in the cyanobacterium *Synechocystis* sp. PCC 6803. *Microb Cell Fact* **17**: 48.
- Ma, A.T., Schmidt, C.M., and Golden, J.W. (2014) Regulation of gene expression in diverse cyanobacterial species by using theophylline-responsive riboswitches. *Appl Environ Microbiol* **80**: 6704–6713.
- Markley, A.L., Begemann, M.B., Clarke, R.E., Gordon, G.C., and Pfleger, B.F. (2015) Synthetic biology toolbox for controlling gene expression in the cyanobacterium *Synechococcus* sp. strain PCC 7002. *ACS Synth Biol* **4**: 595–603.
- Mermet-Bouvier, P., and Chauvat, F. (1994) A conditional expression vector for the cyanobacteria *Synechocystis* sp. strains PCC6803 and PCC6714 or *Synechococcus* sp. strains PCC7942 and PCC6301. *Curr Microbiol* **28**: 145–148.
- Mukherjee, B., Madhu, S., and Wangikar, P.P. (2019) The role of systems biology in developing non-model cyanobacteria as hosts for chemical production. *Curr Opin Biotechnol* **64**: 62–69.
- Nakahira, Y., Ogawa, A., Asano, H., Oyama, T., and Tozawa, Y. (2013) Theophylline-dependent riboswitch as a novel genetic tool for strict regulation of protein expression in Cyanobacterium *Synechococcus elongatus* PCC 7942. *Plant Cell Physiol* **54**: 1724–1735.
- Ng, A.H., Berla, B.M., and Pakrasi, H.B. (2015) Fine-tuning of photoautotrophic protein production by combining promoters and neutral sites in the Cyanobacterium *Synechocystis* sp. strain PCC 6803. *Appl Environ Microbiol* **81**: 6857–6863.
- Oliver, N.J., Rabinovitch-Deere, C.A., Carroll, A.L., Nozzi, N.E., Case, A.E., and Atsumi, S. (2016) Cyanobacterial metabolic engineering for biofuel and chemical production. *Curr Opin Chem Biol* **35**: 43–50.
- Pinto, F., Pacheco, C.C., Oliveira, P., Montagud, A., Landels, A., Couto, N., *et al.* (2015) Improving a *Synechocystis*-based photoautotrophic chassis through systematic genome mapping and validation of neutral sites. *DNA Res* **22**: 425–437.
- Polyviou, D., Hitchcock, A., Baylay, A.J., Moore, C.M., and Bibby, T.S. (2015) Phosphate utilization by the globally important marine diazotroph *Trichodesmium*. *Environ Microbiol Rep* **7**: 824–830.
- Ruffing, A.M., Jensen, T.J., Strickland, L.M., and Santos-Merion, L.M. (2016) Genetic tools for advancement of *Synechococcus* sp. PCC 7002 as a cyanobacterial chassis. *Microb Cell Fact* **15**: 190.
- Selão, T.T., Włodarczyk, A., Nixon, P.J., and Norling, B. (2019) Growth and selection of the cyanobacterium *Synechococcus* sp. PCC 7002 using alternative nitrogen and phosphorus sources. *Metab Eng* **54**: 255–263.
- Sengupta, A., Pakrasi, H.B., and Wangikar, P.P. (2018) Recent advances in synthetic biology of cyanobacteria. *Appl Microbiol Biotechnol* **102**: 5457–5471.
- Tan, X., Liang, F., Cai, K., and Lu, X. (2013) Application of the FLP/FRT recombination system in cyanobacteria for construction of markerless mutants. *Appl Microbiol Biotechnol* **97**: 6373–6382.
- Ueno, K., Tsukakoshi, K., and Ikebukuro, K. (2018) Riboregulator elements as tools to engineer gene expression in cyanobacteria. *Appl Microbiol Biotechnol* **102**: 7717–7723.
- Vasudevan, R., Gale, G.A.R., Schiavon, A.A., Puzorjov, A., Malin, J., Gillespie, M.D., *et al.* (2019) CyanoGate: a modular cloning suite for engineering cyanobacteria based on the plant mocl syntax. *Plant Physiol* **180**: 39–55.
- Wang, B., Eckert, C., Maness, P.C., and Yu, J. (2018) A genetic toolbox for modulating the expression of heterologous genes in the Cyanobacterium *Synechocystis* sp. PCC 6803. *ACS Synth Biol* **7**: 276–286.
- Wendt, K.E., Ungerer, J., Cobb, R.E., Zhao, H., and Pakrasi, H.B. (2016) CRISPR/Cas9 mediated targeted mutagenesis of the fast growing cyanobacterium *Synechococcus elongatus* UTEX 2973. *Microb Cell Fact* **15**: 115.
- Xia, P.F., Ling, H., Foo, J.L., and Chang, M.W. (2019) Synthetic biology toolkits for metabolic engineering of cyanobacteria. *Biotechnol J* **14**: e1800496.
- Xu, Y., Alvey, R.M., Byrne, P.O., Graham, J.E., Shen, G., and Bryant, D.A. (2011) Expression of genes in cyanobacteria: adaptation of endogenous plasmids as platforms for high-level gene expression in *Synechococcus* sp. PCC 7002. *Methods Mol Biol* **684**: 273–293.
- Yao, L., Cengic, I., Anfelt, J., and Hudson, E.P. (2016) Multiple gene repression in cyanobacteria using CRISPRi. *ACS Synth Biol* **5**: 207–212.
- Yu, J., Liberton, M., Cliften, P.F., Head, R.D., Jacobs, J.M., Smith, R.D., *et al.* (2015) *Synechococcus elongatus* UTEX 2973, a fast growing cyanobacterial chassis for biosynthesis using light and CO₂. *Sci Rep* **5**: 8132.