

Canterbury Research and Theses Environment

# Canterbury Christ Church University's repository of research outputs

http://create.canterbury.ac.uk

Please cite this publication as follows:

Lockwood, P., Pittock, L., Lockwood, C., Jeffery, C. and Piper, K. (2015) Intraorbital foreign body detection and localisation by radiographers: a preliminary JAFROC observer performance study. Radiography, 2015. pp. 1-5. ISSN 1078-8174.

Link to official URL (if available):

http://dx.doi.org/10.1016/j.radi.2015.06.005

This version is made available in accordance with publishers' policies. All material made available by CReaTE is protected by intellectual property law, including copyright law. Any use made of the contents should comply with the relevant law.

Contact: create.library@canterbury.ac.uk





# Canterbury Christ Church University's repository of research outputs

http://create.canterbury.ac.uk

Please cite this publication as follows:

Lockwood, Paul, Pittock, Lisa, Lockwood, Claire, Jeffery, Christopher and Piper, Keith (2015) Intraorbital foreign body detection and localisation by radiographers: a preliminary JAFROC observer performance study. Radiography, 2015. pp. 1-5. ISSN 1078-8174

Link to official URL (if available):

http://www.sciencedirect.com/science/article/pii/S1078817415000802

This version is made available in accordance with publishers' policies. All material made available by CReaTE is protected by intellectual property law, including copyright law. Any use made of the contents should comply with the relevant law.

Contact: create.library@canterbury.ac.uk



# Intraorbital foreign body detection and localisation by radiographers: a preliminary JAFROC observer performance study

P. Lockwood<sup>a</sup>, L. Pittock<sup>b</sup>, C. Lockwood<sup>c</sup>, C. Jeffery<sup>d</sup>, K. Piper<sup>e</sup>

Allied Health Department, Canterbury Christ Church University, Medway Campus, Rowan Williams Building, 30 Pembroke Court, Chatham Maritime, ME4 4UF, UK

#### Key words

Radiographer, Intra Orbital Foreign Body, Jackknife free-response receiver operating characteristic, Observer Performance, Magnetic Resonance Imaging.

Word count 2656

# Abstract

Introduction - The purpose of this study was to run a preliminary investigation to establish if a short course of learning would increase radiographers' performance in intraorbital foreign body (IOFB) detection and localisation on pre-magnetic resonance imaging (MRI) orbital computed radiographs (CR).

Method - A multi-reader multi-case (MRMC) human observer study was performed. Fifteen radiographers from 5 hospitals participated. Each radiographer reviewed a pre- and post-training image bank and was instructed to identify the presence or absence of IOFBs, indicating the lesion location on each case whilst scoring the detection using a confidence index on a 5-point scale, for 30 orbital radiographs. The results were analysed using a Jackknife free-response receiver operating characteristic (JAFROC2 equal weighted) methodology.

Results -The performance of the radiographers demonstrated a statistically significant difference after a short period of training in the detection of IOFBs on orbital radiographs (F (1,14)= 12.99, df = 14.0, p = 0.0029). The JAFROC2 analysis averaged figure of merit (FOM) for the radiographers was 0.818 (95% CI 0.769, 0.867) pre-training and 0.920 (95% CI 0.891, 0.950) post-training.

Conclusion - These results suggest that with a short programme of learning in image interpretation for IOFBs in orbital radiographs, radiographers should be able to achieve a high level of accuracy in the identification and localisation of IOFBs prior to MRI examination.

# Highlights

- Using a MRMC study of 15 radiographers from 5 hospitals
- We evaluated radiographers' performance in IOFB detection and localisation
- The examination incorporated a pre- and post- image bank of IOFB examples
- The exam reports were analysed using JAFROC2 methodology
- Radiographers after a short course of learning achieved high levels of accuracy

#### Introduction

Since the first reported cases of serious eye injuries from metallic intraorbital foreign bodies (IOFBs) by Kelly et al<sup>1</sup>. The risk from injuries such as rupture, haemorrhage or blindness have formed the basis for stringent screening protocols for pre- magnetic resonance imaging (MRI) examinations. There has since been numerous reported cases<sup>2, 3, 4, 5</sup> that have helped develop health and safety protocols to screen patients with suspected metallic IOFBs before they enter the controlled area of the MRI department. Current policies applied in National Health Service (NHS) practice conform to recent guidelines by the Society and College of Radiographers (SCoR) and the British Association of Magnetic Resonance Radiographers (BAMRR)<sup>6</sup>, the Medicines and Healthcare products Regulatory Agency(MHRA)<sup>7</sup>, the European Union Physical Agents Directive (EUPAD)<sup>8</sup> and the Ionising Radiation (Medical Exposure) Regulations (IR(ME)R Schedule 1<sup>9,10</sup>. All of which aim to ensure no individual (patient, staff, or visitor) enters an MRI controlled area with a metallic IOFB. This process involves verbal screening, written questionnaires, review of previous imaging (if available) and in uncertain cases pre-MRI orbital radiographes.

MRI radiographers have the potential to role extend and develop their clinical practice in a service improvement capacity to streamline their diagnostic imaging pathways. Under IR(ME)R<sup>10</sup> legislation MRI practitioners have the capacity through local level agreement to develop a scope of entitlement to identify themselves through departmental protocols, referral criteria, job descriptions and appropriate continuing profession development (CPD) training and audit activities to be recognised as the referrer for pre-MRI orbital radiographs. The relevant skills and knowledge to extend radiographers' scope of practice for the rationale to refer patients for orbital radiographs and the interpretation of imaging for the purposes of excluding the presence of metallic IOFBs are contained as part of the protocols in the employers' procedures under IR(ME)R<sup>10</sup>. The demonstrable knowledge would need to show an awareness of the circumstances leading to a penetrating injury resulting in a retained IOFB, with regular audit of the suitability and impact of referrals<sup>11</sup>. Whilst documenting a record of a sufficient amount of test film viewings from a bank of suitable radiographs, and a period of clinical image interpretations to demonstrate a level of competence. Specifically IR(ME)R<sup>10</sup> advises that each medical exposure has a clinical evaluation of the image, which is recorded for audit purposes.

Radiographers have previously evidenced the ability to interpret a wide range of current radiographic investigations<sup>12, 13, 14, 15, 16, 17, 18</sup>. Role extension into referral and interpretation of pre-MRI orbital radiographic imaging would benefit the workflow practice of busy MRI departments and waiting time initiatives, as well increasing radiographer responsibilities in a modern radiology service.

#### **Aims and Objectives**

The purpose of this study was to run a preliminary investigation to establish if a short one day intensive course of lectures, question and answer sessions and interactive case reviews would increase radiographers' performance in IOFB detection and localisation on pre-MRI orbital computed radiographs. The training reviewed orbital and facial bone anatomy, physiology, pathology, image interpretation and search strategies. Whilst reviewing published cases of IOFB injuries, and discussing the current legal perspectives and NHS polices and guidelines to prevent IOFB injuries.

#### Methodology

The study followed a multiple reader multiple case (MRMC) retrospective preliminary study of 15 radiographers from 5 hospitals using a Jackknife free-response receiver operating characteristic

(JAFROC2) methodology<sup>19</sup>. The observers reviewed a pre- and post-training image bank and were instructed to identify the presence or absence of IOFBs, recording the lesion location (LL) on each case and scoring the detection using a confidence index on a 5-point scale for 30 orbital radiographs. The study was approved by an ethics and governance panel, and all observers gave informed written consent.

The sample of participants recruited conformed to the criteria of band 5 and 6 radiographers working within MRI departments of southern United Kingdom (UK) hospitals, all had MRI experience ranging from 3 to 9 years, and some had previous postgraduate training in MRI, but no radiographers had any plain film reporting experience. The sample size for this preliminary study followed Obuchowski<sup>21, 22</sup> tables for receiver operating characteristic (ROC) studies of observers sampled in medical imaging studies, and for external validity purposes<sup>23</sup> would not be representative of being proportional to the general population.

The images were obtained by permission of local NHS trusts from an anonymised retrospective digital teaching library (DTL) used by the university for research and teaching that conformed to section 33 of the UK Data Protection Act<sup>24</sup>, the Cosson and Willis<sup>25</sup> guidance from the National Information Governance Board for Health and Social Care, and the General Medical Council<sup>26</sup>.

Orbital radiographs used included letter box collimated under tilted occipito-mental (OM) with orbitomeatal baseline raised 10 degrees less than for a standard OM (to provide a circular appearance of the orbits, unlike the oval OM standard view). Some of the images used included secondary supplementary views of eyes up or down or lateral side to side.

Images rejected from the study bank included those with the petrous ridge superimpositioned upon the lower orbital margin, poor positioning or rotation of the facial bones, over tilted projections, and the presence of artefacts on the imaging plate.

A sampling bias of pathology was reduced as near as possibly by using an appropriately wide range, amount and size of IOFBs as was feasible that conformed to textbook examples and conspicuity. A suitable range of subtle IOFBs as discriminatory examples ensured a fair process of image interpretation<sup>22, 27</sup>. Where there were multiple IOFBs on an image each lesion was given equal weighting factor for the JAFROC2 analysis, as each lesion was deemed to have an equal importance and risk factor for potential injury. JAFROC2 methodology compares the performance of readers interpreting the same bank of cases at two different intervals (pre-testing was at the start of the study day, post-testing at the end of the study day). The methodology was intended to investigate questions such as whether or not teaching image interpretation improves diagnostic performance in the detection and localisation of IOFBs on orbital radiographs.

Each case had a triple reader retrospective approach of interpretation by three independent and blinded reviewers<sup>27</sup> (a consultant radiologist and two reporting radiographers) to determine concordance for the reference standard, and reduce potential bias (internal validity) of inter-rater disagreement of same case reviews <sup>28, 29, 30</sup>.

The images were evaluated on Liquid Crystal Display (LCD) image monitors with a resolution of 1280 x 1024, calibrated to the Digital Imaging and Communications in Medicine (DICOM) part 14 Greyscale Standard Display Function (GSDF) with VeriLUM software<sup>32</sup>. Quality checks were performed on the monitors prior to each test with a standard diagnostic imaging Society of Motion Picture and Television Engineers (SMPTE) reference pattern for spatial uniformity of luminance and temporal luminance stability complying with Royal College of Radiologists (RCR) guidelines<sup>31</sup>. The cases were viewed using commercially available software (K-PACS<sup>33</sup>) intended for displaying DICOM

images. This allowed the observers to alter the computed radiograph (CR) window width and window level, pan and zoom the image, and measure region of interests (ROI) with the ability to display multiple images from a case for comparison views. An observing environment was chosen specifically for this study that reduced the possibility of disturbance of concentration, by controlling the lighting, noise and interruptions<sup>27</sup>.

The JAFROC2 (equal weighted) methodology allowed the readers to make as many or as few responses per images, and the paradigm allowed multiple IOFBs per case if required. When the observer reviewed each case they labelled the LL ROI (or quadrants in JAFROC2 terminology) and rated the degree of suspicion using a confidence index score, in this study 1-5. A score of 1 equated to high confidence that the ROI in question did not have a lesion (normal), with grading up to a score of 5 inferred the reader was highly confident the ROI in question did have a lesion (similar to a Likert scale).

The reader's scores were marked as either LL i.e. the mark of a lesion was within an acceptance region to the real lesion, or a non-lesion location (NLL). Acceptance regions of lesions are varied in peer review literature, this study has followed the Chakraboty<sup>34</sup> recommendation of a maximum diameter of a lesion or 3 mm. Acceptance radiuses are a controversial topic in observer performance studies and Chakraborty<sup>34, 35</sup> considers the question of what is the maximum inaccuracy with no clinical impact that would be acceptable in a study scenario.

The JAFROC2 (equal weighted)<sup>19</sup> analysis produces a figure of merit (FOM) metric which determines the measure of the observer's detectability, using the number of LLs compared to the total known number of lesions (lesion localisation fraction (LLF), and NLLs relative to the total number of cases in the image bank (non-lesion localisation fraction (NLF). Specifically JAFROC2 (equal weighted)<sup>20</sup> methodology was used as NLL on non-diseased cases are counted towards false positives factor (FPF), which takes into account satisfaction of search errors (the amount of NLL marks on diseased cases).

With all statistical reasoning of data there is a need to estimate the variability of the results, in this study the JAFROC2<sup>19</sup> variance component uses analysis of variance (ANOVA) with the results displayed as 95% confidence intervals (95% CI) which JAFROC2 uses to qualify each statistic as the trapezoidal area under the curve (AUC). This is a non-parametric bootstrapping method to obtain the FOM<sup>35</sup> to bench mark the performance of the observational task. Although the reader performance in decision making is subjective, the FOM is an objective measure of performance which rewards correct answers and penalizes incorrect answers using a scalar unidirectional measure of 0-1 (0 as an incorrect decision and 1 as a correct measure against the reference standard of case answers). The FOM is equal to a Wilcoxon test statistic<sup>36</sup> for two samples (pre- and post-training image bank results) and JAFROC2 determines if the difference is sufficiently bigger evaluated against the statistical variability of chance<sup>35</sup>. JAFROC2 also calculates an *F* statistic,  $n_N$  (numerator) and  $n_D$  (denominator) degrees of freedom (*df*), at the desired probability level<sup>38</sup> used in small sample size studies, and the *p* value determines the quantified degree of unlikelihood<sup>37</sup>.

# Results

The performance of the radiographers demonstrated a statistically significant difference after a short period of training. The detection of IOFBs in CR images demonstrated an increased performance after training (F (1,14) = 12.99, df = 14.0, p = 0.0029, Table 1, 2,and 3). The JAFROC2 analysis average FOM for the radiographers was 0.818 (95% CI 0.769,0.867) pre-training and 0.920 (95% CI 0.891,0.950) post-training.

The overall observer's performance improved with training, showing 11 radiographers increased the amount of correct LL IOFBs recorded from the pre-training bank to the post-training bank, and 4 radiographers scored the same amount of LLs in the pre- and post-training image banks. The variation of the observed distribution of NLL scores between the pre- and post-training banks ranged from an increase of NLLs for 6 radiographers, no difference for 7 radiographers and 2 radiographers reduced their NLL scores with training (Table 1).

# Discussion

The results are consistent to previous studies showing the ability of radiographers to improve their performance of interpretation after a short course of formal training<sup>39, 40</sup>. The radiographers in this study produced a mean FOM of 0.920 (standard error 0.013, *df* 14, 95% CI 0.891,0.950). A literature search to find similar studies of radiographers' abilities in the accuracy of IOFB identification and localisation found no comparable figures. Historic studies have shown a range of predicated observer sensitivity scores of plain film radiographs to identify IOFB's. Bryden et al<sup>41</sup> have shown that observers able to identify IOFBs using radiographic (not CR) images to a 69% sensitivity on x-rays.

The appearance of an IOFB radiographically is a debatable issue. The question of IOFBs to be detectable on CR must take into consideration the size and shape of the metallic IOFB. It has been shown in previous studies<sup>11</sup> that metallic fragments as small as 0.5x0.2mm are not visible on radiographs to observers, and additionally glass and wooden IOFBs have poorer sensitivity on radiographs<sup>44, 45</sup>.

Other additional examinations have been shown as alternative non-ionising examinations of the orbits prior to MRI scanning, ultrasound has proven to be exceptionally sensitive (93%) to IOFBs in screening<sup>41, 44</sup>, although orbital cavity fat has shown to produce reflective artefacts that impact the evaluation of retained intra and extra ocular particles<sup>41</sup>. Additionally there are multiple studies that have researched computed tomography (CT) as highly accurate in determining IOFBs<sup>43, 46</sup>, with sensitivity for metallic IOFB detection at  $100\%^{43, 47}$  (65% sensitivity <0.06mm<sup>3</sup>, 100% sensitivity >0.06mm<sup>3, 48</sup>). Although the increased radiation dose to the radiosensitive orbital contents must be considered before imaging.

It could be argued that there is substantial controversy regarding the necessity of screening patients prior to MR procedures. Various studies have questioned the adequacy of the screening process including the questions asked within pre-MRI questionnaires<sup>11</sup>, and have implied that some clinical sites do not use orbital radiographs when screening patients. The implications for this variation in practice are that some centres may deem the subjection of ionising radiation unjustified, whereas others might see this as an unsafe practice according to their safety protocols and guidance from other countries<sup>49</sup>.

Indeed a study by Williamson et al<sup>50</sup> discussed the risk of injury from an MRI scan to patients with a history of a metallic IOFB, and suggested a figure of one in several thousand will actually incur an injury, and thus routine screening of at risk patients would seem unnecessary. A balanced argument would state that although it has been shown that a questionnaire may not provide an adequate patient history to reduce the risk of harm<sup>11, 51</sup>, and that there are many factors that impact on the level of risk from ferromagnetic IOFB movements (location, size, geometry, metallic properties, MRI field strength, and period of time the IOFB has been in situ<sup>50</sup>), a proven risk has been demonstrated in peer reviewed papers <sup>1, 2, 3, 4, 5</sup>. The majority of NHS facilities have established strict screening criteria for possible intraocular and IOFBs in line with current guideline recommendations<sup>6</sup>.

Radiographic imaging appears to be the acceptable technique<sup>6</sup> for identifying or excluding an intraocular or peri-orbital metallic foreign body that may present a potential hazard to the patient entering the MRI environment.

The value of short training courses for radiographers appears appropriate in this setting to compliment continuous professional development requirements of radiographer's registration to the Health and Care Professions Council and the SCoR. Furthermore it supports research and initial modelling of future role extension and development of radiographer's ability in service improvement within MRI departments. The efficacy of running short courses has the ability to produce change and has been shown to be beneficial in improving radiographic interpretation<sup>52, 53</sup>.

The results of this research, although significant in findings, constitute a small sample and are unrepresentative of the population of radiographers. Further research is recommended and using the preliminary study's JAFROC2<sup>19</sup> data analysis allowed an Obuchowski-Rockette Dorfman-Berbaum-Metz (OR-DBM) method to perform sample size estimation for a future phase 2 study. The OR-DBM method is considered the standard to reasonably calculate how many readers and how many cases are needed to get a 0.05 Alpha, with a desired power of .80 (1-beta). The variance component covariance estimates for sample size ANOVA estimated a future study of 30 observers and 45 cases would give a prediction of observer performance at 0.8.

## Conclusions

These results suggest that with a short programme of learning in image interpretation for IOFBs in orbital radiographs, radiographers were able to achieve a high level of accuracy (0.920 FOM (95% CI 0.891,0.950) in the identification and localisation of IOFBs prior to MRI examination. A significance of p = 0.0029 was demonstrated between the pre and post-training bank results signifying a potential for inexperienced radiographers to improve image interpretation with training. Further study in this field of training radiographers in the interpretation of pre-MRI orbital radiograph should be completed.

#### References

1. Kelly W M, Paglen P G, Pearson J A, San Diego A G, Soloman M A. Ferromagnetism of intraocular foreign body causes unilateral blindness after MR study. American Journal of Neuroradiology 1986; Mar-Apr;7(2):243-5.

2. Kulshrestha M, Misson G. Magnetic resonance imaging and the dangers of orbital foreign bodies. The British Journal of Ophthalmology 1995;79.12:1149.

3. Wheatcroft S, Benjamin L. Magnetic resonance imaging and the dangers of orbital foreign bodies. British Journal of Ophthalmology 1996;80(12):1116.

4. Ta C N, Bowman RW. Hyphaema caused by a metallic foreign body during magnetic resonance imaging. American Journal of Ophthalmology 2000; 129:533-534.

5. Vote B J, Simpson A J. X-ray turns a blind eye to ferrous metal. Clinical and Experimental Ophthalmology 2001;29:262–264.

6. Society and College of Radiographers and the British Association of Magnetic Resonance Radiographers. Safety in Magnetic Resonance Imaging. London: The Society and College of Radiographers, 2013.

7. Medicines and Healthcare products Regulatory Agency (MHRA) Device Bulletin Safety Guidelines for Magnetic Resonance Imaging Equipment in Clinical Use DB2007 (03) 2007. http://www.mhra.gov.uk/Publications/Safetyguidance/DeviceBulletins/CON2033018

8. DIRECTIVE 2004/40/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 29 April 2004 on the minimum health and safety requirements regarding the exposure of workers to the risks arising from physical agents (electromagnetic fields) (18th individual Directive within the meaning of Article 16(1) of Directive 89/391/EEC) Official Journal of the European Union L184/1-L184/7, 24th May 2004.

9. Society and College of Radiographers. IR(ME)R 2000 and IR(ME) Amendment Regulations 2006 & 2011. London: The Society and College of Radiographers; 2012.

10. Great Britain. Ionising Radiation (Medical Exposure) Regulations 2000 (IRMER). London: Department of Health, 2012.

11. Bailey W, Robinson L. Screening for intra-orbital metallic foreign bodies prior to MRI: Review of the evidence. Radiography 2007;13.1:72-80.

12. Price R, Le Masurier S. Longitudinal changes in extended roles in radiography: a new perspective. Radiography 2007;13(1):18–29.

13. Robinson P, Culpan G, Wiggins M. Interpretation of selected accident and emergency radiographic examinations by radiographers: a review of 11000 cases. British Journal of Radiology 1999;72:546–551.

14. Piper K, Paterson A, Godfrey R. Accuracy of radiographers' reports in the interpretation of radiographic examinations of the skeletal system: a review of 6796 cases. Radiography 2005;11 (1):27–34.

15. Smith S, Reeves P. The extension of the role of the diagnostic radiographer in the UK National Health Service over the period 1995–2009. European Journal of Radiography 2009;1.4:108-114.

16. Elliott L. Radiographer reporting in the nuclear medicine department: a learning curve?." Radiography 2003;9.3:247-251.

17. Clarke R, Allen D, Arnold P, Snaith B. Implementing radiographic CT head reporting: The experiences of students and managers. Radiography 2014;20.2:117-120.

18. Piper K, Buscall K. MRI reporting by radiographers: The construction of an objective structured examination. Radiography 2008;14.2:78-89.

19. Chakraborty D. JAFROC v4 software (Version 4.2.1); 2014. Available at: <a href="http://www.devchakraborty.com/">http://www.devchakraborty.com/</a>

20. Chakraborty D. New developments in observer performance methodology in medical imaging. Seminars in Nuclear Medicine 2011; 41(6): 401-418.

21. Obuchowski N A. Sample size tables for receiver operating characteristic studies. American Journal of Roentgenology 2000; 175(3):603-608.

22. Obuchowski N A. How many observers are needed in clinical studies of medical imaging?. American Journal of Roentgenology 2004;182(4):867-869.

23. Robinson P J, Wilson D, Coral A, Murphy A, Verow P. Variation between experienced observers in the interpretation of accident and emergency radiographs. The British Journal of Radiology 1999; 72(856):323-330.

24. Great Britain. Data Protection Act. Chapter 29. London: HMSO; 1998.

25. Cosson P, Willis N. Digital teaching library (DTL) development for radiography education. Radiography 2011;18(2):112-116.

26. General Medical Council. Supplementary Guidance: making and using visual and audio recordings of patients. London. General Medical Council; 2013.

27. Brealey S. Measuring the effects of image interpretation: an evaluative framework. Clinical Radiology 2001;56(5):341-347.

28. Berlin L. Accuracy of diagnostic procedures: has it improved over the past five decades?. American Journal of Roentgenology 2007;188.5:1173-1178.

29. Berlin L. Radiologic errors, past, present and future. Diagnosis 2014;1.1:79-84.

30. Lee C S, Nagy P G, Weaver S J, Newman-Toker D E. Cognitive and system factors contributing to diagnostic errors in radiology. American Journal of Roentgenology 2013; 201(3):611-617.

31. IMAGE Smith, Inc. (VeriLUM (Version 5.2.1); 2008. Available at: http://verilum.software.informer.com/

32. The Royal College of Radiologists. Picture archiving and communication systems (PACS) and guidelines on diagnostic display devices, second edition. London: The Royal College of Radiologists; 2012.

33. IMAGE Information Systems. K-PACS (Version 1.6.0); 2008. Available at: http://www.k-pacs.net/10555.html

34. Chakraborty D, Yoon H J, Mello-Thoms C. Spatial localization accuracy of radiologists in freeresponse studies: inferring perceptual FROC curves from mark-rating data. Academic radiology2007;14.1:4-18.

35. Chakraborty D P. New developments in observer performance methodology in medical imaging." In Seminars in nuclear medicine 2011;vol. 41, no. 6; 401-418.

36. Obuchowski N A, Lieber M L, Powell K A. Data analysis for detection and localization of multiple abnormalities with application to mammography. Academic radiology 2000;7.7:516-525.

37. Penedo M, Souto M, Tahoces P G, Carreira J M, Villalón J, Porto G, Seoane C, Vidal J J, Berbaum K S, Chakraborty D, Fajardo L L. Free response receiver operating characteristic evaluation of lossy JPEG2000 and object based set partitioning in hierarchical trees compression of digitized mammograms. Radiology 2005;237:450–457.

38. Glanz S A. Primer of Biostatistics, 3rd Edition, New York: McGraw-Hill; 1992, p 44.

39. Hardy M, Culpan G. Accident and emergency radiography: A comparison of radiographer commenting and 'red dotting'. Radiography 2007;13.1:65-71.

40. Piper K, Paterson A. Initial image interpretation of appendicular skeletal radiographs: a comparison between nurses and radiographers. Radiography 2009;15.1:40-48.

41. Bryden F M, Pyott A A, Bailey M, McGhee C N. Real time ultrasound in the assessment of intraocular FB. Eye 1990;4:727.

42. Saeed A, Cassidy L, Malone D E, Beatty S. Plain X-ray and computed tomography of the orbit in cases and suspected cases of intraocular foreign body. Eye 2008;22.11:1373-1377.

43. Bray L C, Griffiths P G. The value of plain radiography in suspected intraocular foreign body. Eye 1991;5(6):751-4.

44. Lagalla R, Manfre L, Caronia A, Bencivinni F, Duranti C, Ponte F. Plain film, CT and MRI sensibilityin the evaluation of intraorbital foreign bodies in an in vitro model of the orbit and in pig eyes. European radiology 2000;10.8:1338-1341.

45. Fielding J A. The assessment of ocular injury by ultrasound. Clinical Radiology 2004;59(4):301.

46. Pinto, Antonio, et al. "Role of computed tomography in the assessment of intraorbital foreign bodies." Seminars in Ultrasound, CT and MRI. Vol. 33. No. 5. WB Saunders, 2012.

47. Guneric U, Maden A, Kaynak S, Pimar T. Magnetic resonance imaging and computed tomography in the detection and localisation of intraocular foreign bodies. Documenta Ophthalmologica 1992; 81:369–78.

48. Mester V, Kuhn F. Intraocular foreign bodies. Ophthalmology Clinics of North America 2002;15.2 :235-242.

49. Kanal E. Barkovich A J, Bell C, Borgstede J P, Bradley Jr W G, Froelich J W, Gimbel J , Gosbee J W, Kuhni-Kaminski r T, Larson P A, Lester J W, Nyenhuis J, Schaefer d J, Sebek R N, Weinreb J, Wiloff B L, Woods T O, Lucey L, Hernandez D. ACR Guidance Document on MR Safe Practices: Journal of Magnetic Resonance Imaging 2013;37:501–530.

50. Williamson MR, Espinosa MC, Boutin RD, Orrison WW Jr, Hart BL, Kelsey CA. Metallic foreign bodies in the orbits of patients undergoing MR imaging: prevalence and value of radiography and CT before MR, American Journal of Roentgenology 1994; Apr;162(4):981-3.

51. Elmquist C, Shellock F G, Stoller D. Screening adolescents for metallic foreign bodies before MR procedures. Journal of Magnetic Resonance Imaging 1995;5(6):784–785.

52. Mackay S J. The impact of a short course of study on the performance of radiographers when highlighting fractures on trauma radiographs: "The Red Dot System". The British Journal of Radiology 2006; 79 (942): 468-472.

53. McConnell J R, Webster AJ. Improving radiographer highlighting of trauma films in the accident and emergency department with a short course of study-an evaluation. The British journal of radiology 2000; 73(870): 608-612.

#### Acknowledgements

The authors would like to thank all the radiographers that participated in the study.

#### Funding

There were no financial conflicts of interest.

## **Conflicts of Interest / Affiliations**

Authors a and b are senior lecturers on the MSc Clinical Reporting and BSc Radiography programmes at Canterbury Christ Church University. Author c is a reporting radiographer at Kings College Foundation NHS Trust, London. Author d is a senior lecturer on the MSc Medical Imaging and BSc Radiography programmes at Canterbury Christ Church University. Author e is the programme director for the MSc Clinical Reporting programme at Canterbury Christ Church University.



**Table 1.** JAFROC FOM for pre-training (dark blue bars) and post-training (light blue bars) for each observer.

**Table 2.** Observer averaged pre-training (dark blue bar) and post-training (light blue bar) with 95% confidence intervals.





Analysis	Statistical	Radiographer Mean	Radiographer Mean
	significance	Pre-training	Post-training
		FOM (95% CI)	FOM (95% CI)
JAFROC2	F (1,14) = 12.99	0.818	0.920
Equal Weighting	<i>P</i> < 0.0029	(0.769,0.867)	(0.891,0.950)