



# CREaTE

Canterbury Research and Theses Environment

Canterbury Christ Church University's repository of research outputs

<http://create.canterbury.ac.uk>

Please cite this publication as follows:

Woznitza, N. (2014) Commentary: radiographer reporting. *Journal of Medical Radiation Sciences*, 61 (2). pp. 66-68.

Link to official URL (if available):

<http://dx.doi.org/10.1002/jmrs.51>

This version is made available in accordance with publishers' policies. All material made available by CReaTE is protected by intellectual property law, including copyright law. Any use made of the contents should comply with the relevant law.

Contact: [create.library@canterbury.ac.uk](mailto:create.library@canterbury.ac.uk)



## COMMENTARY

**Radiographer reporting**Nick Woznitza, BSc, PgDip, PgCert<sup>1,2</sup><sup>1</sup>Radiology Department, Homerton University Hospital, London, United Kingdom<sup>2</sup>Allied Health Department, Canterbury Christ Church University, Kent, United Kingdom*Journal of Medical Radiation Sciences* 61 (2014) 66–68

doi: 10.1002/jmrs.51

Radiographers worldwide are integral to the diagnostic pathway and are optimally placed to provide expert comment on radiographs. By nature, the radiographer is the first health care professional to view each diagnostic image, which has been acquired by a focus on the patient. Radiographers are in a unique position to communicate their professional observations directly with the treating clinician in a timely manner and thereby have a significant influence on patient care. Currently, advanced practitioner roles, which incorporate radiographer reporting, are limited to the United Kingdom (UK).<sup>1</sup> The changing nature of health care worldwide has seen several countries including Canada,<sup>2</sup> Australia,<sup>3</sup> Norway<sup>4</sup> and Denmark<sup>5</sup> develop models of advanced radiographer practice which includes definitive clinical reporting.

Swinburne first raised the possibility of trained radiographers expanding their role to incorporate preliminary image interpretation,<sup>6</sup> although the pioneering work of Berman et al.<sup>7</sup> is seen as the origin of radiographer preliminary image interpretation. The proposed system of work required radiographers to highlight abnormal trauma skeletal radiographs by placing a 'red dot' on the image, which indicated to the casualty officer the possible presence of significant pathology. This method has been shown to reduce diagnostic errors in the Emergency Department.<sup>7</sup> In 2006, the Society and College of Radiographers, while recognising the benefits of the 'red dot' system of preliminary radiograph interpretation, also identified several weaknesses, which includes the ambiguity of an absence of a 'red dot'.<sup>8</sup> Preliminary clinical evaluation (PCE) builds on abnormality detection by radiographers, as PCE requires a concise written statement which localises and describes the pertinent findings.<sup>1</sup> The provision of a written interpretation directs the treating clinician to the area(s) of concern and removes many of the ambiguity of the 'red dot' system, such as cases with multiple abnormalities, incorrect interpretation of abnormalities on an abnormal image and

communication of uncertainty in the radiographer decision.<sup>1,9</sup> A survey undertaken of UK radiology departments in 2008 found a significant majority provide a system of radiographer abnormality detection for skeletal trauma imaging; most still use the 'red dot', while some provide a PCE or a hybrid system.<sup>10</sup>

In parallel to this expansion of radiographer practice, the role of the advanced practitioner has been developed in the United Kingdom which incorporates the provision of definitive clinical reports by appropriately trained radiographers.<sup>1</sup> The performance of radiographers in interpreting skeletal radiographs at the end of an accredited postgraduate training program was promising with high levels of sensitivity (91.6–96.7%) and specificity (92.1–94.0%) reported.<sup>11</sup> A large multi-centre clinical evaluation, consisting of 7179 cases conducted across four sites in the United Kingdom, demonstrated very high levels of accuracy, sensitivity and specificity, 99.1%, 97.6% and 99.3%, respectively, for skeletal trauma reports produced by trained reporting radiographers.<sup>12</sup> A subsequent meta-analysis conducted by Brealey et al.<sup>13</sup> examined the performance of radiographer reporting for 28,900 plain imaging examinations and provided the definitive evidence that trained reporting radiographers can provide clinical reports on skeletal radiographs at a level comparable to consultant radiologists.

Trained radiographers now provide definitive clinical reports on skeletal radiographs throughout the U.K., with 59 (41%) of 143 departments providing this service in 2012.<sup>14</sup> In response to evolving service needs, radiographer reporting has expanded in scope beyond skeletal trauma. There is a growing body of evidence that supports trained radiographers who can provide definitive clinical reports for chest radiographs,<sup>15,16</sup> magnetic resonance imaging (MRI) lumbar spine and knee examinations,<sup>17</sup> and mammograms.<sup>18</sup> Multidisciplinary team working, which incorporates radiographer reporting, has been highlighted in recent a

joint publication by the Royal College of Radiologists and Society and College of Radiographers as one method to deliver an effective, efficient and patient focused radiology service.<sup>19</sup>

The interesting article by Neep et al.<sup>20</sup> explores the confidence of a cohort of Australian radiographers in the provision of both radiographer abnormality detection (red dot) and PCE (radiographer comment). Based on the results of a cross-sectional survey, they note that radiographers report high confidence when participating in abnormality detection systems, but lower confidence and perceived accuracy is suggested for PCEs. The authors of the study hypothesise that this may be due to the prior educational support and experience of the radiographers.

The results of Neep et al.<sup>20</sup> are similar to the findings of Coleman and Piper<sup>21</sup> who found that radiographers reported lower confidence when interpreting a bank of trauma radiographs when compared to emergency nurse practitioners (ENP) and junior medical staff. Although the radiographers reported lower confidence, the radiographers had the highest average score (28.5/40) for the image bank of all professional groups, significantly higher than the ENPs (21/40;  $P < 0.01$ ) and junior casualty medical staff (21.5/40;  $P = 0.02$ ).<sup>21</sup> The radiographers were also the only group whose confidence correlated with accuracy ( $r = 0.51$ ;  $P = 0.02$ ).<sup>18</sup> Lower radiographer confidence in the provision of PCEs was identified in the analysis conducted at a multi-site NHS Trust, with education and training highlighted as potential barriers to improved confidence.<sup>22</sup> Uptake of PCE by radiology departments in the United Kingdom remains patchy, with authors speculating whether definitive clinical reporting by radiographers has helped or hindered this progression.<sup>10</sup>

Education and training, which incorporates tutorials and feedback, improves radiographer image interpretation performance.<sup>23</sup> This is true for both definitive clinical reporting<sup>13</sup> and preliminary radiographer interpretations.<sup>9</sup> The magnitude of improvement has been shown to be greater for radiographer abnormality detection when compared to PCE,<sup>9,24</sup> but some of this difference may be due to the inherent ambiguity in the red dot/abnormality detection system.

The College of Radiographers, in conjunction with the U.K. regulatory body (Health and Care Professions Council), have mandated that image interpretation training is included as part of pre-registration.<sup>1</sup> Online resources, which include adult and paediatric skeletal and adult chest radiograph interpretation, have been developed at a national level to support undergraduate students and practitioners in the provision of radiographer PCE.<sup>25</sup> A recent report has highlighted

examples of Australian trained radiographers who have taken up advanced practitioner/reporting radiographer roles in the United Kingdom with appropriate education and support.<sup>26</sup> This reflects well on the potential to develop radiographer reporting in Australia.

The promising work of Neep et al.<sup>20</sup> produced some valuable findings and, together with evidence available from the United Kingdom,<sup>11,13,21</sup> suggest potentially that Australian radiographers may be able to offer a positive contribution to the trauma diagnostic pathway. Accuracy of radiographer image interpretation and confidence in participating in PCE and definitive reporting will improve with appropriate education and training.

## Conflicts of Interest

The author declares no conflict of interest.

## References

1. College of Radiographers. Preliminary Clinical Evaluation and Clinical Reporting by Radiographers: Policy and Practice Guidance. College of Radiographers, London, U.K., 2013.
2. Canadian Association of Medical Radiation Technologists. Advanced Practice in Medical Radiation Technology A Canadian Framework. Canadian Association of Medical Radiation Technologists, Canada, 2014.
3. Advanced Practice Advisory Panel. Proposed Pathway to Advanced Practice. Australian Institute of Radiography, Australia, 2014.
4. Hofmann B, Vikestad KG. Accuracy of upper abdominal ultrasound examinations by sonographers in Norway. *Radiography* 2013; **19**: 186–9.
5. Buskov L, Abild A, Christensen A, Holm O, Hansen C, Christensen H. Radiographers and trainee radiologists reporting accident radiographs: a comparative plain film-reading performance study. *Clin Radiol* 2013; **68**: 55–8.
6. Swinburne K. Pattern recognition for radiographers. *Lancet* 1971; **297**: 589–90.
7. Berman L, de Lacey G, Twomey E, Twomey B, Welch T, Eban R. Reducing errors in the accident department: A simple method using radiographers. *Br Med J* 1985; **290**: 421–2.
8. College of Radiographers. Medical Image Interpretation & Clinical Reporting by Non-Radiologists: The Role of the Radiographer. College of Radiographers, London, U.K., 2006.
9. Hardy M, Culpan G. Accident and emergency radiography: A comparison of radiographer commenting and 'red dotting'. *Radiography* 2007; **13**: 65–71.
10. Snaith B, Hardy M. Radiographer abnormality detection schemes in the trauma environment. An assessment of current practice. *Radiography* 2008; **14**: 277–81.

11. Piper K, Paterson A, Godfrey R. Accuracy of radiographers' reports in the interpretation of radiographic examinations of the skeletal system: A review of 6796 cases. *Radiography* 2005; **11**: 27–34.
12. Piper K, Paterson A, Ryan C. The Implementation of a Radiographic Reporting Service for Trauma Examinations of the Skeletal System in 4 NHS Trusts. NHS Executive South Thames, UK, 1999.
13. Brealey S, Scally A, Hahn S, Thomas N, Godfrey C, Coomasamy A. Accuracy of radiographer plain radiograph reporting in clinical practice: A meta-analysis. *Clin Radiol* 2005; **60**: 232–41.
14. Society & College of Radiographers. Scope of radiographic practice survey 2012. Society and College of Radiographers, London, U.K., 2012.
15. Piper K, Cox S, Paterson A, Thomas A, Thomas N, Jeyagopal N, et al. Chest reporting by radiographers: Findings of an accredited postgraduate programme. *Radiography* 2014; **20**: 94–9.
16. Woznitza N, Piper K, Burke S, Patel K, Amin S, Grayson K, et al. Adult chest radiograph reporting by radiographers: preliminary data from an in-house audit programme. *Radiography* 2014; doi: 10.1016/j.radi.2014.03.002.
17. Brealey S, Piper K, King D, Bland M, Caddick J, Campbell P, et al. Observer agreement in the reporting of knee and lumbar spine magnetic resonance (MR) imaging examinations: Selectively trained MR radiographers and consultant radiologists compared with an index radiologist. *Eur J Radiol* 2013; **82**: e597–605.
18. Wivell G, Denton ER, Eve CB, Inglis JC, Harvey I. Can radiographers read screening mammograms? *Clin Radiol* 2003; **58**: 63–7.
19. Royal College of Radiologists, Society & College of Radiographers. Society & College of Radiographers. Royal College of Radiologists and the Society and College of Radiographers, London, U.K., 2012.
20. Neep M, Steffens T, Owen R, McPhail S. A survey of radiographers' confidence and self-perceived accuracy in frontline image interpretation and their continuing educational preferences. *J Med Radiat Sci* 2014; **61**: 69–77.
21. Coleman L, Piper K. Radiographic interpretation of the appendicular skeleton: A comparison between casualty officers, nurse practitioners and radiographers. *Radiography* 2009; **15**: 196–202.
22. Lancaster A, Hardy M. An investigation into the opportunities and barriers to participation in a radiographer comment scheme, in a multi-centre NHS trust. *Radiography* 2012; **18**: 105–8.
23. Mackay S. The impact of a short course of study on the performance of radiographers when highlighting fractures on trauma radiographs: "The Red Dot System". *Br J Radiol* 2006; **79**: 468–72.
24. Smith TN, Traise P, Cook A. The influence of a continuing education program on the image interpretation accuracy of rural radiographers. *Rural Remote Health* 2009; **9**: 1145.
25. e-Learning for Healthcare. Interpretation of Radiological Images. UK: NHS Health Education England; [Updated 2014; cited 2014 March 16]. Available from: <http://www.e-lfh.org.uk/programmes/image-interpretation/>.
26. Ricote L. Advanced Practice within Medical Imaging – Towards an Australian Model. The Winston Churchill Memorial Trust of Australia, Adelaide, Australia, 2009.

Nick Woznitza  
Reporting Radiographer, Radiology Department,  
Homerton University Hospital, Homerton Row,  
London E9 6SR, United Kingdom.  
Tel: +44 208 510 7105; Fax: +44 208 510 7107;  
E-mail: [nicholas.woznitza@nhs.net](mailto:nicholas.woznitza@nhs.net)