

Anthocyanin pigments in strawberry

Fátima Lopes da Silva¹, María Teresa Escribano-Bailón, José Joaquín Pérez Alonso, Julián C. Rivas-Gonzalo, Celestino Santos-Buelga^{*}

Facultad de Farmacia, Laboratorio de Nutrición y Bromatología, Universidad de Salamanca, Campus Miguel de Unamuno s/n, E-37007 Salamanca, Spain

Received 3 November 2004; received in revised form 29 September 2005; accepted 29 September 2005

Abstract

The anthocyanin composition was analysed in strawberry fruits from five different cultivars (cv. *Eris*, *Oso Grande*, *Carisma*, *Tudnew* and *Camarosa*). Twenty-five defined anthocyanin pigments were detected, most of them containing Pelargonidin (Pg) as aglycone; some cyanidin (Cy) derivatives were also found. Glucose and rutinose were the usual substituting sugars, although arabinose and rhamnose were also tentatively identified; some minor anthocyanins showed acylation with aliphatic acids. A relevant aspect was the detection of anthocyanin-derived pigments, namely 5-carboxypyranopelargonidin-3-glucoside and four condensed pigments containing C–C linked anthocyanin (Pg) and flavanol (catechin and afzelechin) residues. Total anthocyanin content ranged between 200 and 600 mg kg⁻¹, with Pg 3-gluc constituting 77–90% of the anthocyanins in the strawberry extracts followed by Pg 3-rut (6–11%) and Cy 3-gluc (3–10%). A notable variability was found among the anthocyanin concentrations in samples of a same variety and harvest, indicating a strongly influence of the degree of maturity, edaphic-climatic factors and post-harvest storage.

© 2005 Swiss Society of Food Science and Technology. Published by Elsevier Ltd. All rights reserved.

Keywords: Anthocyanins; Strawberry; Pelargonidin; Cyanidin; Anthocyanin-derived pigments

1. Introduction

Strawberry fruits (*Fragaria × ananassa* Duch.) have been shown to possess high in vitro antioxidant activity that has been positively correlated with the content of polyphenolic compounds and, specifically, anthocyanins, the type of polyphenols quantitatively most important in strawberry (Heinonen, Meyer, & Frankel, 1998; Wang & Jiao, 2000; Wang & Lin, 2000). The anthocyanin composition in strawberry has been the object of various studies, but is still not fully characterized regarding minor pigments. Strawberry anthocyanins derive from pelargonidin (Pg) and cyanidin (Cy) aglycones (Fig. 1a) (Mazza & Miniati, 1993). The major anthocyanin in the fruits is Pg 3-glucoside (Pg 3-gluc), as firstly identified by Robinson and Robinson (1931). In smaller proportions the presence of Cy 3-glucoside (Cy 3-gluc) seems also constant in all varieties

(Bridle & Garcia-Viguera, 1997; Hong & Wrolstad, 1990a; Lukton, Chichester, & MacKiney, 1955) and Pg 3-rutinoside (Pg 3-rut) is also commonly found (Bakker, Bridle, & Bellworthy, 1994; Co & Markakis, 1968; Hong & Wrolstad, 1990b). Furthermore, Pg 3-arabinoside (Fiorini, 1995; Goiffon, Mouly, & Gaydou, 1999) and Cy 3-rutinoside (Bridle & Garcia-Viguera, 1997) have been cited in some strawberry cultivars, as well as various acylated anthocyanins. In particular, Pg 3-(6-malonylglucoside) was unequivocally identified by Tamura, Takada and Yoshida (1995) and indicated as one of the main pigments in several Japanese cultivars, comprising 5–30% of total anthocyanin content (Tamura et al., 1995; Yoshida, Koyama, & Tamura, 2002). Other acylated anthocyanins also reported in strawberry are Pg 3-acetylglucoside (Hong & Wrolstad, 1990b) and Pg succinylglucoside (Bakker et al., 1994).

In a previous study by our group (Lopes-da-Silva, de Pascual-Teresa, Rivas-Gonzalo, & Santos-Buelga, 2002), the anthocyanin composition in strawberries of cv. *Camarosa* was analysed using electrospray ionization mass spectrometry ESI-MS coupled to HPLC. In addition to the major anthocyanins (i.e. Pg 3-gluc, Pg 3-rut and Cy 3-gluc)

^{*}Corresponding author. Fax: +34923 294 515.

E-mail address: csb@usal.es (C. Santos-Buelga).

¹Current address: Escola Superior Agrária de Bragança, Campus de Santa Apolónia, P-5301-855 Bragança, Portugal.




Fig. 1. Structures of pigments found in strawberry. (a) Anthocyanin aglycones; (b) 5-carboxypyranopelargonidin 3-glucoside (Andersen et al., 2004); and (c) anthocyanin-flavanol condensed pigments (Fossen et al., 2004).

12 minor anthocyanins were detected although identity could be only assigned to five of them as Pg 3-acetylglucoside, Cy 3-rutinoside, Pg 3-malylglucoside, Pg diglucoside, and Cy 3-malonylglucosyl-5-glucoside, the three latter not being described previously in strawberry.

Quite recently, small amounts of some anthocyanin-related pigments have also been detected and identified in strawberries, including 5-carboxypyranopelargonidin 3-glucoside (Fig. 1b) (Andersen, Fossen, Torskangerpoll, Fossen, & Hauge, 2004) and four purple anthocyanin-flavanol complexes consisting of pelargonidin 3-glucoside C–C linked to (epi)catechin and (epi)afzelechin moieties (Fig. 1c) (Fossen, Rayyan, & Andersen, 2004). That was the first evidence of the occurrence in a natural plant source of this type of condensed pigments, whose formation was associated to reactions taking place during maturation and ageing of red wines (Jurd, 1969; Salas et al., 2004; Somers, 1971; Vivar-Quintana, Santos-Buelga, Francia-Aricha, & Rivas-Gonzalo, 1999). Further evidence about the presence

of anthocyanin–flavanol condensed pigments in strawberry fruits and other plants has also been recently contributed by our group (González-Paramás et al., 2005).

The aim of the present work is to update the knowledge about strawberry anthocyanins, for which the anthocyanin composition, qualitative and quantitative, has been analysed in strawberry fruits from five different cultivars, using HPLC coupled to diode array and MS detection.

2. Materials and methods

2.1. Samples

Strawberries (*Fragaria × ananassa* Duch.) from five selected cultivars (cv. *Camarosa*, *Carisma*, *Eris*, *Oso Grande* and *Tudnew*) grown at an experimental station at Instituto de la Grasa-CSIC in Seville (Spain) and picked at commercial maturity were collected in years 2001 and

2002. After harvest, fruits were washed in water and frozen and stored at -35°C until analysis.

2.2. Sample preparation

Frozen strawberries (40–50 g) were homogenized in MeOH containing 0.1% HCl, kept overnight (~ 14 h) at $3\text{--}5^{\circ}\text{C}$ and later filtered through a Büchner funnel under vacuum. The solid residue was exhaustively washed with methanol and the filtrates obtained were centrifuged (4000g, 15 min, 2°C) and the solid residue further submitted to the same process the number of times necessary to complete extraction of the colour. Hydro-alcoholic phases were combined, water was added, the supernatant was concentrated under vacuum in a rotary evaporator at $<30^{\circ}\text{C}$ to total evaporation of the methanol; the aqueous extract obtained was washed with *n*-hexane to remove liposoluble substances. An aliquot (2 ml) of the aqueous phase was carefully deposited onto a C-18 SepPak[®] Vac 3cc cartridge (Waters); sugars and more polar substances were removed by passing 15 ml of ultrapure water and anthocyanin pigments further eluted with 5 ml of MeOH:0.1% TFA (95:5). The methanolic extract was concentrated under vacuum in a rotary evaporator at $<30^{\circ}\text{C}$ after adding some water. The aqueous extract was collected, its volume completed to 2 ml with ultrapure water and filtered through a $0.45\text{-}\mu\text{m}$ membrane filter for HPLC analysis. For each strawberry variety and year of harvest, three independent extracts were prepared that were purified and analysed separately.

2.3. HPLC-DAS-MS analyses

Analyses were performed in a Hewlett-Packard 1100 series liquid chromatograph. Separation was achieved on a $5\text{-}\mu\text{m}$ AQUA[®] C18 $150\text{ mm} \times 4.6\text{ mm}$ column (Phenomenex[®], Torrance, CA) thermostatted at 35°C . Solvents used were: (A) 0.1% trifluoroacetic acid in water, and (B) HPLC-grade acetonitrile, establishing the following gradient: isocratic 10%B for 5 min, 10–15%B over 15 min, isocratic 15%B for 5 min, 15–18%B over 5 min, and 18–35%B over 20 min, using a flow rate of 0.5 ml min^{-1} . Double on-line detection was carried out in a diode array spectrophotometer (DAS), using 520 nm as the preferred wavelength, and in a mass spectrometer (MS) connected to the HPLC system via the DAS cell outlet.

The mass spectrometer was a Finnigan LCQ (San Jose, CA) equipped with an ESI source and an ion trap mass analyser, which were controlled by the LCQ Xcalibur software. Nitrogen was used as both auxiliary and sheath gas at flow rates of 6 and 1.21 min^{-1} , respectively. The capillary voltage was 4 V and the capillary temperature 195°C . Spectra were recorded in positive ion mode between m/z 150 and 1500. The MS detector was programmed to perform a series of three consecutive scans: a full scan, a zoom scan of the most abundant ion in the

first scan and an MS–MS scan of the most abundant ion, using a normalized collision energy of 45%.

2.4. Quantification

The three major anthocyanins in strawberry were quantified from the areas of their chromatographic peaks recorded at 520 nm by comparison with calibration curves obtained with external standards of Cy 3-gluc (for cyanidin-based anthocyanins) and of Pg 3-gluc (for pelargonidin-based anthocyanins). Strawberry extracts were analysed in triplicate.

3. Results and discussion

3.1. Pigment identification

In the different strawberry varieties analysed 25 anthocyanin pigments were detected for which suitable information concerning their UV-vis or mass spectral characteristics could be obtained. Fig. 2 shows the HPLC anthocyanin profiles in the samples of the five strawberry cultivars analysed (i.e. cv. *Tudnew*, *Carisma*, *Camarosa*, *Eris* and *Oso Grande*). Peak data obtained in the HPLC-DAS-MS analyses (retention time in the HPLC system, λ_{max} in the visible region, molecular ion and main fragments observed in MS²) are summarized in Table 1, together with the strawberry varieties in which each peak was detected. In addition to the compounds indicated in that table, other very minor pigments were also detected although no good absorption or mass spectra could be obtained to allow speculation about their identity.

Pelargonidin shows a characteristic UV-vis spectrum shape with λ_{max} in the visible region at lower wavelengths (about 500 nm) than other common anthocyanins and an additional maximum about 430 nm (Fig. 3). Based on it the peaks corresponding to Pg-derived anthocyanins could be easily assigned in the chromatograms. It is necessary to take into account that during gradient HPLC run a progressive bathochromic shift in the visible λ_{max} of the anthocyanin peaks is produced with the increase in the percentage of acetonitrile in the mobile phase (Hebrero, Santos-Buelga, & Rivas-Gonzalo, 1988). Thus, λ_{max} of the peaks of the Pg-based anthocyanins vary from 500 nm (peak 6) to 508 nm (peak 23). The presence of Pg as anthocyanidin in those peaks was further confirmed by their mass spectra, which showed an MS² signal at m/z [M]⁺ 271. In addition, five peaks (5, 8, 9, 18 and 24) were identified as Cy derivatives based on the presence of a signal at m/z [M]⁺ 287 in their MS² spectra. Mass spectral characteristics also allowed to assign five other peaks to anthocyanin-derived pigments as discussed below.

Major peaks in the HPLC chromatograms in all samples corresponded to Pg 3-gluc (peak 10), Pg 3-rut (peak 12) and Cy 3-gluc (peak 5). Besides them, compounds 1, 2, 3, 19 and 21 were also found in all the samples analysed. Peak 21 would correspond to Pg 3-acetylglucoside, as previously




Fig. 2. Chromatograms (zoom) recorded at 520 nm showing the anthocyanin profiles of the strawberry samples. (a) *cv. Tudnew*; (b) *cv. Carisma*, (c) *cv. Camarosa*, (d) *cv. Eris*, (e) *cv. Oso Grande*. Full chromatogram is shown in the small window. See Table 1 for peak identification.

suggested (Lopes-da-Silva et al., 2002). Its presence in strawberry had also been indicated by Hong and Wrolstad (1990b). Peak 19 was relatively important in the samples of *cv. Tudnew* (Fig. 2a). Its UV-vis (λ_{\max} at 504 nm) and mass

spectra (molecular ion at m/z 519 releasing a unique MS^2 fragment at m/z 271 corresponding to Pg) allowed possible identity as Pg 3-malonylglucoside, previously identified in strawberry by Tamura et al. (1995). The equivalent Cy

Table 1
Retention time (Rt), wavelengths of maximum absorption in the visible region (λ_{\max}), mass spectral data, tentative identification of anthocyanin pigments and strawberry varieties in which they were detected

Peak	Rt (min)	λ_{\max} (nm)	Molecular ion [M^+] (m/z)	MS ² (m/z)	Tentative identification ^a	Strawberry variety ^b
1	12.3	515	721	559, 407, 313, 271	Catechin-(4→8)-Pg 3-glucoside	Ca, Cr, Er, Os, Tu
2	14.0	517	721	559, 407, 313, 271	Epicatechin-(4→8)-Pg 3-glucoside	Ca, Cr, Er, Os, Tu
3	17.7	515	705	543, 407, 313, 271	(Epi)afzelechin-(4→8)-Pg 3-glucoside	Ca, Cr, Er, Os, Tu
4	19.3	518	851	543, 407, 313, 271	(Epi)afzelechin-(4→8)-Pg 3-rutinoside	Ca, Cr, Tu
5	20.3	515	449	287	Cy 3-glucoside	Ca, Cr, Er, Os, Tu
6	21.5	500	433	271	Pg-3-galactoside	Ca, Tu
7	21.5	500	595	433, 271	Pg 3,5-diglucoside	Ca, Cr, Er, Os
8	21.8	515	595	449, 287	Cy 3-rutinoside	Ca, Cr, Er, Os
9	23.0	524	697	535, 449, 287	Cy 3-malonylglucosyl-5-glucoside	Ca
10	23.8	502	433	271	Pg 3-glucoside	Ca, Cr, Er, Os, Tu
11	24.8	492	501	339	5-carboxypyranopelargonidin-3-glucoside	Cr, Os, Tu
12	25.5	503	579	433, 271	Pg 3-rutinoside	Ca, Cr, Er, Os, Tu
13	29.3	503	549	271	Pg 3-malylglucoside	Cr
14	31.6	504	422	331	Unknown	Ca, Er, Tu
15	32.6	503	607	271	Pg-dissaccharide (hexose + pentose) acylated with acetic acid	Ca, Cr, Er
16	32.9	503	607	271	Pg-dissaccharide (hexose + pentose) acylated with acetic acid	Ca
17	33.3	503	403	271	Pg 3-arabinoside	Cr, Tu
18	33.5	n.a.	535	287	Cy 3-malonylglucoside	Ca
19	35.5	504	519	271	Pg 3-malonylglucoside	Ca, Cr, Er, Os, Tu
20	38.4	n.a.	563	271	Pg dirhamnoside?	Ca, Cr
21	39.9	504	475	271	Pg 3-acetylglucoside	Ca, Cr, Er, Os, Tu
22	40.6	504	533	271	Pg 3-succinylglucoside? Pg 3-methylmalonylglucoside?	Os, Tu
23	45.3	508	503	271	Pg 3-succinylarabinoside? Pg malonylrhamnoside? (Pg 3-methylmalonylarabinoside?)	Ca, Cr, Os, Tu
24	46.2	n.a.	549	287	Cy 3-succinylglucoside? Cy malylrhamnoside? (Cy 3-methylmalonylglucoside?)	Ca, Tu
25	47.1	n.a.	517	271	Pg 3-diacetylglucoside? (Pg succinylrhamnoside?) (Pg 3-methylmalonylrhamnoside?)	Ca, Cr, Tu

^aPg, pelargonidin; Cy, cyanidin; n.a., not available.

^bCa, *Camarosa*; Cr, *Carisma*; Er, *Eris*; Os, *Oso Grande*; Tu, *Tudnew*.




Fig. 3. UV-vis spectra of pelargonidin 3-glucoside (—), cyanidin 3-glucoside (---) and peak 11 (5-carboxypyranopelargonidin 3-glucoside) (· · ·) recorded with the HPLC diode array detector.

3-malonylglucoside (peak 18) was also detected in our samples, as a very small peak in the chromatograms of cv. *Camarosa*. It was assigned according to its mass (molecular ion at m/z 535, MS² fragment at m/z 287) and HPLC retention characteristics, eluting immediately before Pg 3-malonylglucoside. Further confirmation of the identity of compounds 18 and 19 was provided for comparison of their chromatographic and absorption and mass spectral characteristics with those of Cy 3-(6''-malonylglucoside) and Pg 3-(6''-malonylglucoside) previously identified in our laboratory in purple corn (de Pascual-Teresa, Santos-Buelga, & Rivas-Gonzalo, 2002) and available in our anthocyanin library. In our knowledge, Cy 3-malonylglucoside had not been previously reported in strawberry.

Compounds 1–4 were assigned to condensed pigments containing C–C linked anthocyanin (Pg) and flavanol (catechin or afzelechin) residues (Fig. 1c), as discussed elsewhere (González-Paramás et al., 2005). Pigments with the same structural characteristics as compounds 1–3 were

isolated from strawberry and fully characterised using NMR by Fossen et al. (2004). Compound 4 (i.e. (epi)afzelechin-(4→8)-Pg 3-rut) was not reported by those authors, but firstly described by our group (González-Paramás et al., 2005).

Anthocyanins 7, 8, 9, 13, 15 and 16 have been previously detected and tentatively identified in strawberries from cv. *Camarosa* by our group (Lopes-da-Silva et al., 2002). They correspond to: Pg-3,5-digluc (peak 7), Cy 3-rutinoside (8), Cy 3-malonylglucosyl-5-glucoside (9), Pg 3-malylglucoside (13) and two Pg biosides acylated with acetic acid (15 and 16).

Peak 17 was only relevant in the samples of cv. *Carisma*. The molecular ion at m/z 403, releasing an MS² fragment at m/z 271 (Pg), is consistent with a Pg pentoside. Arabinose and xylose are the most common pentose substituents in anthocyanins (Mazza & Miniati, 1993). No actual identification can be made of the sugar residue from the MS spectrum, but Pg 3-arabinoside was reported by some authors (Fiorini, 1995; Goiffon et al., 1999) as the third major anthocyanin in some strawberry varieties. For that reason and since no other pentoside was detected in our samples, peak 17 was tentatively assigned to Pg 3-arabinoside.

Bakker et al. (1994) detected in some strawberry cultivars two pigments whose masses, obtained by FAB-MS, were coherent with the succinyl derivatives of Pg 3-gluc and Cy 3-gluc. Molecular ions that may correspond to these two anthocyanins were also detected in our samples for peaks 22 (m/z at 533) and 24 (m/z at 549), respectively. However, their identity causes some doubt since Cy succinylglucoside should be expected to elute before the corresponding Pg succinylglucoside. A theoretical possibility that would also match these molecular ions is that one or both of them are products derived from anthocyanins acylated with malonic acid by esterification of the free carboxyl unit of the malonyl residue with the methanol used as a solvent for the extraction. Similar reaction was previously demonstrated in our lab to occur in anthocyanins of purple corn (de Pascual-Teresa et al., 2002), where the formation of ethyl derivatives of the malonylglucosides of cyanidin, peonidin and pelargonidin was observed following esterification with ethanol. Even if in that study the process was induced by heating (60 °C), methylation should not be discarded in our strawberry samples. In support of that possibility is the fact that peak 22 is particularly noticeable in the extracts of cv. *Tudnew*, where its potential precursor, Pg 3-malonylglucoside (peak 19), is in relatively higher amounts than in the rest of strawberry varieties analysed.

Similarly, peak 24 might be the corresponding methyl derivative of Cy 3-malonylglucoside, although in this case the reduced levels of the precursor anthocyanin (i.e. peak 18) would not support it. Another structure that also matches the molecular ion of peak 24 is that of a Cy malylrhamnoside. Nevertheless, such identity is much more uncertain taking into account that the corresponding non-

acylated Cy rhamnoside, that would have also been expected to occur in the samples, was not detected.

The molecular ion of peak 23 (m/z at 503) would match either a Pg succinylpentoside or a Pg malonylrhamnoside. The facts that peak 23 appeared in the same samples as peak 17 (assigned to Pg 3-arabinoside) and that Pg rhamnoside was not detected in any sample, seem to support the first identity (i.e. Pg succinylpentoside).

Different possible structures can also be suggested for peak 25, whose molecular ion at m/z 517 would match either a diacetyl derivative of Pg 3-gluc, a Pg succinylrhamnoside or a Pg methylmalonylrhamnoside. All of them would be in agreement with its late elution. Newly, the fact that no Pg rhamnoside was detected might support the first identity. Curiously, peak 20 showed a molecular ion (m/z at 563) that coincides with that of a Pg aglycone bearing two rhamnose substituents, identity that would also agree with its lower polarity when compared to Pg 3-gluc and Pg 3-rut. As for peaks 23 and 25, the non-detection of a Pg monorhamnoside would raise some doubt about the dirhamnosyl nature of peak 20.

The possibility that any of the peaks 20, 23 or 25 contain a *p*-coumaroyl acyl residue instead of a rhamnose was discarded. No shoulder at 310–330 nm was observed in the absorption spectra of any of the anthocyanins detected in the samples analysed, thus indicating that no hydroxycinnamic acids but only aliphatic acids are involved in acylation of strawberry anthocyanins.

Finally, three other minor pigments (peaks 6, 11 and 14) were also detected in some samples. Mass characteristics of peak 6 correspond to a Pg hexoside (molecular ion at m/z 433, MS² fragment at m/z 271). Taking into account that galactose is the only relevant hexose found in anthocyanins besides glucose (Mazza & Miniati, 1993) and the retention characteristics of the peak, eluting before Pg 3-glucoside, it was tentatively associated to Pg 3-galactoside (galactosides elute before the equivalent glucosides; *Escribano-Bailón, Santos-Buelga, Alonso, & Salinas, 2002*). Peak 11 showed a molecular ion at m/z 501 that released an MS² fragment at m/z 339 (–162 amu, loss of a glucose moiety) and a characteristic UV-vis spectrum (Fig. 3) with λ_{\max} in the visible region at 492 nm, hypsochromically shifted with regard to that of Pg, and an additional maximum at 335 nm. These mass and spectral features allowed assignment as 5-carboxypyranopelargonidin 3-glucoside (Fig. 1b), a pigment recently identified in strawberry by *Andersen et al. (2004)*. Such a structure is similar to that of the pyruvic acid adducts of anthocyanins formed in red wines during winemaking and ageing (*Mateus, de Pascual-Teresa, Rivas-Gonzalo, Santos-Buelga, & de Freitas, 2002*).

Mass spectrum of peak 14 (Fig. 4) revealed a molecular ion at m/z 422 that released a unique MS² fragment at m/z 331, a mass that might correspond to the anthocyanidin malvidin (Mv). However, the small difference of mass between the molecular ion and the MS² fragment (92 amu) does not allow identification of the peak as a Mv glycoside;




Fig. 4. Mass (molecular ion and MS²) and UV-vis spectra of peak 14 recorded with the HPLC diode array detector.




Fig. 5. Ion chromatogram extracted for m/z 422 corresponding to a strawberry extract from cv. *Tudnew*.

on the other hand, the even molecular ion (m/z 422) may suggest that the lost fragment contains N or S. Furthermore, another compound with the same mass characteristics (molecular ion at m/z 422 and MS² fragment at m/z 331) was found in the same strawberry samples when a search was made in the mass chromatograms (Fig. 5), pointing out that two isomeric structures of the compound exist. The peak of the second compound is overlapped by that of Pg 3-rutinoside and for that reason it was not detected in the DAD chromatograms. Mass characteristics of peak 14 are quite unusual for an anthocyanin, although its UV-vis spectrum (Fig. 4) is in support of its anthocyanin nature. In any case, λ_{\max} in the visible region (504 nm) is

not characteristic of a Mv-derived anthocyanin (λ_{\max} 525–530 nm) but rather of a Pg-derived or a C-4-substituted anthocyanin. Thus, the existence of malvidin as anthocyanidin in compound 14 is doubtful and no definitive structure could be matched with its spectral and mass features.

3.2. Content and distribution of anthocyanins

Table 2 shows the individual concentrations of the three major anthocyanins (i.e. Cy 3-gluc, Pg 3-gluc, Pg 3-rut) in the extracts of the five strawberry varieties in the two harvests analysed. Total anthocyanins would range between 200 and 600 mg kg⁻¹ fresh weight. In general, these anthocyanin contents are roughly similar to those reported by other authors. Values of total anthocyanins ranging 150–350 mg kg⁻¹ were collected by Clifford (2000) from the literature.

In our study notable variability was found among samples of each variety and year, suggesting that edaphic-climatic factors and degree of maturity have a strong influence on the anthocyanin levels. Nevertheless, some influence of the strawberry cultivar may also exist. Meyers, Watkins, Pritts, and Hai-Liu (2003), in a study carried out with eight different strawberry varieties, reported average concentrations of 414 mg kg⁻¹, with strong differences among cultivars, the richest one

Table 2
Anthocyanin contents in samples from five strawberry cultivars harvested in 2001 and 2002

		Cy 3-gluc ^a	Pg 3-gluc ^b	Pg 3-rut ^b
<i>Camarosa</i>	2001	25±6	261±36	43±7
	2002	41±13	384±88	55±9
<i>Carisma</i>	2001	10±3	242±49	15±5
	2002	13±5	314±112	24±4
<i>Eris</i>	2001	27±8	185±66	13±5
	2002	23±4	163±22	14±3
<i>Oso Grande</i>	2001	24±3	289±34	29±2
	2002	11±1	162±6	16±2
<i>Tudnew</i>	2001	24±9	468±124	33±13
	2002	28±4	401±47	31±1

Concentrations (mean ± s.d.; n = 3) are in fresh weight.

^aExpressed as mg Cy 3-gluc.kg⁻¹.

^bExpressed as mg Pg 3-gluc.kg⁻¹.

(*Earliglow*) having twice the concentration than the poorest (*Allstar*). None of the varieties studied by those authors coincides with the ones analysed by us. Concentrations of 185.0±15.0 and 840.2±5.7 mg kg⁻¹ were determined in samples of *Oso Grande* and *Camarosa*, respectively, by Garcia-Viguera, Zafrilla, and Tomas-Barberan (1998). The high levels found in *Camarosa* by those authors were explained by the more intense pigmentation of the inner tissues of the fruit than found in other varieties. Such differences were also observed in the *Camarosa* samples here analysed, as well as in those of cv. *Tudnew*, varieties that showed the highest concentrations among those studied. By contrast, the faintest inner pigmentation and the lowest concentrations were observed in the samples of cv. *Eris*. Anthocyanin content in *Camarosa* strawberries was also determined by Castro, Gonçalves, Teixeira, and Vicente (2002) obtaining values closer to those found in our samples (482±14 mg kg⁻¹). It should also be taken into account that differences in concentration found for a same variety by different authors might also be due to the use of different extraction solvents.

Regarding anthocyanin distribution, Pg 3-gluc is the predominant compound in the strawberry extracts (83% of total anthocyanins on average), usually followed by Pg 3-rut (8%) and Cy 3-gluc (7%). Percentages of 89–95% for Pg 3-gluc and 4–11% for Cy 3-gluc were found by Goiffon et al. (1999) in a study carried out with five strawberry varieties different to those analysed by us. Curiously, this author does not report the presence of Pg 3-rut, but indicates Pg 3-arabinoside to be the third anthocyanin in the varieties *Senga sengana* and *El santa*.

In our study, the samples of *Camarosa* were those with lower percentages of Pg 3-gluc (77–78%) and the highest ones of Pg 3-rut (11–13%), whilst samples of *Carisma* possessed the highest proportions of Pg 3-gluc (88–89%) and the lowest ones of Cy 3-gluc (3–4%). *Eris* was the only variety that showed greater levels of Cy 3-gluc (12%) than Pg 3-rut (6–7%). The proportions found among these three

anthocyanins were fairly consistent for the different samples of the same variety (Table 2), suggesting a characteristic anthocyanin distribution. A similar conclusion was obtained by Garcia-Viguera et al. (1998) who also found similar proportions to us in cv. *Camarosa* and *Oso Grande*.

According to Tamura et al. (1995) Pg 3-malonylglucoside is one of the main anthocyanins in strawberry comprising 30%, 25%, and 12% of the total anthocyanin content in cv's *Nyoho*, *Himesodachi* and *Reiko*, respectively. Proportions between 5% and 24% of this anthocyanin are also determined by Yoshida et al. (2002) in 11 Japanese strawberry cultivars. In our case, no important levels of acylated anthocyanins were found in the samples analysed. The highest percentages were determined in the extracts of cv. *Tudnew* where Pg 3-malonylglucoside and Pg 3-acetylglucoside represented about 1% of total anthocyanins. At this respect, it is pertinent to indicate that extraction with solvents containing HCl may result in pigment degradation due to the concentration of HCl, and this is one of the reasons why anthocyanins containing labile aliphatic acyl moieties may have been overlooked (Strack & Way, 1989). In assays carried out in our laboratory with grape anthocyanins no relevant decrease in the levels of anthocyanins acylated with acetic acid was observed when 0.1% HCl in methanol was used for extraction, as in the present study. We found that a critical point, even when low HCl percentages are used, is the evaporation phase, where an increase in the concentration of HCl is produced as a result of the removal of the organic solvent. For that reason, in this study, water was added prior to solvent evaporation to prevent the increase in the concentration of HCl during evaporation. In spite of this, some cleavage of the aliphatic acyl residues cannot be discarded with subsequent underestimation of the corresponding acylated anthocyanins.

4. Conclusions

The use of optimized HPLC conditions coupled to diode array and mass detection allowed us to detect up to 25 different anthocyanin pigments in strawberry fruits from five different varieties (cv. *Eris*, *Oso Grande*, *Carisma*, *Tudnew* and *Camarosa*). Most anthocyanins showed Pg as aglycone, although some Cy derivatives were also present. Glucose was the most usual substituting sugar, but rutinose and possibly arabinose and rhamnose have been found, as well as acylation with different aliphatic acids (e.g. malic, malonic, succinic or acetic acids). A relevant aspect was the confirmation in the strawberry fruits of the presence of anthocyanin-derived pigments, namely a Pg-derived pyranoanthocyanin and four condensed products containing C–C linked anthocyanin (Pg) and flavanol (catechin and afzelechin) residues, whose detection in natural plant sources was very recently made. Up to then, this type of compounds was associated with reactions taking place during processing and ageing of red wines. In the five

strawberry varieties studied, Pg 3-gluc is always the predominant anthocyanin, usually followed by Pg 3-rut and Cy 3-gluc. These three compounds represented more than 95% of total anthocyanins in strawberry. *Tudnew* and *Camarosa* were the varieties with the higher anthocyanin content among those analysed.

Acknowledgements

Financial support of *Junta de Castilla y León* (Spain) through the grant ref. SA086/01 is greatly acknowledged. Author Lopes da Silva was funded by European Union and Government of Portugal through the Programme PRODEP III (2001–2003). Strawberry samples were kindly provided by Dr. Luis Carlos Sanz Martinez, Scientific Researcher at *Instituto de la Grasa-CSIC* in Seville (Spain).

References

- Andersen, Ø. M., Fossen, T., Torskangerpoll, K., Fossen, A., & Hauge, U. (2004). Anthocyanin from strawberry (*Fragaria ananassa*) with the novel aglycone, 5-carboxypyranopelargonidin. *Phytochemistry*, *65*, 405–410.
- Bakker, J., Bridle, P., & Bellworthy, S. J. (1994). Strawberry juice colour: a study of the quantitative and qualitative pigment composition of juices from 39 genotypes. *Journal of the Science of Food and Agriculture*, *64*, 31–37.
- Bridle, P., & Garcia-Viguera, C. (1997). Analysis of anthocyanins in strawberries and elderberries. A comparison of capillary zone electrophoresis and HPLC. *Food Chemistry*, *59*, 299–304.
- Castro, I., Gonçalves, O., Teixeira, J. A., & Vicente, A. A. (2002). Comparative study of a *Selva* and *Camarosa* strawberries from the commercial market. *Journal of Food Science*, *67*, 2132–2137.
- Clifford, M. N. (2000). Anthocyanins: nature, occurrence and dietary burden. *Journal of the Science of Food and Agriculture*, *80*, 1063–1072.
- Co, H., & Markakis, P. (1968). Flavonoid compounds in strawberry fruits. *Journal of Food Science*, *33*, 281–283.
- de Pascual-Teresa, S., Santos-Buelga, C., & Rivas-Gonzalo, J. C. (2002). LC-MS analysis of anthocyanins from purple corn cob. *Journal of the Science of Food and Agriculture*, *82*, 1003–1006.
- Escribano-Bailón, M. T., Santos-Buelga, C., Alonso, G. L., & Salinas, M. R. (2002). Anthocyanin composition of *Coriaria myrtifolia* L. fruit. *Phytochemical Analysis*, *13*, 354–357.
- Fiorini, M. (1995). Preparative high-performance liquid chromatography for the purification of natural anthocyanins. *Journal of Chromatography A*, *692*, 213–219.
- Fossen, T., Rayyan, S., & Andersen, Ø. M. (2004). Dimeric anthocyanins from strawberry (*Fragaria ananassa*) consisting of pelargonidin 3-glucoside covalently linked to four flavan-3-ols. *Phytochemistry*, *65*, 1421–1428.
- Garcia-Viguera, C., Zafrilla, P., & Tomas-Barberan, F. T. (1998). The use of acetone as an extraction solvent for anthocyanins from strawberry fruits. *Phytochemical Analysis*, *9*, 274–277.
- Goiffon, J. P., Mouly, P. P., & Gaydou, E. M. (1999). Anthocyanic pigment determination in red fruit juices, concentrated juices and syrups using liquid chromatography. *Analytica Chimica Acta*, *382*, 39–50.
- González-Paramás, A. M., Lopes da Silva, F., Martín-López, P., Macz-
Pop, G., González-Manzano, S., Alcalde-Eon, C., Pérez-Alonso, J. J.,
et al. (2005). Flavanol-anthocyanin condensed pigments in natural plant sources. *Food Chemistry* Available online since 16/02/2005.
- Hebrero, E., Santos-Buelga, C., & Rivas-Gonzalo, J. C. (1988). HPLC-DAS identification of anthocyanins of *Vitis vinifera* variety Tempranillo. *American Journal of Enology and Viticulture*, *39*, 227–233.
- Heinonen, I. M., Meyer, A. S., & Frankel, E. N. (1998). Antioxidant activity of berry phenolics on human low-density lipoprotein and liposome oxidation. *Journal of Agricultural and Food Chemistry*, *46*, 4107–4112.
- Hong, V., & Wrolstad, R. E. (1990a). Characterization of anthocyanin-containing colorants in fruit juices by HPLC-photodiode array detection. *Journal of Agricultural and Food Chemistry*, *38*, 698–707.
- Hong, V., & Wrolstad, R. E. (1990b). Use of HPLC separation/photodiode array detection for characterization of anthocyanins. *Journal of Agricultural and Food Chemistry*, *38*, 708–715.
- Jurd, L. (1969). Review of polyphenol condensation reactions and their possible occurrence in the aging of wines. *American Journal of Enology and Viticulture*, *20*, 191–195.
- Lopes-da-Silva, F., de Pascual-Teresa, S., Rivas-Gonzalo, J. C., & Santos-Buelga, C. (2002). Identification of anthocyanin pigments in strawberry (*cv Camarosa*) by LC using DAD and ESI-MS detection. *European Food Research and Technology*, *214*, 248–253.
- Lukton, A., Chichester, C. O., & MacKiney, G. (1955). Characterization of a second pigment in strawberries. *Nature*, *176*, 790.
- Mateus, N., de Pascual-Teresa, S., Rivas-Gonzalo, J., Santos-Buelga, C., & de Freitas, V. (2002). Structural diversity of anthocyanin-derived pigments in port wines. *Food Chemistry*, *76*, 335–342.
- Mazza, G., & Miniati, E. (1993). *Anthocyanins in fruits, vegetables and grains*. Boca Raton, FL: CRC Press Inc.
- Meyers, K. J., Watkins, C. B., Pritts, M. P., & Hai-Liu, R. (2003). Antioxidant and antiproliferative activities of strawberries. *Journal of Agricultural and Food Chemistry*, *51*, 6887–6892.
- Robinson, G. M., & Robinson, M. A. (1931). Survey of anthocyanins I. *Biochemical Journal*, *25*, 1687–1705.
- Salas, E., Atanasova, V., Poncet-Legrand, C., Meudec, E., Mazauric, J. P., & Cheyner, V. (2004). Demonstration of the occurrence of flavanol-anthocyanin adducts in wine and model solutions. *Analytica Chimica Acta*, *513*, 325–332.
- Somers, T. C. (1971). The polymeric nature of wine pigments. *Phytochemistry*, *10*, 2175–2186.
- Strack, D., & Way, V. (1989). Anthocyanins. In J. B. Harborne (Ed.), *Methods in Plant Biochemistry*, Vol. I (pp. 325–356). New York: Academic Press.
- Tamura, H., Takada, M., & Yoshida, Y. (1995). Pelargonidin 3-O-(6-O-malonyl-β-D-glucopyranoside) in *Fragaria x ananassa* Duch cv Nyoho. *Bioscience, Biotechnology and Biochemistry*, *59*, 1157–1158.
- Vivar-Quintana, A. M., Santos-Buelga, C., Francia-Aricha, E., & Rivas-Gonzalo, J. C. (1999). Formation of anthocyanin-derived pigments in experimental red wines. *Food Science and Technology International*, *5*, 347–352.
- Wang, S. Y., & Jiao, H. (2000). Scavenging capacity of berry crops on superoxide radicals, hydrogen peroxide, hydroxyl radicals and singlet oxygen. *Journal of Agricultural and Food Chemistry*, *48*, 5677–5684.
- Wang, S. Y., & Lin, H. S. (2000). Antioxidant activity in fruits and leaves of blackberry, raspberry and strawberry varies with cultivar and developmental stage. *Journal of Agricultural and Food Chemistry*, *48*, 140–146.
- Yoshida, Y., Koyama, N., & Tamura, H. (2002). Color and anthocyanin composition of strawberry fruit: changes during fruit development and differences among cultivars, with special reference to the occurrence of pelargonidin 3-malonylglucoside. *Journal of the Japanese Society for Horticultural Science*, *71*, 355–361.