


Redox-Active Ligands

International Edition: DOI: 10.1002/anie.201908744
German Edition: DOI: 10.1002/ange.201908744**[3+2] Fragmentation of a Pentaphosphido Ligand by Cyanide****

Christian M. Hoidn, Thomas M. Maier, Karolina Trabitsch, Jan J. Weigand, and Robert Wolf*

Abstract: The activation of white phosphorus (P_4) by transition-metal complexes has been studied for several decades, but the functionalization and release of the resulting (organo)-phosphorus ligands has rarely been achieved. Herein we describe the formation of rare diphosphan-1-ide anions from a P_5 ligand by treatment with cyanide. Cobalt diorganopentaphosphido complexes have been synthesized by a stepwise reaction sequence involving a low-valent diimine cobalt complex, white phosphorus, and diorganochlorophosphanes. The reactions of the complexes with tetraalkylammonium or potassium cyanide afford a cyclotriphosphido cobaltate anion **5** and 1-cyanodiphosphan-1-ide anions $[R_2PPCN]^-$ (**6-R**). The molecular structure of a related product **7** suggests a novel reaction mechanism, where coordination of the cyanide anion to the cobalt center induces a ligand rearrangement. This is followed by nucleophilic attack of a second cyanide anion at a phosphorus atom and release of the P_2 fragment.

Over the past few decades, studies on the activation of white phosphorus have produced numerous early and late-transition-metal polyphosphido complexes.^[1] The functionalization of the polyphosphorus ligands in such compounds with electrophiles and/or nucleophiles is potentially an elegant and atom-economical route to unique phosphorus compounds. However, such transition-metal-mediated P_4 functionalizations and, in particular, the release of useful phosphorus building blocks from the metal are generally difficult to achieve because of the low reactivity of many known polyphosphido compounds toward electrophiles.^[2] Rare examples of successful electrophilic functionalization reactions are shown in Scheme 1a. Peruzzini, Stoppioni, and co-workers reported the methylation of a *cyclo*- P_3 ligand at Group 9 metal cations,^[3] Cummins and co-workers described



Scheme 1. a) Examples of established functionalizations of polyphosphorus ligands: i) + CF_3SO_3Me or + $[Me_3O]BF_4/-Me_2O$; $[M] = [M(\text{triphos})]$ ($M = Co, Rh, Ir$, triphos = $CH_3C(CH_2PPh_2)_3$); ii) + $AsCl_3$ or + $SbCl_3/-NaCl$, $-[Nb]Cl_2(thf)$; $[Nb] = [Nb(ODipp)_3]$ (Dipp = 2,6-*iPr*₂C₆H₃); iii) + $[(L')SiCl]/-[Zr]Cl_2$, $L' = PhC(NtBu)_2$; $[Zr] = [(C_5H_3tBu)_2Zr]$; iv) + $R'_2PCL/-KCl$, $-[Ga]$; $[Ga] = [Ga(CH(CMeNDipp)_2)]$, $[Co] = [(^{Mes}BIAN)Co]$, BIAN = 1,2-bis(arylimino)acenaphthene; $R' = Cy, iPr, tBu$. b) Ligand functionalization and subsequent fragmentation reactions affording new phosphorus compounds ($L = PHDI = \text{bis}(2,6\text{-diisopropylphenyl})\text{phenanthrene-9,10\text{-diimine}}$; $R = Cy, tBu, Ph, Mes, N(iPr)_2$).

the remarkable synthesis of EP_3 ($E = As, Sb$) molecules from $[Nb(ODipp)_3(\eta^3-P_3)]^-$ (Dipp = 2,6-*iPr*₂C₆H₃) and ECl_3 ,^[4] and quite recently Scheer and co-workers disclosed the synthesis of a phosphorus-silicon analogue of benzene from a tetraphosphidozirconium complex.^[5] Recent studies from our groups have shown that the heterodinuclear cobalt-gallium species **A** can be successfully applied for the construction of novel alkyl-substituted pentaphosphido complexes **B**.^[6] However, the synthesis of **A** is cumbersome, and the reaction properties of **B**-type species have, therefore, not been explored.


Here we present a much more facile synthesis of diorganopentaphosphido ligands and their fragmentation into P_2 and P_3 units (Scheme 1b). The bis(2,6-diisopropylphenyl)phenanthrene-9,10-diimine (PHDI) complex $[K(18c-6)(thf)_{1.5}][(\text{PHDI})Co(\eta^4-1,5\text{-cod})]$ ($[K(18c-6)]\mathbf{1}$; 18c-6 = [18]crown-6, cod = cycloocta-1,5-diene) enables the straightforward, high-yielding synthesis of an anionic *cyclo*- P_4 cobalt

[*] C. M. Hoidn, T. M. Maier, K. Trabitsch, Prof. Dr. R. Wolf
University of Regensburg
Institute of Inorganic Chemistry
93040 Regensburg (Germany)
E-mail: robert.wolf@ur.de
Prof. Dr. J. J. Weigand
TU Dresden, Faculty of Chemistry and Food Chemistry
01062 Dresden (Germany)
E-mail: jan.weigand@tu-dresden.de

[**] A previous version of this manuscript has been deposited on a preprint server (<https://doi.org/10.26434/chemrxiv.8868701.v1>).

Supporting information and the ORCID identification numbers for some of the authors of this article can be found under: <https://doi.org/10.1002/anie.201908744>.

© 2019 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited, and is not used for commercial purposes.



Scheme 2. Synthesis of compounds [K(18c-6)]1–3 (18c-6 = [18]crown-6); reagents and by-products: a) + 18c-6/–1,5-cod; b) + P₄/–1,5-cod; c) + [W(CO)₅(thf)]/–thf; yields: [K(18c-6)]1 76%, [K(18c-6)]2 80%, [K(18c-6)]3 32%.


complex **2**, which is readily functionalized with dialkyl-, diaryl-, and diaminyldiphosphino groups. Cyanide anions cleave the resulting pentaphosphorus unit into a *cyclo*-P₃ complex **5** and 1-cyanodiphosphano-1-ide anions [R₂PPCN][–] (**6-R**).^[7]

Our investigation commenced with the synthesis of the new α -diimine complex [K(18c-6)]**1** by ligand exchange of the 1,5-cod in [K(thf)_{0.2}][Co(η^4 -1,5-cod)₂] with PHDI followed by the addition of [18]crown-6 (Scheme 2, reaction a). Filtration and crystallization from THF/*n*-hexane afforded analytically pure, dark-green crystals of [K(18c-6)]**1** in good yield. The NMR spectra of **1** compare well with those of the related 1,2-bis(arylimino)acenaphthene (^ABIAN) complexes K[Co(η^4 -1,5-cod)(^ABIAN)] (Ar = Dipp, Mes).^[6,8]

Next, the reaction of [K(18c-6)]**1** with white phosphorus was examined (Scheme 2, reaction b). Monitoring of the reaction by ³¹P{¹H} NMR spectroscopy showed quantitative conversion into [K(18c-6)][(PHDI)Co(η^4 -*cyclo*-P₄)] ([K(18c-6)]**2**) at room temperature. Crystallization from toluene/*n*-hexane gave [K(18c-6)]**2** as dark-turquoise crystals in up to 80% yield (corresponding to > 1.5 g of pure product). The ³¹P{¹H} NMR spectrum of [K(18c-6)]**2** in [D₈]THF shows a sharp singlet at $\delta = 136.5$ ppm (compare with the chemical shifts of $\delta = 175.2$ ppm observed for [(η^5 -Cp^{*t*Bu₃})Co(η^4 -P₄)]^[12] (Cp^{*t*Bu₃} = C₅H₂*t*Bu₃) and $\delta = 114.1$ ppm for [(η^5 -Cp^{Ar})Fe(η^4 -P₄)][–] (Cp^{Ar} = C₅(C₆H₄-4-Et)₅).^[10] A single-crystal X-ray diffraction (XRD) analysis clearly revealed the presence of a bidentate PHDI ligand and a terminal η^4 -coordinated *cyclo*-P₄ unit. A more detailed interpretation of the structure was unfortunately prevented by heavy disorder within the *cyclo*-P₄ unit (see Figure S57 in the Supporting Information for further details). However, the single-crystal X-ray diffraction analysis of an adduct [K(18c-6)][(PHDI)Co(μ^2 : η^1 , η^4 -P₄)W(CO)₅] ([K(18c-6)]**3**; Scheme 2, reaction c), formed quantitatively from [K(18c-6)]**2** and [W(CO)₅(thf)] according to monitoring by ³¹P NMR spectroscopy, revealed an ordered structure (see Figure S58). An almost square, η^4 -coordinated *cyclo*-P₄ unit similar to those in related complexes is observed.^[9–12] The P–P bond lengths (2.132(4) to 2.173(5) Å, mean: 2.147(7) Å) are in-between the values expected for P–P (2.22 Å) and P=P bonds (2.04 Å) and suggest the presence of a (*cyclo*-P₄)^{2–} dianion.^[11e,13] The C–C and C–N bond lengths in the PHDI backbone (C–C 1.42(1) Å; C–N 1.36(1) Å) indicate the presence of PHDI in its radical anionic form.^[14] Further details of the characterization of [K(18c-6)]**3**, including a discussion of the NMR data, are given in the Supporting Information.

In general, mononuclear *cyclo*-P₄ complexes are still surprisingly rare. Besides two recently published iron complexes,^[9,10] only examples with early transition metals (e.g. V, Nb, Ta, Mo) are known to date.^[11] A recently reported, neutral *cyclo*-P₄ cobalt sandwich complex [(η^5 -Cp^{*t*Bu₃})Co(η^4 -P₄)] described by Scheer and co-workers is unstable at room temperature and only accessible in low yields after column chromatography.^[12] In contrast, [K(18c-6)]**2** is indefinitely stable at room temperature, both as a solid and in solution.

This finding suggests [K(18c-6)]**2** to be an excellent precursor for P–P bond-forming reactions with diorganochlorophosphanes. Reactions with R₂PCl (R = Cy, *t*Bu, Ph, Mes, N(*i*Pr)₂; Scheme 3) proceeded quantitatively (³¹P NMR



Scheme 3. Functionalization of the P₄ unit in **2** by various chlorophosphanes to afford the pentaphosphido complexes **4-R**; reagents and by-products: + R₂PCl/–KCl, –18c-6; yields range from 33 to 77%.

monitoring) to afford pentaphosphido complexes [(PHDI)Co(η^4 -P₅R₂)] (**4-R**), which were isolated as dark-blue crystals in up to 77% yield by a convenient work-up procedure. Analogous reactions with K₂[Co₂(μ^2 : η^4 , η^4 -P₄)(^{Dipp}BIAN)₂] were observed to lead only to unproductive outer-sphere electron-transfer reactions.^[8]

Complexes **4-R** are isostructural and resemble those of the recently reported species [(^{Mes}BIAN)Co(η^4 -P₅R₂)] (**B**, R = *i*Pr, *t*Bu, Cy; see Scheme 1 a).^[6] The molecular structure of **4-tBu** is shown as an example in Figure 1 a; the remaining structures with R = Cy, Ph, Mes, and NiPr₂ are given in the Supporting Information. The η^4 -*cyclo*-P₅R₂ ligands show an envelope conformation with P–P bond lengths ranging from 2.1197(2) to 2.182(1) Å, which indicates some delocalized character.^[13] The C–C and C–N distances in **4-R** suggest the presence of PHDI[–] radical anions.^[6,14] The ³¹P{¹H} NMR spectra recorded in C₆D₆ show an AMM'XX' spin system in each case (see Figure 1 b for the spectrum of **4-tBu** as an example).^[6] The chemical shift of the tetracoordinate phosphorus atoms P_A varies with the substituent (e.g. $\delta =$





Figure 1. a) Solid-state molecular structure of $[(\text{PHDI})\text{Co}(\eta^3\text{-P}_5\text{tBu}_2)]$ (**4-tBu**). Hydrogen atoms are omitted for clarity and thermal ellipsoids are drawn at the 60% probability level. Selected bond lengths [Å] and angles [°]: P1–P2 2.1410(9), P1–P5 2.1596(8), P2–P3 2.132(1), P3–P4 2.1394(9), P4–P5 2.1487(9), Co1–P1 2.3687(7), Co1–P2 2.3463(7), Co1–P3 2.3236(7), Co1–P4 2.3928(7), Co1–N1 1.893(2), Co1–N2 1.896(2), C1–N1 1.360(3), C2–N2 1.360(3), C1–C2 1.427(3); P1–P2–P3 103.63(3), P2–P3–P4 105.11(4), P3–P4–P5 100.41(4), P4–P5–P1 95.69(3), P5–P1–P2 100.63(3). b) Section of the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **4-tBu** with nuclei assigned to an AMM'XX' spin system; experimental (top); simulation (bottom): $\delta(\text{P}_A) = 161.7$ ppm, $\delta(\text{P}_{MM'}) = 91.4$ ppm, $\delta(\text{P}_{XX'}) = -171.6$ ppm, $^1J_{AX} = ^1J_{AX'} = -396.3$ Hz, $^1J_{MX} = ^1J_{MX'} = -404.1$ Hz, $^1J_{MM'} = -383.6$ Hz, $^2J_{MX} = ^2J_{MX'} = 35.9$ Hz, $^2J_{AM} = ^2J_{AM'} = 10.9$ Hz, $^2J_{XX'} = -4.9$ Hz; the spectra of the residual compounds **4-R** are very similar (see the Supporting Information for further details); [Co] = (PHDI)Co.

161.7 ppm for **4-tBu** versus $\delta = 84.3$ ppm for **4-Mes**), while the remaining data are similar to those observed for complexes **B** shown in Scheme 1.

Having substantial quantities of *cyclo*- P_5R_2 complexes **4-R** ($\text{R} = \text{Cy}, \text{tBu}, \text{Ph}, \text{Mes}, \text{N}(\text{iPr})_2$) in hand, reactions of these compounds with nucleophiles were examined in the hope of releasing the pentaphosphorus fragment. Although initial reactions with carbon monoxide, cyclohexylisocyanide, and *N,N'*-diisopropyl-4,5-dimethylimidazolin-2-ylidene gave intractable mixtures, the reaction with cyanide salts $[\text{M}]\text{CN}$ ($[\text{M}] = [\text{nBu}_4\text{N}]^+, [\text{Et}_4\text{N}]^+, [\text{K}(\text{18c-6})]^+$; Scheme 4, reaction a) was far more promising.^[15] An immediate color change from dark blue to dark purple-blue occurred upon addition of the cyanide salt. Monitoring the reaction by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy revealed the clean formation of $[(\text{PHDI})\text{Co}(\eta^3\text{-P}_3)(\text{CN})]^-$ (**5**; singlet at $\delta = -193.2$ ppm), which is a very rare example of an anionic *cyclo*- P_3 cobalt complex.^[16] The tetra-*n*-butylammonium salt $[\text{nBu}_4\text{N}]\text{5}$ can be isolated in 40–64% yield by crystallization from toluene. Compared to related neutral species such as $[(\text{ArNC})_3\text{Co}(\eta^3\text{-P}_3)]$ ($\text{Ar} = 2,6\text{-}(\text{Mes})_2\text{C}_6\text{H}_3$, $\delta = -276$ ppm),^[17] and $[(\text{CH}_3\text{C}(\text{CH}_2\text{PPh}_2)_3)\text{Co}(\eta^3\text{-P}_3)]$ ($\delta = -278$ ppm),^[18] the $^{31}\text{P}\{^1\text{H}\}$ NMR resonance of $[\text{nBu}_4\text{N}]\text{5}$ ($\delta = -193.2$ ppm) is shifted downfield. In addition,



Scheme 4. a) Fragmentation and b) rearrangement of the pentaphosphido ligand in **4-R** depending on the substituent R. Reagents and by-products: a) for $\text{R} = \text{Cy}, \text{tBu}, \text{Ph}, \text{N}(\text{iPr})_2$: + 2 equiv $[\text{M}]\text{CN}$; b) only for $\text{R} = \text{Mes}$: + 1 equiv $[\text{M}]\text{CN}$ ($[\text{M}] = [\text{nBu}_4\text{N}]^+, [\text{Et}_4\text{N}]^+, [\text{K}(\text{18c-6})]^+$). I and II are two conceivable resonance structures of anions **6-R**.

the formation of the new cyanodiphoshanide anions $[\text{R}_2\text{PPCN}]^-$ (**6-R**, $\text{R} = \text{Cy}, \text{tBu}, \text{Ph}, \text{N}(\text{iPr})_2$) was observed, which were identified by the characteristic set of doublets ($^1J_{\text{PP}} = 261$ to 278 Hz) in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectra.

The reaction thus results in an unusual fragmentation of the polyphosphido ligand into a P_3 and a P_2 unit. To our knowledge, there is only one related reaction involving the [3+2] fragmentation of a P_5 species in the literature, which has a completely different outcome. As reported by Weigand and co-workers, the reaction of the $[\text{P}_5\text{DippCl}]^+$ cation (Dipp = 2,6-diisopropylphenyl) with *N,N'*-bis(2,6-diisopropylphenyl)-4,5-dichloroimidazolin-2-ylidene (IPrCl_2) affords a triphosphaallyl cation $[\text{P}_3(\text{IPrCl}_2)_2]^+$ and a neutral P_2 species $[(\text{IPrCl}_2)\text{PP}(\text{Cl})\text{Dipp}]$.^[19]

$[\text{nBu}_4\text{N}]\text{6-tBu}$ can be isolated as a light-purple solid in 40% yield by extracting the crude reaction mixture with cyclohexane/*n*-hexane (3:2 v/v).

Although X-ray quality crystals of $[\text{nBu}_4\text{N}]\text{5}$ and $[\text{nBu}_4\text{N}]\text{6-tBu}$ could not be obtained so far, the single-crystal X-ray structures of the two potassium salts $[\text{K}(\text{18c-6})]\text{5}$ and $[\text{K}(\text{18c-6})]\text{6-tBu}$ (Figure 2) have been obtained from separate single crystals grown from the same reaction mixture (attempts to separate these compounds on a preparative scale have so far been unsuccessful). The molecular structure of $[\text{K}(\text{18c-6})]\text{5}$ features a cobalt atom coordinated by a cyanide anion, a formally neutral PHDI ligand [C1–N1 1.338(2), C2–N2 1.344(2), and C1–C2 1.453(2) Å],^[20] and a η^3 -coordinated *cyclo*- P_3 ring.

The mean P–P (2.1361(8) Å) and Co–P (2.300(5) Å) bond lengths in $[\text{K}(\text{18c-6})]\text{5}$ are similar to those in neutral CoP_3 complexes.^[16–18,21] The Co–C (1.909(2) Å) and C≡N (1.153(2) Å) bond lengths of the cyanide ligand as well as the CN stretching frequency ($\bar{\nu} = 2068$ cm^{-1}) are typical for cyanide-cobalt complexes.^[22] The solid-state molecular structure of $[\text{K}(\text{18c-6})]\text{6-tBu}$ shows an almost linear, phosphanyl-substituted PCN moiety (P1–C1–N1 178.4(1)°) with a P–P distance of 2.1895(4) Å, that is, close to a typical single bond.^[13] The




Figure 2. Solid-state molecular structures of $[K(18c-6)][(PHDI)Co(\eta^3-P_3)(CN)]$ (left), $[K(18c-6)]5$ and $[K(18c-6)][(tBu_2PPCN)]$ (right, $[K(18c-6)]6-tBu$). Hydrogen atoms, solvent molecules, and disorder are omitted for clarity and thermal ellipsoids are drawn at the 50% probability level. Selected bond lengths [Å] and angles [°]: $[K(18c-6)]5$: P1–P2 2.1256(8), P1–P3 2.1228(7), P2–P3 2.1599(7), Co1–P1 2.2780(5), Co1–P2 2.3200(5), Co1–P3 2.3027(5), Co1–N1 1.908(1), Co1–N2 1.906(2), Co1–C3 1.909(2), C1–N1 1.338(2), C2–N2 1.344(2), C3–N3 1.153(2), C1–C2 1.453(2), K1...N3 2.714(2); P1–P2–P3 59.38(2), P1–P3–P2 59.51(2), P2–P1–P3 61.12(2), Co1–C3–N3 177.0(2); $[K(18c-6)]6-tBu$: P1–P2 2.1895(4), P1–C1 1.763(1), C1–N1 1.160(2), K1...N1 2.828(1); P1–C1–N1 178.4(1), C1–P1–P2 92.43(4).

structural motif is reminiscent of the anions $[P(CN)_2]^-$ and $[PhPCN]^-$ reported by Schmidpeter et al.^[23–25] The IR spectrum shows the characteristic ν_{CN} stretch at $\tilde{\nu} = 2049\text{ cm}^{-1}$, which is lower than those of known dicyanophosphanides $[P(CN)_2]^-$ ($\tilde{\nu} = 2120, 2113\text{ cm}^{-1}$)^[23,24] and other trivalent monocyano phosphanes ($\tilde{\nu} \approx 2160\text{ cm}^{-1}$).^[26]

Recently, Borger, Grützmacher, and co-workers described related cyanodiphosphanides $[(NHP)PCN]$ with bulky *N*-heterocyclic phosphonium (NHP) substituents.^[7] These species feature a similar CN stretching frequency ($\tilde{\nu} = 2087\text{ to }2046\text{ cm}^{-1}$) and similar bond parameters as $[K(18c-6)]6-tBu$.^[7,23–25] Analogous to the compounds reported by Grützmacher and co-workers, the electronic structure of **6-R** may be described as both a cyanophosphanide ($R'P^--C\equiv N$, **I**, $R' = PR_2$) and a phosphaketeneimide ($R'P=C=N^-$, **II**, $R' = PR_2$) resonance structure (c.f. Scheme 4, pathway a). A natural resonance analysis at the B3LYP/6-31G+* level shows that the phosphaketeneimide form plays only a minor role in the electronic ground state of **6-tBu** (**I**: 69% versus **II**: 23%). Analogous calculations on related cyanate and phosphoethynolate anions ($O^--C\equiv X$ versus $O=C=X^-$, $X = N, P$) gave much higher contributions of the ketene form in NCO^- (33%) and PCO^- (40%).^[27]

Remarkably, a different product, $[Et_4N][(PHDI)Co(\eta^3-P_4PMe_2)(CN)]$ ($[Et_4N]7$), was isolated when the mesityl-substituted complex **4-Mes** was reacted with $[Et_4N]CN$ (one equiv, Scheme 4, pathway b). Monitoring the reaction by $^{31}P\{^1H\}$ NMR spectroscopy showed the reaction reaches full conversion within two hours at -30°C in MeCN; the main product, $[Et_4N]7$, was identified by an AB_2CD spin system ($\delta = 32.8, 59.2, 94.0, \text{ and } 193.7\text{ ppm}$). Deep-purple crystals were isolated in 38% yield from toluene/THF after work-up at low temperature ($< -30^\circ\text{C}$).^[28] A single-crystal XRD

structure (Figure 3) shows a bent P_4 ring coordinating to a cobalt center through three P atoms with P3–P4 and P4–P5 bond lengths of 2.1667(9) and 2.172(1) Å, respectively, thus indicating some delocalized character. The remaining P–P bond lengths (2.2416(9), 2.241(1), and 2.2381(9) Å) are typical for single bonds.^[13,29] The structural parameters of the PHDI ligand (C1–N1 1.328(4), C2–N2 1.335(3), and C1–C2 1.457(4) Å) are similar to those of $[K(18c-6)]5$, which suggests the presence of a neutral PHDI molecule.^[20]




Figure 3. Solid-state molecular structure of $[Et_4N][(PHDI)Co(\eta^3-P_4PMe_2)(CN)]$ ($[Et_4N]7$). Hydrogen atoms are omitted for clarity and thermal ellipsoids are drawn at the 50% probability level. Selected bond lengths [Å] and angles [°]: P1–P2 2.2416(9), P2–P3 2.241(1), P2–P5 2.2381(9), P3–P4 2.1667(9), P4–P5 2.172(1), Co1–P3 2.3356(8), Co1–P4 2.2671(8), Co1–P5 2.3174(8), Co1–N1 1.921(2), Co1–N2 1.927(2), Co1–C3 1.925(3), C1–N1 1.328(4), C2–N2 1.335(3), C3–N3 1.144(4), C1–C2 1.457(4); P1–P2–P3 94.77(4), P1–P2–P5 98.13(3), P2–P3–P4 90.36(4), P3–P4–P5 83.68(4), P4–P5–P2 90.28(4), P5–P2–P3 80.52(3).

The mechanism for the formation of $[Et_4N]7$ is proposed to involve an attack of the cyanide anion on the cobalt center, which induces a rearrangement to a phosphanyl-substituted *cyclo-P*₄ ring. Note that $[Et_4N]7$ is not converted into P_3 and P_2 products (**5** and hypothetical **6-Mes**) by the addition of a second equivalent of cyanide. Presumably, the bulky mesityl substituents prevents the CN^- anion from approaching P2. It is very likely that intermediates similar to $[Et_4N]7$ are formed in the fragmentation reactions of **4-R** with smaller substituents.

In conclusion, neutral pentaphosphido complexes **4-R** with an unprecedented range of alkyl, aryl, and amino substituents are readily accessible from $[K(18c-6)]1$, P_4 , and R_2PCl in two steps. The pentaphosphorus ligands of **4-R** undergo a remarkable [3+2] fragmentation upon reaction with two equivalents of cyanide, thereby forming the anionic cyclotriphosphido cobalt complex **5** and phosphanyl-substituted cyanodiphosphanides **6-R**. The reaction of **4-Mes** with cyanide affords $[Et_4N]7$, which contains a rearranged P_5Mes_2 ligand. The structure of this complex suggests that cyclo-tetraphosphido complexes are key intermediates en route to anions **5** and **6-R**. The results of this work show that diimine cobalt complexes are excellent platforms for studying the degradation of polyphosphorus ligands with inorganic nucleophiles. An extension of this approach to a wider range of polyphosphides and other nucleophiles should give rise to other unusual phosphorus compounds. In addition, reactions

of $[K(18c-6)]_2$ and $[nBu_4N]_5$ with electrophiles will likely give rise to unusual polyphosphanes and polyphosphanido complexes. Investigations into this research area are currently underway.

Acknowledgements

Financial support by the Deutsche Forschungsgemeinschaft (WE4621/3-1 and WO1496/7-1), the European Research Council (CoG 772299), the Stiftung der Deutschen Wirtschaft (sdw, fellowship to C.M.H.), and the Fonds der Chemischen Industrie (fellowship to T.M.M.) is gratefully acknowledged. We thank Dr. Stefanie Gärtner for crystallographic assistance as well as Dr. Daniel J. Scott and Dr. John Kelly (all University of Regensburg) for helpful comments on the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Keywords: cobalt · cyanides · phosphorus · pnictides · redox-active ligands

How to cite: *Angew. Chem. Int. Ed.* **2019**, *58*, 18931–18936
Angew. Chem. **2019**, *131*, 19107–19112

- [1] a) B. M. Cossairt, N. A. Piro, C. C. Cummins, *Chem. Rev.* **2010**, *110*, 4164–4177; b) M. Scheer, G. Balázs, A. Seitz, *Chem. Rev.* **2010**, *110*, 4236–4256; c) M. Caporali, L. Gonsalvi, A. Rossin, M. Peruzzini, *Chem. Rev.* **2010**, *110*, 4178–4235.
- [2] a) A. Velian, C. C. Cummins, *Chem. Sci.* **2012**, *3*, 1003; b) P. Barbaro, A. Ienco, C. Mealli, M. Peruzzini, O. J. Scherer, G. Schmitt, F. Vizza, G. Wolmershäuser, *Chem. Eur. J.* **2003**, *9*, 5195–5210; c) S. Pelties, A. W. Ehlers, R. Wolf, *Chem. Commun.* **2016**, *52*, 6601–6604; d) E. Mädl, M. V. Butovskii, G. Balázs, E. V. Peresypkina, A. V. Virovets, M. Seidl, M. Scheer, *Angew. Chem. Int. Ed.* **2014**, *53*, 7643–7646; *Angew. Chem.* **2014**, *126*, 7774–7777; e) P. Barbaro, C. Bazzicalupi, M. Peruzzini, S. Seniori Costantini, P. Stoppioni, *Angew. Chem. Int. Ed.* **2012**, *51*, 8628–8631; *Angew. Chem.* **2012**, *124*, 8756–8759.
- [3] G. Capozzi, L. Chiti, M. Di Vaira, M. Peruzzini, P. Stoppioni, *J. Chem. Soc. Chem. Commun.* **1986**, 1799–1800.
- [4] B. M. Cossairt, M.-C. Diawara, C. C. Cummins, *Science* **2009**, *323*, 602.
- [5] A. E. Seitz, M. Eckhardt, A. Erlebach, E. V. Peresypkina, M. Sierka, M. Scheer, *J. Am. Chem. Soc.* **2016**, *138*, 10433–10436.
- [6] C. G. P. Ziegler, T. M. Maier, S. Pelties, C. Taube, F. Hennersdorf, A. W. Ehlers, J. J. Weigand, R. Wolf, *Chem. Sci.* **2019**, *10*, 1302–1308.
- [7] Z. Li, J. E. Borger, F. Müller, J. R. Harmer, C.-Y. Su, H. Grützmacher, *Angew. Chem. Int. Ed.* **2019**, *58*, 11429–11433; *Angew. Chem.* **2019**, *131*, 11551–11555.
- [8] S. Pelties, T. Maier, D. Herrmann, B. de Bruin, C. Rebreyend, S. Gärtner, I. G. Shenderovich, R. Wolf, *Chem. Eur. J.* **2017**, *23*, 6094–6102.
- [9] A. Cavailé, N. Saffon-Merceron, N. Nebra, M. Fustier-Boutignon, N. Mézailles, *Angew. Chem. Int. Ed.* **2018**, *57*, 1874–1878; *Angew. Chem.* **2018**, *130*, 1892–1896.
- [10] U. Chakraborty, J. Leil, B. Mühldorf, M. Bodensteiner, S. Pelties, R. Wolf, *Dalton Trans.* **2018**, *47*, 3693–3697.
- [11] a) O. J. Scherer, R. Winter, G. Wolmershäuser, *Z. Anorg. Allg. Chem.* **1993**, *619*, 827–835; b) O. J. Scherer, J. Vondung, G. Wolmershäuser, *Angew. Chem. Int. Ed. Engl.* **1989**, *28*, 1355–1357; *Angew. Chem.* **1989**, *101*, 1395–1397; c) M. Herberhold, G. Frohmader, W. Milius, *J. Organomet. Chem.* **1996**, *522*, 185–196; d) K. A. Mandla, C. E. Moore, A. L. Rheingold, J. S. Figueroa, *Angew. Chem. Int. Ed.* **2019**, *58*, 1779–1783; *Angew. Chem.* **2019**, *131*, 1793–1797; e) K. A. Mandla, M. L. Neville, C. E. Moore, A. L. Rheingold, J. S. Figueroa, *Angew. Chem. Int. Ed.* **2019**, *58*, 15329–15333; *Angew. Chem.* **2019**, *131*, 15473–15477.
- [12] F. Dielmann, A. Timoshkin, M. Piesch, G. Balázs, M. Scheer, *Angew. Chem. Int. Ed.* **2017**, *56*, 1671–1675; *Angew. Chem.* **2017**, *129*, 1693–1698.
- [13] For calculated single- and double-bond lengths, see P. Pyykkö, M. Atsumi, *Chem. Eur. J.* **2009**, *15*, 12770–12779.
- [14] B. Gao, X. Luo, W. Gao, L. Huang, S.-m. Gao, X. Liu, Q. Wu, Y. Mu, *Dalton Trans.* **2012**, *41*, 2755–2763.
- [15] For the use of cyanide to release polyphosphorus molecules from a copper(I) iodide matrix, see A. Pfützer, M. F. Bräu, J. Zweck, G. Brunklaus, H. Eckert, *Angew. Chem. Int. Ed.* **2004**, *43*, 4228–4231; *Angew. Chem.* **2004**, *116*, 4324–4327.
- [16] While this manuscript was under review, a related anionic complex $(MeNHC)_2P[\eta^5-Cp^{tBu3}Co(\eta^3-P_3)]$ ($MeNHC = 1,3,4,5$ -tetramethylimidazol-2-ylidene) was published: M. Scheer, M. Piesch, S. Reichl, M. Seidl, G. Balázs, *Angew. Chem. Int. Ed.* **2019**, <https://doi.org/10.1002/anie.201908397>; *Angew. Chem.* **2019**, <https://doi.org/10.1002/ange.201908397>.
- [17] C. Chan, A. E. Carpenter, M. Gembicky, C. E. Moore, A. L. Rheingold, J. S. Figueroa, *Organometallics* **2019**, *38*, 1436–1444.
- [18] M. Di Vaira, L. Sacconi, P. Stoppioni, *J. Organomet. Chem.* **1983**, *250*, 183–195.
- [19] M. H. Holthausen, S. K. Surmiak, P. Jerabek, G. Frenking, J. J. Weigand, *Angew. Chem. Int. Ed.* **2013**, *52*, 11078–11082; *Angew. Chem.* **2013**, *125*, 11284–11288.
- [20] a) R. van Belzen, R. A. Klein, W. J. J. Smeets, A. L. Spek, R. Benedix, C. J. Elsevier, *Recl. Trav. Chim. Pays-Bas* **1996**, *115*, 275–285; b) R. van Belzen, R. A. Klein, H. Kooijman, N. Veldman, A. L. Spek, C. J. Elsevier, *Organometallics* **1998**, *17*, 1812–1825; c) W. W. Kramer, L. A. Cameron, R. A. Zarkesh, J. W. Ziller, A. F. Heyduk, *Inorg. Chem.* **2014**, *53*, 8825–8837; d) G. A. Abakumov, N. O. Druzhkov, T. N. Kocherova, K. A. Kozhanov, A. v. Murugova, E. N. Egorova, *Dokl. Chem.* **2016**, *467*, 109–112.
- [21] a) F. Ceconi, P. Dapporto, S. Midollini, L. Sacconi, *Inorg. Chem.* **1978**, *17*, 3292–3294; b) C. A. Ghilardi, S. Midollini, A. Orlandini, L. Sacconi, *Inorg. Chem.* **1980**, *19*, 301–306.
- [22] a) A search of the Cambridge Crystal Structure Database (CCSD), version 5.39 update 4, 14/05/2019, revealed 96 cobalt complexes bearing only one terminal cyanide ligand with a mean Co–C distance of 1.900 Å (median 1.886 Å) and a mean C≡N distance of 1.134 Å (median 1.141 Å); b) P. Rigo, A. Turco, *Coord. Chem. Rev.* **1974**, *13*, 133–172.
- [23] A. Schmidpeter, G. Burget, F. Zwaschka, W. S. Sheldrick, *Z. Anorg. Allg. Chem.* **1985**, *527*, 17–32.
- [24] W. S. Sheldrick, J. Kroner, F. Zwaschka, A. Schmidpeter, *Angew. Chem. Int. Ed. Engl.* **1979**, *18*, 934–935; *Angew. Chem.* **1979**, *91*, 998–1000.
- [25] A. Schmidpeter, K.-H. Zirzow, G. Burget, G. Huttner, I. Jibril, *Chem. Ber.* **1984**, *117*, 1695–1706.
- [26] a) S. Holand, F. Mathy, *Organometallics* **1988**, *7*, 1796–1801; b) A. Fischer, I. Neda, P. G. Jones, R. Schmutzler, *Phosphorus Sulfur Silicon Relat. Elem.* **1993**, *83*, 135–148; c) A. Vollbrecht, I. Neda, A. Fischer, P. G. Jones, R. Schmutzler, *Phosphorus Sulfur Silicon Relat. Elem.* **1995**, *107*, 69–78.
- [27] S. Alidori, D. Heift, G. Santiso-Quinones, Z. Benkő, H. Grützmacher, M. Caporali, L. Gonsalvi, A. Rossin, M. Peruzzini, *Chem. Eur. J.* **2012**, *18*, 14805–14811.

- [28] $[\text{Et}_4\text{N}]7$ decomposes to $[\text{Et}_4\text{N}]2$ and Mes_2PH in solution at room temperature over several days. These species were identified by their $^{31}\text{P}\{^1\text{H}\}$ NMR shifts: R. A. Bartlett, M. M. Olmstead, P. P. Power, G. A. Sigel, *Inorg. Chem.* **1987**, *26*, 1941–1946.
- [29] Related phosphoallylic *cyclo-P₄R* ligands are found in oligonuclear Cr, Mo, and W carbonyl complexes: a) M. Scheer, U. Becker, J. Magull, *Polyhedron* **1998**, *17*, 1983–1989; b) M. Scheer, M. Dargatz, P. G. Jones, *J. Organomet. Chem.* **1993**, *447*, 259–264.

Manuscript received: July 14, 2019
Accepted manuscript online: October 1, 2019
Version of record online: November 7, 2019