

**SUPERCRITICAL CARBON DIOXIDE
EXTRACTION OF MANGO SEED KERNEL FAT
BLENDED WITH PALM OIL MID-FRACTION
AND PALM STEARIN TO FORMULATE COCOA
BUTTER REPLACERS**

MD. JAHURUL HAQUE AKANDA

**UNIVERSITI SAINS MALAYSIA
2013**

**SUPERCRITICAL CARBON DIOXIDE EXTRACTION OF MANGO SEED
KERNEL FAT BLENDED WITH PALM OIL MID-FRACTION AND PALM
STEARIN TO FORMULATE COCOA BUTTER REPLACERS**

by

MD. JAHURUL HAQUE AKANDA

**Thesis submitted in fulfilment of the requirements
for the degree of
Doctor of Philosophy**

November 2013

ACKNOWLEDGEMENTS

Firstly, praise to God, Almighty Allah, who created and control the universe. This humble work could reach the present form through His blessing. My profound gratitude and appreciation are assigning to my supervisor Professor Ir. Mohd Omar Ab. Kadir and my field-supervisor Professor Md. Zaidul Islam Sarker for their insightful guidance's, encouragements and important supports throughout this work. I would like to express my thanks to Professor Nik Norulaini Nik Ab. Rahman for her valuable advices and comments throughout my Ph.D. program. I wish to acknowledge University Postgraduate Research Grant Scheme (PRGS, No. 1001/PTEKIND/845026) for the financial support and Universiti Sains Malaysia for the fellowship provided to me.

I would like to acknowledge Sime Darby Research Sdn Bhd for giving me chance to do some analysis of this thesis. Nur Azwani Ab. Karim, Scientist, Sime Darby Research Sdn Bhd is gratefully appreciated. I wish to express my warmest thanks to all staff of the School of Industrial Technology at Universiti Sains Malaysia for their kind help and cooperation. My special thanks to Mr. Mazlan Mohamed, Mr. Ravi Vinayagamuertty, Mr. Khairul, Mr. Shamsul Zoolkiffli, Ms. Noraida Bukhari, Mr. Azmaizan Yaakub for their accommodating the chemical analysis lab-work throughout this work. I also would like to thank my colleagues in the school of industrial Technology and specially Sahena, Shalima, Zainul, Sohrab, Moftah, Mark and other lab mate for the pleasant time and for sharing the lovely moments.

Finally, I would like to acknowledge my family: my parents Kalim Uddin and Mrs. Jhora Begum, my wife Maksuda, and son Mausooof for their love, support and kind encouragement.

TABLE OF CONTENTS

Acknowledgement	ii
Table of Contents	iii
List of Tables	xii
List of Figures	xv
List of Abbreviations	xvii
List of Appendices	xix
Abstrak	xx
Abstract	xxii
CHAPTER 1 – INTRODUCTION	
1.1 Research Background	1
1.2 Problem Statement	4
1.3 Objectives of Study	5
1.4 Research Contributions	6
1.5 Scope of Study	7
1.6 Limitation of Study	8
1.7 Overview of Thesis	9
CHAPTER 2 - LITERATURE REVIEW	
2.1 Introduction	10
2.2 Cocoa Butter	10
2.2.1 Background of Cocoa	10
2.2.2 Cocoa Production	10

2.2.3	Cocoa Butter Extraction	12
2.2.4	Composition of Natural Cocoa Butter	14
2.2.5	Physico-chemical Properties of Natural Cocoa Butter	15
2.2.6	Cocoa Butter Replacer (CBR)	17
2.2.6.1	Cocoa Butter Equivalent (CBE)	17
2.2.6.2	Cocoa Butter Substitute (CBS)	17
2.2.7	Alternative Sources of Cocoa Butter	17
2.2.7.1	Cocoa Butter Equivalent (CBE) from MSKF	18
2.2.7.2	Cocoa Butter from Palm Oil and Palm Kernel Oil	19
2.2.7.3	Cocoa Butter Replacers from Kokum Butter	24
2.2.7.4	Cocoa Butter Extender from Sal Fat	27
2.2.7.5	Cocoa Butter Substitute from Shea Butter	28
2.2.7.6	Cocoa Butter Equivalent from Illipé Butter	29
2.2.7.7	Cocoa Butter from Other Vegetable Oils	29
2.3	Palm Oil	30
2.3.1	Background	30
2.3.2	Extraction Process	31
2.3.3	Palm Oil Mid-Fraction (POMF) and Palm Stearin (PS)	31
2.3.4	Major Constituents of POMF and PS	32
2.3.5	Major Uses of Palm Oil	32
2.4	Mango (<i>Mangifera indica</i> L.)	35
2.4.1	Background	35
2.4.2	Mango Production	35
2.4.3	Extraction Process	36
2.4.4	Mango Seed Kernel Fat (MSKF)	36

2.4.5	Application of MSKF	37
2.5	Supercritical Carbon Dioxide (SC-CO ₂) Extraction	38
2.6	Advantages and Economy of SC-CO ₂ as a Solvent	39
2.7	Prospect and Limitation of SC-CO ₂ in the Oil Industry	42
2.8	Conventional Versus SC-CO ₂ Extraction Method	43
2.9	Research Gap to be Fulfilled by This Work	44

CHAPTER 3 – CONVENTIONAL AND SUPERCRITICAL CARBON DIOXIDE EXTRACTION OF MANGO SEED KERNEL FAT FROM DIFFERENT MANGO VARIETIES FOR SCREENING COCOA BUTTER ANALOGY FATS

3.1	Introduction	45
3.2	Material and Methods	46
3.2.1	Experimental Samples	46
3.2.2	Preparation of Mango Seed Kernel	46
3.2.3	Determination of Moisture Content in Mango Seed Kernel	47
3.2.4	Extraction of Fat from Mango Seed Kernel by Soxhlet Method	47
3.2.4.1	Method	47
3.2.4.2	Expression of Results	48
3.2.5	Supercritical Carbon Dioxide (SC-CO ₂) Extraction	49
3.2.5.1	Method	49
3.2.5.2	Continuous and Soaking Method	50
3.2.5.3	Expression of Results	50
3.2.6	Fatty Acid Methyl Esters Preparation	50
3.2.6.1	Reagents	51

3.2.6.2	Apparatus	51
3.2.6.3	Method	51
3.2.7	Determination of FAMES by Gas Chromatography	51
3.2.7.1	Materials	51
3.2.7.2	Instrument	52
3.2.7.3	Method	52
3.2.7.4	Expression of Results	52
3.2.8	Determination of Physico-chemical Properties of MSKF	53
3.2.8.1	Determination of Iodine Value	53
3.2.8.1.1	Reagents	53
3.2.8.1.2	Apparatus	54
3.2.8.1.3	Method	54
3.2.8.1.4	Expression of Results	54
3.2.8.2	Determination of Saponification Value (SPV)	55
3.2.8.2.1	Reagents	55
3.2.8.2.2	Apparatus	55
3.2.8.2.3	Method	56
3.2.8.2.4	Expression of Results	56
3.2.8.3	Determination of Slip Melting Point (SMP)	56
3.2.8.3.1	Apparatus	57
3.2.8.3.2	Method	57
3.2.8.4	Determination of Free Fatty Acid (FFA)	58
3.2.8.4.1	Reagents	58
3.2.8.4.2	Apparatus	58
3.2.8.4.3	Method	59

3.2.8.4.4	Expression of Results	59
3.3	Results and Discussion	59
3.3.1	Moisture Content	59
3.3.2	Total Fat in Different MSKs	60
3.3.3	Effects of SC-CO ₂ Methods on the Yield of MSKF	61
3.3.4	Fatty Acid Compositions	63
3.3.5	Physico-chemical Properties of SC-CO ₂ Extracted MSKF	70
3.3.5.1	Iodine Value	70
3.3.5.2	Saponification Value	70
3.3.5.3	Acid Value	71
3.3.5.4	Slip Melting Point	72
3.4	Conclusions	73

CHAPTER 4 – OPTIMIZATION OF SUPERCRITICAL CARBON DIOXIDE
EXTRACTION PARAMETERS EXTRACTING MANGO SEED KERNEL
FAT FROM MANGO SEED KERNEL USING RESPONSE SURFACE
METHODOLOGY

4.1	Introduction	75
4.2	Material and Methods	76
4.2.1	Experimental Samples	76
4.2.2	Preparation of Mango Seed Kernel	76
4.2.3	Determination of Moisture Content in Mango Seed Kernel	76
4.2.4	Extraction of Fat from Mango Seed Kernel by Soxhlet Method	77
4.2.5	Response Surface Methodology-Central Composite Design	77
4.2.5.1	Experimental Design	77

4.2.5.2	Expression of Results	78
4.3	Results and Discussion	78
4.3.1	Fitting the Model	78
4.3.2	Optimization and Effect of SC-CO ₂ Extraction Parameters on the Yield	81
4.4	Conclusions	84
CHAPTER 5 – COCOA BUTTER REPLACERS FROM SUPERCRITICAL CARBON DIOXIDE (SC-CO ₂) EXTRACTED MANGO SEED KERNEL FAT BLENDED WITH PALM OIL MID-FRACTION AND PALM STEARIN		
5.1	Introduction	86
5.2	Material and Methods	88
5.2.1	Materials and Chemicals	88
5.2.2	Supercritical Carbon Dioxide Extraction	89
5.2.3	Blending of Fat	89
5.2.4	Analysis of Fatty Acid Compositions of Each Blend	90
5.2.5	Determination of Physico-chemical Properties	90
5.2.6	Statistical Analysis	91
5.3	Results and Discussion	91
5.3.1	Fatty Acid Compositions of Blends	91
5.3.2	Physico-chemical Properties of Blends	96
5.3.2.1	Iodine value	96
5.3.2.2	Slip Melting Point	99
5.3.2.3	Saponification Value	101

5.3.2.4	Acid Value	102
5.3.3	Correlation Between the Fatty Acid Constituents of all Blends and Their Properties by Multiple Regression Equation	103
5.4	Conclusions	107
CHAPTER 6 – CHARACTERIZATION OF CRYSTALLIZATION AND MELTING PROFILES OF BLENDS OF MANGO SEED KERNEL FAT AND PALM OIL MID-FRACTION AS COCOA BUTTER REPLACERS USING DIFFERENTIAL SCANNING CALORIMETRY AND PULSE NUCLEAR MAGNETIC RESONANCE		
6.1	Introduction	109
6.2	Materials and Methods	111
6.2.1	Materials	111
6.2.2	Determination of Triglycerides by HPLC	112
6.2.3	Determination of Solid Fat Content (SFC) by Pulsed Nuclear Magnetic Resonance (pNMR)	112
6.2.4	Analysis of Crystallization and Melting Characteristics by Differential Scanning Calorimetry (DSC)	113
6.2.5	Crystal Morphological Study by Polarized Light Microscopy	113
6.2.6	Statistical Analysis	114
6.3	Results and Discussion	114
6.3.1	Triglyceride Compositions	114
6.3.2	Effect of Blending on DSC Melting Thermograms	116
6.3.3	Effect of Blending on DSC Crystallization Thermograms	120
6.3.4	Solid Fat Content	124

6.3.5	Relationships between SFC, Fatty Acids, and Triglycerides of Blends of MSKF and POMF	126
6.3.6	Morphological Study	126
6.3.7	Correlation between the Triglycerides and SFC of the Blends Using Multiple Regression Equation	129
6.4	Conclusions	130
CHAPTER 7 - HARD COCOA BUTTER REPLACERS FROM MANGO SEED KERNEL FAT AND PALM STEARIN		
7.1	Introduction	132
7.2	Material and Methods	133
7.2.1	Blending of MSKF and PS	133
7.2.2	Determination of Triglycerides (TGs) by HPLC	134
7.2.3	Determination of Solid Fat Content (SFC) by pNMR	134
7.2.4	Analysis of crystallization and Melting Characteristics by DSC	134
7.2.5	Crystal Morphological Study by Polarized Light Microscopy	134
7.2.6	Statistical Analysis	134
7.3	Results and Discussion	135
7.3.1	Triglyceride Compositions	135
7.3.2	Blending Effect on DSC Melting Thermograms	137
7.3.3	Blending Effect on DSC Crystallization Thermograms	139
7.3.4	Solid Fat Content	142
7.3.5	Morphological Study	144
7.4	Conclusions	146

CHAPTER 8 - CONCLUSIONS AND RECOMMENDATIONS

8.1	Conclusions	147
8.2	Recommendations	149
	REFERENCES	150
	APPENDICES	167
	LIST OF PUBLICATIONS	184

LIST OF TABLES

		Page
Table 2.1	Production of cocoa beans in the world (thousands of tonnes).	12
Table 2.2	The major fatty acid composition (%) of natural cocoa butter produced from various countries.	15
Table 2.3	Chemical characteristics of commercial CB from different countries.	16
Table 2.4	Fatty acid constituents for blends 1-10 (referred to f-PKO-3 and f-PKO-4 as a blend component) and commercial CB.	21
Table 2.5	TG compositions of POMF, CB and CBEs produced by inter-esterification reaction.	24
Table 2.6	Fatty acid and TG compositions of hard butter from kokum fat and phulwara butter.	27
Table 2.7	Fatty acids of sal fat from different studies.	28
Table 2.8	Major fatty acid compositions of shea butter from different studies.	29
Table 2.9	Fatty acid compositions of mango seed kernel fat from different studies.	37
Table 2.10	Examples of substances used as supercritical solvents and its corresponding critical temperature and pressure.	41
Table 3.1	Total fats extracted from MSK using Soxhlet method.	60
Table 3.2	Total fats extracted from MSK using continuous and soaking techniques of SC-CO ₂ extraction.	62
Table 3.3	Fatty acid composition in fats extracted from various MSK using soaking technique of SC-CO ₂ extraction at 44.2 MPa, 70.2° C and 3.4 ml/min.	68
Table 3.4	Fatty acid composition in fats extracted from varieties MSK using Soxhlet method.	69
Table 3.5	Physico-chemical properties of MSKFs extracted using soaking technique of SC-CO ₂ extraction.	73

Table 4.1	Levels and ranges of independent variables used in the RSM design.	77
Table 4.2	Experimental and predicted MSKF yield (%) of CCD using SC-CO ₂ .	79
Table 4.3	Estimated regression coefficients, corresponding t and p-values and analysis of variance for the final regression model of MSKF yield.	80
Table 5.1	Fatty acid compositions (%) of SC-CO ₂ extracted MSKF, POMF, PS, and commercial CB.	89
Table 5.2	Blending ratios (%) of SC-CO ₂ extracted MSKF and POMF in 100 g (w/w).	90
Table 5.3	Blending ratios (%) of SC-CO ₂ extracted MSKF and PS in 100 g (w/w).	90
Table 5.4	Fatty acid compositions (%) of SC-CO ₂ extracted MSKF and POMF blends at various ratios 5 to 50% of POMF, and commercial CB.	92
Table 5.5	Fatty acid compositions (%) of SC-CO ₂ extracted MSKF and PS blends at various ratios 5 to 50% of PS, and commercial CB.	95
Table 5.6	IV, SMP, SPV, and AV determined (Referred to SC-CO ₂ extracted MSKF and POMF as a blending components according to the blending ratios in Table 5.2) for all blends and commercial CB.	97
Table 5.7	IV, SMP, SPV, and AV determined (referred to SC-CO ₂ extracted MSKF and PS as a blending components according to the blending ratios in Table 5.3) for all blends and commercial CB.	98
Table 5.8	Multiple regression coefficients of fatty acid constituents of MSKF: POMF blends, E is constant, and R ² is coefficient determination.	105
Table 5.9	Multiple regression coefficients of fatty acid constituents of MSKF: PS blends, E is constant, and R ² is coefficient determination.	105
Table 6.1	Melting characteristics of pure MSKF, POMF and their blends and commercial CB.	118
Table 6.2	Crystallization characteristics of pure MSKF, POMF and their	122

	blends and commercial CB.	
Table 6.3	Multiple regression coefficients of triglyceride profiles of MSKF: POMFblends, G is constant, and R^2 is coefficient determination.	129
Table 7.1	Melting characteristics of blends of MSKF and PS, and commercial CB.	137
Table 7.2	Crystallization characteristics of MSKF and PS blends, and commercial CB.	142

LIST OF FIGURES

	Page	
Figure 2.1	Flow diagram showing the usage of palm oil and palm kernel oil.	34
Figure 3.1	Flow chart of MSKF extraction and CBRs preparation.	48
Figure 3.2	Typical GC chromatogram of fatty acid methyl esters of Apple (a), Arumanis (b), Chokonan (c), Elephant (d), Sala (e), and Waterlily (f) mango seed kernel fats.	65
Figure 4.1	Generated response surface plot showing the effects of pressure and temperature on the fat yield at a fixed CO ₂ flow rate of 2.5 ml/min.	82
Figure 4.2	Generated response surface plot showing the effect of pressure and CO ₂ flow rate on the fat yield at a fixed temperature of 60 °C.	83
Figure 4.3	Generated response surface plot showing the effect of temperature and CO ₂ flow rate on the fat yield at a fixed pressure of 35 MPa.	84
Figure 5.1	Residual of experimental and calculated properties of blends (referred to Table 5.6) versus fatty acid constituents of all blends (referred to Table 5.4).	107
Figure 5.2	Residual of experimental and calculated properties of blends (referred to Table 5.7) versus fatty acid constituents of all blends (referred to Table 5.5).	107
Figure 6.1	Triglyceride compositions of blends of MSKF and POMF (area %, mean of three replications, standard deviation (±) ranges from 0.46 to 1.20).	115
Figure 6.2	DSC melting curves of pure POMF and MSKF.	117
Figure 6.3	DSC melting curves of blends of MSKF and POMF. Melting curves of blend 1 to 10 represent fat blend containing 50 to 95% of MSKF, respectively.	120
Figure 6.4	DSC crystallization curves of pure POMF and MSKF.	121
Figure 6.5	DSC crystallization curves of blends of MSKF and POMF. Crystallization curves of blend 1 to 10 represent fat blend containing 50 to 95% of MSKF, respectively.	123

Figure 6.6	Solid Fat Content (%), each blends represents the average of triplicates and standard deviation (\pm) ranges from 0.84 to 1.75) of Blends 1 to 10 of MSKF and POMF.	125
Figure 6.7	Polarized light microphotographs (40X lens) of blends of MSKF and POMF crystallized at 21-23 °C for 48 h. (a) 85%, (b) 80%, (c) 75%, and (d) 70% MSKF.	128
Figure 6.8	Residual of experimental and calculated SFC (%) of blends versus blend number.	130
Figure 7.1	Triglyceride compositions of MSF and PS blends (area %, mean of three replications, standard deviation (\pm) ranges from 0.51 to 1.40).	136
Figure 7.2	DSC melting curves of blends of MSKF and PS. melting curves of blend 1 to 10 represent fat blend containing 50 to 95% of MSKF, respectively.	138
Figure 7.3	DSC crystallization curves of blends of MSKF and PS. crystallization curves of blend 1 to 10 represent fat blend containing 50 to 95% of MSKF, respectively.	141
Figure 7.4	Solid fat content (%), each blend represents the average of triplicates and standard deviation (\pm) ranges from 0.77 to 1.55) of blends 1 to 10 of MSF and PS.	144
Figure 7.5	Polarized light microphotographs (40X lens) of blends of MSKF and PS crystallized at 21-23 °C for 48 h. (a) 95%, (b) 90%, (c) 85%, (d) 80% MSKF.	145

LIST OF ABBREVIATIONS AND SYMBOLS

ANOVA	Analysis of variance
AR	Analytical reagent
AV	Acid value
CB	Cocoa butter
CBE	Cocoa butter equivalent
CBEXs	Cocoa butter extenders
CBIs	Cocoa butter improvers
CBR	Cocoa butter replacer
CBS	Cocoa butter substitute
C14	Myristic acid
C15	Pentadecanoic acid
C16	Palmitic acid
C16:1	Palmitoleic acid
C17	Heptadecanoic acid
C18:0	Stearic acid
C18:1	Oleic acid
C18:2	Linoleic acid
C18:3	Linolenic acid
C20	Arachidic acid
C20:1	11-eicosenoic acid
C22	Behenic acid
C24	Lignoceric acid
°C	Degree celsius

DOE	Design of experiment
DSC	Differential scanning calorimetry
FAME	Fatty acid methyl ester
FID	Flame ionization detector
GC	Gas chromatography
GC/MS	Gas chromatography/mass spectrometry
HPLC	High performance liquid chromatography
IV	Iodine value
MPa	Mega pascal
MSKF	Mango seed kernel fat
NMR	Nuclear magnetic resonance
PKO	Palm kernel oil
PLM	Plan polarized light microscopy
POMF	Palm oil mid-fraction
POP	1,3-dipalmitoyl-2-oleoyl-glycerol
PORIM	Palm oil research institute malaysia
POS	1-palmitoyl-3-stearoyl-2-oleoyl-glycerol
PS	Palm stearin
RSM	Response surface methodology
SC-CO ₂	Supercritical carbon dioxide
SFC	Solid fat content
SMP	Slip/sharp melting point
SOS	1,3-distearoyl-2-oleoyl-glycerol
SV	Saponification value
TG	Triglyceride

LIST OF APPENDICES

	Page	
Appendix A.1	Mango fruits and mango seed kernel powder used in this study.	168
Appendix A.2	MSKFs extracted from different varieties using soaking technique of SC-CO ₂ extraction at 35 MPa, 72° C and 3.4 ml/min.	169
Appendix A.3	SFE instrument used in this study	170
Appendix A.4	Schematic diagram of SC-CO ₂ extraction apparatus	170
Appendix B	Contour plots of oil yield (%) vs pressure (MPa), Temperature (°C) and flow rate (ml/min).	171
Appendix C.1:	Blends of MSKF and POMF at various ratios 5 to 50% of POMF.	172
Appendix C.2:	Blends of MSKF and PS at various ratios 5 to 50% of POMF.	172
Appendix C.3:	Recommended blends of MSKF and POMF containing 65 to 90% MSKF	173
Appendix C.4:	Recommended blends of MSKF and PS containing 65 to 90% MSKF.	173
Appendix D.1:	GC chromatogram of fatty acid methyl esters of MSKF and POMF blends at various ratios 5 to 50% of POMF.	177
Appendix D.2:	GC chromatogram of fatty acid methyl esters of MSKF and PS blends (1 to 10) at various ratios 5 to 50% of PS.	180
Appendix E.1:	Typical HPLC chromatogram of MSKF and POMF blends identifying the TG profiles by comparing the elution times (in minute) of standard.	181
Appendix E.2:	Typical HPLC chromatogram of MSKF and PS blends identifying the TG profiles by comparing the elution times (in minute) of standard.	182
Appendix F	Polarized light microphotographs (40X lens) of blends of MSKF and POMF crystallized at 21-23 °C for 48 h. (Blend 1) 5% POMF, (Blend 2) 10% POMF, (Blend 7) 35% POMF, (Blend 8) 40% POMF, and (Blend 9) 45% POMF and (Blend 10) 50% POMF.	183

**PENGEKSTRAKAN LEMAK ISIRONG BIJI MANGGA MENGGUNAKAN
KARBON DIOKSIDA LAMPAU GENTING SEBATI DENGAN MINYAK
SAWIT PECAHAN PERTENGAHAN DAN STEARIN PALMA UNTUK
MEMBENTUK PENGANTI MENTEGA KOKO**

ABSTRAK

Mangga (*Mangifera tindan* L.) ialah sejenis buah tropika tahunan yang penting. Isirung biji mangga adalah hasil sampingan perindustrian yang mengandungi mentega koko hampir serupa dengan lemak. Pada masa ini tidak terdapat laporan atas pengekstrakan lemak isirung biji mangga menggunakan karbon dioksida lampau genting (BLG-CO₂). Tujuan kajian ini adalah untuk mencampurkan ekstrak lemak isirung biji mangga (LIBM) dengan fraksi pertengahan minyak sawit (FPMS) dan stearin sawit (SS) untuk menjanakan rumusan pengganti mentega koko (PMK). Ciri fiziko kimia, termal, kandungan lemak pepejal dan morfologi campuran LIBM: FPMS dan LIBM:SS dijalankan dengan menggunakan teknik kromatografi dan termal berbeza. Pengoptimuman parameter pengekstrakan SC-CO₂ lemak isirung biji mangga (LIBM) dari isirung biji mangga (IBM) dijalankan menggunakan kaedah permukaan gerak balas rekabentuk komposit berpusat. Pembolehubah yang dipertimbangkan dalam kajian itu ialah tekanan (20-50 MPa), suhu (40-80 °C), dan kadar aliran CO₂ (1-4 ml/min). Hasil lemak yang dioptimumkan telah diramalkan sebanyak 11.29% pada 44.2 MPa, suhu 72.2 °C dan kadar aliran CO₂, 3.4 ml/min yang nilai hampir sama dengan hasil lemak pengekstrakan dengan Soxhlet. Campuran yang mengandungi 70 hingga 85% MSKF mempunyai sifat fiziko-kimia, profil asid lemak, nilai iodin (IV), takat lebur 'slip' (SMP), nilai saponifikasi (SPV) dan nilai asid (AV) yang hampir sama dengan

mentega koko komersial. Hasil kajian menunjukkan bahawa julat kandungan trigliserida (TG) utama bagi semua campuran adalah POP: 11-38.8%, SOS: 22.1-36.9%, dan POS: 15.4-16.2%,. Ciri daripada pencairan menunjukkan satu lengkung tunggal dengan hanya satu titik maksima dan satu bahu kecil untuk campuran no. 3 hingga 6 (LIBM: FPMS) dan 1 hingga 4 (LIBM: SS). Campuran sama mempunyai 70 hingga 95% LIBM yang menunjukkan corak penghabluran serupa dengan satu lengkung tunggal yang mempunyai satu puncak maksimum pada suhu-suhu 10.17, 10.58, 11.54 , dan 11.66 °C. Kandungan lemak pepejal (KLP) campuran no. 1 hingga 5 didapati hamper sama dengan mentega koko (MK) komersial pada suhu 10 hingga 20 °C. Pada 35 °C, KLP kesemua campuran mempamerkan ciri seperti MK. Mikroskop cahaya terkutub digunakan untuk memantau mikrostruktur rangkaian hablur semua campuran itu. Mikrostruktur hablur campuran no. 3 hingga 6 dan 1 hingga 4 ialah sferulit yang terdiri daripada jarum seperti hablur bersinar dan bercabang menuju ke luar daripada teras pusat, serupa dengan MK. Kajian mendedahkan bahawa PMK yang berciri mesra alam boleh disediakan dengan pengadunan LIBM dengan FPMS tempatan and SS dan boleh digunakan secara langsung dalam produk coklat.

SUPERCRITICAL CARBON DIOXIDE EXTRACTION OF MANGO SEED KERNEL FAT BLENDED WITH PALM OIL MID-FRACTION AND PALM STEARIN TO FORMULATE COCOA BUTTER REPLACERS

ABSTRACT

Mango (*Mangifera indica* L.) is an important annual tropical fruit. Mango seed kernel (MSK) which is industrial by-product contains considerable amount of cocoa butter analogy fats. It is remarkable that no reports on the extraction of the mango seed kernel fat (MSKF) using supercritical carbon dioxide (SC-CO₂) have been published. The aim of this work is to blend SC-CO₂ extracted MSKF with palm oil mid-fraction (POMF) and palm stearin (PS) to formulate new cocoa butter replacers (CBRs). The physico-chemical properties, thermal properties, solid fat content (SFC) and morphology for the blends of MSKF: POMF and MSKF: PS conducted using different chromatographic and thermal techniques. Optimization of the SC-CO₂ extraction parameters of MSKF from MSK were conducted using central composite design (CCD) of response surface methodology (RSM). The variables considered in the study are pressure (20-50 MPa), temperature (40-80 °C), and CO₂ flow rate (1-4 ml/min). The optimized fat yield was predicted to be 11.29% at 44.2 MPa, 72.2 °C and CO₂ flow rate of 3.4 ml/min which was close to the fat yield (11.7%) of Soxhlet extraction. The blends containing 70 to 85% of MSKF had physico-chemical properties like fatty acid profiles, iodine value (IV), slip melting point (SMP), saponification value (SPV) and acid value (AV) close to that of commercial CB. Results showed that the major TG ranges in all blends were POP 11-38.8%, SOS 22.1-36.9%, and POS 15.4-16.2%, respectively. The melting

behaviour indicated a single curve with only one maximum and one small shoulder for blends no. 3 to 6 (MSKF: POMF) and 1 to 4 (MSKF: PS). The same blends having 70 to 95% of MSKF indicated similar crystallization pattern with a single curve having one maximum peak heights at 10.17, 10.58, 11.54, and 11.66 °C temperatures. The SFC of blends no. 1 to 5 was found to be close to that of commercial CB at temperatures 10 to 20 °C. At 35 °C the SFC of all blends were depicting that of CB. Polarized light microscopy was used to monitor microstructure of crystal network of the all blends. The microstructure of the crystals of blends no. 3 to 6 and 1 to 4 was spherulites consisting of needle like crystals radiating and branching outward from the central nuclei, similar to that of CB. The studies revealed that environmental friendly CBRs could be prepared by blending MSKF with local POMF and PS and could be used directly in chocolate products.

CHAPTER 1

INTRODUCTION

1.1 Research background

Mango (*Mangifera indica* L.) fruit belongs to the family of *Anacardiaceae*. It is the most important popular seasonal tropical fruit around the world and in some countries it is known as the “king of fruits”. As it is a seasonal fruit, about 20% of mango fruits are processed for the purpose of making puree, leather, pickles, canned slices, jam, jelly, and chutney (Loelillet, 1994). Moreover, mango juice is the predominant fruit juice especially in tropical and subtropical areas. It has been well acknowledged that mango fruit and its processed products have gained popularity in both the national and international market due to their nutritional quality, delicious, succulence, sweet taste, and delicate flavor. In the industry, only large edible portion of mangoes are used. Consequently, considerable amounts of mango seeds are discarded as agricultural wastes which are produced either by industrial processing or after consumption of the fruit, amounting 35 to 60% of the total fruit weight (Puravankara et al., 2000). Depending on the mango varieties, the seed represents from 10 to 25% of the weight of the total fruit and the kernel inside the seed represents from 45 to 85% of the seed and about 20% of the whole fruit (Hemavathy et al., 1988; Arogba, 1997; Solís-Fuentes and Durán-de-Bazúa, 2011).

Mango seed kernels (MSK) contain considerable amounts of cocoa butter analogy fats. On a dry basis, MSK contains about 7.1 to 15% crude fat (Ali et al., 1985; Hemavathy et al., 1988; Abdalla et al., 2007; Gunstone, 2011). It is the rich

sources of palmitic (C_{16:0}), stearic (C_{18:0}) and oleic acids (C_{18:1}) containing about 3 to 18%, 24 to 57% and 34 to 56%. The other major fatty acids are linoleic, C_{18:2} (up to 13%), linolenic, C_{18:3} (0.6 to 1.0%), arachidic, C_{20:0} (1.7 to 2.6%), and myristic, C₁₄ (0.5 to 8%) acids. This group of fats received considerable attention for their particular physical and chemical properties which make them very suitable for the fabrication of confectionary. Therefore, the lipid composition of MSK has drawn immense research interest because of their potential application in the confectionery industry as a source of cocoa butter alternatives (Lakshminarayana et al., 1983; Ali et al., 1985; Hemavathy et al., 1988; Muchiri et al., 2012).

Currently, mango seed kernel fats (MSKF) are extracted and fractionated using solvents and their quality evaluated by many researchers in the literature (Gaydou and Bouchet, 1984; Ali et al., 1985; Abdalla et al., 2007). These organic solvent has adverse effects and it is non-selective, costly and requires long extraction time. Therefore, it is essential to develop improved method for MSKF extraction. Moreover, greater concern over the disposal of such toxic organic solvents and their effect on the human health as well as on the environment has led to a move towards cleaner extraction methods such as SC-CO₂ extraction (Staby and Mollerup, 1993; Hultin, 1994). Until now, no studies have been conducted for the extraction of MSKF using SC-CO₂.

Food industries are keen to find for alternative fats to cocoa butter (CB) from natural matrices that are denoted as cocoa butter replacers (CBRs), cocoa butter equivalents (CBEs), and cocoa butter substitutes (CBSs) fat. CB is a natural fat obtained from cocoa seeds (*Theobroma cacao*). It is commonly used as an essential

major ingredient of chocolate and other confectionary products due to its specific physical and chemical properties. CB is solid at room temperature (below 25 °C) and at body temperature (~37 °C) it is liquid. CB mainly consists of C₁₆, C_{18:0}, C_{18:1}, and C_{18:2} acids but low amount of C₁₂ and C₁₄ acids. CB can crystallize into several polymorphic forms, having α , γ , β' , and β crystals, with melting points of 17, 23, 26, and 35–37 °C, respectively. In the chocolate production, only β crystal is used because it has a high melting point. This crystal structure confers chocolate products an excellent quality in terms of sheen, snap, and smooth texture. In addition, CB exhibits resistance to fat bloom, arising from changes in the fat in the chocolate during storage. This can be seen as undesirable white or streaky grey-white spots on the chocolate surface.

CB is highly appreciated and is expensive compared with all other vegetable fats and oils because of its specific characteristics. Another important reason is that cocoa beans contain low amount of CB (Zaidul et al., 2007a). Moreover, cocoa has only been cultivated in a few countries (Moreton, 1988; Hassan et al., 1995). Therefore, many researchers have produced cocoa butter alternative fats either by fractionation and blending or enzymatic interesterification of palm kernel oil (PKO) and palm oil (PO) (Bloomer et al., 1990; Hashimoto et al., 2001; Undurraga et al., 2001; Calliauw et al., 2005; Zaidul et al., 2007a), mango seed fat (Lakshminarayana et al., 1983; Ali et al., 1985; Jiménez- Bermúdez et al., 1995; Solís-Fuentes, 1998), kokum butter (Reddy and Prabhakar, 1994; Maheshwari and Reddy, 2005), sal fat (Reddy and Prabhakar, 1989; Gunstone, 2011), shea butter (Olajide et al., 2000), and illipé fat (Gunstone, 2011).

1.2 Problem statement

CB is the most important class of fats with demonstrated health benefits, and is used entirely for making chocolate and other confectionery products (Wollgast and Anklam, 2000; Kim et al., 2011). Moreover, it is a highly demanded fat among tropical plant fats, but its world production is decreasing by the day due to the cultivation difficulties, weather event, low productivity, and pest attacks (Bootello et al., 2012). As only a few countries cultivate cocoa, supply can be unstable. In recent years, cocoa price is increasing due to increasing consumption of chocolate products that contain higher level of CB (Afoakwa, 2010). Therefore, the food industries are in need for CB alternatives that might have similar quality of natural CB to meet the demand of premium grade and relatively cheaper CB overcoming the uncertainty in the supply of original CB due to its less existibility in the cocoa beans. Thus, interests of CB analogy fats from various natural sources are increasing.

Like CB, MSKF is a natural fat containing high saturated and monounsaturated fatty acids. These fatty acid constituents and other physico-chemical properties of MSKF make it a valuable fat that is comparable to that of commercial CB (Muchiri et al., 2012). In the industry, only large edible portion of mangoes are used or consumed. As a result considerable amounts of seeds are discarded as industrial waste or raised as by-products. These by-products particularly come from the tropical or subtropical fruit processing industries which have adverse effects on the environments, human health as well as creating both disposal and pollution problems. Due to increasing popularity and production of mangoes worldwide, disposal amounting 35 to 60% of the fruit weight represents a growing problem as the plant material is prone to microbial spoilage (Larrauri et al., 1996).

Moreover, drying cost, storage, and shipment of these by- products are economically limiting factors (Schieber et al., 2001).

Traditional solvent extraction is the current method for the extraction of MSKF (Lakshminarayana et al., 1983; Ali et al., 1985; Solis-Fuentes and Durán-de-Bazúa, 2004; Abdalla et al., 2007). In many countries, health and safety regulations are getting stricter in addressing environmental problems created by the use of organic solvents and these issues are forcing the industries to search for alternative processing methods. The solvents are unsafe to handle and unacceptable as it is harmful to human health and the environment, restricting its use in the food, cosmetic and pharmaceutical industries (Liu et al., 2009a). Moreover, there is an increasing concern of the health and safety hazards associated with the use of organic solvents often introduces contaminants into the MSKF that must be removed later. The major demerits of solvent extraction methods are (a) they require high temperature that affects nutritional quality of the products, (b) they are heat sensitive labile natural compounds, and (c) they contain toxic solvent left in the final products which have diverse adverse human health effects (Staby and Mollerup, 1993; Hultin, 1994).

1.3 Objectives of Study

The objectives of the study are:

1. To extract mango seed kernel fat from different varieties of mango seed kernel using conventional and SC-CO₂ methods for screening cocoa butter analogy fats.

2. To optimize the SC-CO₂ extraction parameters extracting CB analogy fat from MSK using response surface methodology (RSM).
3. To formulate CBRs from SC-CO₂ extracted MSKF blended with palm oil mid-fraction (POMF) and palm stearin (PS).
4. To characterize the physico-chemical, thermal properties and solid fat content for the blends of MSKF: POMF and MSKF: PS as CBRs using DSC and pNMR.

1.4 Research Contributions

This work does indeed offer some significant contributions to the current body of knowledge on research involving SC-CO₂ extraction and CBRs formulation. The physico-chemical properties of the SC-CO₂ extracted MSKF are ideal for use in CBRs formulation in food industry in particular chocolate industry. According to the Malaysian Palm Oil Council (MPOC), POMF and PS are obtained by fractionation of palm oil. PS is always traded at a discount compared to any other palm oil products; making it a cost effective ingredient for various applications. Blending of MSKF with POMF and PS can lower the costs comparative to commercial CB, while having the desired taste of CB. SC-CO₂ extraction and direct blending processes have the additional advantage that no chemical process is involved, consistent with the consumer trend toward natural and chemical free products.

The results of this study suggest that certain blends of MSKF and PS could be used to prepare CBRs without altering the physical and chemical properties of the product significantly. These blends could be used for the production of high temperature resistant hard butter for countries with hot climate. Moreover, blends of

MSKF and PS could solve the tempering difficulties faced by chocolate manufacturers in tropical countries like Malaysia or in countries with a moderate climate during the summer season. This blending method is a feasible alternative to expand the use of Malaysian POMF, and PS through the development of new CBRs.

1.5 Scope of Study

The following scopes of the study were performed in order to achieve the objectives:

- Examination of the effect of continuous and soaking techniques of SC-CO₂ extraction on mango seed kernel fat yield.
- Examination of the effect of SC-CO₂ extraction parameters on mango seed kernel fat yield.
- Comparison the fatty acids content in mango seed kernel fat obtained by SC-CO₂ and Soxhlet methods.
- Determination of the physico-chemical properties such as fatty acid constituents, iodine value, saponification vale, slip/sharp melting point, and acid value of SC-CO₂ extracted mango seed kernel fat.
- Comparison of the physico-chemical properties of SC-CO₂ extracted mango seed kernel fat with that of commercial CB.
- Preparation of cocoa butter replacers by blending of SC-CO₂ extracted MSKF with POMF and PS.
- Determination of the physico-chemical properties such as fatty acid and triglyceride compositions, iodine value, saponification value, acid value, slip/sharp melting point, solid fat content, and melting and crystallization profiles of MSKF: POMF and MSKF: PS blends.

- Examination of the effect of blending on physico-chemical properties in the formulated blends.
- Comparison the physico-chemical properties of all blend with that of commercial CB.

1.6 Limitation of Study

As discussed in earlier sections, SC-CO₂ could be advantageous for extracting of MSK fat and be blended again in desired ratios to formulate cocoa butter alternative fats according to the market demand. Supercritical CO₂ being a cheaper recyclable solvent and it will not be costly to produce the desired products and bear no threat to health and environment. The major hurdle to the application of SC-CO₂ in fats and oils industry is its instrumentation to an industrial scale. Another problem is the maintenance of the SC-CO₂ equipment which is somewhat costly as it goes frequent gas-leakage during high pressure operation. Attention should be paid while constructing the heavy duty equipment for use in the industry. It is also important to keep in mind during construction of the equipment to link a supercritical fluid chromatography to SC-CO₂ equipment to be used in special purpose to concentrate desired components. By this time SC-CO₂ has been established to industrial scale in essential oil extraction from different spices and in extraction of some pharmaceutical or phytochemicals from different plant parts. Once the industry adopts SC-CO₂ equipment in fats and oils extraction, the process can appear cost-effective or more profitable than the conventional methods.

1.7 Overview of Thesis

This thesis consists of eight chapters which begins with the introduction as the first chapter. The introduction describes in detail the background of the research. In this chapter, the problem statement, research objectives, research contributions, scope and limitation of the study have been identified and described. Chapter 2, covers the literature review regarding the application of SC-CO₂ in the extraction of fats and oils from various plant sources presented in detail to further understand the principles of this research. Blending of fats and oils to obtain CBRs is also reviewed as well as appraisal on MSKF, CB and cocoa butter alternative fats. In Chapter 3, screening of cocoa butter analogy fats from different varieties of MSK are presented and discussed. In Chapter 4 presents and discusses the optimization of the SC-CO₂ extraction parameters of CB analogy fat from MSK using RSM are presented and discussed. In Chapter 5, SC-CO₂ extracted MSKF blended with palm oil mid-fraction and palm stearin and investigated their physico-chemical properties. In Chapter 6 and 7, the thermal properties and SFC of MSKF: POMF and MSKF: PS blends are discussed. The conclusion of this research and some recommendations for the future work are addressed in Chapter 8.

CHAPTER 2

LITERATURE REVIEW

2.1 Introduction

In this chapter, literature reviews regarding the application of SC-CO₂ in the extraction of fats and oils from various plant sources are presented. Blending of those fats and oils to obtain CBRs are also reviewed. Moreover, reviews on mango seed kernel fat (MSKF), cocoa butter and cocoa butter alternative fats are summarised.

2.2 Cocoa Butter

2.2.1 Background of Cocoa

The generic name of cocoa is *Theobroma* belonging to the family of Sterculiaceae, also called “Food of God”. It contains about 30 to 50 beans, covered with pulp. About 500 years ago, cocoa beans were originated from Latin America, and within a few years it spread to Europe. From there it was then distributed throughout the world (ICCO, 2000).

2.2.2 Cocoa Production

The major cocoa beans growing countries in the world are Ivory Coast, Ghana, Indonesia, Cameroon, Nigeria, Brazil, Ecuador, Dominican Republic, and Malaysia, contributing almost 90% of world production (ICCO, 2009/2010; FAO, 2012). The world total cocoa beans productions in season 2007/08 to 2009/10 are shown in Table 2.1. The Global cocoa beans production declined at 3.613 million

tonnes in 2009/2010 season, while in 2007/2008 it was 3.752 million tonnes (Table 2.1). In season 2007/2008 to 2009/2010, the cocoa beans production declined by 3% in Africa, while it increased by 1.9% and 1.8% in the Americas and in Asia and Oceania to 14.4% and 17.5%, respectively. Africa is still the largest cocoa producing region, contributing 68.0% of the total world production followed by Asia and Oceania and the Americas in 2009/2010.

In Malaysia, cocoa beans production has also declined gradually. According to the Malaysian Cocoa Board, cocoa beans production in 2011 was 15000 tons, while in 2007 it was 35180 tons (MCB, 2011). The International Cocoa Organization (2009/2010) reported that weather has a great impact on the cocoa beans production. de Magalhães et al. (2011) reported that about 71% of the cocoa consumed around the world comes from the Africa's, especially from Ivory coast, Ghana and Nigeria. In the last decade, the consumption of cocoa has increased while its production has been declining day by day (Jonfia-Essien et al., 2008).

Table 2.1: Production of cocoa beans in the world (thousands of tonnes)

	2007/2008		2008/2009		2009/2010	
Africa	2693	71.8%	2518	69.9%	2458	68.0%
Ivory Coast	1382		1222		1242	
Ghana	729		662		632	
Nigeria	230		250		240	
Cameroon	185		227		190	
Others	166		158		154	
America	469	12.5%	488	13.5%	522	14.4%
Brazil	171		157		161	
Ecuador	118		134		160	
Others	180		197		201	
Asia and Oceania	591	15.8%	599	16.6%	633	17.5%
Indonesia	485		490		535	
Papua New Guinea	52		59		50	
Others	55		50		48	
World Total	3752	100.0%	3605	100.0%	3613	100.0%

Source: ICCO Quarterly Bulletin of Cocoa Statistics, Vol. xxxvi, No. 4, Cocoa year 2009/2010.

2.2.3 Cocoa Butter Extraction

The cocoa seeds are referred to as cocoa beans consist about 85% cotyledon (nib) and 15% shell. The nibs contain about 55% fat. The nibs ground to a paste are called cocoa liquor or mass, some are directly used in chocolate. Cocoa beans are natural oil seeds like palm kernel, groundnut, sesame seed, or any other oil seeds. Processing of cocoa beans for getting the fats or oils is not same like other oilseed due to the unique physico-chemical properties of cocoa beans, especially the fats (Adeyeye et al., 2010). Presently, several methods are employed for the extraction of CB from mass or liquor, or from other sources including hydraulic press, mechanical press, screw presses, SC-CO₂, and solvent extractions (Asep et al., 2008; Nair, 2010). Recently, Nair (2010) reported that the hydraulic press and screw presses are not successful methods for the extraction of CB. The major demerits of hydraulic press, mechanical press, screw presses, and solvent extractions are a) they require high

temperature that affects nutritional quality of the CB, b) they are heat sensitive labile natural compounds, and c) they contain toxic solvent left in the final products which have diverse adverse human health effects (Staby and Mollerup, 1993; Hultin, 1994).

Solvent extraction with hexane has been widely used to extract the CB as well as fats and oils from the oil-contained sources. However, there is an increasing concern of the health and safety hazards associated with the use of organic solvents, while expression by hydraulic method often introduces contaminants into the CB that must be removed later.

Meanwhile, several studies have been carried out for the extraction of CB from cocoa liquor and shell by SC-CO₂ at different temperatures and pressures (Li and Hartland, 1992; McHugh and Krukonis, 1994; Li and Hartland, 1996; Rossi, 1996; Asep et al., 2008). The studied pressure and temperature ranges from 15 to 40 MPa and 40 to 80 °C. The yield of CB extracted with SC-CO₂ at 30-40 MPa and 50-80 °C depend on the degree of disruption of lipid bearing cells reported by McHugh and Krukonis (1994), Rossi (1996). At higher pressure, the yield of CB was reported to be higher and temperature range being studied. Moreover, in their study, the triglycerides in terms of fatty acid compositions of extracted CB were within the required range and retained the aroma of the residue. Recently, Li and Hartland (1996), Rossi (1996), Asep et al. (2008), and Jinap et al. (2013) studied the effect of co-solvent such as ethanol on SC-CO₂ extraction of CB. Their results showed that addition of co-solvent improve the efficiency of the CB extraction. By adding co-solvent such as ethanol (20-25%, w/w), the solubility of CB increased and maximum

yield was also reported by Li and Hartland (1992), Asep et al. (2008), and Jinap et al. (2013).

2.2.4 Composition of Natural Cocoa Butter

CB obtained from cocoa beans and on dry weight basis accounts for 50 to 57% and is responsible for the melting properties of chocolate (Steinberg et al., 2003). The major fatty acids of CB are C₁₆ (25 to 33.7%), C_{18:0} (33.7 to 40.2%), C_{18:1} (26.3 to 35%) and C_{18:2} (1.7 to 3%) which contribute about 98% of the total fatty acids (Bracco, 1994; Asep et al., 2008). The fatty acid compositions of CB differ depending on the country of origin. The key fatty acid compositions of CB from different countries are shown in Table 2.2. 1,3-dipalmitoyl-2-oleoylglycerol (POP), 1-palmitoyl-2-oleoyl-3-stearoyl-glycerol (POS) and 1,3-distearoyl-2-oleoyl-glycerol (SOS) are the three main triglycerides (TGs) account for 92 to 96% of total lipid composition of CB (Lehrian and Keeney, 1980; D'Alonzo et al., 1982; Davis and Dimick, 1989; Lipp and Anklam, 1998; Lipp et al., 2001; Asep et al., 2008). Among these three TGs, POS is the major leading TG component present in CB with range 42.5 to 46.4% yield followed by SOS (27.8 to 33.0%) and POP (18.9 to 22.6%) (Asep et al., 2008).

Table 2.2: The major fatty acid composition (%) of natural cocoa butter produced from various countries.

Country	Fatty acid (%)				References
	C _{18:0}	C _{18:1}	C ₁₆	C _{18:2}	
Ivory Coast	36.9	32.9-33	25.8-26.6	2.6-2.8	Davis and Dimick (1989); Lipp and Anklam (1998)
Ghana	36.69-37.6	32.7-32.99	25.3-25.46	2.51-2.8	Spangenberg and Dionisi (2001); Lipp and Anklam (1998)
Indonesia	36.88-37.3	33.06-34.3	24.1-25.13	2.5-2.7	Spangenberg and Dionisi (2001); Lipp and Anklam (1998)
Brazil	33.3-33.8	34.5-36.5	25.1-27.9	3.5-3.6	Lehrian and Keeney (1980); Lipp and Anklam (1998)
Ecuador	34.62-36	34.6-34.91	25.2-25.6	2.6-3.04	Spangenberg and Dionisi (2001); Lipp and Anklam (1998)
Malaysia	36-37.4	33.5-34	24.9-26	2.6-3.0	Kheiri (1982); Lipp and Anklam (1998)

2.2.5 Physico-chemical Properties of Natural Cocoa Butter

The physico-chemical properties of cocoa beans and the characterization of cocoa matrix are greatly influenced by processing method (Hoskin and Dimick, 1984, 1994; Jinap, 1994; Amin et al., 1997, 1998, 2002; Puziah et al., 1998). The properties of CB fat are the properties of the mixture of TGs. CB is commercially available fat that contains significantly higher amount of saturated acids, leading to TGs of POS, SOS, and POP. Due to such types of TGs in CB, it confers the short melting behavior which is appreciated by consumers. CB has relatively low and sharp melting point which ranges from 27 to 35 °C. At room temperature, CB is hard and brittle and its hardness depends on the solid fat content (SFC) and it might be 0% just above 37 °C temperature. Furthermore, the nature of crystalline lattice also renders the hardness/consistency of the CB (Kheiri, 1982).

The other physico-chemical properties of CB such as IV and SPV are shown in Table 2.3. Iodine values have great impact on oil quality; it indicates the degree of

unsaturation in the fats or oils. Typically, at room temperature fats are liquid because of the higher unsaturated fatty acid components, while it is solid at lower unsaturated fatty acid components. Chaiseri and Dimick (1989) reported that the higher IV contributes to the softness of CB. They also stated that higher IV content CB is softer than the lower IV content CB. The commercial CB from different countries shows different IV that ranges from 34.40 to 38.65.

The SPV indicates that the average chain length of fatty acid present in fat. If the SPV value of the fat is high, then the chain length of fatty acid will be shorter, and vice versa. The AV is defined as the weight in milligrams of potassium hydroxide necessary to neutralize the free fatty acid present in 1 g of fat and it is used to quantify the free fatty acids present in fats or oils. The cloud point (CP) is related to the unsaturation of oil, that is, the unsaturation of oil is higher, when its CP is low.

Table 2.3: Chemical characteristics of commercial CB from different countries.

Country	Iodine value	Saponification value
Bolivia	36.02	195.43
Brazil	37.46	195.07
Colombia	36.56	195.75
Ecuador	36.68	195.85
Peru	37.94	195.92
Costa Rica	36.64	195.27
Dominican Republic	36.72	194.92
Mexico	35.79	193.72
Panama	36.86	196.71
Ivory Coast	35.54	193.58
Nigeria	37.33	193.62
Malaysia	34.74	194.36

Source: Chaiseri and Dimick, 1989

2.2.6 Cocoa Butter Replacer (CBR)

The CBR or cocoa butter alternatives are defined as non-lauric fats that could replace CB either partially or completely in the chocolate or other food products (Kheiri, 1982). The fatty acid compositions of CBR are similar to that of commercial CB with more or less similar TGs structure. It should be cheaper than that of commercial CB. CBR can be divided into the following groups.

2.2.6.1 Cocoa Butter Equivalent (CBE)

CBEs are vegetable fats which have similar physical and chemical characteristics like CB. Therefore, CBEs can be mixed with CB in any amount without changing the behavior of the final product. The major fatty acids contained in CBEs are C₁₆, C_{18:0}, and C_{18:1} acids, which are similar to that of CB. CBEs are divided into two subgroups, namely cocoa butter extenders (CBEXs) and cocoa butter improvers (CBIs) (Lipp and Anklam, 1998). CBEXs cannot be mixed with CB in every proportion, while CBIs are similar to CBEs, contain higher level of solid TGs, and because of this characteristic it is commonly used for improving soft CB.

2.2.6.2 Cocoa Butter Substitute (CBS)

CBSs are lauric and myristic plant fats (containing lauric and myristic acids) with some physical similarities to CB, but chemically they are completely different. Therefore, they are suitable for wholly replacement of CB.

2.2.7 Alternative Sources of Cocoa Butter

Many researches have carried out work for the production of CBRs, CBEs and CBSs from various natural sources. All of these are obtained from fats of natural

plant, such as PKO, PO, MSKF, kokum butter, sal fat, shea butter, illipé butter, soya oil, rape seed oil, cotton oil, ground nut oil, and coconut oil. Replacing the CB either partially or wholly with other natural fats has been investigated due to the technological and economical advantages. These methods include chemical or enzymatic fractionation or SC-CO₂ extraction of fats and their blends from various sources.

2.2.7.1 Cocoa Butter Equivalent (CBE) from Mango Seed Kernel Fat (MSKF)

Up till now, lots of research have been conducted for the extraction and fractionation of various MSKF as their lipid compositions as well as unique physical and chemical properties are similar to that of CB (Narashima-Char et al., 1977; Lakshminarayana et al., 1983; Ali et al., 1985; Dhinigra and Kapoor, 1985; Ali and Dimick, 1994; Jiménez-Bermúdez et al., 1995; Solís-Fuentes, 1998). Recently, Kaphueakngam et al. (2009) produced CBE by blending mango seed almond fat (MAF) with POMF. Seven blends of MAF-POMF with different ratios (100/0, 90/10, 80/20, 70/30, 60/40, 50/50, 40/60, 30/70, 20/80, 10/90 and 0/100) have been studied using various techniques. Their results showed that C₁₆, C_{18:0}, and C_{18:1} acids were predominant fatty acids in all blends, which were similar to commercial CB components. They reported that the major fatty acids, the melting behavior and slip melting point of the 80/40 (%wt) blend resemble to that of commercial CB.

In another study, Solís-Fuentes and Durán-de-Bazúa (2004) studied the thermal behaviour of MAF and its mixtures with CB. They reported that the fatty acid constituents and the physico-chemical properties of MAF are similar to that of commercial CB. They also showed that the MAF curves for solid–liquid phase

change and the solid fat content of MAF profiles are great similar to that of CB. The properties of MAF, thermal conduct and the presence of α and β crystalline forms in MAF made it a suitable fat like CB. The iso-solid diagrams showed the compatibility between the mixture of MAF and CB, even better than that of mixtures of CB with milk fat, lauric fats, or hydrogenated cottonseed oil (Solís-Fuentes and Durán-de-Bazúa, 2004).

2.2.7.2 Cocoa Butter from Palm Oil and Palm Kernel Oil (PKO)

The level of fatty acids such as C_{12} and C_{14} are present in CB as trace or very low amount while the amount of C_{16} , $C_{18:0}$, and $C_{18:1}$ acids are high. On the other hand, PKO contains a high level of C_{12} constituent but low in $C_{18:0}$ and $C_{18:1}$ constituents compared with commercial CB. Although PKO contains high level of C_{12} and C_{14} acids, it is widely used as a suitable raw material in confectionery (Pantzaris and Ahmad, 2001). To produce CBRs blend components, Zaidul et al. (2006) successfully fractionated the PKO to reduce its C_{12} content and increase $C_{18:0}$ and $C_{18:1}$ constituents using SC-CO₂. About 28% of C_{12} yield was reduced in fraction 4, while 31% of $C_{18:1}$ constituent increased in yield which is closer to the fatty acid composition of CB. The authors extracted PKO into four fractions and observed that at higher temperature (80°C) the total PKO yield increased with increasing pressures. At higher pressure of 48.3 MPa and temperature of 80°C, the highest yield reported 99.6%. They also reported that the highest level of short chain fatty acid such as C_8 , C_{10} and C_{12} constituents was found in fraction 1, while longer chain fatty acid such as C_{16} , $C_{18:0}$ and $C_{18:1}$ constituents were seen in fraction 4. The lower melting point was reported in fraction 4 than in other studied fractions. The authors fractionated

PKO to produce triglycerides in terms of fatty acid constituents that could be used in CBRs blend.

In another study, Zaidul et al. (2007a) produced CBRs by blending of SC-CO₂ extracted PKO fractions with conventionally extracted PO and commercial C_{18:0} and C_{18:1} constituents at various ratios. They also fractionated PKO into four fractions using SC-CO₂ and are thus called f-PKO-1, f-PKO-2, f-PKO-3 and f-PKO-4. To obtain CBRs, the authors blended the f-PKO-3 and f-PKO-4 fraction as blending components (denoted as low lauric and high oleic acid constituents) with PO and commercial fatty acids such as C_{18:0} and C_{18:1} constituents at different ratios and labeled as blends 1 to 10.

The fatty acid compositions of different blends of f-PKO-3, f-PKO-4, PO, commercial C_{18:0} and C_{18:1} constituents are shown in Table 2.4. In both blends (f-PKO-3 and f-PKO-4) 1 to 10 low levels of C₈ and C₁₀ were found while large amounts of the longer chain fatty acid such as C₁₆, C_{18:0}, C_{18:1} and C_{18:2} were reported which are closer to that of commercial CB. The authors reported that increasing the amount of f-PKO (f-PKO-3 and f-PKO-4), more than 50% increased shorter chain fatty acid and decreased longer chain fatty acid constituents. Moreover, their results showed that the physico-chemical properties like SMP, SFC, IV, SPV and AV of blends 2 to 10 were closer to that of commercial CB.

Table 2.4: Fatty acid constituents for blends 1-10 (referred to f-PKO-3 and f-PKO-4 as a blend component) and commercial CB. (Zaidul et al., 2007a).

f-PKO-3									f-PKO-4								
Blend number	Fatty acid constituents (%)								Blend number	Fatty acid constituents (%)							
	C ₈	C ₁₀	C ₁₂	C ₁₄	C ₁₆	C ₁₈	C _{18:1}	C _{18:2}		C ₈	C ₁₀	C ₁₂	C ₁₄	C ₁₆	C ₁₈	C _{18:1}	C _{18:2}
1	0.1	0.1	1.8	1.3	28.7	25.2	34.6	8.1	1	0.0	0.0	1.4	1.1	28.8	25.2	35.2	8.2
2	0.1	1.1	3.6	1.9	27.0	19.9	37.8	8.5	2	0.0	0.0	2.8	1.7	27.8	20.5	38.4	8.7
3	0.1	1.1	5.1	2.6	27.5	19.9	35.4	8.2	3	0.0	0.0	4.2	2.3	28.1	20.4	36.5	8.4
4	0.2	0.2	7.2	2.7	27.1	19.6	34.9	8.0	4	0.1	0.1	5.6	2.9	26.7	20.4	35.8	8.3
5	0.3	0.3	9.0	4.0	24.6	19.3	34.6	7.8	5	0.1	0.1	7.1	3.4	25.2	20.3	35.5	8.2
6	0.3	0.3	10.8	4.7	23.2	19.1	33.9	7.6	6	0.1	0.1	8.5	4.0	23.8	20.2	35.2	8.0
7	0.4	0.4	12.6	5.4	21.4	18.8	33.5	7.4	7	0.1	0.1	9.8	4.5	22.7	19.9	34.9	7.9
8	0.4	0.4	14.3	6.1	20.1	18.5	32.9	7.2	8	0.1	0.1	11.3	5.1	20.9	19.8	34.8	7.8
9	0.5	0.5	16.1	6.7	18.4	18.2	32.5	7.0	9	0.1	0.1	12.1	5.6	20.2	19.7	34.4	7.7
10	0.5	0.5	17.9	7.4	16.7	17.9	32.2	6.8	10	0.1	0.1	14.0	6.2	18.2	19.5	34.2	7.6
CB ^a	0.0	0.0	Trace	0.7	25.2	35.5	35.2	3.2	CB ^a	0.0	0.0	Trace	0.7	25.2	35.5	35.2	3.2

^aKheiri (1982) and Pease (1985)

Many other researchers have produced CBS from the fractionation of PKO in the literature (Hashimoto et al., 2001; Calliau et al., 2005). Calliau et al. (2005) developed a two-stage dry static fractionation method for the fractionation of PKO. They also studied single-stage static fractionation method. When both methods were compared, the major advantage of the two-stage method was that it produces a higher yield of good quality palm kernel stearin (PKS) and reduces the additional hydrogenation that would be advantageous for the production of CB like fat. By using this method, they produced CBS from PKO without hydrogenation. They also reported that about 30% of PKS produced by the two-stage process needs to be hydrogenated for use as CBS. Ali (1996) produced CBEs by blending of sal fat with co-fractionated PO. Their results showed that the TG composition and solidification characteristics were similar to that of the Malaysian CB. They also reported that the co-fractionation method increases the compatibility between CBE triglyceride components.

Due to the absence of any waxy taste, *trans* fatty acids and low level of linoleic acid in POMF, Samsudin and Rahim (1996) used it in white chocolate formulation. Two types of POMF (POMF I, a commercial sample and POMF II, from a laboratory-scale acetone fractionation of POMF I) and Malaysian CB were used in white chocolate formulation. They investigated the effect of the tempering process and bloom resistance of the produced white chocolate products. They found that the tempering time to produce a well-tempered chocolate using POMF I was longer than that using POMF II, while the time to produce a well-tempered CB chocolate increased with increasing tempering temperature.

Since 1980s the modification of fats through interesterification reactions catalyzed by enzyme has been widely studied. It is a method which has several advantages (lower energy consumption, absence of isomerization by products, and better control of products) over the conventional methods of the chemical interesterification (Undurraga et al., 2001). Many researchers produced CBEs through enzymatic interesterification of POMF in the literature (Bloomer et al., 1990; Undurraga et al., 2001). To produce CBEs, Bloomer et al. (1990) studied the interesterification reaction of POMF using lipase enzyme as catalysts. They studied the purity of the product and the influence of the solvent concentration on the reaction rate at different temperatures. Their results showed that the solvent concentration and temperature have great effect on the reaction rate. The optimum temperature was 40 °C and the solvent concentration was between 1 to 1.15 grams of solvent/gram of substrate reported in that study. They also showed that above this temperature the rate of interesterification reduced and the product purity decreased with increase of solvent.

In another study, Undurraga et al. (2001) produced CBEs through enzymatic interesterification of POMF with stearic acid in a solvent free system using Novo lipase Lipozyme™ as a catalyst. Their study performed in batch and in a continuous packed bed reactor. They investigated the effect of different parameters such as stearic acid-POMF ratio, enzyme-substrate ratio, and humidity of the enzyme preparation on productivity. Their results showed that the highest specific productivity obtained in shake flask was 0.0393 g/Batch Interesterification Unit (BIU) h at a stearic acid-POMF ratio of 1.6 and enzyme-substrate ratio of 23 BIU/g. On the other hand, the highest mass productivity observed was 1.54 g/g·h, using an

enzymic load of 73 BIU/g in the continuous packed bed reactor. They reported that the thermograms of their products obtained by differential scanning calorimetry (DSC) were similar to that of CB, but exhibited several distinct peaks due to the presence of diglycerides and trisaturated TGs. They also reported that POMF could be used as a suitable raw material for interesterification with stearic acid using Lipozyme™, leading to the production of CBEs whose composition resembles closely to that of CB. Table 2.5 shows the composition of POMF, pure CB, and CBEs which were produced in their study. In comparison with all extraction methods, SC-CO₂ is feasible in terms of quality product and has the potential to produce a higher yield and good quality CBRs blends.

Table 2.5: TG compositions of POMF, CB and CBEs produced by interesterification reaction.

Sample	POP	POS	SOS	Other TGs	DGs
PMF	74.3	14.3	2.0	9.4	-
CB	23.4	42.8	27.5	3.6	2.7
CBE	23.4	38.5	20.2	8.2	9.7

Source: Undurraga et al., 2001

2.2.7.3 Cocoa Butter Replacers from Kokum Butter

Kokum butter (*Garcinia indica* Choisy Syn *Brindonia indica*) belongs to the family of Guttiferae. It is found in the western peninsular coastal regions and the states of Maharashtra, Goa, Karnataka and Kerala India and parts of Eastern India in the states of West Bengal, Assam, and North Eastern Hill regions and other parts of peninsular India. It is a small evergreen tree and is also known as mangosteen, goa butter tree, kokum butter tree (Baliga et al., 2011). The mature tree yields fruits annually and takes five months to complete its fruiting process and by March to May