

Comparative analysis of bacterial communities associated with healthy and diseased corals in the Indonesian sea

by Agus Trianto

Submission date: 27-Jan-2020 09:47AM (UTC+0700)

Submission ID: 1246803518

File name: 2019-Peer_J_Karimunjawa_Coral.pdf (7.13M)

Word count: 8755

Character count: 48771

Comparative analysis of bacterial communities associated with healthy and diseased corals in the Indonesian sea

Wuttichai Mhuantong^{1,*}, Handung Nuryadi^{2,*}, Agus Trianto², Agus Sabdono², Sithichoke Tangphatsornruang³, Lily Eurwilaichitr¹, Pattanop Kanokratana¹ and Verawat Champreda¹

¹ Biorefinery and Bioproduct Technology Research Group, National Center for Genetic Engineering and Biotechnology, Pathum Thani, Thailand

² Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia

³ National Omics Center, National Center for Genetic Engineering and Biotechnology, Pathum Thani, Thailand

* These authors contributed equally to this work.

ABSTRACT

Coral reef ecosystems are impacted by climate change and human activities, such as increasing coastal development, overfishing, sewage and other pollutant discharge, and consequent eutrophication, which triggers increasing incidents of diseases and deterioration of corals worldwide. In this study, bacterial communities associated with four species of corals: *Acropora aspera*, *Acropora formosa*, *Cyphastrea* sp., and *Isopora* sp. in the healthy and disease stages with different diseases were compared using tagged 16S rRNA sequencing. In total, 59 bacterial phyla, 190 orders, and 307 genera were assigned in coral metagenomes where *Proteobacteria* and *Firmicutes* were predominated followed by *Bacteroidetes* together with *Actinobacteria*, *Fusobacteria*, and *Lentisphaerae* as minor taxa. Principal Coordinates Analysis (PCoA) showed separated clustering of bacterial diversity in healthy and infected groups for individual coral species. *Fusibacter* was found as the major bacterial genus across all corals. The lower number of *Fusibacter* was found in *A. aspera* infected with white band disease and *Isopora* sp. with white plaque disease, but marked increases of *Vibrio* and *Acrobacter*, respectively, were observed. This was in contrast to *A. formosa* infected by a black band and *Cyphastrea* sp. infected by yellow blotch diseases which showed an increasing abundance of *Fusibacter* but a decrease in WH1-8 bacteria. Overall, infection was shown to result in disturbance in the complexity and structure of the associated bacterial microbiomes which can be relevant to the pathogenicity of the microbes associated with infected corals.

Subjects Biodiversity, Microbiology

Keywords Bacterial diversity, Coral, Metagenome, Next generation sequencing, Pathogenicity, 16S rRNA

INTRODUCTION

Coral reefs contain important ecological space harboring a diversity of marine organisms and represents the core element of the complex marine ecosystem. Southeast Asia contains the largest area of coral reefs, accounting for 34% of the world's total (Wilkinson, 2008).

Submitted 15 March 2019
Accepted 1 November 2019
Published 19 December 2019

Corresponding author
Pattanop Kanokratana,
pattanopk@biotec.or.th

Academic editor
Linda Kelly

Additional Information and
Declarations can be found on
page 15

DOI 10.7717/peerj.8137

© Copyright
2019 Mhuantong et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

The reefs play crucial roles in ecology such as protecting the coastline from erosion, providing habitats for marine organisms, and supplying nutrients for the complex marine food chains. The coral reef ecosystem is estimated to provide USD 29.8 billion per year of economic benefit worldwide from various sectors, e.g., fisheries, tourism, coastal protection, and biodiversity (Cesar, Burke & Pet-Soede, 2003). From a scientific point of view, corals are also considered a rich source of unique and unexplored biosynthetic products (Radjasa et al., 2011). In this ecosystem, diverse symbioses exist with complex interactive dependences between corals and associated communities of eukaryotic and prokaryotic microorganisms (Blackall, Wilson & Van Oppen, 2015). For example, dinoflagellate endosymbionts, Symbiodiniaceae (LaJeunesse et al., 2018) utilizes light energy (Brodersen et al., 2014) and secrete fixed carbon to the coral host (Burriesci, Raab & Pringle, 2012). This symbiosis creates a highly complex and unique ecosystem.

In the last few decades, coral reefs have been facing world-wide crisis related to coral degradation driven by complicated phenomena including anthropogenic stresses and natural factors (Wilkinson, 2008). It is estimated that the world has collectively lost 19% of the original area of coral reefs while 15% are seriously endangered with expected damage within the next 10–20 years and 20% are under threat of loss in 20–40 years (Wilkinson, 2008). Degradation of coral reefs ecosystem can lead to massive loss of marine biodiversity. One of the serious threats to coral reefs ecosystem comes from various coral diseases (Harvell et al., 2007; Hughes et al., 2003). The occurrence of these diseases has been increasing dramatically during the last decade due to the rising sea-water temperature as a consequence of climate change (Rosenberg & Ben-Haim, 2002). Many of these diseases are prevalent in reef builder corals (Order Scleractinia), leading to the deterioration of entire reefs structures (Harvell et al., 2007; Rosenberg et al., 2007). Among them, black band, white band, white plague, white pox, brown band, red band, and yellow band syndromes are the main threats on corals in the Caribbean, Indo-Pacific, and Great Barrier Reef.

Coral diseases can be triggered by infection of pathogenic agents involving specific groups of bacteria, fungi, and viruses (Bourne et al., 2009; Harvell et al., 2007). Displacement of primary symbiotic microorganisms with other members has been shown to be related to the appearance of the disease signs. For example, the yellow blotch disease is caused by infection of *Vibrio* spp. which targets endosymbiotic zooxanthellae rather than the coral tissue, resulting in a decrease in chlorophyll concentration (Cervino et al., 2004). Bacteria are considered as the major infectious microorganisms in corals. So far, only a limited number of bacteria have been isolated and identified as causative agents of coral diseases by Koch's postulates, including cyanobacteria for black band disease, *Vibrio* sp and *Acrobacter* sp for white band disease, *Vibrio* spp for yellow blotch disease and *Serratia marcescens* for white pox (Sheridan et al., 2013). However, the symbiotic nature of bacteria to corals makes them unculturable using current isolation techniques. Therefore, identification of the causative microbes by conventional culture-dependent approaches may not be able to give the complete scenario of coral associated microbial communities upon infection.

Culture-independent analysis of environmental communities through phylogenetic molecular markers allows direct study of the uncultured microorganisms, which can make up to 99% of the total diversity in many ecosystems (Schloss & Handelsman, 2005). Several

studies have shown that culture-independent approaches based on bacterial molecular markers have been performed to explore the microbial communities associated with healthy and corals under stresses in the Caribbean and Indo-Pacific regions and their roles in pathogenesis (Gignoux-Wolfsohn & Vollmer, 2015; Meyer et al., 2017; Pootakham et al., 2018). However, variation in the taxonomic composition of the microbial consortia related to existing coral diseases can depend on different geographical, seasonal, and physical factors of the coral habitats (Sokolow, 2009; Woo et al., 2017). In order to gain more information on the roles of uncultured bacteria on coral diseases in Southeast Asia, the bacterial communities associated with corals collected from Indonesian sea, which is considered one of the world's most important coral habitat, in healthy and diseased stages affected by black band disease (BBD), white band disease (WBD), white plaque disease (WPD), and yellow blotch disease (YBD) were explored in this study using tagged 16S rRNA sequencing on a next-generation sequencing platform. Our work compares coral-associated bacteria in different host species and shows shifts in the bacterial community structure during the diseased stage. Our findings expand the current understanding on the microbiology of these prevalent coral diseases.

MATERIALS AND METHODS

Sample collection and DNA extraction

Coral specimens were taken from the Indonesian sea in Central Java (Cemara Kecil (S05°49'57,7"E110°22'50,5") and Karimunjawa (S05°50'12,9"E110°25'14,1")) (Fig. S1) during September, 2015 from a depth of about 3–5 m, with the permission from the Balai Taman Nasional Karimunjawa (the approval number 934 /BTNKJ-1.6 /SIMAKSI /2015). Triplicate samples (10 g each) of fragments from healthy and lesion fragments from diseased corals were collected from individual coral species including *Acropora aspera*, *Acropora Formosa*, *Cyphastrea* sp. , and *Isopora* sp. Healthy and infected coral were collected from different colonies of same species on same reefs. Coral fragments were placed in sterile bottles filled with seawater obtained from the same sampling area during transportation. The samples were rinsed using sterilized sea water and preserved at –80 °C upon arrival at the laboratory (Wegley et al., 2007). Physicochemical properties of the seawater including temperature, salinity, dissolved oxygen and pH were measured using Lutron YK-2005WA (Lutron Electronic Enterprise, Taipei, Taiwan).

The coral specimens were grounded in a mortar prior to metagenomic DNA extraction using the SDS-based DNA extraction procedure (Zhou, Bruns & Tiedje, 1996), with slight modifications (Kanokratana et al., 2004). Briefly, 1 g of grounded sample was mixed vigorously for 30 min at 37 °C with 2.7 ml DNA extraction buffer (100 mM Tris-HCl (pH 8.0), 100 mM sodium EDTA (pH 8.0), 100 mM sodium phosphate (pH 8.0), 1.5 M NaCl, 1% (wt/vol) cetyltrimethylammonium bromide (CTAB) and 20 µl of proteinase K (10 mg/ml). Subsequently, 300 µl of 20% [wt/vol] SDS was added and incubated at 65 °C for 2 h with gentle mixing every 15 min. After 2 h of incubation samples were centrifuged at 6,000× g for 10 min at room temperature. The resulting supernatants were mixed with an equal volume of chloroform: iso-amyl alcohol (24: 1, vol/vol). The aqueous phase was transferred to a new

tube after centrifugation at $6,000 \times g$ for 10 min. To the aqueous phase, 0.6 volume of iso-propanol was added and the mixture was left at room temperature for 1 h, followed by centrifugation ($14,000 \times g$, 20 min). Then, the DNA pellet was washed with ice-cold 70% (vol/vol) ethanol and resuspended in double-distilled water. Healthy and diseased coral fragments were extracted separately. The integrity of the extracted DNA was verified by agarose gel electrophoresis. The concentration of extracted DNA was quantified using a NanoDrop ND-1000 spectrophotometer (Peqlab Biotechnologie GmbH, Erlangen, Germany).

16S rRNA amplification and sequencing

The purified metagenomic DNA was used as the template for amplification of the partial 16S rRNA gene using universal bacterial primers E785F (5'-GGATTAGATACCCTGGTAGTCC-3') and E1081R (5'-CTCACGRCACGAGCTGACG-3') encompassing the 5 and 6 hypervariable regions in prokaryotic 16S rRNA gene. Each pair of primers was attached with a specific 8-bp barcode sequence at 5' end (Meyer, Stenzel & Hofreiter, 2008) and used for amplification of each metagenomic DNA sample in order to generate tagged 16S rRNA amplicons. Polymerase chain reactions were performed using Phusion DNA polymerase (Thermo Scientific, Espoo, Finland) on a MyCycler thermocycler for 25 cycles of denaturation at 98°C for 10 s, annealing at 69°C for 30 s, and extension at 72°C for 30 s. The amplicons were purified using QIAGEN PCR purification kit (QIAGEN, Hilden, Germany) and were quantified using NanoDrop ND-1000 Spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). Three independent PCR amplicons obtained from the individual coral metagenomic sample were performed. Sequencing was done using the ION PGM™ platform (Life Technologies, Carlsbad, CA, USA) following the manufacturer's recommended protocols.

Data cleaning and analyses

Data analysis was done in QIIME version 1.9.1 (Caporaso et al., 2010) mainly for clustering sequences into operational taxonomic units (OTUs), classifying taxonomy and calculating diversity index. First, the raw sequences were initially demultiplexed into specific groups based on the tagged sequences and trimmed off low-quality score reads (Phred \leq Q20), barcodes, and primers. Chimeric sequences and amplification errors were filtered by UCHIME (Edgar et al., 2011) using the referenced dataset from the Ribosomal Database Project (Cole et al., 2014). The OTUs were clustered from the remaining sequences with an "open-reference" method using UCLUST (Edgar, 2010) at 97% similarity. The representative sequence of each cluster was assigned for taxonomy against the Greengene database (McDonald et al., 2012). In order to compare the diversity among the samples, alpha diversity index including observed OTUs, Shannon-Weaver index, and the Chao1 richness estimator was calculated using a cutoff of 21,089 reads per sample, which was the minimal number of reads per sample obtained. The difference of microbial communities among samples was measured using weighted UniFrac distance (Lozupone et al., 2011) and Principal Coordinates Analysis (PCoA). The differences of bacterial abundance, OTU number, and diversity index between healthy and infected corals were statistically tested

Table 1 Sampling locations and sea water characteristics.

Sampling location	Temperature (°C)	pH	Salinity ^a (PSU)	Dissolve oxygen (mg/l)	Turbidity (m)	Collected coral name
Ujung gelam (S05°49'57.7", E110°22'50.5")	30.1 ± 0.4	7.3 ± 0.1	32.0 ± 0.4	6.4 ± 0.6	5 ± 0	<i>Acropora formosa</i>
Cemara Kecil (S05°50'12.9", E110°25'14.1")	28.3 ± 0.2	7.8 ± 0.1	34.3 ± 0.3	6.2 ± 0.3	2 ± 0	<i>Acropora aspera</i> <i>Cyphastrea</i> sp. <i>Isopora</i> sp.

Notes.^aPractical salinity unit.

using White's non-parametric *t*-test (White, Nagarajan & Pop, 2009) implemented in STAMP (Parks et al., 2014).

RESULTS

Sample characterization

Fragments of both healthy and diseased corals were collected from four major coral species in the sea of Central Java, Indonesia. Physicochemical characteristics of seawater in the sampling areas are described in Table 1. Both of the sampling sites have the same coral reef topology. Those sites are covered by various coral species from many lifeforms, such as branching, massive, submassive and encrusting. The seawater in Ujung Gelam had slightly higher temperature (30.1 °C) and dissolved oxygen (6.4 mg/l) but with lower pH (7.3) and salinity (32.0 PSU) than that in Cemara Kecil (28.3 °C, 6.2 mg/l, 7.8, and 34.3 PSU, respectively). The diseased corals were affected by different diseases commonly found in the Indo-Pacific sea. The physical appearances of the healthy and diseased corals are shown in Fig. 1 and Fig. S2. *A. aspera* infected by white band disease (*A. aspera*-WBD) showed symptom with the appearance of the white band between healthy and dead coral tissues. *A. formosa* infected by black band diseases (*A. formosa*-BBD) showed lesion as a reddish to dark-colored band. *Cyphastrea* sp. infected by yellow blotch disease (*Cyphastrea* sp.-YBD) which resulted in a band of yellow tissue around the enlarging sediment patch. *Isopora* sp. infected by white plaque disease (*Isopora* sp.-WPD) showed progressing band of bleached coral tissue followed by necrotic tissue starting from the base of the branch.

Overview of sequencing dataset and alpha diversity analysis

The structure of bacterial communities associated with healthy and infected corals were investigated based on 16S rRNA sequences analyzed by the ION PGM platform. A total of 1,788,418 raw reads were obtained with an average read length of 260 base pairs. After the data cleaning process, 89.25% of raw reads (1,596,127 reads) were obtained ranging from 21,453 to 107,722 reads per sample. There was no significant difference of the average sequence number obtained from individual coral species in healthy and infected stages (ANOVA; $P > 0.05$). Total numbers of sequencing read for all samples are presented in Table 2 and Table S1.

Analysis of microbial community by alpha and beta diversity suggested the inequality of diversity among coral samples (Table 2). Overall, there was no significant difference in bacterial diversity between the two sampling sites or among the coral species. According




Figure 1 Appearance of coral specimens in healthy and infected stages; (A) black band disease, (B) white band disease, (C) white plaque disease, (D) yellow blotch disease.

Full-size [DOI: 10.7717/peerj.8137/fig-1](https://doi.org/10.7717/peerj.8137/fig-1)

46 QIIME, the data set covered 96.8–99.1% of the bacterial diversity at 97% similarity. The number of OTUs at the 97% similarity ranged from 837 to 3,081 OTUs/ sample, which was lowest and highest in sample *A. aspera* (Aa-3) and *Isopora* sp. (Is-WPD-1), respectively. The highest total number of OTUs was found in sample *A. formosa* (Af-1) (4,440) while the lowest was observed in *Cyphastrea* sp. (Cs-YBD-1) (1,648) as estimated by the richness estimator, Chao1. In order to reduce the bias of unequal number of reads among the individual samples, normalization was performed by a subsampling method using a cutoff level of the minimal number of reads in an individual sample obtained (21,089 reads). Shannon's diversity indexes from OTU table (97% similarity) indicated significantly reduced diversity of bacteria in the diseased *A. formosa* and *Cyphastrea* sp. than healthy ones (P -value < 0.05); however, there was no significant difference in bacterial diversity between the healthy and diseased corals for species *A. aspera* and *Isopora* sp. (P -value > 0.05) (Fig. 2A). The shared OTU analysis revealed that the number of common OTUs in both healthy and diseased coral species ranging from 24.31–31.25%. A substantial fraction of OTUs in the healthy corals was replaced by newly introduced OTUs upon infection. This resulted in the presence of unique OTUs related to causative 71 opportunistic bacteria in all diseased corals, accounting for 28.78–49.51% of the total diversity of the coral-associated bacteria in the diseased stage (Fig. 2B).

The coverage of the dataset to the total diversity was assessed by rarefaction curve as shown in Fig. S3. For *A. formosa* and *Cyphastrea* sp., similar slopes were found for the healthy and infected corals, suggesting saturation of OTUs from both groups. In contrast, the slopes for the healthy *A. aspera* and *Isopora* sp. were substantially higher than that of

Table 2 Diversity estimates of bacterial 16S rRNA gene in sequencing datasets of microbial communities associated with healthy and infected corals.

Coral species	Sampling site	Condition	Sample ID	No. of sequences after QC	Observed OTUs	Chao1	Shannon
<i>Acropora aspera</i>	Cemara Kecil	Healthy	Aa-1	102,886	1,732	3,096	5.5581
			Aa-2	70,966	1,386	2,666	5.1340
			Aa-3	24,556	837	1,913	5.0301
		Infected with white band disease	Aa-WBD-1	61,634	2,249	3,609	5.9713
			Aa-WBD-2	107,722	1,948	3,337	5.4652
			Aa-WBD-3	27,701	1,363	2,539	6.1005
<i>Acropora formosa</i>	Karimunjawa	Healthy	Af-1	94,076	2,431	4,440	6.0624
			Af-2	88,344	2,300	4,177	6.0045
			Af-3	27,979	1,344	2,474	6.3581
		Infected with black band disease	Af-BBD-1	65,206	1,740	3,434	5.4460
			Af-BBD-2	73,163	2,222	4,401	5.5205
			Af-BBD-3	21,453	1,140	2,568	5.6828
<i>Cyphastrea</i> sp.	Cemara Kecil	Healthy	Cs-1	55,264	1,793	3,479	6.1171
			Cs-2	33,475	1,376	2,958	5.9524
			Cs-3	77,176	2,212	3,764	6.2162
		Infected with yellow blotch disease	Cs-YBD-1	23,850	925	1,648	5.3805
			Cs-YBD-2	63,703	1,987	3,769	5.5155
			Cs-YBD-3	56,042	1,704	3,533	5.2922
<i>Isopora</i> sp.	Cemara Kecil	Healthy	Is-1	55,471	1,592	3,324	5.6822
			Is-2	103,771	2,033	3,611	5.6433
			Is-3	102,458	2,252	3,555	6.0461
		Infected with white Plaque disease	Is-WPD-1	89,080	3,081	3,849	6.8630
			Is-WPD-2	78,463	1,965	3,305	6.1575
			Is-WPD-3	91,688	2,026	3,541	6.0657

the infected groups while the higher variation of OTUs was found in the infected corals compared to the healthy corals. The results thus suggested a high microbial adaptation of the bacteria associated with *A. aspera* and *Isopora* sp. in the infected stage.

Taxonomic assignment of coral-associated microbial communities

Taxonomic classification by QIIME revealed that totally 59 bacterial phyla, 190 orders, and 307 genera were assigned in coral metagenomes. Only a small fraction (1.11%) of the total sequences could not be classified into any known phyla and was labeled as an unassigned sequence. According to the total dataset, the overall bacterial communities in the corals was dominated by sequences affiliated to the phylum *Firmicutes*, followed by *Proteobacteria* and *Bacteroidetes* as the major phyla across all samples, respective in their relative abundance, except in *A. aspera* where these three phyla showed relatively balanced distribution. Other phyla including *Actinobacteria*, *Fusobacteria*, and *Lentisphaerae* were present in lower abundances. Variation in their relative abundance was found depending on coral species and health status (Fig. 3). At the class level, *Clostridia* were a predominated class in *Firmicutes* while *Delta-* and *Gammaproteobacteria* were majorly presented in




Figure 2 Comparison of Shannon's diversity indexes (A) and number of shared/unique OTUs (B) between healthy and infected conditions in each coral species. Coral species marked with asterisk represented significant difference (White's non-parametric t-test) between healthy and infected corals.

Full-size [DOI: 10.7717/peerj.8137/fig-2](https://doi.org/10.7717/peerj.8137/fig-2)

phylum *Proteobacteria* in all coral samples, except in *Isopora* sp.-WPD. *Bacteroidia* and *Flavobacteria* existed as a major class in *Bacteroidetes* in all samples.

Comparative analysis based on coral's health status indicated that relative abundance of *Proteobacteria* was markedly increased in diseased corals *A. aspera*-WBD (50.18%) and *Isopora* sp.-WPD (39.14%) but slightly decreased in the other two coral species. With a higher number of *Delta*- and *Gammaproteobacteria* in *A. aspera*-WBD compared to the healthy one, a significantly decreasing number of *Flavobacteria* (from 14.48% to 2.44%) in infected *A. aspera*-WBD was clearly observed while an abundance of *Clostridia* was not significantly different (P -value > 0.05) in both healthy and infected groups. Higher relative abundances of *Delta*-, *Epsilon*- and *Gammaproteobacteria* in *Isopora* sp.-WPD were observed in agreement with the reduction of *Clostridia* when compared to the healthy corals. The ratio of *Clostridia* was significantly higher in *A. formosa* -BBD and *Cyphastrea* sp.-YBD compared to the healthy ones.

Focusing at the deeper taxonomic level, different relative abundances with significant variations in some bacterial genus were observed depending on coral species and health status (Fig. 4). *Fusibacter* belonging to phylum *Firmicutes* was predominantly present across all samples except *A. aspera*-WBD, where *Vibrio* was a major genus with approximately two times higher than the *Fusibacter*. This was related to the marked decreasing relative




Figure 3 Relative abundance of bacterial distribution profiles at class level within the coral metagenomic samples obtained from healthy and diseased coral species. Bacterial taxa labelled with coral species; Aa, Af, Cs, and Is indicated significantly difference (White's non-parametric t-test) between healthy and infected corals.

Full-size [DOI: 10.7717/peerj.8137/fig-3](https://doi.org/10.7717/peerj.8137/fig-3)




Figure 4 Taxonomic classification of bacterial diversity at the genus level. Major bacterial genera were selected and shown as (A) a heat map based on the coral species and their health status and (B–E) difference in mean proportions healthy and infected corals.

Full-size [DOI: 10.7717/peerj.8137/fig-4](https://doi.org/10.7717/peerj.8137/fig-4)

abundance of *Fusibacter* in *A. aspera*-WBD along with a significantly increased ratio of *Vibrio* (and *Acrobacter* but not significant). A lower number of *Fusibacter* in infected corals was also found in the *Isopora* sp.-WPD, but the number of *Vibrio* was not changed significantly (P -value > 0.05) regardless of the coral health. Relative abundance of *Arcobacter* in class *Epsilonproteobacteria* was significantly higher (P -value < 0.05) in *Isopora* sp.-WPD but was not different in the other groups of diseased coral. On the other hand, the amount of *Fusibacter* found in the *A. formosa*-BBD and *Cyphastrea* sp.-YBD was higher than that found in the healthy ones (P -value < 0.05). This was associated with the increase of genus WH1-8 within phylum *Firmicutes* in the *A. aspera*-WBD and *Isopora* sp.-WPD samples.




Figure 5 Weighted UniFrac PCoA plot showing the differences between bacterial communities based on (A) the coral health status and (B) coral species.

Full-size [DOI: 10.7717/peerj.8137/fig-5](https://doi.org/10.7717/peerj.8137/fig-5)

Correlation of bacterial diversity and beta diversity analysis

The beta diversity of bacterial communities related to the coral health status was measured using the weighted Unifrac method. Principal Coordinates Analysis (PCoA) based on weighted distance metric indicated that the bacterial diversity in healthy and diseased coral samples were clustered separately from each other for individual coral species (Fig. 5A). This suggested alteration of bacterial diversity when the corals were infected by any of these diseases. Distribution of bacteria was also clustered according to health status of corals regardless of the coral species, i.e., clusters between (1) healthy *A. formosa* and *Cyphastrea* sp., (2) infected *A. formosa*-BBD and *Cyphastrea* sp.-YBD, (3) healthy *A. aspera* and *Isopora* sp., and (4) infected *A. aspera*-WBD and *Isopora* sp.-WPD (Fig. 5B). Moreover, the result from PCoA plot conforms to the UPGMA tree constructed from weighted UniFrac distances (Fig. 6). The UPGMA tree topology showed clearly separated clusters between healthy and infected coral samples including (1) healthy *A. aspera* and infected *A. aspera*-WBD, (2) healthy *Isopora* sp. and infected *Isopora* sp.-WPD, and (3) healthy *A. formosa*, *Cyphastrea* sp. and infected *A. formosa*-BBD, *Cyphastrea* sp.-YBD. As individual coral species used in this study displayed different symptoms of diseases, the results suggested that when different coral species containing similar bacterial profiles were infected with different diseases, bacterial communities tend to alter to the similar direction. For example, bacterial profiles between *A. formosa* and *Cyphastrea* sp. were closed in the health status. However, when they were infected by BBD (*A. formosa*-BBD) and YBD (*Cyphastrea* sp.-YBD), bacterial communities in these two coral species were changed to the same direction and clustered together.

Changes in bacterial diversity in the healthy and infected corals are shown by a box plot of Phylogenetic Diversity (PD) Whole Tree. According to the results (Fig. S4), there is no significant difference in bacterial diversity measured by PD between healthy and infected corals (P -value > 0.05). Interestingly, the variation of PD values in diseased corals tended




Figure 6 Jackknifed UPGMA tree of all coral samples based on weighted UniFrac distance matrix. The jackknifed bootstrapped tree was illustrated with internal nodes colored; red for 75–100% support, yellow for 50–75%, green for 25–50%, and blue for < 25% support.

Full-size [DOI: 10.7717/peerj.8137/fig-6](https://doi.org/10.7717/peerj.8137/fig-6)

to be higher than the healthy coral possibly because the diseased coral is more susceptible to colonization by opportunistic bacteria. This was associated with higher or lower in diversity of the associated bacteria. Infection of *A. aspera* with WBD resulted in a marked increase in diversity of the bacterial communities. This situation was also observed in *Isopora* sp. and *Isopora* sp.-WPD. Decrease in complexity of the communities was found in *A. formosa*-BBD and *Cyphastrea* sp.-YBD. The results thus suggest alteration in the community structure of coral associated bacteria upon infection.

DISCUSSION

Coral diseases are the result of complex interactions between host, causative agents, and environment (Martin, Meek & Willeberg, 1987; Sunagawa, Woodley & Medina, 2010).

They are characterized by a shift in microbial communities in coral mucus and tissue. However, causes and consequences of this phenomenon to pathogenesis is usually not fully understood due to the complexity and dynamics of the associated bacteria as well as effects of abiotic factors. According to our study, shifts in bacterial communities were found in all taxonomic levels upon infection of all coral species by all diseases. WBD is an important coral disease causing loss of corals in many regions of the world. It has been reported to exhibit high host specificity, particularly *Acropora* species, including *A. cervicornis* and *A. palmata* in the Caribbean (Kline & Vollmer, 2011). Based on the findings in our work, *A. aspera* showed a reduced diversity index for its associated bacteria upon infection by WBD. This was related to the increasing abundance of all sub-phyla of *Proteobacteria* along with the decreases in *Firmicutes* (*Clostridia*) and *Bacteroidetes* (*Flavobacteria*) with the presence of *Fusibacteria* found only in the diseased stage. This phenomenon occurred along with a significant increase in *Vibrio* and WH1-8 and decreasing *Fusibacter* upon infection. The causative agent of WBD is currently unknown but has been shown to possibly cause by bacteria according to a study on *A. cervicornis* and *A. palmata* using antibiotic treatment (Sweet, Croquer & Bythell, 2014). *Vibrio* spp. were consistently found in association with both healthy and diseased corals (Cumming et al., 2008; Mouchka, Hewson & Harvell, 2010). Comparison of bacterial diversity in healthy and WBD-infected *A. palmata* based on 16S rRNA showed decreasing relative abundance of *Betaproteobacteria* and *Actinomycetes* with increasing abundance of *Planctomycetes* and *Cyanobacteria* (Pantos & Bythell, 2006). A putative pathogen *V. charcharii* has been identified and partially proven as the causative agent for WBD type II based on Henle-Koch's postulate (Kline & Vollmer, 2011). However, the causative bacteria for WBD type I has not yet been identified.

BBD is a polymicrobial coral disease which is considered one of the most virulent disease of scleractinian corals caused by polymicrobial factors which result in massive destruction of framework-building corals worldwide. Although factors affecting susceptibility of corals to BBD are still not fully understood, a few works showed that BBD pathogenesis is linked to nutrient enrichment, elevated temperature and light intensity. A recent study on Caribbean corals showed that the microbial communities in healthy corals were dominated by *Gammaproteobacteria*, particularly *Halomonas* spp. while the microbiome of BBD consortia were more variable and diverse (Meyer et al., 2017). Studies using a culture-independent molecular approach showed a diverse microbial community classified into four functional groups, including photoautotrophs (*Cyanobacteria*), sulfate-reducers (*Desulfovibrio*), sulfide oxidizers (*Beggiatoa*) and organo-heterotrophs (*Vibrio*) (Sere et al., 2016). Among them, *Desulfovibrio* spp. and *Vibrio coralliilyticus* were suspected as the primary pathogens; however, without proven by Henle Koch's postulate (Sere et al., 2016). Basically, the complex microbial consortia act to produce highly concentrated sulfide specifically by the promoted *Deltaproteobacteria* under anoxic conditions beneath the BBD mat that are lethal to coral tissue (Sato et al., 2017). The accumulation of sulfide underneath the BBD mat was partially due to the lack of sulfur oxidizers which contributes to the lethality of the disease (Meyer et al., 2017). Meta-analysis of published clonal library studies of BBD microbial communities showed that, with few exceptions, the microbial species composition of BBD communities did not correlate with the species of the host corals

with the domination of OTU of *Roseofilum reptotaenium* in over 70% of the samples (Miller & Richardson, 2011). In addition, three OTUs of Bacteroidetes and Alpha-proteobacteria were present in 13% of the samples with other OTUs found in <7% of the samples.

According to our study, decreasing relative abundance of *Proteobacteria*, particularly the subphyla *Epsilon* and *Gammaproteobacteria*, was found with increasing *Firmicutes* (*Clostridia*) in the *A. formosa*-BBD. This was related to increasing abundance of *Fusibacter* and WH1-8 bacteria upon the infection. Increasing abundance of *Fimicutes*, *Cytophaga-Flexibacter-Bacteroidetes* (CFB) and *Deltaproteobacteria* in the infectious mat was identified in a pioneered molecular analysis work using Terminal Fragment Length Polymorphism (T-RFLP) (Frias-Lopez et al., 2004). The results showed that the composition of the infectious bacterial mat was not related to the species of coral being infected. Instead, differences in the mat composition appear to be linked to the species of cyanobacteria dominant in the infection. However, the presence of cyanobacteria in *A. formosa*-BBD could not be detected due to the specificity of primers used for amplification (Nübel, Garcia-Pichel & Muyzer, 1997). No increase in *Desulfovibrio* was observed in the 16S rRNA sequencing analysis of *A. formosa*-BBD in this study. According to a previous study (Meyer et al., 2016), *Desulfovibrioprofundus* which is thought to be responsible the production of H₂S was detected in only 5% of the clone libraries analyzed. These discrepancies could be due to technical issues (e.g., amplification biases and low coverage of microbiome in clonal libraries), seasonal and/or regional differences in the BBD composition or function-based (rather than taxonomic-based) of the BBD community. Variations have been detected in bacterial communities associated across geographical regions and between sympatric coral species (Sere et al., 2016). High variability in the BBD bacterial communities in different geographical areas and coral species suggested that this disease derives from an earlier infection, which aids subsequent infection of opportunistic microorganisms such as cyanobacteria (Sere et al., 2016). Recent analysis of metagenomes from Caribbean and Pacific BBD mat revealed five metagenome-assembled genomes of *Roseofilum*, *Proteobacteria* and *Bacteroidetes* which are proposed to play symbiotic interaction (Meyer et al., 2017). However, the mechanisms of BBD development remain unclear, and no primary pathogens have yet been identified. The difference on the results observed in this study to previous works could be due to the difference of environmental factors such as climatic condition and location (Mouchka, Hanson & Harvell, 2010).

YBD infection in corals resulted in a decrease in microbial diversity index in *Cyphastrea* sp.. This was along with reduction in the relative abundance of *Proteobacteria* and *Bacteroidetes* (*Cytophaga*) and increase in *Firmicutes* (*Bacilli*) along the infection. *Vibrio alginolyticus* has been identified as a causative agent for YBD (Cervino et al., 2004b). Four *Vibrio* species (*V. rotiferianus*, *V. harveyi*, *V. alginolyticus*, and *V. proteolyticus*) were identified as causative agents in YBD in Caribbean corals through a series of infection and isolation experiments (Cervino et al., 2004b; Cuning et al., 2008). It is suggested that this *Vibrio* consortium infected the coral's symbiotic algae and resulted in degradation of zooxanthellae leading to the pale-yellow bands observed on infected Caribbean and Indo-Pacific corals (Cervino et al., 2008; Cuning et al., 2008). However, the pathogenesis mechanism during YBD infection is still poorly understood. A survey of *Vibrio* species

associated with healthy corals and YBD infected corals were conducted using a culture-based approach (Cunning et al., 2008). The results showed a shift from isolates taxonomically affiliated with *V. fortis* dominate in healthy corals to those related to *V. harveyi*, a known marine pathogen in diseased corals. However, this study did not find any *Vibrio* species that are always present in YBD lesion but not in healthy corals. Although *Vibrio* have been reported as pathogenic bacteria for various coral diseases, it should be noted that they are a part of common bacterial taxa found in coral microbiomes (Bourne & Munn, 2005; Daniels et al., 2011; Gray et al., 2011; Nithyanand & Pandian, 2009). The higher overall abundance of *Vibrio* in the healthy corals than the infected corals as found in our study could be expected due to the taxonomic refinement of the partial rDNA sequence was analyzed to the genus level only, not specifically to the specific group of pathogenic *Vibrio* species. The conflicting phenomenon on the higher abundance of *Vibrionales* in healthy corals has been previously reported in analysis of symbionts in corals infected by WPD (Cardenas et al., 2012; Kellogg et al., 2013).

WPD has been reported to affect more than 40 coral species (Sunagawa et al., 2009; Weil, Smith & Gil-Agudelo, 2006). However, identification of its causative agent is still problematic, suggesting a complex etiological phenomenon involving alterations in the dynamic interaction between environmental factors, and symbiotic microbiomes. According to our study, reduction in the relative abundance of all subphyla of *Proteobacteria* was found in *Isopora* sp. upon infection by WPD with increasing abundance of *Firmicutes* (*Clostridia*) and changes in the phylum *Bacteroidetes* with decreasing *Cytophaga* and increasing *Flavobacteria* along with reduction in *Lentisphaeria*. This change in the bacterial community structure led to an increase in *Acrobacter* and WH1-8 bacteria along with the decrease in *Clostridia* and *Fusibacter* at the genus level. Increasing diversity and a shift in bacterial community structure in *Montastraea faveolata* infected by WPS Type II has been shown using high-density 16S rRNA gene microarray and clone library sequencing (Sunagawa et al., 2009). Accumulation of various known bacterial families known as coral pathogens including *Alteromonadaceae* and *Vibrionaceae* has been found. However, the primary pathogen *Aurantimonas coralicida* (Denner et al., 2003) previously proven by Koch's postulate was not detected in this molecular study (Sunagawa et al., 2009). Analysis of differentially abundant OTUs in Caribbean coral species *Orbicella faceolata* and *O. franksi* showed marked differences in bacterial communities in the healthy and diseased coral samples but not between coral species (Sunagawa, Woodley & Medina, 2010). The subsequent comparison in Indo-Pacific coral species (*Pavonaduerdeni* and *Poriteslutea*) showed distinct bacterial community patterns associated with ocean basin, coral species and health status. Increasing bacterial richness was found in the diseased samples suggesting the role of opportunistic bacteria during pathogenesis. These studies showed microbial community patterns related to WPD that are consistent over coral species and oceans, irrespective of the putative underlying pathogens (Sunagawa, Woodley & Medina, 2010). This includes various taxa involving *Proteobacteria*, *Bacteroidetes*, *Cyanobacteria*, and *Firmicutes*. A higher abundance of known coral pathogens, e.g., *Alteromonadaceae*, *Rhodobacteraceae*, and *Vibrionaceae* have been reported in *Pavona duerdeni* and *Porites lutea* infected by WPD using 16S rRNA gene microarray (Roder et

al., 2014). A significant increase in *Alphaproteobacteria* and a concomitant decrease in the *Beta-* and *Gammaproteobacteria* were also observed in WPD-affected reef-building corals *Diploriastrigose* and *Siderastrea sidereal* using culture-dependent methods and pyrosequencing of 16S rRNA sequences (Cardenas et al., 2012). Significant shifts were also found for the orders *Rhizobiales*, *Caulobacteriales*, *Burkholderiales*, *Rhodobacteriales*, *Aletheromonadales*, and *Xanthomonadales*, suggesting roles of these bacteria on pathogenesis.

CONCLUSION

Our findings showed structural alteration of microbiomes associated with important reef-building corals in Indonesian sea revealed by a culture-independent molecular analysis approach. Coral disease pathogenesis led to disturbance of the complex microbial communities and shifts in the overall bacterial community structures as shown by well separated clustering of the healthy and infected coral microbiomes with significant changes in certain key bacterial species. This study provides insights on understanding the functions of associated symbiont bacteria in pathogenesis. Further in-depth analysis on the role of microbiomes on coral health is warrant.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. Philip J. Shaw for manuscript proofreading and comments.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This project was financially supported by the National Center for Genetic Engineering and Biotechnology, National Science and Technology Development Agency, Thailand. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
National Center for Genetic Engineering and Biotechnology,
National Science and Technology Development Agency, Thailand.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Wuttichai Mhuantong and Pattanop Kanokratana conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

- Handung Nuryadi conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.
- Agus Trianto, Agus Sabdono and Lily Eurwilaichitr conceived and designed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Sithichoke Tangphatsornruang analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Verawat Champreda conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Field sampling was approved by the Balai Taman Nasional Karimunjawa (934 /BTNKJ-1.6 /SIMAKSI /2015).

Data Availability

The following information was supplied regarding data availability:

The 16S rRNA gene sequences are deposited in NCBI Sequence Read Archive (SRA) with the accession number SRP071125.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8137#supplemental-information>.

REFERENCES

- Blackall LL, Wilson B, Van Oppen MJ. 2015. Coral—the world’s most diverse symbiotic ecosystem. *Molecular Ecology* 24(21):5330–5347 DOI 10.1111/mec.13400.
- Bourne DG, Garren M, Work TM, Rosenberg E, Smith GW, Harvell CD. 2009. Microbial disease and the coral holobiont. *Trends in Microbiology* 17(12):554–562 DOI 10.1016/j.tim.2009.09.004.
- Bourne DG, Munn CB. 2005. Diversity of bacteria associated with the coral *Pocillopora damicornis* from the Great Barrier Reef. *Environmental Microbiology* 7(8):1162–1174 DOI 10.1111/j.1462-2920.2005.00793.x.
- Brodersen KE, Lichtenberg M, Ralph PJ, Kuhl M, Wangpraseurt D. 2014. Radiative energy budget reveals high photosynthetic efficiency in symbiont-bearing corals. *Journal of the Royal Society Interface* 11(93):20130997 DOI 10.1098/rsif.2013.0997.
- Burriesci MS, Raab TK, Pringle JR. 2012. Evidence that glucose is the major transferred metabolite in dinoflagellate-cnidarian symbiosis. *Journal of Experimental Biology* 215(pt 19):3467–3477 DOI 10.1242/jeb.070946.
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Pena AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE,

- Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunencko T, Zaneveld J, Knight R. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7(5):335–336 DOI 10.1038/nmeth.f.303.
- Cardenas A, Rodriguez RL, Pizarro V, Cadavid LF, Arevalo-Ferro C. 2012. Shifts in bacterial communities of two Caribbean reef-building coral species affected by white plague disease. *ISME Journal* 6(3):502–512 DOI 10.1038/ismej.2011.123.
- Cervino JM, Hayes RL, Goreau TJ, Smith GW. 2004a. Zooxanthellae regulation in yellow blotch/band and other coral diseases contrasted with temperature related bleaching: in situ destruction vs expulsion. *Symbiosis* 37:63–85.
- Cervino JM, Hayes RL, Polson SW, Polson SC, Goreau TJ, Martinez RJ, Smith GW. 2004b. Relationship of *Vibrio* species infection and elevated temperatures to yellow blotch/band disease in Caribbean corals. *Applied and Environmental Microbiology* 70(11):6855–6864 DOI 10.1128/AEM.70.11.6855-6864.2004.
- Cervino JM, Thompson FL, Gomez-Gil B, Lorence EA, Goreau TJ, Hayes RL, Winiarski-Cervino KB, Smith GW, Hughen K, Bartels E. 2008. The *Vibrio* core group induces yellow band disease in Caribbean and Indo-Pacific reef-building corals. *Journal of Applied Microbiology* 105(5):1658–1671 DOI 10.1111/j.1365-2672.2008.03871.x.
- Cesar H, Burke L, Pet-Soede L. 2003. *The economics of worldwide coral reef degradation*. Cesar Environmental Economics Consulting (CEEC).
- Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, Brown CT, Porras-Alfaro A, Kuske CR, Tiedje JM. 2014. Ribosomal Database Project: data and tools for high throughput rRNA analysis. *Nucleic Acids Research* 42(Database issue):D633–D642 DOI 10.1093/nar/gkt1244.
- Cunning JR, Thurmond JB, Smith GW, Weil E, Ritchie KB. 2008. Proceedings of the 11th international coral reef symposium. In: *A survey of Vibrios associated with healthy and Yellow Band diseased Montastraea faveolata*. 206–210.
- Daniels CA, Zeifman A, Heym K, Ritchie KB, Watson CA, Berzins I, Breitbart M. 2011. Spatial heterogeneity of bacterial communities in the mucus of *Montastraea annularis*. *Marine Ecology Progress Series* 426:29–40 DOI 10.3354/meps09024.
- Denner EB, Smith GW, Busse HJ, Schumann P, Narzt T, Polson SW, Lubitz W, Richardson LL. 2003. *Aurantimonas corallicida* gen. nov. sp. nov. the causative agent of white plague type II on Caribbean scleractinian corals. *International Journal of Systematic and Evolutionary Microbiology* 53(pt 4):1115–1122 DOI 10.1099/ijs.0.02359-0.
- Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26(19):2460–2461 DOI 10.1093/bioinformatics/btq461.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27(16):2194–2200 DOI 10.1093/bioinformatics/btr381.

- Frias-Lopez J, Klaus JS, Bonheyo GT, Fouke BW. 2004. Bacterial community associated with black band disease in corals. *Applied and Environmental Microbiology* 70(10):5955–5962 DOI 10.1128/AEM.70.10.5955-5962.2004.
- Gignoux-Wolfsohn SA, Vollmer SV. 2015. Identification of candidate coral pathogens on white band disease-infected staghorn coral. *PLOS ONE* 10(8):e0134416 DOI 10.1371/journal.pone.0134416.
- Gray MA, Stone RP, McLaughlin MR, Kellogg CA. 2011. Microbial consortia of gorgonian corals from the Aleutian islands. *FEMS Microbiology Ecology* 76(1):109–120 DOI 10.1111/j.1574-6941.2010.01033.x.
- Harvell D, Jordán-Dahlgren E, Merkel S, Rosenberg E, Raymundo L, Smith G, Weil E, Willis B. 2007. Coral disease, environmental drivers, and the balance between coral and microbial associates. *Oceanography* 20(1):172–195 DOI 10.5670/oceanog.2007.91.
- Hughes TP, Baird AH, Bellwood DR, Card M, Connolly SR, Folke C, Grosberg R, Hoegh-Guldberg O, Jackson JB, Kleypas J, Lough JM, Marshall P, Nystrom M, Palumbi SR, Pandolfi JM, Rosen B, Roughgarden J. 2003. Climate change, human impacts, and the resilience of coral reefs. *Science* 301(5635):929–933 DOI 10.1126/science.1085046.
- Kanokratana P, Chanapan S, Pootanakit K, Eurwilaichitr L. 2004. Diversity and abundance of bacteria and archaea in the Bor Khlueng hot spring in Thailand. *Journal of Basic Microbiology* 44(6):430–444 DOI 10.1002/jobm.200410388.
- Kellogg CA, Piceno YM, Tom LM, DeSantis TZ, Gray MA, Zawada DG, Andersen GL. 2013. Comparing bacterial community composition between healthy and white plague-like disease states in *orbicella annularis* using PhyloChip™ G3 microarrays. *PLOS ONE* 8(11):e79801 DOI 10.1371/journal.pone.0079801.
- Kline DI, Vollmer SV. 2011. White Band Disease (type I) of endangered caribbean acroporid corals is caused by pathogenic bacteria. *Scientific Reports* 1:7 DOI 10.1038/srep00007.
- LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, Santos SR. 2018. Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Current Biology* 28:2570–2580 DOI 10.1016/j.cub.2018.07.008.
- Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R. 2011. UniFrac: an effective distance metric for microbial community comparison. *ISME Journal* 5(2):169–172 DOI 10.1038/ismej.2010.133.
- Martin SW, Meek AH, Willeberg P. 1987. *Veterinary epidemiology principles and methods*. Iowa State University Press.
- McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL, Knight R, Hugenholtz P. 2012. An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME Journal* 6(3):610–618 DOI 10.1038/ismej.2011.139.

- Meyer JL, Gunasekera SP, Scott RM, Paul VJ, Teplitski M. 2016.** Microbiome shifts and the inhibition of quorum sensing by Black Band Disease cyanobacteria. *ISME Journal* **10**(5):1204–1216 DOI [10.1038/ismej.2015.184](https://doi.org/10.1038/ismej.2015.184).
- Meyer JL, Paul VJ, Raymundo LJ, Teplitski M. 2017.** Comparative metagenomics of the polymicrobial black band disease of corals. *Frontiers in Microbiology* **8**:618.
- Meyer M, Stenzel U, Hofreiter M. 2008.** Parallel tagged sequencing on the 454 platform. *Nature Protocols* **3**(2):267–278 DOI [10.1038/nprot.2007.520](https://doi.org/10.1038/nprot.2007.520).
- Miller AW, Richardson LL. 2011.** A meta-analysis of 16S rRNA gene clone libraries from the polymicrobial black band disease of corals. *FEMS Microbiology Ecology* **75**(2):231–241 DOI [10.1111/j.1574-6941.2010.00991.x](https://doi.org/10.1111/j.1574-6941.2010.00991.x).
- Mouchka ME, Hewson I, Harvell CD. 2010.** Coral-associated bacterial assemblages: current knowledge and the potential for climate-driven impacts. *Integrative and Comparative Biology* **50**(4):662–674 DOI [10.1093/icb/icq061](https://doi.org/10.1093/icb/icq061).
- Nithyanand P, Pandian SK. 2009.** Phylogenetic characterization of culturable bacterial diversity associated with the mucus and tissue of the coral *Acropora digitifera* from the Gulf of Mannar. *FEMS Microbiology Ecology* **69**(3):384–394 DOI [10.1111/j.1574-6941.2009.00723.x](https://doi.org/10.1111/j.1574-6941.2009.00723.x).
- Nübel U, Garcia-Pichel F, Muyzer G. 1997.** PCR primers to amplify 16S rRNA genes from cyanobacteria. *Applied and Environmental Microbiology* **63**(8):3327–3332.
- Pantos O, Bythell JC. 2006.** Bacterial community structure associated with white band disease in the elkhorn coral *Acropora palmata* determined using culture-independent 16S rRNA techniques. *Diseases of Aquatic Organisms* **69**(1):79–88 DOI [10.3354/dao069079](https://doi.org/10.3354/dao069079).
- Parks DH, Tyson GW, Hugenholtz P, Beiko RG. 2014.** STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics* **30**(21):3123–3124 DOI [10.1093/bioinformatics/btu494](https://doi.org/10.1093/bioinformatics/btu494).
- Pootakham W, Mhuantong W, Putchim L, Yoocha T, Sonthirod C, Kongkachana W, Sangsrakru D, Naktang C, Jomchai N, Thongtham N, Tangphatsornruang S. 2018.** Dynamics of coral-associated microbiomes during a thermal bleaching event. *Microbiologyopen* **7**(5):e00604 DOI [10.1002/mbo3.604](https://doi.org/10.1002/mbo3.604).
- Radjasa OK, Vaske YM, Navarro G, Vervoort HC, Tenney K, Linington RG, Crews P. 2011.** Highlights of marine invertebrate-derived biosynthetic products: their biomedical potential and possible production by microbial associates. *Bioorganic and Medicinal Chemistry* **19**(22):6658–6674 DOI [10.1016/j.bmc.2011.07.017](https://doi.org/10.1016/j.bmc.2011.07.017).
- Roder C, Arif C, Bayer T, Aranda M, Daniels C, Shibl A, Chavanich S, Voolstra CR. 2014.** Bacterial profiling of White Plague Disease in a comparative coral species framework. *ISME Journal* **8**(1):31–39 DOI [10.1038/ismej.2013.127](https://doi.org/10.1038/ismej.2013.127).
- Rosenberg E, Ben-Haim Y. 2002.** Microbial diseases of corals and global warming. *Environmental Microbiology* **4**(6):318–326 DOI [10.1046/j.1462-2920.2002.00302.x](https://doi.org/10.1046/j.1462-2920.2002.00302.x).
- Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I. 2007.** The role of microorganisms in coral health, disease and evolution. *Nature Reviews. Microbiology* **5**(5):355–362 DOI [10.1038/nrmicro1635](https://doi.org/10.1038/nrmicro1635).

- Sato Y, Ling EY, Turaev D, Laffy P, Weynberg KD, Rattei T, Willis BL, Bourne DG. 2017. Unraveling the microbial processes of black band disease in corals through integrated genomics. *Scientific Reports* 7:40455 DOI 10.1038/srep40455.
- Schloss PD, Handelsman J. 2005. Metagenomics for studying unculturable microorganisms: cutting the Gordian knot. *Genome Biology* 6(8):229 DOI 10.1186/gb-2005-6-8-229.
- Sere M, Wilkinson DA, Schleyer MH, Chabanet P, Quod JP, Tortosa P. 2016. Characterisation of an atypical manifestation of black band disease on *Porites lutea* in the Western Indian Ocean. *PeerJ* 4:e2073.
- Sheridan C, Kramarsky-Winter E, Sweet M, Kushmaro A, Leal MC. 2013. Diseases in coral aquaculture: causes, implications and preventions. *Aquaculture* 396-399:124–135 DOI 10.1016/j.aquaculture.2013.02.037.
- Sokolow S. 2009. Effects of a changing climate on the dynamics of coral infectious disease: a review of the evidence. *Diseases of Aquatic Organisms* 87(1-2):5–18 DOI 10.3354/dao02099.
- Sunagawa S, DeSantis TZ, Piceno YM, Brodie EL, DeSalvo MK, Voolstra CR, Weil E, Andersen GL, Medina M. 2009. Bacterial diversity and White Plague Disease-associated community changes in the Caribbean coral *Montastraea faveolata*. *ISME Journal* 3(5):512–521 DOI 10.1038/ismej.2008.131.
- Sunagawa S, Woodley CM, Medina M. 2010. Threatened corals provide underexplored microbial habitats. *PLOS ONE* 5(3):e9554 DOI 10.1371/journal.pone.0009554.
- Sweet MJ, Croquer A, Bythell JC. 2014. Experimental antibiotic treatment identifies potential pathogens of white band disease in the endangered Caribbean coral *Acropora cervicornis*. *Proceedings of the Royal Society B: Biological Sciences* 281(1788):20140094 DOI 10.1098/rspb.2014.0094.
- Wegley L, Edwards R, Rodriguez-Brito B, Liu H, Rohwer F. 2007. Metagenomic analysis of the microbial community associated with the coral *Porites astreoides*. *Environmental Microbiology* 9(11):2707–2719 DOI 10.1111/j.1462-2920.2007.01383.x.
- Weil E, Smith G, Gil-Agudelo DL. 2006. Status and progress in coral reef disease research. *Diseases of Aquatic Organisms* 69(1):1–7 DOI 10.3354/dao069001.
- White JR, Nagarajan N, Pop M. 2009. Statistical methods for detecting differentially abundant features in clinical metagenomic samples. *PLOS Computational Biology* 5(4):e1000352 DOI 10.1371/journal.pcbi.1000352.
- Wilkinson C. 2008. *Status of coral reefs of the world: 2008*. Townsville, Australia: Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre.
- Woo S, Yang SH, Chen HJ, Tseng YF, Hwang SJ, Palmas SDe, Denis V, Imahara Y, Iwase F, Yum S, Tang SL. 2017. Geographical variations in bacterial communities associated with soft coral *Scleronephthya gracillimum*. *PLOS ONE* 12(8):e0183663 DOI 10.1371/journal.pone.0183663.
- Zhou J, Bruns MA, Tiedje JM. 1996. DNA recovery from soils of diverse composition. *Applied and Environmental Microbiology* 62(2):316–322.

Comparative analysis of bacterial communities associated with healthy and diseased corals in the Indonesian sea

ORIGINALITY REPORT

19%

SIMILARITY INDEX

%

INTERNET SOURCES

19%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Julie L Meyer, Sarath P Gunasekera, Raymond M Scott, Valerie J Paul, Max Teplitski. "Microbiome shifts and the inhibition of quorum sensing by Black Band Disease cyanobacteria", The ISME Journal, 2015 2%

Publication
- 2** Suhong Fu, Ming Lei, Yongqun Zhang, Zhaomin Deng, Jing Shi, Doudou Hao. "De novo transcriptome analysis of Tibetan medicinal plant *Dysphania schraderiana*", Genetics and Molecular Biology, 2019 1%

Publication
- 3** Anny Cárdenas, Luis M Rodriguez-R, Valeria Pizarro, Luis F Cadavid, Catalina Arévalo-Ferro. "Shifts in bacterial communities of two caribbean reef-building coral species affected by white plague disease", The ISME Journal, 2011 1%

Publication
- 4** Roder, Cornelia, Chatchanit Arif, Camille Daniels, Ernesto Weil, and Christian R. 1%

Voolstra. "Bacterial profiling of White Plague Disease across corals and oceans indicates a conserved and distinct disease microbiome", *Molecular Ecology*, 2014.

Publication

5

"Coral Reefs: An Ecosystem in Transition", Springer Nature, 2011

Publication

6

Wirulda Pootakham, Wuttichai Mhuantong, Thippawan Yoocha, Lalita Putchim et al. "High resolution profiling of coral-associated bacterial communities using full-length 16S rRNA sequence data from PacBio SMRT sequencing system", *Scientific Reports*, 2017

Publication

7

Michael Sweet, Deborah Burn, Aldo Croquer, Peter Leary. "Characterisation of the Bacterial and Fungal Communities Associated with Different Lesion Sizes of Dark Spot Syndrome Occurring in the Coral *Stephanocoenia intersepta*", *PLoS ONE*, 2013

Publication

8

Shinichi Sunagawa, Todd Z DeSantis, Yvette M Piceno, Eoin L Brodie et al. "Bacterial diversity and White Plague Disease-associated community changes in the Caribbean coral *Montastraea faveolata*", *The ISME Journal*,

1%

1%

1%

1%

2009

Publication

9

Pattanop Kanokratana, Tanaporn Uengwetwanit, Ukrit Rattanachomsri, Benjarat Bunternngsook et al. "Insights into the Phylogeny and Metabolic Potential of a Primary Tropical Peat Swamp Forest Microbial Community by Metagenomic Analysis", *Microbial Ecology*, 2010

Publication

1%

10

Boonyapakron, Katewadee, Kusol Pootanakit, Duriya Chantasingh, Kanyawim Kirtikara, and Lily Eurwilaichitr. "Cloning and expression of xylanase 10 from *Cryptovalsa mangrovei* (BCC7197) in *Pichia pastoris* : FULL LENGTH RESEARCH PAPER", *DNA Sequence*, 2005.

Publication

<1%

11

"Submarine Geomorphology", Springer Science and Business Media LLC, 2018

Publication

<1%

12

H. L. BUSS. "The coupling of biological iron cycling and mineral weathering during saprolite formation, Luquillo Mountains, Puerto Rico", *Geobiology*, 10/2005

Publication

<1%

13

Wong, Sandi, W. Zac Stephens, Adam R. Burns, Keaton Stagaman, Lawrence A. David,

<1%

Brendan J. M. Bohannon, Karen Guillemin, and John F. Rawls. "Ontogenetic Differences in Dietary Fat Influence Microbiota Assembly in the Zebrafish Gut", mBio, 2015.

Publication

14

"Coral Reefs of the Gulf", Springer Science and Business Media LLC, 2012

<1%

Publication

15

Chandan Badapanda. "Advancing Our Understanding of the Soil Microbial Communities Using QIIME Software: A 16S Data Analysis Pipeline", Journal of Applied Biotechnology & Bioengineering, 2017

<1%

Publication

16

Anne-Leila Meistertzheim, Maggy M. Nugues, Gaëlle Quéré, Pierre E. Galand. "Pathobiomes Differ between Two Diseases Affecting Reef Building Coralline Algae", Frontiers in Microbiology, 2017

<1%

Publication

17

"Coral Reefs of the Red Sea", Springer Science and Business Media LLC, 2019

<1%

Publication

18

Sarath P. Gunasekera, Julie L. Meyer, Yousong Ding, Khalil A. Abboud et al. "Chemical and Metagenomic Studies of the Lethal Black Band Disease of Corals Reveal Two Broadly

<1%

Distributed, Redox-Sensitive Mixed Polyketide/Peptide Macrocycles", Journal of Natural Products, 2019

Publication

19

J. Frias-Lopez, J. S. Klaus, G. T. Bonheyo, B. W. Fouke. "Bacterial Community Associated with Black Band Disease in Corals", Applied and Environmental Microbiology, 2004

<1%

Publication

20

Sweet, Michael J., John C. Bythell, and Maggy M. Nugues. "Algae as Reservoirs for Coral Pathogens", PLoS ONE, 2013.

<1%

Publication

21

Dada, Nsa, Estelle Jumas-Bilak, Sylvie Manguin, Razak Seidu, Thor-Axel Stenström, and Hans J Overgaard. "Comparative assessment of the bacterial communities associated with Aedes aegypti larvae and water from domestic water storage containers", Parasites & Vectors, 2014.

<1%

Publication

22

Sung-Suk Suh, Mirye Park, Jinik Hwang, Sukchan Lee, Sang Hyun Moh, Ki Hwan Park, Taek-Kyun Lee. "Characterization of bacterial communities associated with seasonal water masses from Tongyoung in South Sea of Korea", Ocean Science Journal, 2014

<1%

Publication

23 "Oceanographic and Biological Aspects of the Red Sea", Springer Science and Business Media LLC, 2019 <1%
Publication

24 Elham Karimi, Beate M Slaby, André R Soares, Jochen Blom, Ute Hentschel, Rodrigo Costa. "Metagenomic binning reveals versatile nutrient cycling and distinct adaptive features in alphaproteobacterial symbionts of marine sponges", FEMS Microbiology Ecology, 2018 <1%
Publication

25 Coral Health and Disease, 2004. <1%
Publication

26 Greta S. Aeby, Thierry M. Work, Christina M. Runyon, Amanda Shore-Maggio et al. "First Record of Black Band Disease in the Hawaiian Archipelago: Response, Outbreak Status, Virulence, and a Method of Treatment", PLOS ONE, 2015 <1%
Publication

27 Juliet Motteram, Isabell Kufner, Siân Deller, Frédéric Brunner, Kim E. Hammond-Kosack, Thorsten Nürnberger, Jason J. Rudd. "Molecular Characterization and Functional Analysis of , the Sole NPP1 Domain–Containing Protein, from the Fungal Wheat Leaf Pathogen ", Molecular Plant-Microbe Interactions, 2009 <1%

28

"Encyclopedia of Modern Coral Reefs", Springer Nature, 2011

Publication

<1%

29

E Eljarrat, J Caixach, J Rivera. "Determination of PCDDs and PCDFs in different animal feed ingredients", Chemosphere, 2002

Publication

<1%

30

Haohao Sun, Xiwei He, Lin Ye, Xu-Xiang Zhang, Bing Wu, Hongqiang Ren. "Diversity, abundance, and possible sources of fecal bacteria in the Yangtze River", Applied Microbiology and Biotechnology, 2016

Publication

<1%

31

Y. Sato, D. G. Bourne, B. L. Willis. "Effects of temperature and light on the progression of black band disease on the reef coral, *Montipora hispida*", Coral Reefs, 2011

Publication

<1%

32

Tomoko Hashimoto-Yasuda, Makoto Ikenaga, Susumu Asakawa, Han-Yong Kim, Masumi Okada, Kazuhiko Kobayashi, Makoto Kimura. "Effect of Free-Air CO Enrichment (FACE) on Methanogenic Archaeal Communities Inhabiting Rice Roots in a Japanese Rice Field ", Soil Science and Plant Nutrition, 2005

Publication

<1%

33

H. J. Liew, G. De Boeck, C. M. Wood. "An in vitro study of urea, water, ion and CO₂/HCO₃-transport in the gastrointestinal tract of the dogfish shark (*Squalus acanthias*): the influence of feeding", *Journal of Experimental Biology*, 2013

Publication

<1%

34

Tagliavia, Marcello, Enzo Messina, Barbara Manachini, Simone Cappello, and Paola Quatrini. "The gut microbiota of larvae of *Rhynchophorus ferrugineus* Oliver (Coleoptera: Curculionidae)", *BMC Microbiology*, 2014.

Publication

<1%

35

Caroline S Fortunato, Alexander Eiler, Lydie Herfort, Joseph A Needoba, Tawnya D Peterson, Byron C Crump. "Determining indicator taxa across spatial and seasonal gradients in the Columbia River coastal margin", *The ISME Journal*, 2013

Publication

<1%

36

Netta Kooperman, Eitan Ben-Dov, Esti Kramarsky-Winter, Zeev Barak, Ariel Kushmaro. "Coral mucus-associated bacterial communities from natural and aquarium environments", *FEMS Microbiology Letters*, 2007

Publication

<1%

37

Jamie M. McDevitt-Irwin, Melissa Garren, Ryan

McMinds, Rebecca Vega Thurber, Julia K. Baum. "Variable interaction outcomes of local disturbance and El Niño-induced heat stress on coral microbiome alpha and beta diversity", *Coral Reefs*, 2019

Publication

<1%

38

LD Mydlarz, CS Couch, E Weil, G Smith, CD Harvell. "Immune defenses of healthy, bleached and diseased *Montastraea faveolata* during a natural bleaching event", *Diseases of Aquatic Organisms*, 2009

Publication

<1%

39

Mathieu G. Séré, Pablo Tortosa, Pascale Chabanet, Jean Turquet, Jean-Pascal Quod, Michael H. Schleyer. "Bacterial Communities Associated with Porites White Patch Syndrome (PWPS) on Three Western Indian Ocean (WIO) Coral Reefs", *PLoS ONE*, 2013

Publication

<1%

40

Lawler, Stephanie N., Christina A. Kellogg, Scott C. France, Rachel W. Clostio, Sandra D. Brooke, and Steve W. Ross. "Coral-Associated Bacterial Diversity Is Conserved across Two Deep-Sea *Anthothela* Species", *Frontiers in Microbiology*, 2016.

Publication

<1%

41

"Reproductive Sciences in Animal

<1%

Conservation", Springer Science and Business Media LLC, 2014

Publication

42

Julie L. Meyer, Valerie J. Paul, Laurie J. Raymundo, Max Teplitski. "Comparative Metagenomics of the Polymicrobial Black Band Disease of Corals", *Frontiers in Microbiology*, 2017

Publication

<1%

43

LL Richardson, AW Miller, E Broderick, L Kaczmarzsky, M Gantar, D Stanic, R Sekar. "Sulfide, microcystin, and the etiology of black band disease", *Diseases of Aquatic Organisms*, 2009

Publication

<1%

44

Den Uyl, Paul A., Laurie L. Richardson, Sunit Jain, and Gregory J. Dick. "Unraveling the Physiological Roles of the Cyanobacterium *Geitlerinema* sp. BBD and Other Black Band Disease Community Members through Genomic Analysis of a Mixed Culture", *PLoS ONE*, 2016.

Publication

<1%

45

Cristina Becerra-Castro, Ana Rita Lopes, Sara Teixeira, M. Elisabete F. Silva et al. "Characterization of bacterial communities from Masseiras, a unique Portuguese greenhouse agricultural system", *Antonie van Leeuwenhoek*,

<1%

2017

Publication

46

Shang Wang, Weiguo Hou, Hailiang Dong, Hongchen Jiang et al. "Control of Temperature on Microbial Community Structure in Hot Springs of the Tibetan Plateau", PLoS ONE, 2013

Publication

<1%

47

MG Séré, P Chabanet, J Turquet, JP Quod, MH Schleyer. "Identification and prevalence of coral diseases on three Western Indian Ocean coral reefs", Diseases of Aquatic Organisms, 2015

Publication

<1%

48

Willner, Dana, Joshua Daly, David Whiley, Keith Grimwood, Claire E. Wainwright, and Philip Hugenholtz. "Comparison of DNA Extraction Methods for Microbial Community Profiling with an Application to Pediatric Bronchoalveolar Lavage Samples", PLoS ONE, 2012.

Publication

<1%

49

Zeber-Lubecka, Natalia, Maria Kulecka, Filip Ambrozkiwicz, Agnieszka Paziewska, Milosz Lechowicz, Ewa Konopka, Urszula Majewska, Maria Borszewska-Kornacka, Michal Mikula, Bozena Cukrowska, and Jerzy Ostrowski. "Effect of Saccharomyces boulardii and Mode of Delivery on the Early Development of the Gut

<1%

Microbial Community in Preterm Infants", PLoS ONE, 2016.

Publication

50

G. M. Cook, J. P. Rothenberger, M. Sikaroodi, P. M. Gillevet, E. C. Peters, R. B. Jonas. "A comparison of culture-dependent and culture-independent techniques used to characterize bacterial communities on healthy and white plague-diseased corals of the *Montastraea annularis* species complex", Coral Reefs, 2013

<1%

Publication

51

Loredana Stabili, Frine Cardone, Pietro Alifano, S. Maurizio Tredici, Stefano Piraino, Giuseppe Corriero, Elda Gaino. "Epidemic Mortality of the Sponge *Ircinia variabilis* (Schmidt, 1862) Associated to Proliferation of a *Vibrio* Bacterium", Microbial Ecology, 2012

<1%

Publication

52

Andrian P. Gajigan, Leomir A. Diaz, Cecilia Conaco. " Resilience of the prokaryotic microbial community of to elevated temperature ", MicrobiologyOpen, 2017

<1%

Publication

53

"The Cnidaria, Past, Present and Future", Springer Nature, 2016

<1%

Publication

54

Meng, Fangang, Yuan Wang, Li-Nan Huang, Jie

Li, Feng Jiang, Shiyu Li, and Guang-Hao Chen. "A novel nonwoven hybrid bioreactor (NWHBR) for enhancing simultaneous nitrification and denitrification", *Biotechnology and Bioengineering*, 2013.

Publication

<1%

55

Cornelia Roder, Chatchanit Arif, Till Bayer, Manuel Aranda et al. "Bacterial profiling of White Plague Disease in a comparative coral species framework", *The ISME Journal*, 2013

Publication

<1%

56

Microbial Factories, 2015.

Publication

<1%

57

Katie L. Barott. "Microbial diversity associated with four functional groups of benthic reef algae and the reef-building coral *Montastraea annularis* : Microbial diversity on benthic algae and corals", *Environmental Microbiology*, 05/2011

Publication

<1%

58

Godwin, Scott, Elizabeth Bent, James Borneman, and Lily Pereg. "The Role of Coral-Associated Bacterial Communities in Australian Subtropical White Syndrome of *Turbinaria mesenterina*", *PLoS ONE*, 2012.

Publication

<1%

59

Kostić, Tanja, Michael Ellis, Maggie R. Williams,

Tiffany M. Stedtfeld, John B. Kaneene, Robert D. Stedtfeld, and Syed A. Hashsham. "Thirty-minute screening of antibiotic resistance genes in bacterial isolates with minimal sample preparation in static self-dispensing 64 and 384 assay cards", Applied Microbiology and Biotechnology, 2015.

Publication

<1%

60

Jorge H. Pinzón C., Joshua Beach-Letendre, Ernesto Weil, Laura D. Mydlarz. "Relationship between Phylogeny and Immunity Suggests Older Caribbean Coral Lineages Are More Resistant to Disease", PLoS ONE, 2014

Publication

<1%

61

Raeid M. M. Abed. "Long-term compositional changes after transplant in a microbial mat cyanobacterial community revealed using a polyphasic approach", Environmental Microbiology, 2/2001

Publication

<1%

62

Cai, Yuanfeng, Fanxiang Kong, Limei Shi, and Yang Yu. "Spatial heterogeneity of cyanobacterial communities and genetic variation of Microcystis populations within large, shallow eutrophic lakes (Lake Taihu and Lake Chaohu, China)", Journal of Environmental Sciences, 2012.

Publication

<1%

63

Linda Wegley. "Metagenomic analysis of the microbial community associated with the coral *Porites astreoides*", *Environmental Microbiology*, 7/21/2007

Publication

<1%

64

"Coral Bleaching", Springer Science and Business Media LLC, 2018

Publication

<1%

65

Shan Yang, Wei Sun, Cen Tang, Liling Jin, Fengli Zhang, Zhiyong Li. "Phylogenetic Diversity of Actinobacteria Associated with Soft Coral *Alcyonium graclimum* and Stony Coral *Tubastraea coccinea* in the East China Sea", *Microbial Ecology*, 2013

Publication

<1%

66

Luba Arotsker, Nachshon Siboni, Eitan Bendov, Esti Kramarsky-Winter, Yossi Loya, Ariel Kushmaro. " sp. as a potentially important member of the Black Band Disease (BBD) consortium in sp. corals ", *FEMS Microbiology Ecology*, 2009

Publication

<1%

67

"Posters", *FEMS Microbiology Letters*, 2003

Publication

<1%

68

Samodha C. Fernando, Jia Wang, Kimberly Sparling, Gizele D. Garcia et al. "Microbiota of the Major South Atlantic Reef Building Coral

<1%

Mussismilia", Microbial Ecology, 2014

Publication

69

Blackall, Linda L., Bryan Wilson, and Madeleine J. H. van Oppen. "Coral-the world's most diverse symbiotic ecosystem", *Molecular Ecology*, 2015.

Publication

70

Lucas H Sampaio, Mariane MA Stefani, Regiane M Oliveira, Ana LM Sousa, Greg C Ireton, Steven G Reed, Malcolm S Duthie. "Immunologically reactive *M. leprae* antigens with relevance to diagnosis and vaccine development", *BMC Infectious Diseases*, 2011

Publication

71

Yan-Ying Zhang, Juan Ling, Qing-Song Yang, You-Shao Wang et al. "The diversity of coral associated bacteria and the environmental factors affect their community variation", *Ecotoxicology*, 2015

Publication

72

Beneficial Microorganisms in Multicellular Life Forms, 2011.

Publication

73

The Prokaryotes, 2013.

Publication

74

Elizabeth Pleshe, John Truesdell, Robert T. Batey. " Structure of a class II TrmH tRNA-

<1%

<1%

<1%

<1%

<1%

<1%

modifying enzyme from ", Acta
Crystallographica Section F Structural Biology
and Crystallization Communications, 2005

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On