

Incidence of end-stage renal disease in patients with type 1 diabetes

Jaakko Helve, MD ^{1,2}, Reijo Sund, DSocSc ^{3,4}, Martti Arffman, MSocS ^{4,5}, Valma Harjutsalo, PhD ^{6,7}, Per-Henrik Groop, MD ^{1,2,6,7,8}, Carola Grönhagen-Riska, MD ¹ and Patrik Finne, MD ^{1,2}

¹Finnish Registry for Kidney Diseases, Helsinki, Finland

²Abdominal Center Nephrology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

³Department of Social Research, Centre for Research Methods, University of Helsinki, Helsinki, Finland

⁴Institute of Clinical Medicine, University of Eastern Finland, Kuopio, Finland

⁵National Institute for Health and Welfare (THL), Helsinki, Finland

⁶Diabetes and Obesity Research Program, Research Programs Unity, University of Helsinki, Helsinki, Finland

⁷Folkhälsan Institute of Genetics, Folkhälsan Research Center Biomedicum Helsinki, Finland

⁸Baker IDI Heart & Diabetes Institute, Melbourne, Australia

Corresponding author:

Jaakko Helve

Finnish Registry for Kidney Diseases

Kumpulantie 1 A, 6th floor

FI-00520 Helsinki

Finland

Tel +358-9-43422760

Fax +358-9-45410075

Jaakko.Helve@Helsinki.fi

Running title: Incidence of ESRD in patients with T1D

Manuscript word count: 2781

Number of tables: 2

Number of figures: 2

Abstract

OBJECTIVE To investigate how risk of end-stage renal disease among patients with type 1 diabetes has changed over time, and further how the risk is affected by age, sex, and time period of diagnosis of diabetes.

RESEARCH DESIGN AND METHODS A cohort including all patients younger than 30 years diagnosed with type 1 diabetes in Finland in 1965–2011 was followed until start of renal replacement therapy, death, or end of follow-up at the end of 2013. Altogether 29 906 patients were included. The main outcome was cumulative risk of end-stage renal disease, accounting for death as a competing risk.

RESULTS The patients were followed up for a median of 20 years. During 616 403 patient-years, 1 543 end-stage renal disease cases and 4 185 deaths were recorded. The cumulative risk of end-stage renal disease was 2.2% after 20 years, and 7.0% after 30 years from the diabetes diagnosis. The relative risk of end-stage renal disease was 0.13 [95% CI 0.08–0.22) among patients diagnosed in 1995–2011 compared to those diagnosed in 1965–1979. Patients younger than 5 years at the time of diagnosis had the lowest risk of end-stage renal disease after diagnosis. If the cumulative risk of end-stage renal disease was estimated from time of birth, the patients aged 5–9 years at diabetes diagnosis were at highest risk.

CONCLUSIONS The cumulative risk of end-stage renal disease has decreased markedly during the past five decades. This highlights the importance of modern treatment of diabetes and diabetic nephropathy.

Introduction

It is well known that diabetic nephropathy is associated with increased morbidity and mortality in patients with type 1 diabetes, and end-stage renal disease (ESRD) increases the mortality markedly (1-4). However, the risk of diabetic nephropathy has decreased during the past decades probably because of improvements in glucose and blood pressure control (5-7). In 2005 we showed that the risk of ESRD among patients with type 1 diabetes had diminished during the past 4 decades and was only 7.8% at 30 years after the diabetes diagnosis (8). A more recent study on Swedish patients with type 1 diabetes reported an even lower cumulative incidence of ESRD (9).

Many studies have reported better renal prognosis if type 1 diabetes is diagnosed before puberty (8-10), but the results on the effect of sex have been conflicting (8,9,11,12). The incidence of ESRD starts to increase after 15 years and continues to increase up to 30 years from the diabetes diagnosis, and it has been suggested that new treatments have postponed the development of ESRD (8,13). The use of multiple injection insulin therapy, angiotensin converting enzyme inhibitors, angiotensin 2 receptor blockers, and statins has become increasingly common from the 1990's, and therefore only now we are able to see a possible effect of long-term use of these medications on the risk of ESRD.

It is of note that compared to previous reports our study population is comprehensive with an almost 50 year study period and complete coverage of patients with type 1 diabetes in Finland. The aim of this nationwide population-based study was to investigate how the trends in risk of ESRD have changed in patients with type 1 diabetes during last 5 decades.

Research Design and Methods

Study population

Patients with type 1 diabetes were identified from the FinDM II study (14), which has collected information on patients with diabetes in Finland. A primary source of this information is the register of entitlements to special reimbursement for medicines maintained by the Finnish Social Insurance Institution. In Finland insulin therapy has been fully reimbursed since 1964 for patients diagnosed with type 1 diabetes, and therefore the coverage of patients with type 1 diabetes in the register is complete. We selected all patients with type 1 diabetes, who had started insulin therapy before the age of 30 years between 1965 and 2011. For this population, we also obtained data on purchases of other medications from the Finnish Social Insurance Institution prescription database for the period between 1993 and 2011. Thus, patients using metformin (n=264) or other oral medication (n=49) used for type 2 diabetes within one year from the start of insulin therapy were excluded from the study. In addition, those patients having a diagnosis of secondary diabetes (n=221) in the Finnish Care Register were excluded. After these exclusions a total of 29 906 patients with type 1 diabetes were included in the study. ESRD was defined as onset of renal replacement therapy (RRT) based on information from the Finnish Registry for Kidney Diseases, which has almost full coverage (97-99%) of patients starting RRT (dialysis or kidney transplantation) in Finland since 1965 until the end of 2013. Information on deaths from 1965 until the end of 2013 was obtained from the Cause of Death Register maintained by Statistics Finland. Linkage between different registries was possible because of the Finnish system of unique personal identification numbers for all citizens. Notably, all data in this study were obtained from registries fully financed by the Finnish government. The ethics committee of the Finnish National Institute for Health and Welfare has approved the use of patient data for the FinDM II study, and the patients in the Finnish Registry for Kidney Diseases provided written

informed consent for use of their data for research purposes, and therefore separate approval by an ethics committee was not needed for this observational study.

Statistical methods

Patients were followed from the start of insulin therapy marking the onset of diabetes until the start of RRT, death, or end of follow up on 31 December 2013. Death is a competing risk event for ESRD as deceased patients are no longer at risk of ESRD. Therefore, the cumulative risk of ESRD was calculated using a method that takes into account the effect of death as a competing risk event. All-cause mortality was assessed using Kaplan-Meier survival probabilities. Cox proportional hazards- model was used to assess the relative risks of ESRD and death. Effect of age at diagnosis, time period of diagnosis, sex and the time-dependent variable of ESRD were estimated using Cox regression model.

The R statistical software 3.2.2 (The R Foundation for Statistical Computing, Vienna, Austria; available at <http://www.r-project.org>) and SAS statistical software, release 9.3 (SAS Institute Inc., Cary, NC, USA) were used to perform the statistical analyses.

Results

Altogether 29 906 patients aged under 30 years were diagnosed with type 1 diabetes in Finland in 1965-2011, of which 17 365 were men (58%). Median age at diabetes diagnosis was 12.5 years, and median follow-up time after diagnosis of diabetes was 20.3 years (Table 1). During 616 403 patient-years of follow-up, 1 543 ESRD cases and 4 185 deaths were recorded.

The incidence rate of ESRD started to rise 15 years after the diabetes diagnosis, increased up to 25 years from diagnosis, and thereafter reached a plateau and remained at the same level until the end of the 45-year follow-up in patients diagnosed 1965-69. However, the incidence rate at 16-20 years from the diabetes diagnosis has become lower over time (Figure 1). The cumulative incidence of ESRD was 2.2% (95% CI 2.0 to 2.4%) after 20 years, and 7.0% (95% CI 6.6 to 7.4%) after 30 years among all patients in the study cohort. However, if including only those diagnosed with diabetes after 1980, the cumulative incidence was only 1.3% (95% CI 1.1 to 1.5%) and 4.4% (95% CI 3.8 to 5.0%), respectively. Figure 2 shows the cumulative incidence of ESRD according to sex and age at diabetes diagnosis. The cumulative incidence of ESRD after 30 years was higher among men 7.7% (95% CI 7.1 to 8.3%) than women 6.0% (95% CI 5.4 to 6.6%). Patients diagnosed with type 1 diabetes before the age of five years had the lowest cumulative incidence of ESRD, but otherwise there was no association with age at diagnosis and risk of ESRD. The results were similar if including only the patients diagnosed with diabetes after 1980. The cumulative incidence was the second lowest in the age-group 5-9-years until approximately 25 years of follow-up, after which the incidence started to catch up and even overtake the cumulative incidence in the age groups 10-29 years.

Although the cumulative risk of ESRD, as assessed from time of diagnosis of diabetes, was the lowest among patients diagnosed under the age of five, the cumulative risk of ESRD as assessed from time of birth was 12.1% in men and 13.2% in women at age 50, compared to 3.5% and 2.1%, respectively, among those diagnosed with diabetes at age 25-29 years (Figure 2).

In multivariable analysis (Table 2), the patients diagnosed with type 1 diabetes before the age of 5 years had significantly lower risk of ESRD, whereas there was no difference in the risk between the older age group. The risk of ESRD was 26% lower among women than men. A

diabetes diagnosis in 1965-79 was associated with the highest risk of ESRD, but the prognosis has improved continuously during later time periods.

Only a small portion of the deceased patients had ESRD. The cumulative all-cause-mortality was 7.0% (95% CI 6.7 to 7.4%) at 20 years and 12.5% (95% CI 12.0 to 13.0%) at 30 years after the diabetes diagnosis. Hereby, the cumulative mortality was markedly higher than the cumulative incidence of ESRD. This highlights the importance of death as a competing risk event that reduces the risk of ever developing ESRD. Consequently, the adjusted relative risk of death in patients with type 1 diabetes and ESRD was 10.2 (95% CI 9.4 to 11.1) compared to other patients with type 1 diabetes, showing the considerable impact of ESRD on mortality in these patients. The relative risk of death was 34% lower in women than men. The risk of death increased with older age at diabetes diagnosis, while a later time period of diagnosis was associated with lower risk of death (Table 2).

Conclusions

We here showed that the risk of ESRD has decreased continuously over time and that the progression of diabetic nephropathy and renal failure is slower than before among patients with type 1 diabetes. However, type 1 diabetes is still a notable cause of ESRD, and comprised 14% of all patients who entered RRT in Finland in 2011-2015 (15). Within 30 years from the diagnosis of type 1 diabetes 7.0% of the patients developed ESRD, and the risk of death among these patients was 10 times as high as in other patients with type 1 diabetes. Although ESRD developed more slowly, the life-time risk of ESRD was the highest if diabetes was diagnosed at younger age. This nationwide study covering almost 50 years and 29 906 patients is the largest one to study the incidence in patients with type 1 diabetes.

We were able to perform this study, because of the national registries in Finland, which cover practically all patients with type 1 diabetes, all patients with ESRD, and all deaths during the past decades. Unique personal identification numbers for all citizens enabled linkage between these registries. For this reason, we could avoid selection bias, and assess how the incidence of ESRD has changed by time. It is of note that the incidence of type 1 diabetes in Finland is the highest in the world (16), and a lot of efforts and resources have consequently been invested in the treatment and study of these patients. It is therefore possible that this could have led to better prognosis of the patients with type 1 diabetes in Finland compared to many other countries (17). In addition, the Finnish population is almost entirely Caucasian and genetically homogenous. For these reasons, our results may not be directly generalizable to other parts of the world. In regression analyses it has to be taken into account that Kaplan-Meier curves according to age groups do not fulfil the proportional hazards assumption. Shapes of the curves are different if diabetes is diagnosed before or after ten-years of age, probably because of the influence of puberty.

In 2005 we showed that the cumulative incidence of ESRD in patients with type 1 diabetes was 7.8% after 30 years (8), which was lower than previously reported (18). Later a large population-based study from Sweden (n=11 681) reported an even lower cumulative incidence of ESRD, 3.3% after 30 years (9). The patient selection and study design was similar to ours, but the study period started later and hereby the follow-up period was shorter. In contrast a study from Pittsburgh, United States, showed a markedly higher incidence rate of ESRD for patients diagnosed 1965-1980. The cumulative incidence after 30 years was 13.7% for men, and 21.0% for women, but the incidence was even higher if diabetes was diagnosed 1950-1964, namely 43.4% for men and 24.6% for women, respectively (12). At least part of these differences can be explained by an earlier study period and a different patient selection. Another study from United States reported a decline in the cumulative incidence rates of

ESRD. If type 1 diabetes was diagnosed between 1975 and 1979, the 20-year cumulative incidence was 3.6% (19). Notably studies from Europe, Canada, and Australia have also reported declining incidence rates as well as an increase in the age at start of RRT in patients with type 1 diabetes (7,15,20,21), which is in line with our results. However, the incidence of ESRD caused by type 1 diabetes in the United States has not decreased but rather increased over the past 20 years, although also there it nowadays occurs at slightly older ages (22). Importantly, there was no risk reduction of ESRD despite of efficient renoprotective medication (23). Furthermore, it must be kept in mind that the prognosis of patients with type 1 diabetes and diabetic nephropathy is still poor in the developing countries (24).

There is strong evidence that better glucose control as well as treatment of hypertension and dyslipidemia decrease the risk of diabetic nephropathy and ESRD in patients with type 1 diabetes (7,25-31). The use of multiple insulin injections became more common in the 1990s, and was followed by the development of rapid-acting and long-acting insulin regimens, which enabled patients with type 1 diabetes to maintain a more stable blood glucose control. Also the use and variety of medication for dyslipidemia and hypertension has increased from 1990s. For instance angiotensin converting enzyme inhibitors and angiotensin receptor blockers have become a mainstay in the treatment of diabetic nephropathy during recent decades. As our study showed, the incidence of ESRD after 30 years has decreased over time. However, because there has been a broad use of these medications for only less than 20 years, we could expect to see a further decrease in the incidence of ESRD in the future. Although the risk of ESRD cannot be fully eliminated, the development of this devastating complication can probably be moved forward (32). Postponing kidney failure and the start of RRT start will increase quality of life, and reduce medical expenses.

Previous reports have showed conflicting results on how sex affects the incidence of ESRD among patients with type 1 diabetes. Many studies have reported that men more frequently

develop diabetic nephropathy (2,30,33), but the incidence of ESRD seems more complex. Costacou et al showed that the incidence of ESRD was higher among men if diabetes was diagnosed in 1950-1964, but if diabetes was diagnosed 1965-1980, the male excess was eliminated (12). There is evidence that the risk of ESRD is equal in men and women if diabetes is diagnosed during childhood, but if diabetes develops after puberty, the risk of ESRD is higher among men (9,11). This suggests a role of sex hormones. Supporting this theory, we showed in this study that the risk of ESRD is higher among men only if diabetes is diagnosed after puberty.

Diabetic nephropathy, especially when it proceeds to ESRD, is associated with an increased risk of premature death (1,2,33). Patients with type 1 diabetes and ESRD are estimated to have 18- to 30-fold higher standardized mortality ratio compared to the general population (3,4). Our study also showed a 10 fold-risk among type 1 diabetes patients with compared those without ESRD. However, the survival of patients with type 1 diabetes on RRT has improved during recent decades (34).

In conclusion, the risk of ESRD in patients with type 1 diabetes has decreased over time. Females and those diagnosed with type 1 diabetes at younger age are at lower risk, although the lifetime risk of ESRD is the highest among patients diagnosed with diabetes before the age of 10 years. Because modern treatment of diabetes with multiple insulin injections, renin-angiotensin system inhibitors, and statin therapy has been mainstay only for less than 20 years, there is hope that the cumulative incidence of ESRD will continue to decrease in the future.

Acknowledgments. We acknowledge support from the Board of the Finnish Registry for Kidney Diseases, and all the nephrologists and staff in all Finnish central hospitals which have reported to the Finnish Registry for Kidney Diseases: Helsinki University Central Hospital, Turku University Central Hospital, Satakunta Central Hospital, Kanta-Häme Central Hospital, Tampere University Central Hospital, Päijät-Häme Central Hospital, Kymenlaakso Central Hospital, Etelä-Karjala Central Hospital, Mikkeli Central Hospital, Itä-Savo Central Hospital, Pohjois-Karjala Central Hospital, Kuopio University Central Hospital, Keski-Suomi Central Hospital, Etelä Pohjanmaa Central Hospital, Vaasa Central Hospital, Keski-Pohjanmaa Central Hospital, Oulu University Central Hospital, Kainuu Central Hospital, Länsi-Pohja Central Hospital, Lappi Central Hospital and Åland Central Hospital.

Funding. Finnish Medical Society, Munuaissäätiö, and Finnish Academy. The sponsors had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; or preparation, review, and approval of the manuscript.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Authors Contributions. JH contributed to the literature search, study design, data collection, data interpretation, and writing. MA and RS contributed to the study design, data collection, data analysis, data interpretation, figures, and writing. VH contributed to the data collection, data interpretation, and writing. PHG and CGR contributed to the study design, data collection, data interpretation, and writing. PF contributed to the literature search, study design, data collection, data interpretation, and writing. All authors have seen and approved the final version of the article.

Figure 1. Incidence of Renal Replacement Therapy after Diagnosis of Type 1 Diabetes

Figure 2. Cumulative Incidence of End-Stage Renal Disease after Type 1 Diabetes Diagnosis and from Birth among Male and Female According to Age at Diagnosis of Diabetes

References

1. Astrup AS, Tarnow L, Rossing P, Pietraszek L, Riis Hansen P, Parving HH. Improved prognosis in type 1 diabetic patients with nephropathy: A prospective follow-up study. *Kidney Int* 2005 Sep;68(3):1250-1257
2. Andersen AR, Christiansen JS, Andersen JK, Kreiner S, Deckert T. Diabetic nephropathy in type 1 (insulin-dependent) diabetes: An epidemiological study. *Diabetologia* 1983 Dec;25(6):496-501
3. Groop PH, Thomas MC, Moran JL, Waden J, Thorn LM, Makinen VP, Rosengard-Barlund M, Saraheimo M, Hietala K, Heikkila O, Forsblom C, FinnDiane Study Group. The presence and severity of chronic kidney disease predicts all-cause mortality in type 1 diabetes. *Diabetes* 2009 Jul;58(7):1651-1658
4. Orchard TJ, Secrest AM, Miller RG, Costacou T. In the absence of renal disease, 20 year mortality risk in type 1 diabetes is comparable to that of the general population: A report from the pittsburgh epidemiology of diabetes complications study. *Diabetologia* 2010 Nov;53(11):2312-2319
5. Bojestig M, Arnqvist HJ, Hermansson G, Karlberg BE, Ludvigsson J. Declining incidence of nephropathy in insulin-dependent diabetes mellitus. *N Engl J Med* 1994 Jan 6;330(1):15-18

6. Hovind P, Tarnow L, Rossing K, Rossing P, Eising S, Larsen N, Binder C, Parving HH. Decreasing incidence of severe diabetic microangiopathy in type 1 diabetes. *Diabetes Care* 2003 Apr;26(4):1258-1264
7. Otani T, Yokoyama H, Ohashi Y, Uchigata Y. Improved incidence of end-stage renal disease of type 1 diabetes in japan, from a hospital-based survey. *BMJ Open Diabetes Res Care* 2016 Apr 11;4(1):e000177
8. Finne P, Reunanen A, Stenman S, Groop PH, Gronhagen-Riska C. Incidence of end-stage renal disease in patients with type 1 diabetes. *Jama* 2005 Oct 12;294(14):1782-1787
9. Mollsten A, Svensson M, Waernbaum I, Berhan Y, Schon S, Nystrom L, Arnqvist HJ, Dahlquist G, Swedish Childhood Diabetes Study Group, Diabetes Incidence Study in Sweden, Swedish Renal Registry. Cumulative risk, age at onset, and sex-specific differences for developing end-stage renal disease in young patients with type 1 diabetes: A nationwide population-based cohort study. *Diabetes* 2010 Jul;59(7):1803-1808
10. Svensson M, Nystrom L, Schon S, Dahlquist G. Age at onset of childhood-onset type 1 diabetes and the development of end-stage renal disease: A nationwide population-based study. *Diabetes Care* 2006 Mar;29(3):538-542
11. Harjutsalo V, Maric C, Forsblom C, Thorn L, Waden J, Groop PH, FinnDiane Study Group. Sex-related differences in the long-term risk of microvascular complications by age at onset of type 1 diabetes. *Diabetologia* 2011 Aug;54(8):1992-1999
12. Costacou T, Fried L, Ellis D, Orchard TJ. Sex differences in the development of kidney disease in individuals with type 1 diabetes mellitus: A contemporary analysis. *Am J Kidney Dis* 2011 Oct;58(4):565-573

13. Toppe C, Mollsten A, Schon S, Jonsson A, Dahlquist G. Renal replacement therapy due to type 1 diabetes; time trends during 1995-2010--a swedish population based register study. *J Diabetes Complications* 2014 Mar-Apr;28(2):152-155
14. Sund R, Koski S. FinDM II. On the register-based measurement of the prevalence and incidence of diabetes and its long-term complications. A technical report. Tampere: the Finnish Diabetes Association. 2009. Available at http://www.diabetes.fi/files/1167/DehkoFinDM_Raportti_ENG.pdf. Last viewed 9 Nov 2017.
15. Finnish Registry for Kidney Diseases, Report 2012. Available at http://www.muma.fi/files/2154/Finnish_Registry_for_Kidney_Diseases_2014.pdf Last viewed 9 Nov 2017.
16. Karvonen M, Viik-Kajander M, Moltchanova E, Libman I, LaPorte R, Tuomilehto J. Incidence of childhood type 1 diabetes worldwide. diabetes mondiale (DiaMond) project group. *Diabetes Care* 2000 Oct;23(10):1516-1526
17. Asao K, Sarti C, Forsen T, Hyttinen V, Nishimura R, Matsushima M, Reunanen A, Tuomilehto J, Tajima N, Diabetes Epidemiology Research International Mortality Study Group. Long-term mortality in nationwide cohorts of childhood-onset type 1 diabetes in japan and finland. *Diabetes Care* 2003 Jul;26(7):2037-2042
18. Krolewski M, Eggers PW, Warram JH. Magnitude of end-stage renal disease in IDDM: A 35 year follow-up study. *Kidney Int* 1996 Dec;50(6):2041-2046
19. Nishimura R, Dorman JS, Bosnyak Z, Tajima N, Becker DJ, Orchard TJ, Diabetes Epidemiology Research International Mortality Study, Allegheny County Registry.

- Incidence of ESRD and survival after renal replacement therapy in patients with type 1 diabetes: A report from the allegheny county registry. *Am J Kidney Dis* 2003 Jul;42(1):117-124
20. Assogba FG, Couchoud C, Hannedouche T, Villar E, Frimat L, Fagot-Campagna A, Jacquelinet C, Stengel B, French Renal Epidemiology and Information Network Registry. Trends in the epidemiology and care of diabetes mellitus-related end-stage renal disease in france, 2007-2011. *Diabetologia* 2014 Apr;57(4):718-728
21. ESRD Incidence Study Group, Stewart JH, McCredie MR, Williams SM. Divergent trends in the incidence of end-stage renal disease due to type 1 and type 2 diabetes in europe, canada and australia during 1998-2002. *Diabet Med* 2006 Dec;23(12):1364-1369
22. Krolewski AS, Bonventre JV. High risk of ESRD in type 1 diabetes: New strategies are needed to retard progressive renal function decline. *Semin Nephrol* 2012 Sep;32(5):407-414
23. Rosolowsky ET, Skupien J, Smiles AM, Niewczas M, Roshan B, Stanton R, Eckfeldt JH, Warram JH, Krolewski AS. Risk for ESRD in type 1 diabetes remains high despite renoprotection. *J Am Soc Nephrol* 2011 Mar;22(3):545-553
24. Bentata Y, Haddiya I, Latrech H, Serraj K, Abouqal R. Progression of diabetic nephropathy, risk of end-stage renal disease and mortality in patients with type-1 diabetes. *Saudi J Kidney Dis Transpl* 2013 Mar;24(2):392-402
25. de Boer IH, Afkarian M, Rue TC, Cleary PA, Lachin JM, Molitch ME, Steffes MW, Sun W, Zinman B, Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Research Group. Renal outcomes in

patients with type 1 diabetes and macroalbuminuria. *J Am Soc Nephrol* 2014
Oct;25(10):2342-2350

26. DCCT/EDIC Research Group, de Boer IH, Sun W, Cleary PA, Lachin JM, Molitch ME, Steffes MW, Zinman B. Intensive diabetes therapy and glomerular filtration rate in type 1 diabetes. *N Engl J Med* 2011 Dec 22;365(25):2366-2376
27. de Boer IH, Rue TC, Cleary PA, Lachin JM, Molitch ME, Steffes MW, Sun W, Zinman B, Brunzell JD, Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Study Research Group, White NH, Danis RP, Davis MD, Hainsworth D, Hubbard LD, Nathan DM. Long-term renal outcomes of patients with type 1 diabetes mellitus and microalbuminuria: An analysis of the diabetes control and complications Trial/Epidemiology of diabetes interventions and complications cohort. *Arch Intern Med* 2011 Mar 14;171(5):412-420
28. Forsblom C, Harjutsalo V, Thorn LM, Waden J, Tolonen N, Saraheimo M, Gordin D, Moran JL, Thomas MC, Groop PH, FinnDiane Study Group. Competing-risk analysis of ESRD and death among patients with type 1 diabetes and macroalbuminuria. *J Am Soc Nephrol* 2011 Mar;22(3):537-544
29. Lecaire TJ, Klein BE, Howard KP, Lee KE, Klein R. Risk for end-stage renal disease over 25 years in the population-based WESDR cohort. *Diabetes Care* 2014 Feb;37(2):381-388
30. Raile K, Galler A, Hofer S, Herbst A, Dunstheimer D, Busch P, Holl RW. Diabetic nephropathy in 27,805 children, adolescents, and adults with type 1 diabetes: Effect of diabetes duration, A1C, hypertension, dyslipidemia, diabetes onset, and sex. *Diabetes Care* 2007 Oct;30(10):2523-2528

31. Stadler M, Peric S, Strohner-Kaestenbauer H, Kramar R, Kaestenbauer T, Reitner A, Auinger M, Kronenberg F, Irsigler K, Amiel SA, Prager R. Mortality and incidence of renal replacement therapy in people with type 1 diabetes mellitus--a three decade long prospective observational study in the lainz T1DM cohort. *J Clin Endocrinol Metab* 2014 Dec;99(12):4523-4530
32. Marshall SM. Diabetic nephropathy in type 1 diabetes: Has the outlook improved since the 1980s?. *Diabetologia* 2012 Sep;55(9):2301-2306
33. Borch-Johnsen K, Andersen PK, Deckert T. The effect of proteinuria on relative mortality in type 1 (insulin-dependent) diabetes mellitus. *Diabetologia* 1985 Aug;28(8):590-596
34. Haapio M, Helve J, Groop PH, Gronhagen-Riska C, Finne P. Survival of patients with type 1 diabetes receiving renal replacement therapy in 1980-2007. *Diabetes Care* 2010 Aug;33(8):1718-1723

Table 1. Number of Males and Females Diagnosed as Having Type 1 Diabetes in Finland
According to Age and Time Period of Diagnosis

Age group	Sex	1965-79	1980-1984	1985-89	1990-94	1995-2011	Total
0-4 years	Total	808	328	378	501	2371	4386
	Males	431	184	208	254	1273	2350
	Females	377	144	170	247	1098	2036
5-9 years	Total	1577	590	638	693	3216	6714
	Males	813	313	354	351	1676	3507
	Females	764	277	284	342	1540	3207
10-14 years	Total	2147	663	630	684	3059	7183
	Males	1155	375	368	374	1799	4071
	Females	992	288	262	310	1260	3112
15-19 years	Total	1619	420	374	385	1479	4277
	Males	984	259	232	247	991	2713
	Females	635	161	142	138	488	1564
20-24 years	Total	1414	388	377	328	1040	3547
	Males	839	228	220	194	692	2173
	Females	575	160	157	134	348	1374
25-29 years	Total	1466	401	448	380	1104	3799
	Males	960	276	302	245	768	2551
	Females	506	125	146	135	336	1248
0-29 years	Total	9031	2790	2845	2971	12269	29906
	Males	5182	1635	1684	1665	7199	17365
	Females	3849	1155	1161	1306	5070	12541

Table 2. Relative Risks of ESRD and Death Associated with Sex, Age, and Time Period of Diagnosis of Type 1 Diabetes

Variable		ESRD RR	95% CI	P Value	Death RR	95% CI	P Value
Sex	Male	1			1		
	Female	0.74	0.67-0.82	<.001	0.66	0.62-0.71	<.001
Age (Years)	0-4	1			1		
	5-9	1.83	1.45-2.31	<.001	1.58	1.32-1.89	<.001
	10-14	2.00	1.60-2.51	<.001	2.01	1.70-2.38	<.001
	15-19	1.90	1.50-2.40	<.001	2.85	2.41-3.38	<.001
	20-24	2.02	1.59-2.58	<.001	4.57	3.87-5.39	<.001
	25-29	2.16	1.69-2.75	<.001	6.52	5.54-7.67	<.001
Year of Diabetes Diagnosis	1965-79	1			1		
	1980-84	0.56	0.48-0.67	<.001	0.64	0.57-0.71	<.001
	1985-89	0.40	0.32-0.50	<.001	0.48	0.42-0.55	<.001
	1990-94	0.32	0.23-0.44	<.001	0.47	0.40-0.55	<.001
	1995-2011	0.13	0.08-0.22	<.001	0.27	0.23-0.32	<.001

Abbreviations: ESRD, end-stage renal disease; RR, relative risk; CI, confidence interval