

COMPARISON OF NON-MAXIMAL TESTS FOR ESTIMATING EXERCISE CAPACITY

Reem A. Alajmi, Carl Foster, John P. Porcari, Kimberley Radtke, & Scott Doberstein

Department of Exercise and Sport Science, University of Wisconsin-La Crosse, USA

Original scientific article

DOI: 10.26582/k.52.1.2

Abstract:

Although maximal incremental exercise tests (GXT) are the gold standard for outcome assessment and exercise prescription, they are not widely available in either fitness or clinical exercise programs. This study compared the prediction of VO_2 max in healthy, sedentary volunteers using a non-exercise prediction (Matthews et al., 1999), RPE extrapolation to 19 and 20 and the Rockport Walking Test (RWT), and of ventilatory threshold (VT) using the Talk Test and RPE @ 13,14,15. Subjects performed a treadmill GXT with gas exchange, a submaximal treadmill with RPE and Talk Test, the RWT and Matthews. All methods provided reasonable estimates of both VO_2 max and VT, with correlations of >0.80 and $SEE \sim 1.3$ METs. VO_2 max was best estimated with the extrapolation to RPE=19. VT was intermediate between the TT Last Positive and Equivocal stages and between RPE 13 and 14. Non-maximal evaluation can be used in place of maximal GXT with gas exchange to make reasonable estimates of both VO_2 max and VT.

Key words: *exercise prescription, exercise test, VO_2 max, ventilatory threshold*

Introduction

Maximal graded exercise testing (GXT) has been the standard test performed for exercise evaluation and prescription and prognosis estimation (ACSM, 2017; Mezzani, et al., 2012). Although GXT is the gold standard, it has several limitations. One is that it is primarily performed for diagnostic and prognostic purposes rather than for exercise prescription (Myers, Voodi, Umann, & Froelicher, 2000), and it requires physician's involvement. Also, the use of GXT is limited in individuals who have pain/fatigue or who have contraindications to GXT (Noonan & Dean, 2000). Performing GXT is also limited by its cost and the availability of equipment.

To overcome the limitations of GXT, it would be useful to find simpler, safer, and lower cost alternatives. Several non-maximal tests have been used to estimate exercise capacity and guide exercise training intensity, including the Talk Test (TT), the Rating of Perceived Exertion (RPE), and Rockport 1-mile Walk Test (RWT). However, how well these tests measure the same exercise capacity qualities has not been established yet. Further, as each of these tests was developed in different samples of the population of nominally healthy individuals, the degree to which they are actually interchangeable remains uncertain.

The TT is easily conducted by asking the participants to carry on a conversation or to read a standard

passage at the end of each exercise stage. If the individual can speak comfortably, the exercise intensity is typically below the ventilatory threshold (VT). If they have some difficulties speaking, the exercise intensity is typically above the VT. Numerous studies have demonstrated the validity of TT as a method for exercise prescription by predicting the VT in samples from different populations, including healthy sedentary individuals, healthy active individuals, athletes, and patients with stable cardiovascular disease. The TT has also been used to identify the ischemic threshold in patients who had exertional ischemia (Foster, et al., 2018).

The Rating of Perceived Exertion (RPE) is a well-known method for determining, guiding and monitoring the exercise intensity (Eston, 2012). The RPE assesses how easy or hard a physical activity feels at any point in time. The RPE is based on its' very robust relationship with training intensity and with objective physiological markers (HR, VO_2 , VT, RCT, and blood lactate). The two most commonly used RPE scales are the original scale (Borg 6-20 Category Scale) and the Borg Category-Ratio-10 Scale (CR-10) (Borg, 1982). Both scales are widely used in cardiac rehabilitation and preventive medicine as methods of gauging exercise intensity (Mezzani, et al., 2012). Since HR can be uniquely variable in cardiac patients secondary both to pathology and medication effects, and is also

highly variable amongst healthy persons, the use of RPE is an attractive alternative to HR (Mezzani, et al., 2012). Earlier studies have shown a good correspondence between RPE 13 'somewhat hard' and the VT (Purvis & Cukiton, 1981) and training at RPE=13 has been shown to be both pleasant and effective (Parfitt, Evans, & Eston, 2012).

Because of the strong linear relationship between RPE and VO_2 , the RPE: VO_2 relationship can be utilized to predict the maximal oxygen consumption (VO_{2max}) from non-maximal exercise tests. There are two main procedures that can be used to predict VO_{2max} from linear regression: an estimation procedure and a production procedure (Coquart, Garcin, Parfitt, Tourny-Chollet, & Eston, 2014). The estimation procedure is a passive process in which individuals rate their physical exertion using the RPE scale at the end of each GXT stage. Conversely, the production procedure is an active process in which individuals self-regulate the intensity equivalent to a pre-determined RPE during each exercise test stage. By using either procedure, individual linear regression between RPE: VO_2 can be extrapolated to the theoretical endpoint at RPE 20 (or RPE 19) to predict VO_{2max} (Coquart, et al., 2014b). Several studies have shown RPE to be a valid method to estimate VO_{2max} and to monitor exercise intensity in healthy older adults (Chung, Zhao, Liu, & Quach, 2015), healthy participants (Eston, Lamb, Parfitt, & King, 2005; Eston, Faulkner, Mason, & Parfitt, 2006; Faulkner & Eston, 2007; Faulkner, Lambrick, Parfitt, Rowlands, & Eston, 2009; Lambrick, Faulkner, Rowlands, & Eston, 2009), athletes (Coquart, Eston, Nycz, Grosbois, & Garcin, 2012a), clinical participants (Coquart, et al., 2009), healthy and clinical populations (Al-Rahamneh & Eston, 2011; Al-Rahamneh, Faulkner, Byrne, & Eston, 2011), and in physically active and sedentary individuals (Eston, et al., 2012; Eston, Lambrick, Sheppard, & Parfitt, 2008; Faulkner, Parfitt, & Eston, 2007). RPE is also useful for estimating the anaerobic threshold (e.g. VT) (Parfitt, Evans & Eston, 2012).

The Rockport Walking Test (RWT), also known as 1-Mile Walk Test (1-MWT), is a popular submaximal field test used to predict VO_{2max} among a wide range of age categories and fitness levels (Kline, et al., 1987). The RWT is a brisk walking field test performed on a level surface track, for a distance of 1-mile (1,609 meters). Predicting VO_{2max} using the RWT is dependent on the age, gender, body weight, terminal HR and time required for completion. Dolgener, Hensley, Marsh, and Fielstul (1994) found that the original RWT equation should not be used for young individuals. Thus, an age-specific equation for RWT was established to predict VO_{2max} for young individuals.

Several previous studies have shown the validity of estimating VO_{2max} without the requirement for

exercise testing based on personal information collected from questionnaires (Jackson, et al., 1990; Matthews, Heil, Freedson, & Pastides, 1999). These questionnaire-based models have been shown to be of equivalent accuracy compared to submaximal exercise tests, particularly field tests such as the RWT (Jackson, et al., 1990). As an example of these models, Matthews et al. (1999) developed an equation, including age, age², gender, physical activity status, height, and body mass to predict VO_{2max} . Non-exercise questionnaire-based predictive equations can reasonably categorize individual fitness levels. Physical activity status was assessed using the NASA Physical Activity Rating (PA-R) questionnaire (Matthews, et al. 1999; Ross & Jackson, 1990).

There is a lack of evidence supporting non-maximal or non-exercise tests as a primary outcome measure and comparing these tests simultaneously relative to each other. Because different studies were used to create the predictions of VO_{2max} or VT, there remains an open question of how well they actually correspond with each other, or with objective measurements. Accordingly, the purpose of this study was to compare three non-maximal tests: the TT, the RPE, and the RWT relative to the accuracy of predicting exercise capacity, reflected by VO_{2max} and VT. The secondary aim was to examine the accuracy of questionnaire-based prediction equation compared to maximal as well as non-maximal testing. We hypothesized that there would be no significant difference between the non-maximal tests and the GXT and that the non-maximal methods could be used as primary methods for predicting exercise capacity.

Methods

Participants

The subjects in this study were 20 healthy adults, eight men and 12 women. All were of a low to moderate fitness level (no regular physical activity in the three months prior to the study). Prior to testing, there was an introductory session designed to define participants' characteristics and to familiarize them to the laboratory setting, exercise testing and equipment. The study protocol was approved by the University of Wisconsin-La Crosse Institutional Review Board for the Protection of Human Subjects. Written informed consent was provided by participants. They completed the following assessment forms: (1) Physical Activity Readiness Questionnaire (PAR-Q), (2) AHA/ACSM Health/Fitness Facility Preparticipation Screening Questionnaire to identify contraindications to GXT, and (3) the NASA Physical Activity Rating (PA-R) (Jackson, et al., 1990; Ross & Jackson, 1990) to assess the participants' level of physical activity.

Experimental design

Participants performed four tests, with at least 48 hours between the tests. The procedures included a maximal test, and three non-maximal tests. Test 1 was a maximal incremental exercise test conducted on the treadmill to the point of fatigue using a modified Balke protocol (ACSM, 2017), with respiratory gas exchange measurements using open circuit spirometry using a mixing chamber based metabolic cart (AEI, Bastrop, TX). The treadmill velocity/grade were initially set at 3.0 mph (4.8 kph)/0% grade for 2 minutes and increased by 2.5% grade every two minutes until fatigue. The RPE was reported at the end of each 2-minute stage using the Borg (6-20) scale (Borg, 1982). The gas analyzers were calibrated with reference gases (16% O₂, 5% CO₂) and room air. The pneumotach was calibrated with a 3-L syringe. Gas analysis data were summated every 30 seconds, with the highest 30 second value accepted as VO_{2max}. The VT and RCT were determined using both the V-slope and ventilatory equivalent methods (Foster & Cotter, 2006; Schneider, Phillips, & Stoffolano, 1993). Heart rate

was measured using radiotelemetry (Polar Electro Oy, Port Washington, NY), integrated every 5 seconds.

Test 2 was similar to Test 1, but without respiratory gas exchange measurement. During the last 30 seconds of each stage, the TT was performed by instructing the subjects to read a 101-word paragraph, the “Rainbow Passage”, out loud and respond to the question “Can you speak comfortably?” Possible answers were: “Yes”, which indicated the positive (POS) stage, any equivocal expressions like “Yes, but” or “I am not sure”, which indicated the equivocal (EQ) stage, and “No”, which indicated the negative (NEG) stage of TT. Based on previous studies (Foster, et al., 2018), the highest exercise intensity at which the subject indicates that they can speak comfortably is the last positive (LP) stage, which is typically below the VT. The EQ stage typically approximates the VT. The first NEG stage typically approximates the RCT. The RPE was assessed during the last 10 seconds of each 2-minute stage, immediately following the TT. The participant walked on the treadmill until the NEG stage of TT (or fatigue), and the RPE was recorded until the subject reached or exceeded RPE=15. By using the estimation procedure, RPEs reported throughout the exercise test and VO₂ values calculated from metabolic calculations (ACSM, 2017) (VO₂:RPE) were extrapolated to RPE=19 and RPE=20 to estimate VO_{2max} (Faulkner & Eston, 2007; Lambrick, et al., 2009; Faulkner, et al., 2009) (Figure 1).

For Test 3, the RWT test was performed on an indoor track (8 laps=1 mile). The subjects walked briskly for one mile. At the end of the walk, the time and terminal HR were recorded. VO_{2max} was calculated using the Rockport formula (Kline, et al., 1987):

$$\text{VO}_2\text{max (ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}) = 132.853 + (6.315*\text{gender}) - (0.3877*\text{age}) - (0.0769*\text{weight [pounds]}) - (3.2649*\text{time [min]}) - (0.1565*\text{final heart rate}) \text{ [Equation 1]},$$

where gender = 1 for male, 0 for female.

Additionally, VO_{2max} was calculated using the non-exercise prediction equation of Matthews et al. (1999):

$$\text{VO}_2\text{max (ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}) = (0.133 \times \text{age}) - (0.005 \times \text{age}^2) + (11.403 \times \text{gender}) + (1.463 \times \text{PA-R score}) + (9.17 \times \text{height [m]}) - (0.254 \times \text{body mass [Kg]}) + 34.142 \text{ [Equation 2]},$$

where gender = 1 for male, 0 for female. The PA-R score was obtained from the NASA PA-R questionnaire (Jackson, et al., 1990; Ross & Jackson, 1990).




Figure 1. Example of VO_{2max} prediction using the individual linear regression between RPE ≤ 15 and VO₂ calculated from metabolic calculations (ACSM, 2017) for one subject. The dotted lines represent the extrapolation of the individual RPE ≤ 15: VO₂ linear regression to a.) RPE 19 (predicted VO_{2max}: $1.81(19) - 1.19 = 33.2 \text{ mL}\cdot\text{kg}\cdot\text{min}^{-1}$), and to b.) maximal theoretical RPE (i.e. RPE 20) (predicted VO_{2max}: $1.81(20) - 1.19 = 35 \text{ mL}\cdot\text{kg}\cdot\text{min}^{-1}$).

Statistical analysis

Data were analyzed using the Statistical Package for Social Sciences (SPSS Inc., Chicago, IL, USA). Standard descriptive statistics were used to assess the baseline physical characteristics. Comparisons between the measured and the predicted VO_{2max}

and VT were made using repeated measures analysis of variance (ANOVA). If ANOVA showed a significant difference, *post-hoc* analyses were performed using the Tukey's Honestly Significance Difference (HSD) test. A p value ≤ 0.05 was considered statistically significant. Linear regression analysis between the calculated VO_2 values and $\text{RPE} \leq 15$ values was made for each subject. The resultant equation ($\text{VO}_2\text{max} = a + b [\text{RPE } 19 \text{ or } 20]$) was used to predict VO_2max (Figure 1). Pearson product-moment correlations and standard errors of the estimate (SEE) versus the predicted VO_2max were used to compare values.

Results

The characteristics and maximal exercise test data of the 20 participants are presented in Table 1. $\text{VO}_2\text{@RCT}$ was indeterminate in six subjects; thus, $\text{VO}_2\text{@RCT}$ and $\text{VO}_2\text{@NEG}$ stage of the TT were not analyzed in the study.

Overall, mean measured VO_2max was $38.1 \pm 6.52 \text{ mL} \times \text{kg}^{-1} \times \text{min}^{-1}$ and mean predicted VO_2max was 40.1 ± 6.9 , 40.1 ± 7.03 , 42.6 ± 7.43 , and $40.0 \pm 7.8 \text{ mL} \times \text{kg}^{-1} \times \text{min}^{-1}$ for the Matthews et al. (1999) equation, extrapolation to RPE 19, extrapolation to RPE 20, and the RWT equation, respectively (Table 2, Figure 2). Tukey's *post-hoc* test found that there was no significant difference between the measured and the predicted VO_2max using the Matthews et al. (1999) equation, extrapolation to RPE19 and the RWT equation (Kline, et al., 1987). VO_2max

Table 1. Descriptive characteristics of subjects (N=20)

Variable	Male (n=8)	Female (n=12)
Age (year)	25.2 \pm 7.90	28.3 \pm 9.37
Height (cm)	181.5 \pm 6.42	164.8 \pm 8.67
Weight (kg)	87.3 \pm 11.85	66.9 \pm 14.80
VO_2max (L \cdot min $^{-1}$)	3.67 \pm 0.689	2.33 \pm 0.426
VO_2max (mL \cdot kg $^{-1}$ \cdot min $^{-1}$)	42.1 \pm 5.84	35.5 \pm 5.71
VO_2 @ VT (L \cdot min $^{-1}$)	2.55 \pm 0.754	1.67 \pm 0.373
VO_2 @VT (mL \cdot kg $^{-1}$ \cdot min $^{-1}$)	29.1 \pm 6.38	25.4 \pm 4.54
HRmax (bpm)	175.6 \pm 14.64	183.7 \pm 10.86
HR@VT	140.3 \pm 22.33	150.9 \pm 14.58
RPE@VT	13.6 \pm 1.84	14.0 \pm 1.78
RPEmax	19.2 \pm 0.46	19.0 \pm 0.28

Note. Data are reported as mean \pm standard deviation.

Table 2. Comparison of the measured and the predicted VO_2max values (means \pm SD) in mL \cdot kg $^{-1}$ \cdot min $^{-1}$

Test	Mean \pm SD r R 2
Maximal test (GXT)	38.1 \pm 6.52
Matthews et al. equation ²⁷	40.1 \pm 6.91 0.76 0.57
Extrapolation to RPE 20	42.6 \pm 7.43* 0.84 0.70
Extrapolation to RPE 19	40.1 \pm 7.03 0.85 0.73
RWT equation ²⁴	40.0 \pm 7.82 0.71 0.51

Note. r = Pearson product-moment correlation. R 2 = coefficient of determination. * significantly different than maximal test ($p < .05$).




Figure 2. Mean (\pm standard error, SE) values for the measured VO_2max (mL \cdot kg $^{-1}$ \cdot min $^{-1}$) in relation to the predicted VO_2max (mL \cdot kg $^{-1}$ \cdot min $^{-1}$) using the Matthews et al. (1999) equation, extrapolation to RPE 19 and 20, and the RWT original equation. The dashed line represents a referent to the measured VO_2max .

was significantly less than predicted VO_2max using extrapolation to RPE 20 ($p < .05$).

Figure 3 shows the relationship between the measured and the predicted VO_2max using the non-maximal tests. The R values indicated a moderate to good correlation between the measured and the predicted VO_2max , with values of R 2 = 0.75, 0.85, 0.83, and 0.73 for the Matthews et al. (1999) equation, extrapolation to RPE 19, extrapolation to RPE 20, and the RWT equation, respectively. There was a good correlation between the measured and the predicted VO_2max values using extrapolation to RPE 19 and RPE 20. The predicted VO_2max values using either the Matthews et al. (1999) equation or the RWT equation (Kline, et al., 1987) were moderately correlated with the measured VO_2max .

The means of the VO_2 at VT and VO_2 at the LP and EQ stages of the TT, and at the RPE13, 14, and 15 were compared. Tukey's *post-hoc* test revealed that there was no significant difference between the VO_2 in mL \times kg $^{-1}$ \times min $^{-1}$ at VT (26.9 ± 5.51), the LP (25.7 ± 5.32), and EQ (29.4 ± 5.30) stages of the TT, at RPE13 (25.9 ± 5.07) and RPE14 (28.5 ± 5.26). However, the VO_2 at RPE 15 (31.0 ± 5.55) was significantly higher than VO_2 at the VT (Table 3, Figure 4).

The VO_2 at VT was well correlated with VO_2 at the LP (R = 0.76) and EQ (R = 0.77) stages of the TT, and at the RPE13 (R = 0.73), RPE14 (R = 0.71) and RPE15 (R = 0.70) (Figure 4). Although VO_2 at VT versus VO_2 at EQ stage of the TT and RPE14 slightly overestimated VT, they fit very close to the line of identity. Likewise, VO_2 at VT versus VO_2 at LP stage of the TT and RPE13 fit very close to the line of identity, although they slightly underestimated VT. Therefore, the LP and EQ stages of the TT as well as RPE13 and 14 were relatively accurate predictors of VT.




Figure 3. The relationship between the measured VO_{2max} ($mL \cdot kg^{-1} \cdot min^{-1}$) and the predicted VO_{2max} ($mL \cdot kg^{-1} \cdot min^{-1}$) using the Matthews et al. (1999) equation, extrapolation to RPE19 and RPE20, and the RWT original equation.

Table 3. Means (\pm SD) $VO_{2@VT}$ ($mL \cdot kg^{-1} \cdot min^{-1}$), $VO_{2@LP}$ and EQ stages of the TT, and the $VO_{2@RPE13}$, $RPE14$, and $RPE15$ ($mL \cdot kg^{-1} \cdot min^{-1}$)

	Mean \pm SD	r	R ²
$VO_{2@VT}$	26.9 \pm 5.51		
TT			
VT @ LP	25.7 \pm 5.32	0.76	0.58
VT @ EQ	29.4 \pm 5.30	0.76	0.59
RPE			
VT @ RPE13	25.9 \pm 5.07	0.74	0.54
VT @ RPE14	28.5 \pm 5.26	0.71	0.51
VT @ RPE15	31.0 \pm 5.55*	0.70	0.49

Note. *Significantly different than $VO_{2@VT}$ ($p < .05$).




Figure 4. Mean (\pm SE) values for $VO_{2@}$ at the VT ($mL \cdot kg^{-1} \cdot min^{-1}$) in relation to the $VO_{2@}$ at the LP and EQ stages of the TT, and at the RPE 13,14, and 15. The dashed line represents a referent to the measured VT.




Figure 5. The relationship between the VO_2 at VT (mL·kg⁻¹·min⁻¹) and the VO_2 during the LP and EQ stages of the TT, and the VO_2 at each of RPE 13, RPE 14, and RPE 15 (mL·kg⁻¹·min⁻¹).

Discussion and conclusion

The overall results of the current study suggest that sub-maximal tests, and even non-exercise tests, can be used as relatively accurate methods for predicting VO_2 max and VT, at least in healthy, sedentary individuals. The primary findings showed that there were no significant differences between

the measured and the predicted VO_2 max using the RWT equation (Kline, et al., 1987) and extrapolation to RPE 19. Extrapolation to RPE 20 overestimated VO_2 max by about 12%. The LP and EQ stage of the TT as well as RPE13 and RPE 14 can be used as predictors of VT. The secondary finding of this study found that a non-exercise model using the Mathews et al. (1999) equation could be used to predict VO_2 max and characterize fitness level with an error of about 5%. There were no significant differences between the measured and the predicted VO_2 max using either the extrapolation to RPE 19, the RWT equation (Kline, et al., 1987) or the Mathews et al. (1999) equation.

The results of the current study confirmed that the TT can be used to accurately predict VT. Although VO_2 at VT was not significantly different than VO_2 at either the LP or EQ stages of the TT, the EQ stage more closely correlated with VT. These results are in accordance with previous studies that have also observed that the EQ stage of the TT can be used as a surrogate of VT in different populations (Foster, et al., 2018). We conclude that when subjects were at the LP or EQ stage of the TT, they were very close to VT.

Interestingly, when the 6-20 RPE scale was used in conjunction with the TT, the results showed the comparability of the TT (LP and EQ) and RPE 13 and 14 for predicting VT. The VO_2 at the EQ stage of the TT as well as at RPE 14 were close to the measured VO_2 at VT. Exercising at the LP stage of the TT, as well as RPE 13, slightly underestimated VT. However, exercising at RPE 15 or the NEG stage of the TT significantly overestimated VT. The current results generally support the results of Purvis and Cukiton (1981), which indicated a strong correlation between the VO_2 at VT and VO_2 at RPE 13. The current results indicated that the estimation procedure for VO_2max prediction by extrapolating the measured RPEs ≤ 15 to RPE 19 yielded a prediction that was not significantly different to the measured VO_2max . The findings are in agreement with previous studies in healthy participants (Faulkner, et al., 2009; Faulkner & Eston, 2007; Lambrick, et al., 2009) and competitive cyclists (Coquart, et al., 2012b). However, two studies (Faulkner, et al., 2009; Lambrick, et al., 2009) found that there was no significant difference between the actual and the predicted VO_2max when the VO_2 :RPE ≤ 13 values were extrapolated to either RPE 19 or RPE 20. Conversely, the current study showed that extrapolation to RPE 20 overestimated VO_2max by $4.5 \text{ ml} \times \text{kg} \times \text{min}^{-1}$ (~12%). This finding is in opposition to previous studies in clinical participants (Coquart, et al., 2009), and in healthy and clinical participants (Al-Rahamneh & Eston, 2011; Al-Rahamneh, et al., 2011), which found that extrapolation to RPE=20 yielded the best fit with the measured VO_2max . The opposing finding observed in the current study may be due to the differences in the exercise test protocol and the exercise modality (treadmill vs. cycle ergometer). The other possible explanation is the perceptual end point of extrapolation (RPE 20). Although the RPE 20 is theoretically considered the point of maximal effort or exhaustion, a submaximal RPE, specifically RPE=19, is more often observed (Demello, Cureton, Boineau, & Singh, 1987; Coquart, et al., 2012a; Eston, Faulkner, St Clair Gibson, Noakes, & Parfitt, 2007). In this regard, it has earlier been suggested that the brain increases RPE consistently with the residual time to the end of the test, and that the time to fatigue relates to the reasonably tolerated maximal RPE (i.e. RPE 19 rather than RPE 20) (Coquart, et al, 2012a; Eston, et al., 2007; Noakes, 2004). The current results are in agreement with previous studies (Eston, et al., 2012; Smith, Eston, Norton, & Parfitt, 2015), which revealed that predicting VO_2max by extrapolating RPE: VO_2 values to RPE=19 is more accurate compared to RPE=20.

The current study revealed the predicted VO_2max using the RWT equation was not significantly different than the measured VO_2max . The

results yielded a strong correlation ($R= 0.71$) and low standard error of estimate ($\text{SEE} = 4.6 \text{ ml} \times \text{kg}^{-1} \times \text{min}^{-1}$) between the actual and the predicted VO_2max . The current results are in accordance with the finding of Kline et al. (1987), which showed a strong correlation between the measured and the predicted VO_2max for validation and cross validation groups ($R = 0.88$ and 0.88 , respectively), and the $\text{SEE} = 5.0$ and $4.4 \text{ ml} \times \text{kg}^{-1} \times \text{min}^{-1}$, respectively. The better results from Kline et al. (1987) study may be attributable to a large sample size (343 subjects) and a wider range of subject's VO_2max . Another possible reason could be the testing procedure that allowed the subjects to do several walks with the two fastest walks being used. Although the age-specific equation developed by Dolgener et al. (1994) might be more appropriate for many of the participants in the current study, it is only applicable for a narrow range of ages (18–30 years), which limits its use, and makes the original equation more attractive. However, the original equation includes HR during the last part of the 1-mile walk, which can be highly variable in some conditions, such as in patients on HR altering medications. Therefore, there is a need to find an alternative variable to HR. Additionally, further research needs to be done to find accurate alternative protocols for the RWT. Recently Cress et al. (2015) have shown promising results relative to predicting both VO_2max and VT using 6-minute walk distance and terminal RPE (as a replacement for the HR used in the RWT). Whether this approach will prove useful relative to adapting the RWT remains to be documented.

An impressive secondary result for the current study showed that a non-exercise VO_2max prediction model could be used with reasonable accuracy. The results showed that there was no significant difference between the measured and the predicted VO_2max using the equation of Matthews et al. (1999) ($R = 0.76$, $\text{SEE} = 4.3 \text{ ml} \times \text{kg}^{-1} \times \text{min}^{-1}$). This finding is in accordance with the results of Matthews et al. (1999), who found a stronger correlation but a slightly larger SEE than the current study ($r = 0.86$, $\text{SEE} = 5.7 \text{ ml} \times \text{kg}^{-1} \times \text{min}^{-1}$). The stronger correlation was arguably because of the very large sample size ($N = 799$, healthy adults) with a very wide range of ages (19–79 years). Further studies would be useful in determining the accuracy of using this equation in different populations.

The present study has some technical limitations. For the TT, frequent concerns were the subjects' ability to correctly determine the EQ stage and reading a relatively long and novel passage, requiring about ~30 seconds, the "Rainbow Passage", within a short duration (two minutes) exercise stage. A recent study (Schroeder, Foster, Porcari, & Mikat, 2017) found that longer speech passages, such as the "Rainbow Passage", better predicted VT and RCT, compared to shorter (30

words) speech passages, using the EQ and NEG stages of the TT. Another study (Foster, et al., 2018) pointed out that stage duration affected the power output when trying to identify the VT and RCT, but had no effect on other physiological markers (HR, VO_2 , RPE). Further research is needed to indicate the stage duration that optimally predicts VT and RCT, using shorter and more familiar speech passages. In using the Borg 6-20 RPE scale, some subjects found it difficult to identify appropriate RPE at the end of each stage. For this reason, it might be ideal to have subjects perform several trails in order to optimize the prediction of VO_{2max} . However, both the TT and RPE are attractive because both are very

robust and intuitive. Protocols that require multiple trials to achieve better results would be somewhat self-defeating, as the purpose of sub-maximal tests is to simplify the process of exercise evaluation. The present study compared the accuracy of three non-maximal tests, the TT, RPE extrapolation, and the RWT, for predicting VO_{2max} and VT. Additionally, the non-exercise prediction models are of comparable accuracy to the non-maximal exercise tests when the exercise testing is not feasible. Future studies are required to further support the present results by examining the accuracy of these non-maximal tests in different populations.

References

- Al-Rahamneh, H.Q., & Eston, R.G. (2011). Prediction of peak oxygen consumption from the ratings of perceived exertion during a graded exercise test and ramp exercise test in able-bodied participants and paraplegic persons. *Archives of Physical Medicine and Rehabilitation*, 92, 277-283.
- Al-Rahamneh, H.Q., Faulkner, J.A., Byrne, C., & Eston, R.G. (2011). Prediction of peak oxygen uptake from ratings of perceived exertion during arm exercise in able-bodied and persons with poliomyelitis. *Spinal Cord*, 49(1), 131-135.
- American College of Sports Medicine. (2017). *ACSM's Guidelines for Exercise Testing and Prescription* (10th ed.). Philadelphia, PA: Wolters Kluwer/ Lippincott Williams and Wilkins.
- Borg, G.A. (1982). Psychophysical bases of perceived exertion. *Medicine and Science in Sports and Exercise*, 14(5), 377-381.
- Chung, P.K., Zhao, Y., Liu, J.D., & Quach, B. (2015). A brief note on the validity and reliability of the rating of perceived exertion scale in monitoring exercise intensity among Chinese older adults in Hong Kong. *Perceptual and Motor Skills*, 121(3), 805-809.
- Coquart, J.B., Dufour, Y., Gros Lambert, A., Matran, R., & Garcin, M. (2012a). Relationships between psychological factors, RPE and time limit estimated by teleoanticipation. *The Sport Psychologist*, 26(3), 359-374.
- Coquart, J.B., Eston, R., Nycz, M., Grosbois, J.M., Garcin, M. (2012b). Estimation of maximal oxygen uptake from ratings of perceived exertion elicited during sub-maximal tests in competitive cyclists. *Archives of Medical Science*, 171(2), 165-172.
- Coquart, J.B., Garcin, M., Parfitt, G., Tourny-Chollet, C., & Eston, R.G. (2014). Prediction of maximal or peak oxygen uptake from ratings of perceived exertion. *Sports Medicine*, 44(5), 563-578.
- Coquart, J.B., Lemaire, C., Dubart, A.E., Douillard, C., Luttenbacher, D.P., Wibaux, F., & Garcin, M. (2009). Prediction of peak oxygen uptake from sub-maximal ratings of perceived exertion elicited during a graded exercise test in obese women. *Psychophysiology*, 46(6), 1150-1153.
- Cress, M., Porcari, J.P., Brown, H., Foster, C., Greany, J.F., French, K., Schmidt, K., Donahue, M. (2015). Use of 6-minute walk distance and Rating of Perceived Exertion to predict maximal aerobic capacity and ventilatory threshold in cardiac rehabilitation. *Journal of Cardiopulmonary Rehabilitation and Prevention*, 35, 288S.
- Demello, J.J., Cureton, K.J., Boineau, R.E., & Singh, M.M. (1987). Ratings of perceived exertion at the lactate threshold in trained and untrained men and women. *Medicine and Science in Sports and Exercise*, 19(4), 354-362.
- Dolgener, F.A., Hensley, L.D., Marsh, J.J., & Fielstul, J.K. (1994). Validation of the Rockport Fitness Walking Test in college males and females. *Research Quarterly for Exercise and Sport*, 65(2), 152-158.
- Eston, R. (2012). Use of ratings of perceived exertion in sports. *International Journal of Sports Physiology and Performance*, 7, 175-182.
- Eston, R., Evans, H., Faulkner, J., Lambrick, D., Al-Rahamneh, H., & Parfitt, G. (2012). A perceptually regulated, graded exercise test predicts peak oxygen uptake during treadmill exercise in active and sedentary participants. *European Journal of Applied Physiology*, 112(10), 3459-3468.
- Eston, R.G., Faulkner, J.A., Mason, E.A., & Parfitt, G. (2006). The validity of predicting maximal oxygen uptake from perceptually regulated graded exercise tests of different durations. *European Journal of Applied Physiology*, 97(5), 535-541.
- Eston, R., Faulkner, J., St Clair Gibson, A., Noakes, T., & Parfitt, G. (2007). The effect of antecedent fatiguing activity on the relationship between perceived exertion and physiological activity during a constant load exercise task. *Psychophysiology*, 44(5), 779-786.

- Eston, R.G., Lamb, K.L., Parfitt, G., & King, N. (2005). The validity of predicting maximal oxygen uptake from a perceptually-regulated graded exercise test. *European Journal of Applied Physiology*, 94(3), 221-227.
- Eston, R., Lambrick, D., Sheppard, K., & Parfitt, G. (2008). Prediction of maximal oxygen uptake in sedentary males from a perceptually regulated, sub-maximal graded exercise test. *Journal of Sports Sciences*, 26(2), 131-139.
- Faulkner, J., & Eston, R. (2007). Overall and peripheral ratings of perceived exertion during a graded exercise test to volitional exhaustion in individuals of high and low fitness. *European Journal of Applied Physiology*, 101(5), 613-620.
- Faulkner, J.A., Lambrick, D.M., Parfitt, G., Rowlands, A.V., & Eston, R.G. (2009). Prediction of maximal oxygen uptake from the Åstrand-Ryhming nomogram and ratings of perceived exertion. In G. Atkinson & T. Reilly (Eds.), *Contemporary sport, leisure and ergonomics* (pp. 197-214). London, GB: Routledge.
- Faulkner, J., Parfitt, G., & Eston, R. (2007). Prediction of maximal oxygen uptake from the ratings of perceived exertion and heart rate during a perceptually-regulated sub-maximal exercise test in active and sedentary participants. *European Journal of Applied Physiology*, 101, 397-407.
- Foster, C., & Cotter, H. (2006). Blood lactate, respiratory and heart rate markers of the capacity for sustained exercise. In C. Foster & P.J. Maud (Eds.), *Physiological assessment of human fitness* (pp. 63-75). Champaign, IL: Human Kinetics.
- Foster, C., Porcari, J.P., Doro, K., Dubiel, J., Engen, M., Kolman, D., Ault, S., & Xiong, S. (2018). Exercise prescription when there is no exercise test: The Talk Test. *Kinesiolog*, 50(2), 333-348.
- Jackson, A.S., Blair, S.N., Mahar, M.T., Wier, L.T., Ross, R.M., & Stuteville, J.E. (1990). Prediction of functional aerobic capacity without exercise testing. *Medicine and Science in Sports and Exercise*, 22(6), 863-870.
- Kline, G.M., Porcari, J.P., Hintermeister, R., Freedson, P.S., Ward, A., Mccarron, R.F., ..., & Rippe, J.M. (1987). Estimation of VO_{2max} from a one-mile track walk, gender, age, and body weight. *Medicine and Science in Sports and Exercise*, 19(3), 253-259.
- Lambrick, D.M., Faulkner, J.A., Rowlands, A.V., & Eston, R.G. (2009). Prediction of maximal oxygen uptake from submaximal ratings of perceived exertion and heart rate during a continuous exercise test: The efficacy of RPE 13. *European Journal of Applied Physiology*, 107(1), 1-9
- Matthews, C.E., Heil, D.P., Freedson, P.S., & Pastides, H. (1999). Classification of cardiorespiratory fitness without exercise testing. *Medicine and Science in Sports and Exercise*, 31(3), 486-493.
- Mezzani, A., Hamm, L.F., Jones, A.M., McBride, P.E., Moholdt, T., Stone, J.A., ..., & Williams, M.A. (2012). Aerobic exercise intensity assessment and prescription in cardiac rehabilitation. *Journal of Cardiopulmonary Rehabilitation and Prevention*, 32, 327-350.
- Myers, J., Voodi, L., Umann, T., & Froelicher, V.F. (2000). A survey of exercise testing: Methods, utilization, interpretation, and safety in the VAHCS. *Journal of Cardiopulmonary Rehabilitation*, 20, 251-258.
- Noakes, T.D. (2004). Linear relationship between the perception of effort and the duration of constant load exercise that remains. *Journal of Applied Physiology*, 96(4), 1571-1573.
- Noonan, V., & Dean, E. (2000). Submaximal exercise testing: Clinical application and interpretation. *Physical Therapy*, 80, 782-807.
- Parfitt, G., Evans, H., & Eston, R. (2012). Perceptually regulated training at RPE13 is pleasant and improves physical health. *Medicine and Science in Sports and Exercise*, 44(8), 1613-1618.
- Purvis, J.W., & Cukiton, K.J. (1981). Ratings of perceived exertion at the anaerobic threshold. *Ergonomics*, 24(4), 295-300.
- Ross, R.M., & Jackson, A.S. (1990). *Exercise concepts, calculations, and computer applications*. Carmel, IN: Benchmark Press.
- Schneider, D.A., Phillips, S.E., & Stoffolano, S. (1993). The simplified V-slope method of detecting the gas exchange threshold. *Medicine and Science in Sports and Exercise*, 25(10), 1180-1184.
- Schroeder, M.M., Foster, C., Porcari, J.P., & Mikat, R.P. (2017). Effects of speech passage length on accuracy of predicting metabolic thresholds using the Talk Test. *Kinesiology*, 49, 9-14.
- Smith, A.E., Eston, R.G., Norton, B., & Parfitt, G. (2015). A perceptually-regulated exercise test predicts peak oxygen uptake in older active adults. *Journal of Aging and Physical Activity*, 23(2), 205-211.

Submitted: October 25, 2018
Accepted: November 26, 2019
Published Online First: February 27, 2020

Correspondence to:
Carl Foster, Ph.D.
Department of Exercise and Sport Science
133 Mitchell Hall
University of Wisconsin-La Crosse
La Crosse, WI 54601, USA
Phone: 608-792-2170
E-mail: cfoster@uwlax.edu