

Bounds for Invariance Pressure^{*}

Fritz Colonius^{a,*}, João A. N. Cossich^b, Alexandre J. Santana^c

^a *Institut für Mathematik, Universität Augsburg, Universitätsstr. 14, 86159 Augsburg, Germany*

^b *Department of Mathematics, State University of Maringá, Av. Colombo, 5790, 87020-900 Maringá, Brazil*

^c *Department of Mathematics, State University of Maringá, Av. Colombo, 5790, 87020-900 Maringá, Brazil*

Abstract

This paper provides an upper bound for the invariance pressure of control sets with nonempty interior and a lower bound for sets with finite volume. In the special case of the control set of a hyperbolic linear control system on \mathbb{R}^d this yields an explicit formula. Further applications to linear control systems on Lie groups and to inner control sets are discussed.

Keywords: Invariance pressure, invariance entropy, control sets

1. Introduction

2 The notion of invariance pressure generalizes invariance entropy by adding
3 potentials f on the control range. It has been introduced and analyzed in
4 Colonius, Cossich and Santana [5, 6]. Zhong and Huang [19] show that in-
5 variance pressure can be characterized as a dimension-like notion within the
6 framework due to Pesin. A basic reference for invariance entropy is Kawan's
7 monograph [17]; here also the relation to minimal data rates is explained
8 which gives the main motivation from applications. Further references in-
9 clude the seminal paper Nair, Evans, Mareels and Moran [18] as well as
10 Colonius and Kawan [8] and Da Silva and Kawan [12], [13]. In the latter

^{*}This work was partly supported by Network NT8 OEA ICTP and PICME/CAPES.

^{*}Corresponding author

Email addresses: fritz.colonius@math.uni-augsburg.de (Fritz Colonius),
joaocossich@hotmail.com (João A. N. Cossich), ajsantana@uem.br (Alexandre J.
Santana)

11 paper, robustness properties in the hyperbolic case are proved. Huang and
 12 Zhong [15] show that several generalized notions of invariance entropy fit into
 13 the dimension-theoretic framework.

14 The main results of the present paper are upper and lower bounds for the
 15 invariance pressure of compact subsets K in a control set D with nonvoid
 16 interior and compact closure as well as a formula for the invariance pressure
 17 in the case of hyperbolic linear control systems on \mathbb{R}^d where a unique control
 18 set with nonvoid interior exists. We also give applications for inner control
 19 sets and for certain linear systems on Lie groups. Invariance entropy of these
 20 systems has been analyzed by Da Silva [10].

21 Section 2 collects results on linearization of control systems and on the
 22 notion of invariance pressure. Upper and lower bounds for invariance pressure
 23 are given in Sections 3 and 4, respectively. Section 5 presents a formula for
 24 the invariance pressure of linear control systems on \mathbb{R}^d and Section 6 discusses
 25 applications to linear systems on Lie groups and for inner control sets.

26 2. Preliminaries

27 In this section we first recall basic notions for control systems on manifolds
 28 and their linearization. Then the concepts of invariance pressure and outer
 29 invariance pressure are presented as well as some of their properties.

30 2.1. Control systems and linearization

31 Throughout the paper M will denote a smooth manifold, that is, a con-
 32 nected, second-countable, topological Hausdorff manifold endowed with a C^∞
 33 differentiable structure. A continuous-time **control system** on a smooth
 34 manifold M is a family of ordinary differential equations

$$\dot{x}(t) = F(x(t), \omega(t)), \omega \in \mathcal{U}, \tag{1}$$

35 on M which is parametrized by measurable functions $\omega : \mathbb{R} \rightarrow \mathbb{R}^m$, $\omega(t) \in$
 36 $U \subset \mathbb{R}^m$ almost everywhere, called **controls** forming the set \mathcal{U} of **admis-**
 37 **sible control functions**, where $U \subset \mathbb{R}^m$ is a compact set, the **control**
 38 **range**. The function $F : M \times \mathbb{R}^m \rightarrow TM$ is a C^1 -map such that for each
 39 $u \in U$, $F_u(\cdot) := F(\cdot, u)$ is a smooth vector field on M . For each $x \in M$ and
 40 $\omega \in \mathcal{U}$, we suppose that there exists an unique solution $\varphi(t, x, \omega)$ which is
 41 defined for all $t \in \mathbb{R}$. We usually refer to the solution $\varphi(\cdot, x, \omega)$ as a **tra-**
 42 **jectory** of x with control function ω and write $\varphi_t(x, \omega) = \varphi(t, x, \omega)$ where

43 convenient. We need several notions characterizing controllability properties
 44 of subsets of the state space M of system (1). For $x \in M$ and $t > 0$, the
 45 **set of points reachable from x up to time t** and the **set of points**
 46 **controllable to x within time t** are given by

$$\mathcal{O}_{\leq t}^+(x) := \{y \in M; \text{ there are } s \in [0, t] \text{ and } \omega \in \mathcal{U} \text{ with } \varphi(s, x, \omega) = y\},$$

47 and

$$\mathcal{O}_{\leq t}^-(x) := \{y \in M; \text{ there are } s \in [0, t] \text{ and } \omega \in \mathcal{U} \text{ with } \varphi(s, y, \omega) = x\},$$

48 respectively. The **positive** and **negative orbits from $x \in M$** are

$$\mathcal{O}^+(x) := \bigcup_{t>0} \mathcal{O}_{\leq t}^+(x) \text{ and } \mathcal{O}^-(x) := \bigcup_{t>0} \mathcal{O}_{\leq t}^-(x),$$

49 respectively.

50 A key concept of this paper is presented in the following definition.

51 **Definition 1.** *A subset D of M is a **control set** if*

- 52 (i) *for each $x \in D$, there exists $\omega \in \mathcal{U}$ with $\varphi(\mathbb{R}_+, x, \omega) \subset D$ (controlled*
- 53 *invariance);*
- 54 (ii) *for each $x \in D$ one has $D \subset \overline{\mathcal{O}^+(x)}$ (approximate controllability);*
- 55 (iii) *D is maximal with these properties.*

56 If for all $t > 0$ the sets $\mathcal{O}_{\leq t}^-(x)$ and $\mathcal{O}_{\leq t}^+(x)$ have nonempty interior, we
 57 say that system (1) is **locally accessible from $x \in M$** . We are mainly
 58 interested in control sets with nonvoid interior which are locally accessible
 59 from all $x \in \text{int}D$, since here a general theory can be developed. In particular,
 60 they enjoy the property $\text{int}D \subset \mathcal{O}^+(x)$ for all $x \in D$, cf. Colonius and
 61 Kliemann [9, Lemma 3.2.13].

62 Next we recall some basic concepts and results on linearization of a control
 63 system on a smooth Riemannian manifold (M, g) , cf. Kawan [17].

64 **Definition 2.** *For a control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$ the linearized*
 65 *system is given by*

$$\frac{Dz}{dt}(t) = A(t)z(t) + B(t)\mu(t), \quad \mu \in L^\infty(\mathbb{R}, \mathbb{R}^m), \quad (2)$$

66 where $A(t) := \nabla F_{\omega(t)}(\varphi(t, x, \omega))$ and $B(t) := D_2F(\varphi(t, x, \omega), \omega(t))$.

67 The derivative on the left-hand side of (2) is the covariant derivative of $z(\cdot)$
68 along $\varphi(\cdot, x, \omega)$ and D_2 is the derivative with respect to second component. A
69 solution of (2) corresponding to $\mu \in L^\infty(\mathbb{R}, \mathbb{R}^m)$ with initial value $\lambda \in T_x M$ is
70 a locally absolutely continuous vector field $z = \phi^{x,\omega}(\cdot, \lambda, \mu) : \mathbb{R} \rightarrow TM$ along
71 $\varphi(\cdot, x, \omega)$ with $z(0) = \lambda$, satisfying the differential equation (2) for almost all
72 $t \in \mathbb{R}$.

73 The next proposition presents some properties of linearized systems.

74 **Proposition 3.** *Let $(\omega(\cdot), \varphi(\cdot, x, \omega))$ be a control-trajectory pair with corre-*
75 *sponding linearization (2). Then the following statements hold:*

76 (i) *For all $\tau > 0$ the mapping $\varphi_\tau : M \times L^\infty([0, \tau], \mathbb{R}^m) \rightarrow M, (x, \omega) \mapsto$*
77 *$\varphi(\tau, x, \omega)$ is continuously (Fréchet) differentiable.*

78 (ii) *For every initial value $\lambda \in T_x M$ and every $\mu \in L^\infty(\mathbb{R}, \mathbb{R}^m)$ there*
79 *exists a unique solution $\phi^{x,\omega}(\cdot, \lambda, \mu) : \mathbb{R} \rightarrow TM$ of (2) satisfying*

$$\phi^{x,\omega}(0, \lambda, \mu) = \lambda, \phi^{x,\omega}(t, \lambda, \mu) = D\varphi_t(x, \omega)(\lambda, \mu), t \in \mathbb{R}, \quad (3)$$

80 for $(\lambda, \mu) \in T_x M \times L^\infty(\mathbb{R}, \mathbb{R}^m)$, where D stands for the total derivative of $\varphi_t :$
81 $M \times L^\infty(\mathbb{R}, \mathbb{R}^m) \rightarrow M$ which consists of the derivative $d_x \varphi_t(\cdot, \omega) : T_x M \rightarrow$
82 $T_{\varphi(t,x,\omega)} M$ in the first, and the Fréchet derivative of $\varphi_t(x, \cdot) : L^\infty(\mathbb{R}, \mathbb{R}^m) \rightarrow$
83 $T_{\varphi(t,x,\omega)} M$ in the second component.

84 (iii) *For every $\tau > 0$ the map $\phi^{x,\omega}(\tau, \cdot, \cdot) : T_x M \times L^\infty([0, \tau], \mathbb{R}^m) \rightarrow$*
85 *$T_{\varphi(\tau,x,\omega)} M$ is linear and continuous.*

86 (iv) *For each $t \in \mathbb{R}$ abbreviate $\phi_t^{x,\omega} := \phi^{\varphi(t,x,\omega), \omega(t+\cdot)}$. Then for all $t, s \in \mathbb{R}$,*
87 *$\lambda \in T_x M$ and $\mu \in L^\infty(\mathbb{R}, \mathbb{R}^m)$,*

$$\phi_s^{x,\omega}(t, \phi^{x,\omega}(s, \lambda, \mu), \Theta_s \mu) = \phi^{x,\omega}(t+s, \lambda, \mu),$$

88 and, in particular,

$$\phi_s^{x,\omega}(t, \phi^{x,\omega}(s, \lambda, \mathbf{0}), \mathbf{0}) = \phi^{x,\omega}(t+s, \lambda, \mathbf{0}).$$

89 Now we present the notion of regularity of a control-trajectory pair.

90 **Definition 4.** *Consider some $(x, \omega, \tau) \in M \times \mathcal{U} \times (0, \infty)$ and let $y :=$*
91 *$\varphi(\tau, x, \omega)$. The linearization along $(\omega(\cdot), \varphi(\cdot, x, \omega))$ is **controllable on** $[0, \tau]$*
92 *if for each $\lambda_1 \in T_x M$ and $\lambda_2 \in T_y M$ there exists $\mu \in L^\infty([0, \tau], \mathbb{R}^m)$ with*

$$\phi^{x,\omega}(\tau, \lambda_1, \mu) = \lambda_2.$$

93 *In this case, we say that the control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$ is **regular***
94 *on $[0, \tau]$.*

95 A control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$ is called τ -periodic, $\tau \geq 0$, if
 96 $(\varphi(t + \tau, x, \omega), \omega(t + \tau)) = (\varphi(t, x, \omega), \omega(t))$ for all $t \in \mathbb{R}$, or equivalently
 97 if $\varphi(\tau, x, \omega) = x$ and $\Theta_\tau \omega = \omega$, where $(\Theta_\tau \omega)(t) = \omega(t + \tau)$, $t \in \mathbb{R}$, is the
 98 τ -shift on \mathcal{U} . A periodic regular control-trajectory pair enjoys the property
 99 described in the following proposition (cf. [17, Proposition 1.30]).

100 **Proposition 5.** *Let $(\omega(\cdot), \varphi(\cdot, x, \omega))$ be a τ -periodic control-trajectory pair*
 101 *which is regular on $[0, \tau]$. Then there exists $C > 0$ such that for every $\lambda \in$*
 102 *$T_x M$ there is $\mu \in L^\infty([0, \tau], \mathbb{R}^m)$ with $\phi^{x, \omega}(\tau, \lambda, \mu) = 0_x$ and $\|\mu\|_{[0, \tau]} \leq C|\lambda|$,*
 103 *where $\|\cdot\|_{[0, \tau]}$ denotes the L^∞ -norm.*

104 For a τ -periodic control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$ the Floquet or
 105 Lyapunov exponents are given by

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \|\phi^{x, \omega}(t, \lambda, \mathbf{0})\| = \lim_{n \rightarrow \infty} \frac{1}{n\tau} \log \|\phi^{x, \omega}(n\tau, \lambda, \mathbf{0})\|, \lambda \in T_x M. \quad (4)$$

106 These limits exist and the Lyapunov exponents are denoted by $\rho_1(\omega, x), \dots,$
 107 $\rho_r(\omega, x)$ with $1 \leq r := r(\omega, x) \leq d = \dim M$. The Lyapunov spaces are given
 108 by

$$L_j(\omega, x) = \left\{ \lambda \in T_x M; \lim_{t \rightarrow \pm\infty} \frac{1}{t} \log \|\phi^{x, \omega}(t, \lambda, \mathbf{0})\| = \rho_j(\omega, x) \right\}, j = 1, \dots, r,$$

109 with dimensions $d_j(\omega, x)$. They yield the decomposition

$$T_x M = L_1(\omega, x) \oplus \dots \oplus L_r(\omega, x).$$

110 2.2. Invariance pressure

111 In this subsection we recall the concepts of invariance and outer invariance
 112 pressure introduced in Colonius, Cossich and Santana [5, 6] and some of their
 113 properties.

114 A pair (K, Q) of nonempty subsets of a smooth Riemannian manifold M
 115 is called **admissible** if K is compact and for each $x \in K$ there exists $\omega \in \mathcal{U}$
 116 such that $\varphi(\mathbb{R}_+, x, \omega) \subset Q$. For an admissible pair (K, Q) and $\tau > 0$, a
 117 (τ, K, Q) -**spanning set** \mathcal{S} is a subset of \mathcal{U} such that for all $x \in K$ there is
 118 $\omega \in \mathcal{S}$ with $\varphi(t, x, \omega) \in Q$ for all $t \in [0, \tau]$. Denote by $C(U, \mathbb{R})$ the set of
 119 continuous function $f : U \rightarrow \mathbb{R}$ which we call **potentials**.

120 For a potential $f \in C(U, \mathbb{R})$ denote $(S_\tau f)(\omega) := \int_0^\tau f(\omega(t))dt$ and

$$a_\tau(f, K, Q) := \inf \left\{ \sum_{\omega \in \mathcal{S}} e^{(S_\tau f)(\omega)}; \mathcal{S} (\tau, K, Q)\text{-spanning} \right\}.$$

121 The **invariance pressure** $P_{inv}(f, K, Q)$ of control system (1) is defined by

$$P_{inv}(f, K, Q) := \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log a_\tau(f, K, Q).$$

122 Given an admissible pair (K, Q) such that Q is closed in M and a metric
 123 ϱ on M which is compatible with the Riemannian structure, we define the
 124 **outer invariance pressure** of (K, Q) by

$$P_{out}(f, K, Q) := \lim_{\varepsilon \rightarrow 0} P_{inv}(f, K, N_\varepsilon(Q)),$$

125 where $N_\varepsilon(Q) = \{y \in M; \exists x \in Q \text{ with } \varrho(x, y) < \varepsilon\}$ denotes the ε - neigh-
 126 borhood of Q .

127 Note that $-\infty < P_{out}(f, K, Q) \leq P_{inv}(f, K, Q) \leq \infty$ for every admissible
 128 pair (K, Q) and all potentials f . For the potential $f = \mathbf{0}$, this reduces to the
 129 notion of invariance entropy, $P_{inv}(\mathbf{0}, K, Q) = h_{inv}(K, Q)$ and $P_{out}(\mathbf{0}, K, Q) =$
 130 $h_{out}(K, Q)$, cf. Kawan [17].

131 The next proposition presents some properties of the function $P_{inv}(\cdot, K, Q)$
 132 : $C(U, \mathbb{R}) \rightarrow \mathbb{R}$, cf. [6, Proposition 3.4].

133 **Proposition 6.** *The following assertions hold for an admissible pair (K, Q) ,*
 134 *functions $f, g \in C(U, \mathbb{R})$ and $c \in \mathbb{R}$:*

135 (i) $P_{inv}(f, K, Q) \leq P_{inv}(g, K, Q)$ and $P_{out}(f, K, Q) \leq P_{out}(g, K, Q)$ for
 136 $f \leq g$.

137 (ii) $P_{inv}(f + c, K, Q) = P_{inv}(f, K, Q) + c$.

138 (iii) $h_{inv}(K, Q) + \min_{u \in U} f(u) \leq P_{inv}(f, K, Q) \leq h_{inv}(K, Q) + \max_{u \in U} f(u)$.

139 Proposition 6 (iii) shows, in particular, that $P_{inv}(f, K, Q) < \infty$ if and
 140 only if $h_{inv}(K, Q) < \infty$. For general admissible pairs (K, Q) , one cannot
 141 guarantee the existence of finite (τ, K, Q) -spanning sets \mathcal{S} . The following two
 142 remarks discuss the cardinality of spanning sets and relations to properties
 143 of invariance pressure.

144 **Remark 7.** *If there is no countable (τ, K, Q) -spanning set, then $a_\tau(f, K, Q)$
145 $= \infty$ (see Kawan [17, Example 2.3] for an example). If $P_{inv}(f, K, Q) <$
146 ∞ , then $a_\tau(f, K, Q) < \infty$ for every $\tau > 0$. Hence there is a (τ, K, Q) -
147 spanning set with $\sum_{\omega \in \mathcal{S}} e^{(S_\tau f)(\omega)} < \infty$ implying that there can be only count-
148 ably many summands, i.e., there is a countable (τ, K, Q) -spanning set. On
149 the other hand, if for all $\tau > 0$ there is a countable (τ, K, Q) -spanning set,
150 $a_\tau(f, K, Q) = \infty$ is also possible. If every (τ, K, Q) -spanning set \mathcal{S} contains
151 a finite (τ, K, Q) -spanning subset \mathcal{S}' , then*

$$a_\tau(f, K, Q) = \inf \left\{ \sum_{\omega \in \mathcal{S}} e^{(S_\tau f)(\omega)}; \mathcal{S} \text{ finite and } (\tau, K, Q)\text{-spanning} \right\}.$$

152 *This follows, since all summands satisfy $e^{(S_\tau f)(\omega)} > 0$, and hence the sum-
153 mandants in $\mathcal{S} \setminus \mathcal{S}'$ can be omitted. This situation occurs e.g. if Q is open
154 where compactness of K may be used. For the outer invariance entropy one
155 considers $(\tau, K, N_\varepsilon(Q))$ -spanning sets, $\varepsilon > 0$, and hence here it is also suf-
156 ficient to consider finite $(\tau, K, N_\varepsilon(Q))$ -spanning sets. In the definition of
157 inner invariance pressure of discrete time systems, one considers sets which
158 are $(\tau, K, \text{int}Q)$ -spanning. Here again finite spanning sets are sufficient.*

159 **Remark 8.** *The Lipschitz continuity property*

$$|P_{inv}(f, K, Q) - P_{inv}(g, K, Q)| \leq \|f - g\|_\infty \text{ for } f, g \in C(U, \mathbb{R}),$$

160 *holds if $h_{inv}(K, Q) < \infty$. In fact, as seen in Remark 7, in this case there are
161 for every $\tau > 0$ countable (τ, K, Q) -spanning sets \mathcal{S} with $\sum_{\omega \in \mathcal{S}} e^{(S_\tau f)(\omega)} < \infty$.
162 The arguments used in [5, Proposition 13(iii)] to show Lipschitz continuity
163 under the assumption that finite (τ, K, Q) -spanning sets exist, can be ap-
164 plied in this situation observing that the elementary lemma [5, Lemma 12],
165 on which the proof is based, is valid not only for finite but also for infinite
166 sequences: Let $a_i \geq 0, b_i > 0, i \in \mathbb{N}$. Then for all $n \in \mathbb{N}$*

$$\frac{\sum_{i=1}^n a_i}{\sum_{i=1}^n b_i} \geq \min_{i=1, \dots, n} \frac{a_i}{b_i} \geq \inf_{i \in \mathbb{N}} \frac{a_i}{b_i},$$

167 *and one may take the limit for $n \rightarrow \infty$.*

168 The following proposition shows that in the definition of invariance pres-
169 sure we can take the limit superior over times which are integer multiples of
170 some fixed time step $\tau > 0$.

171 **Proposition 9.** *The invariance pressure satisfies for every $\tau > 0$*

$$P_{inv}(f, K, Q) = \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \log a_{n\tau}(f, K, Q) \text{ for all } f \in C(U, \mathbb{R}). \quad (5)$$

Proof. Let $\tau_k \in (0, \infty), k \in \mathbb{N}$, with $\tau_k \rightarrow \infty$. Then for every $k \geq 1$ there exists $n_k \geq 1$ such that $n_k\tau \leq \tau_k \leq (n_k + 1)\tau$ and $n_k \rightarrow \infty$ for $k \rightarrow \infty$. Since $\tilde{f}(u) := f(u) - \min f, u \in U$, is nonnegative, it follows that $a_{\tau_k}(\tilde{f}, K, Q) \leq a_{(n_k+1)\tau}(\tilde{f}, K, Q)$ and consequently $\frac{1}{\tau_k} \log a_{\tau_k}(\tilde{f}, K, Q)$ is less than or equal to $\frac{1}{n_k\tau} \log a_{(n_k+1)\tau}(\tilde{f}, K, Q)$. Hence

$$\begin{aligned} \limsup_{k \rightarrow \infty} \frac{1}{\tau_k} \log a_{\tau_k}(\tilde{f}, K, Q) &\leq \limsup_{k \rightarrow \infty} \frac{1}{n_k\tau} \log a_{(n_k+1)\tau}(\tilde{f}, K, Q) \\ &= \limsup_{k \rightarrow \infty} \frac{n_k + 1}{n_k} \frac{1}{(n_k + 1)\tau} \log a_{(n_k+1)\tau}(\tilde{f}, K, Q) \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \log a_{n\tau}(\tilde{f}, K, Q). \end{aligned}$$

172 This shows that

$$P_{inv}(f - \min f, K, Q) \leq \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \log a_{n\tau}(f - \min f, K, Q).$$

173 Using $a_{n\tau}(\tilde{f}, K, Q) = a_{n\tau}(f, K, Q) - \min f$ and Proposition 6 (ii) we obtain

$$P_{inv}(f, K, Q) \leq \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \log a_{n\tau}(f, K, Q).$$

174 The converse inequality is obvious. ■

175 For the proof of the following proposition see [6, Corollary 15].

176 **Proposition 10.** *Let K_1, K_2 be two compact sets with nonempty interior*
 177 *contained in a control set $D \subset M$. Then (K_1, Q) and (K_2, Q) are admissible*
 178 *pairs and for all $f \in C(U, \mathbb{R})$ we have*

$$P_{inv}(f, K_1, Q) = P_{inv}(f, K_2, Q).$$

179 **3. An upper bound on control sets**

180 Our goal in this section is to obtain an upper bound for the invariance
 181 pressure of a control set. We consider a smooth control system (1) on a
 182 Riemannian manifold (M, g) under our standard assumptions.

183 In the following theorem, given a periodic control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$, the
 184 different Lyapunov exponents at (x, ω) are denoted by $\rho_1(x, \omega), \dots, \rho_r(x, \omega)$,
 185 $r = r(x, \omega)$, with Lyapunov spaces of dimensions $d_1(x, \omega), \dots, d_r(x, \omega)$, re-
 186 spectively.

187 **Theorem 11.** *Let $D \subset M$ be a control set with nonempty interior and com-
 188 pact closure for control system (1). Then for every compact set $K \subset D$
 189 and every set $Q \supset D$, the pair (K, Q) is admissible and for all potentials
 190 $f \in C(U, \mathbb{R})$ the invariance pressure satisfies*

$$P_{inv}(f, K, Q) \leq \inf_{(T, x, \omega)} \left\{ \sum_{j=1}^{r(x, \omega)} \max\{0, d_j(x, \omega) \rho_j(x, \omega)\} + \frac{1}{T} \int_0^T f(\omega(s)) ds \right\},$$

191 where the infimum is taken over all $(T, x, \omega) \in (0, \infty) \times \text{int}D \times \mathcal{U}$ such that
 192 the control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$ is T -periodic and regular and the
 193 values $\omega(t), t \in [0, T]$, are in a compact subset of $\text{int}U$.

194 **Remark 12.** *For $f \equiv 0$, the statement of the theorem reduces to Kawan [16,
 195 Theorem 4.3],*

$$h_{inv}(K, Q) = P_{inv}(\mathbf{0}, K, Q) \leq \inf_{(T, x, \omega)} \left\{ \sum_{j=1}^{r(x, \omega)} \max\{0, d_j(x, \omega) \rho_j(x, \omega)\} \right\}.$$

196 **Proof.** The theorem will follow by an extension of the proof given in [16,
 197 Theorem 4.3] for invariance entropy. First we briefly sketch the construc-
 198 tion in [16, pp. 740-745], then we indicate the new arguments needed for
 199 invariance pressure.

200 By Proposition 10 one can choose K as an arbitrary compact subset of
 201 D with nonvoid interior. Let $(\omega_0(\cdot), \varphi(\cdot, x_0, \omega_0))$ be a T -periodic and regular
 202 control-trajectory pair as in the statement of the theorem. Then fix real
 203 numbers $\varepsilon > 0$ and

$$S_0 > \sum_{j=1}^r \max(0, d_j \rho_j),$$

204 where $d_j = d_j(x_0, \omega_0)$ and $\rho_j = \rho_j(x_0, \omega_0)$, $j = 1, \dots, r$. An ingenious and
 205 lengthy construction in [16] provides a closed ball $K := \text{cl}(B_{b_0}(x_0)) \subset D$
 206 with radius $b_0 > 0$ centered at x_0 with the following properties: For some
 207 $\tau = kT$, $k \in \mathbb{N}$, and arbitrary $n \in \mathbb{N}$ one finds a set \mathcal{S}_n of $(n\tau, K, Q)$ -spanning
 208 controls $\omega \in \mathcal{S}_n$ satisfying

$$\|\omega - \omega_0\|_{[0, n\tau]} \leq Cb_0\sqrt{d}, \quad (6)$$

209 where $C > 0$ is a constant and $b_0 > 0$ can be taken arbitrarily small (see [16,
 210 formula (4.17)]: the elements of \mathcal{S}_n are n -fold concatenations of the controls
 211 denoted there by u_x). The cardinality $\#\mathcal{S}_n$ of \mathcal{S}_n is bounded by

$$\frac{1}{n\tau} \log \#\mathcal{S}_n \leq S_0 + \varepsilon, \quad (7)$$

212 cf. [16, estimates on middle of p. 745].

213 In order to get a bound for the invariance pressure we need the following
 214 additional arguments: Let $f \in C(U, \mathbb{R})$ be a potential. Since f is defined on
 215 the compact set U , its uniform continuity implies that there exists $\delta > 0$ such
 216 that $|u - v| < \delta$ implies $|f(u) - f(v)| < \varepsilon$. Take $b_0 > 0$ small enough such that
 217 $Cb_0\sqrt{d} < \delta$. By (6) every $\omega \in \mathcal{S}_n$ satisfies $|\omega(t) - \omega_0(t)| \leq \|\omega - \omega_0\|_{[0, n\tau]} < \delta$
 218 for almost all $t \in [0, n\tau]$. Hence it follows that $|f(\omega(t)) - f(\omega_0(t))| < \varepsilon$ for
 219 almost all $t \in [0, n\tau]$.

Now we can estimate

$$\begin{aligned} \frac{1}{n\tau} \log a_{n\tau}(f, K, Q) &\leq \frac{1}{n\tau} \log \sum_{\omega \in \mathcal{S}_n} e^{(S_{n\tau}f)(\omega)} = \frac{1}{n\tau} \log \sum_{\omega \in \mathcal{S}_n} e^{\int_0^{n\tau} f(\omega(t))dt} \\ &= \frac{1}{n\tau} \log \sum_{\omega \in \mathcal{S}_n} e^{\int_0^{n\tau} f(\omega_0(t))dt + \int_0^{n\tau} [f(\omega(t)) - f(\omega_0(t))]dt} \\ &\leq \frac{1}{n\tau} \log \left[\sum_{\omega \in \mathcal{S}_n} e^{\int_0^{n\tau} f(\omega_0(t))dt} \cdot e^{\int_0^{n\tau} \varepsilon dt} \right] \\ &= \frac{1}{n\tau} \log \left(\#\mathcal{S}_n e^{\int_0^{n\tau} f(\omega_0(t))dt} \right) + \frac{1}{n\tau} \log e^{\int_0^{n\tau} \varepsilon dt} \\ &= \frac{1}{n\tau} \log \#\mathcal{S}_n + \frac{1}{n\tau} \int_0^{n\tau} f(\omega_0(t))dt + \varepsilon \\ &< S_0 + \frac{1}{T} \int_0^T f(\omega_0(t))dt + 2\varepsilon. \end{aligned}$$

220 For the last inequality we have used (7) and T -periodicity of ω_0 . By Propo-
 221 sition 9 this implies

$$P_{inv}(f, K, Q) = \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \log a_{n\tau}(f, K, Q) \leq S_0 + \frac{1}{T} \int_0^T f(\omega_0(t)) dt + 2\varepsilon.$$

222 Since $\varepsilon > 0$ can be chosen arbitrarily small and S_0 arbitrarily close to
 223 $\sum_{j=1}^r \max(0, d_j \rho_j)$, the assertion of the theorem follows. ■

224 **Remark 13.** In Kawan [17, Section 5.2] and Da Silva and Kawan [12, Sec-
 225 tion 3.2] one finds more information on regular periodic control-trajectory
 226 pairs.

227 4. A lower bound

228 Again we consider a smooth control system (1) on a Riemannian manifold
 229 (M, g) under our standard assumptions. The next theorem presents a lower
 230 bound for the invariance pressure of admissible pairs (K, Q) .

Theorem 14. *Let (K, Q) be an admissible pair where both K and Q have positive and finite volume. Then for every $f \in C(U, \mathbb{R})$*

$$P_{inv}(f, K, Q) \geq \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \left(\inf_{(x, \omega)} \int_0^\tau f(\omega(s)) ds + \max\{0, \inf_{(x, \omega)} \int_0^\tau \operatorname{div} F_{\omega(s)}(\varphi(s, x, \omega)) ds\} \right),$$

231 where both infima are taken over all $(x, \omega) \in K \times \mathcal{U}$ with $\varphi([0, \tau], x, \omega) \subset Q$.

232 **Proof.** First observe that by Remark 7 we may assume that for all $\tau > 0$
 233 there exists a countable (τ, K, Q) -spanning set, since otherwise $P_{inv}(f, K, Q)$
 234 $= \infty$, and the infimum in $a_\tau(f, K, Q)$ may be taken over all countable
 235 (τ, K, Q) -spanning sets \mathcal{S} . For each ω in a countable (τ, K, Q) -spanning set
 236 \mathcal{S} define

$$K_\omega := \{x \in K; \varphi([0, \tau], x, \omega) \subset Q\}.$$

237 Thus $K = \bigcup_{\omega \in \mathcal{S}} K_\omega$. Since Q is Borel measurable, each set K_ω is measurable
 238 as the countable intersection of measurable sets,

$$K_\omega = K \cap \bigcap_{t \in [0, \tau] \cap \mathbb{Q}} \varphi_{t, \omega}^{-1}(Q).$$

Then

$$\begin{aligned} \text{vol}(Q) &\geq \text{vol}(\varphi_{\tau,\omega}(K_\omega)) = \int_{\varphi_{\tau,\omega}(K_\omega)} \text{dvol} = \int_{K_\omega} |\det \text{d}_x \varphi_{\tau,\omega}| \text{dvol} \\ &\geq \text{vol}(K_\omega) \inf_{(x,\omega)} |\det \text{d}_x \varphi_{\tau,\omega}|, \end{aligned}$$

239 where the infimum is taken over all $(x, \omega) \in K \times \mathcal{U}$ with $\varphi([0, \tau], x, \omega) \subset Q$.

240 Abbreviating with the same infima

$$\alpha(\tau) := \inf_{(x,\omega)} |\det \text{d}_x \varphi_{\tau,\omega}|, \quad \beta(\tau) := \inf_{(x,\omega)} S_\tau(f)(\omega),$$

we find

$$\begin{aligned} e^{\beta(\tau)} \text{vol}(K) &\leq \sum_{\omega \in \mathcal{S}} e^{(S_\tau f)(\omega)} \text{vol}(K_\omega) \leq \sup_{\omega \in \mathcal{S}} \text{vol}(K_\omega) \sum_{\omega \in \mathcal{S}} e^{(S_\tau f)(\omega)} \\ &\leq \frac{\text{vol}(Q)}{\max\{1, \alpha(\tau)\}} \sum_{\omega \in \mathcal{S}} e^{(S_\tau f)(\omega)}. \end{aligned}$$

Since this holds for every countable (τ, K, Q) -spanning set \mathcal{S} , we find

$$\begin{aligned} a_\tau(f, K, Q) &= \inf \left\{ \sum_{\omega \in \mathcal{S}} e^{(S_\tau f)(\omega)}; \mathcal{S} \text{ countable } (\tau, K, Q)\text{-spanning} \right\} \\ &\geq \frac{\text{vol}(K)}{\text{vol}(Q)} e^{\beta(\tau)} \max\{1, \alpha(\tau)\}, \end{aligned}$$

241 Since for each $t \geq 0$ and each control $\omega \in \mathcal{U}$ the map $\varphi_{t,\omega} : M \rightarrow M$ is a
242 diffeomorphism, Liouville's formula shows

$$\log \det \text{d}_x \varphi_{\tau,\omega} = \int_0^\tau \text{div} F_{\omega(s)}(\varphi(s, x, \omega)) ds. \quad (8)$$

243 Now the assertion of the theorem follows from

$$\begin{aligned} P_{inv}(f, K, Q) &= \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \log a_\tau(f, K, Q) \\ &\geq \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} (\beta(\tau) + \log \max\{1, \alpha(\tau)\}) \\ &= \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \left(\inf_{(x,\omega)} \int_0^\tau f(\omega(s)) ds + \max\{0, \inf_{(x,\omega)} \int_0^\tau \text{div} F_{\omega(s)}(\varphi(s, x, \omega)) ds\} \right). \end{aligned}$$

246 ■

247 **5. Linear control systems**

248 In this section we consider linear control systems on \mathbb{R}^d with restricted
 249 controls. Here a unique control set D with nonvoid interior exists and the
 250 previous bounds on the invariance pressure are sharpened to provide a for-
 251 mula for the invariance pressure of D .

252 Linear control systems on \mathbb{R}^d have the form

$$\dot{x}(t) = Ax(t) + B\omega(t), \quad \omega \in \mathcal{U}, \quad (9)$$

253 with $A \in \mathbb{R}^{d \times d}$ and $B \in \mathbb{R}^{d \times m}$ and we suppose that the set \mathcal{U} of control
 254 functions is as for (1).

255 For system (9) there exists a unique control set D with nonvoid interior,
 256 if, without control constraint, the system is controllable (which holds if and
 257 only if $\text{rank}[B, AB, \dots, A^{d-1}B] = d$) and the control range U is a compact
 258 neighborhood of the origin in \mathbb{R}^m . It is convex with $0 \in \text{int}D$, and it is
 259 bounded if and only if A is hyperbolic, i.e., there is no eigenvalue of A with
 260 vanishing imaginary part (cf. Hinrichsen and Pritchard [14, Theorems 6.2.22
 261 and 6.2.23], Colonius and Kliemann [9, Example 3.2.16]). Then the state
 262 space \mathbb{R}^d can be decomposed into the direct sum of the stable subspace E^s
 263 and the unstable subspace E^u which are the direct sums of all generalized
 264 real eigenspaces for the eigenvalues λ with $\text{Re } \lambda < 0$ and $\text{Re } \lambda > 0$, resp. Let
 265 $\pi : \mathbb{R}^d \rightarrow E^u$ be the projection along E^s . We obtain the following estimates,
 266 where λ_j denote the r eigenvalues of A with algebraic multiplicities d_j .

Lemma 15. *Consider a linear control system in \mathbb{R}^d of the form (9) and assume that the pair (A, B) is controllable, that A is hyperbolic and the control range U is a compact neighborhood of the origin. Let D be the unique control set with nonvoid interior. Then for every compact set $K \subset D$ with positive Lebesgue measure every potential $f \in C(U, \mathbb{R})$ satisfies*

$$\begin{aligned} \inf_{(T', x', \omega')} \frac{1}{T'} \int_0^{T'} f(\omega'(s)) ds &\leq P_{\text{inv}}(f, K, D) - \sum_{j=1}^r d_j \max\{0, \text{Re } \lambda_j\} \\ &\leq \inf_{(T, x, \omega)} \frac{1}{T} \int_0^T f(\omega(s)) ds, \end{aligned}$$

267 where the first infimum is taken over all $(T', x', \omega') \in (0, \infty) \times \pi K \times \mathcal{U}$ with
 268 $\pi\varphi([0, T'], x', \omega') \subset \pi D$ and the second infimum is taken over all $(T, x, \omega) \in$
 269 $(0, \infty) \times \text{int}D \times \mathcal{U}$ such that the control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$ is
 270 T -periodic and the values $\omega(t), t \in [0, T]$, are in a compact subset of $\text{int}U$.

271 **Proof.** The hypotheses imply that $0 \in \text{int}D \subset \mathbb{R}^d$ and the Lebesgue mea-
 272 sures of K and D (which coincide with the volumes) are finite and positive.
 273 Theorem 11 yields

$$P_{inv}(f, K, D) \leq \inf_{(T, x, \omega)} \left\{ \sum_{j=1}^{r(x, \omega)} \max\{0, d_j(x, \omega) \rho_j(x, \omega)\} + \frac{1}{T} \int_0^T f(\omega(s)) ds \right\}, \quad (10)$$

274 where the infimum is taken over all $T > 0$ and all $(x, \omega) \in \text{int}D \times \mathcal{U}$ such
 275 that the control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$ is T -periodic and the values
 276 $\omega(t), t \in [0, T]$, are in a compact subset of $\text{int}U$. Note that this control-
 277 trajectory pair is regular, since we assume that (A, B) is controllable. By
 278 Floquet theory it follows (cf. [6, Proposition 20]) that for all T -periodic
 279 $(\omega(\cdot), \varphi(\cdot, x, \omega))$

$$\sum_{j=1}^{r(x, \omega)} \max\{0, d_j(x, \omega) \rho_j(x, \omega)\} = \sum_{j=1}^r \max\{0, d_j \text{Re } \lambda_j\},$$

280 where the sum is over the r eigenvalues λ_j of A with multiplicities d_j . Hence

$$P_{inv}(f, K, D) \leq \inf_{(T, x, \omega)} \frac{1}{T} \int_0^T f(\omega(s)) ds + \sum_{j=1}^r d_j \max\{0, \text{Re } \lambda_j\},$$

281 where the infimum is taken over all (T, x, ω) as in (10). This proves the
 282 second inequality.

Hence it remains to prove the first inequality. By Theorem 14

$$\begin{aligned} & P_{inv}(f, K, D) \\ & \geq \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \left(\inf_{(x, \omega)} \int_0^\tau f(\omega(s)) ds + \max \left\{ 0, \inf_{(x, \omega)} \int_0^\tau \text{div} F_{\omega(s)}(\varphi(s, x, \omega)) ds \right\} \right) \\ & \geq \inf_{(T, x, \omega)} \frac{1}{T} \int_0^T f(\omega(s)) ds + \max \left\{ 0, \inf_{(T, x, \omega)} \frac{1}{T} \int_0^T \text{div} F_{\omega(s)}(\varphi(s, x, \omega)) ds \right\}, \end{aligned}$$

283 where both infima in the second line are taken over all pairs $(x, \omega) \in K \times \mathcal{U}$
 284 with $\varphi([0, \tau], x, \omega) \subset D$ and both infima in the third line are taken over all
 285 $(T, x, \omega) \in (0, \infty) \times K \times \mathcal{U}$ with $\varphi([0, T], x, \omega) \subset D$. In the considered linear
 286 case one has $d_x \varphi_{T, \omega} = A$ and

$$\int_0^T \text{div} F_{\omega(s)}(\varphi(s, x, \omega)) ds = \log \det d_x \varphi_{T, \omega} = T \sum_{j=1}^r d_j \text{Re } \lambda_j,$$

287 where the sum is over the r eigenvalues λ_j of A with multiplicities d_j .

Step 1: Suppose that $\operatorname{Re} \lambda_j > 0$ for all j . Then it follows that

$$P_{inv}(f, K, D) \geq \inf_{(T, x, \omega)} \frac{1}{T} \int_0^T f(\omega(s)) ds + \sum_{j=1}^r d_j \operatorname{Re} \lambda_j,$$

288 where the infimum is taken over all $(T, x, \omega) \in (0, \infty) \times K \times \mathcal{U}$ with
 289 $\varphi([0, T], x, \omega) \subset D$.

290 **Step 2:** Next we treat the general case, where also eigenvalues with neg-
 291 ative real part are allowed. Recall that $\pi : \mathbb{R}^d \rightarrow E^u$ denotes the projection
 292 onto the unstable subspace E^u along the stable subspace E^s .

293 Since these subspaces are A -invariant, this defines a (time-invariant) semi-
 294 conjugacy between system (9) and the system on E^u given by

$$\dot{y}(t) = A|_{E^u} y(t) + \pi B u(t), u \in \mathcal{U}, \quad (11)$$

295 with trajectories $\pi\varphi(\cdot, x', \omega')$, and the sets K and D are mapped to πK and
 296 πD , resp. Then πK and πD have positive volume and form an admissible
 297 pair (cf. Kawan [17, proof of Theorem 3.1]). One easily proves that (cf. [6,
 298 Proposition 10])

$$P_{inv}(f, K, Q) \geq P_{inv}(f, \pi K, \pi Q),$$

299 since every (τ, K, D) -spanning set yields a $(\tau, \pi K, \pi D)$ -spanning set. Simi-
 300 larly as in Step 1, Theorem 14 applied to system (11) implies that

$$P_{inv}(f, \pi K, \pi D) \geq \inf_{(T', x', \omega')} \frac{1}{T'} \int_0^{T'} f(\omega'(s)) ds + \sum_{j=1}^r d_j \max\{0, \operatorname{Re} \lambda_j\},$$

301 where the infimum is taken over all $(T', x', \omega') \in \mathbb{R}_+ \times \pi K \times \mathcal{U}$ with

$$\pi\varphi([0, T'], x', \omega') \subset \pi D.$$

302 ■

303 Next we show that the two infima in the lemma above actually coincide
 304 again using hyperbolicity of A in a crucial way. This provides the announced
 305 formula for the invariance pressure involving the r eigenvalues λ_j of A with
 306 algebraic multiplicities d_j .

307 **Theorem 16.** Consider a linear control system in \mathbb{R}^d of the form (9) and
 308 assume that the pair (A, B) is controllable, the matrix A is hyperbolic and
 309 the control range U is a compact neighborhood of the origin. Let D be the
 310 unique control set with nonvoid interior. Then for every compact set $K \subset D$
 311 with nonempty interior every potential $f \in C(U, \mathbb{R})$ satisfies

$$P_{inv}(f, K, D) = \min_{u \in U} f(u) + \sum_{j=1}^r d_j \max\{0, \operatorname{Re} \lambda_j\}. \quad (12)$$

312 **Proof.** Let $\varepsilon > 0$ and consider $T_0 > 0$ and a control $\omega_0 \in \mathcal{U}$ satisfying

$$\frac{1}{T_0} \int_0^{T_0} f(\omega_0(s)) ds \leq \inf_{(T'', \omega'') \in (0, \infty) \times \mathcal{U}} \frac{1}{T''} \int_0^{T''} f(\omega''(s)) ds + \varepsilon. \quad (13)$$

313 Since f is continuous, there is a control value $u_0 \in U$ with

$$f(u_0) = \min_{u \in U} f(u) = \inf_{(T'', \omega'') \in (0, \infty) \times \mathcal{U}} \frac{1}{T''} \int_0^{T''} f(\omega''(s)) ds, \quad (14)$$

314 where the second equality holds trivially. There is a control ω_1 in the set

$$\operatorname{int}\mathcal{U}_{[0, T_0]} = \{\omega \in L^\infty([0, T_0]; \exists K \subset \operatorname{int}U \text{ compact with } \omega(t) \in K \text{ a.e.}\}$$

315 such that

$$\frac{1}{T_0} \int_0^{T_0} f(\omega_1(s)) ds \leq \frac{1}{T_0} \int_0^{T_0} f(\omega_0(s)) ds + \varepsilon. \quad (15)$$

316 **Claim:** For every $T > 0$ and every control $\omega \in \mathcal{U}$ there exists $x_1 \in \mathbb{R}^d$ with
 317 $\varphi(T, x_1, \omega) = x_1$.

318 In fact, hyperbolicity of A implies that the matrix $I - e^{AT}$ is invertible,
 319 and hence there is a unique solution $x(T, \omega)$ of

$$(I - e^{AT}) x(T, \omega) = \varphi(T, 0, \omega).$$

320 Now the variation-of-constants formula shows the claim:

$$x(T, \omega) = e^{AT} x(T, \omega) + \varphi(T, 0, \omega) = \varphi(T, x(T, \omega), \omega).$$

321 Applying this to T_0 and ω_1 we find a point $x_1 := x(T_0, \omega_1) = \varphi(T_0, x_1, \omega_1)$.
 322 Since $\omega_1 \in \operatorname{int}\mathcal{U}_{[0, T_0]}$ every point in a neighborhood of x_1 can be reached in
 323 time T_0 from x_1 . This follows, since by controllability the map

$$L_\infty([0, T_0], \mathbb{R}^m) \rightarrow \mathbb{R}^d, \omega \mapsto \varphi(T_0, 0, \omega)$$

324 is a linear surjective map, hence maps open sets to open sets, and the same
 325 is true for the map

$$\omega \mapsto \varphi(T_0, x_1, \omega) = e^{AT_1} + \varphi(T_0, 0, \omega).$$

326 Analogously, x_1 can be reached from every point in a neighborhood of x_1 in
 327 time T_0 . Hence in the intersection of these two neighborhoods every point
 328 can be steered in time $2T_0$ into every other point. This shows that x_1 is in
 329 the interior of the (unique) control set D , and the corresponding trajectory
 330 $\varphi(t, x_1, \omega_1), t \in [0, T_0]$, remains in the interior of D . Extending $\omega_1(t), t \in$
 331 $[0, T_0]$, to a T_0 -periodic control, again denoted by ω_1 we find that the control-
 332 trajectory pair $(\omega_1(\cdot), \varphi(\cdot, x_1, \omega_1))$ is T_0 -periodic, the trajectory is contained
 333 in $\text{int}D$ and the values $\omega_1(t), t \in [0, T_0]$, are in a compact subset of $\text{int}U$. By
 334 (13) and (15) it follows that

$$\inf_{(T'', \omega'')} \frac{1}{T''} \int_0^{T''} f(\omega''(s)) ds \geq \frac{1}{T_0} \int_0^{T_0} f(\omega_0(s)) ds - \varepsilon$$

335

$$\geq \frac{1}{T_0} \int_0^{T_0} f(\omega_1(s)) ds - 2\varepsilon \geq \inf_{(T, x, \omega)} \frac{1}{T} \int_0^T f(\omega(s)) ds - 2\varepsilon,$$

336 where the infimum in the last line is taken over all $(T, x, \omega) \in (0, \infty) \times$
 337 $D \times \mathcal{U}$ such that the control-trajectory pair $(\omega(\cdot), \varphi(\cdot, x, \omega))$ is T -periodic,
 338 the trajectory is contained in $\text{int}D$ and the values $\omega(t), t \in [0, T]$, are in a
 339 compact subset of $\text{int}U$.

Together with (14) and the inequalities in Lemma 15 this implies

$$\begin{aligned} \min_{u \in U} f(u) &= \inf_{(T'', \omega'')} \frac{1}{T''} \int_0^{T''} f(\omega''(s)) ds \leq \inf_{(T', x', \omega')} \frac{1}{T'} \int_0^{T'} f(\omega'(s)) ds \\ &\leq P_{inv}(f, K, D) - \sum_{j=1}^r d_j \max\{0, \text{Re } \lambda_j\} \\ &\leq \inf_{(T, x, \omega)} \frac{1}{T} \int_0^T f(\omega(s)) ds \\ &\leq \inf_{(T'', \omega'')} \frac{1}{T''} \int_0^{T''} f(\omega''(s)) ds + 2\varepsilon \\ &= \min_{u \in U} f(u) + 2\varepsilon. \end{aligned}$$

340 Since $\varepsilon > 0$ is arbitrary, assertion (12) follows. ■

341 **Remark 17.** *The proof of the **Claim** above follows arguments in the proof*
 342 *of Da Silva and Kawan [13, Theorem 20].*

343 **Remark 18.** *Theorem 16 improves [6, Theorem 6.2], where it had to be*
 344 *assumed additionally that the minimum of $f(u), u \in U$, is attained in an*
 345 *equilibrium.*

346 6. Further applications

347 In this section, we apply Theorem 11 to linear control systems on Lie
 348 groups and to inner control sets.

349 6.1. Control sets and equilibrium pairs

350 Given a control system (1), a pair $(u_0, x_0) \in U \times M$ is called an **equilib-**
 351 **rium pair** if $F(x_0, u_0) = 0$, or equivalently, $\varphi(t, x_0, \bar{u}_0) = x_0$ for all $t \in \mathbb{R}$,
 352 where $\bar{u}_0(t) \equiv u_0$.

353 If (u_0, x_0) is an equilibrium pair, the linearized system is an autonomous
 354 linear control system in $T_{x_0}M$ and the Lyapunov exponents at (u_0, x_0) in
 355 the direction $\lambda \in T_{x_0}M \setminus \{0_{x_0}\}$ coincide with the real parts of the eigenvalues
 356 of $\nabla F_{u_0}(x_0) : T_{x_0}M \rightarrow T_{x_0}M$. Then regularity, i.e., controllability of the
 357 linearized system, can be checked by Kalman's rank condition.

358 **Corollary 19.** *Let $D \subset M$ be a control set with nonempty interior and let*
 359 *$f \in C(U, \mathbb{R})$. Suppose that there is a regular equilibrium pair $(u_0, x_0) \in$*
 360 *$\text{int}U \times \text{int}D$. Then for every compact set $K \subset D$ and every set $Q \supset D$ we*
 361 *have*

$$P_{inv}(f, K, Q) \leq \sum_{\lambda \in \sigma(\nabla F_{u_0}(x_0))} \max\{0, d_\lambda \text{Re}(\lambda)\} + f(u_0),$$

362 where d_λ is the algebraic multiplicity of the eigenvalue λ in the spectrum
 363 $\sigma(\nabla F_{u_0}(x_0))$.

Proof. Since (u_0, x_0) is a regular equilibrium pair, the control-trajectory pair
 $(\bar{u}_0(\cdot), \varphi(\cdot, x_0, \bar{u}_0))$ is T -periodic and regular for every $T > 0$. By Theorem
 11 we obtain

$$\begin{aligned} P_{inv}(f, K, Q) &\leq \inf_{(T, x, \omega)} \left\{ \sum_{j=1}^{r(x, \omega)} \max\{0, d_j(x, \omega) \rho_j(x, \omega)\} + \frac{1}{T} \int_0^T f(\omega(s)) ds \right\} \\ &\leq \sum_{\lambda \in \sigma(\nabla F_{\omega_0}(x_0))} \max\{0, d_\lambda \text{Re}(\lambda)\} + f(u_0). \end{aligned}$$

364 ■

365 *6.2. Control sets of linear control systems on Lie groups*

366 In this subsection we consider **linear control systems on a connected**
 367 **Lie group** G introduced in Ayala and San Martin [2] and Ayala and Tirao
 368 [4].

369 They are given by a family of ordinary differential equations on G of the
 370 form

$$\dot{x}(t) = \mathcal{X}(x(t)) + \sum_{j=1}^m \omega_j(t) X_j(x(t)), \quad \omega = (\omega_1, \dots, \omega_m) \in \mathcal{U}, \quad (16)$$

371 where the drift vector field \mathcal{X} , called the **linear vector field**, is an infinitesimal
 372 automorphism, i.e., its solutions are a family of automorphisms of the
 373 group, and the X_j are right invariant vector fields. Note that the linear
 374 control systems of the form (9) are a special case with $G = \mathbb{R}^d$.

375 Their controllability properties have been analyzed in Da Silva [11], Ayala,
 376 Da Silva and Zsigmond [3] and Ayala and Da Silva [1]. In particular,
 377 the existence and uniqueness of control sets for general systems of the form
 378 (16) has been analyzed in [3]. If 0 is in the interior of the control range U
 379 and the reachable set $\mathcal{O}^+(e_G)$ from the neutral element e_G is open (this holds
 380 e.g. if $e_G \in \text{int}\mathcal{O}^+(e_G)$), then there exists a control set D containing e_G
 381 in the interior. Sufficient conditions for boundedness and uniqueness of D are
 382 given in [3, Theorem 3.9] and [3, Corollary 3.12], respectively.

383 Along with system (16) comes an associated derivation \mathcal{D} of the Lie algebra
 384 \mathfrak{g} of G which is given by

$$\mathcal{D}(Y) = -\text{ad}(\mathcal{X})(Y) := [\mathcal{X}, Y](e_G).$$

385 **Corollary 20.** *Consider the linear control system (16) on a Lie group G .*
 386 *Suppose that D is a control set with $e_G \in \text{int}D$ and compact closure \overline{D} and*
 387 *let $K \subset D \subset Q$. Let $f \in C(U, \mathbb{R})$ be a potential. If the equilibrium pair*
 388 *$(0, e_G) \in \text{int}U \times \text{int}D$ is regular, then*

$$P_{inv}(f, K, Q) \leq \sum_{\lambda \in \sigma(\mathcal{D})} \max\{0, d_\lambda \text{Re}(\lambda)\} + f(0).$$

389 *If furthermore K has positive Haar measure and $f(0) = \min_{u \in U} f(u)$, then*

$$P_{inv}(f, K, Q) = P_{out}(f, K, Q) = \sum_{\lambda \in \sigma(\mathcal{D})} \max\{0, d_\lambda \text{Re}(\lambda)\} + f(0).$$

390 **Proof.** Note that the right hand side of the system is given by $F(x, u) =$
 391 $\mathcal{X}(x) + \sum_{i=1}^m u_i X_i(x)$ and hence $F_0(x) := F(x, 0) = \mathcal{X}(x)$. Let (ϕ, U) be a
 392 local coordinate neighborhood of e_G and pick a left invariant vector field Y
 393 in the Lie algebra \mathfrak{g} of G . Then we can express \mathcal{X} in terms of (ϕ, U) by

$$\mathcal{X}(h) = \sum_{i=1}^d y_i(h) \frac{\partial}{\partial x_i}.$$

394 Note that $\mathcal{X}(e_G) = 0$ implies $y_i(e_G) = 0$ for every $i \in \{1, \dots, d\}$, hence the
 395 Levi-Civita connection ∇ satisfies

$$(\nabla_{\mathcal{X}} Y)(e_G) = \sum_{i=1}^d y_i(e_G) \left(\nabla_{\frac{\partial}{\partial x_i}} Y \right) (e_G) = 0.$$

396 Since ∇ is symmetric, we have

$$(\nabla_Y F_0)(e_G) = (\nabla_Y \mathcal{X})(e_G) = (\nabla_{\mathcal{X}} Y - [\mathcal{X}, Y])(e_G) = -[\mathcal{X}, Y] = \mathcal{D}(Y).$$

397 Since this holds for every $Y \in \mathfrak{g}$, we have $\nabla F_0(e_G) = \mathcal{D}$. By Corollary 19 we
 398 obtain

$$P_{inv}(f, K, Q) \leq \sum_{\lambda \in \sigma(\mathcal{D})} \max\{0, d_\lambda \operatorname{Re}(\lambda)\} + f(0).$$

399 Now, suppose that K has positive Haar measure. By Da Silva [10, Theorem
 400 4.3], we know that

$$h_{out}(K, Q) \geq \sum_{\lambda \in \sigma(\mathcal{D})} \max\{0, d_\lambda \operatorname{Re}(\lambda)\}.$$

401 Define $\tilde{f}(u) = f(u) - f(0)$, $u \in U$. Since $\tilde{f} \geq 0$ Proposition 6(i) implies that

$$P_{inv}(\tilde{f}, K, Q) \geq P_{out}(\tilde{f}, K, Q) \geq h_{out}(K, Q) \geq \sum_{\lambda \in \sigma(\mathcal{D})} \max\{0, d_\lambda \operatorname{Re}(\lambda)\}.$$

402 Proposition 6(ii) implies $P_{inv}(\tilde{f}, K, Q) = P_{inv}(f, K, Q) - \min f$, hence this
 403 yields

$$P_{inv}(f, K, Q) = P_{out}(f, K, Q) = \sum_{\lambda \in \sigma(\mathcal{D})} \max\{0, d_\lambda \operatorname{Re}(\lambda)\} + \min f.$$

404

405 *6.3. Inner control sets*

406 This section presents an application of Theorem 11 to the class of *inner*
 407 *control sets* as defined (with small changes) in Kawan [17, Definition 2.6].
 408 This nomenclature refers to a control set $D \subset M$ for which there exists a
 409 decreasing family of compact and convex sets $\{U_\rho\}_{\rho \in [0,1]}$ in \mathbb{R}^m (i.e., $U_{\rho_2} \subset U_{\rho_1}$
 410 for $\rho_1 < \rho_2$), such that for every $\rho \in [0, 1]$ system $(1)_\rho$ with control range U_ρ
 411 (instead of U in (1)) has a control set D_ρ with nonvoid interior and compact
 412 closure, and the following conditions are satisfied:

- 413 (i) $U = U_0$ and $D = D_1$;
- 414 (ii) $\overline{D_{\rho_2}} \subset \text{int} D_{\rho_1}$ whenever $\rho_1 < \rho_2$;
- 415 (iii) for every neighborhood W of \overline{D} there is $\rho \in [0, 1)$ with $\overline{D_\rho} \subset W$.

416 We will estimate the outer invariance pressure of the set $Q = \overline{D}$ for the
 417 system with control range $U = U_0$. Note that, in general, D is not a control
 418 set for this system, since we only have $D = D_1 \subset D_0$.

419 **Corollary 21.** *Consider an inner control set D of control system (1). Let*
 420 *$(\omega_0(\cdot), \varphi(\cdot, x_0, \omega_0))$ be a regular T -periodic control-trajectory pair with $x_0 \in \overline{D}$*
 421 *and $\omega_0 \in \mathcal{U}_1$. Then*

$$P_{out}(f, \overline{D}) \leq \sum_{j=1}^r \max\{0, d_j \text{Re } \lambda_j\} + \frac{1}{T} \int_0^T f(\omega_0(s)) ds$$

422 holds, where $\lambda_1, \dots, \lambda_r$ are the Lyapunov exponents at (x_0, ω_0) with corre-
 423 sponding multiplicities d_1, \dots, d_r .

424 **Proof.** Note that the definition of inner control sets implies that for every
 425 $\rho \in [0, 1)$ the set \overline{D} is a compact subset of D_ρ and the pair (\overline{D}, D_ρ) is
 426 admissible. By Theorem 11 it follows that the outer invariance pressure
 427 $P_{out}^\rho(f, \overline{D}, \overline{D_\rho})$ for system $(1)_\rho$ satisfies

$$P_{out}^\rho(f, \overline{D}, \overline{D_\rho}) \leq \sum_{j=1}^r \max\{0, d_j \rho_j\} + \frac{1}{T} \int_0^T f(\omega_0(s)) ds \text{ for all } \rho \in [0, 1).$$

Now for given $\varepsilon > 0$ we may choose $\rho \in [0, 1)$ such that $\overline{D_\rho} \subset N_\varepsilon(\overline{D})$. Then

$$\begin{aligned} P_{out}(f, \overline{D}, N_\varepsilon(\overline{D})) &\leq P_{out}^\rho(f, \overline{D}, N_\varepsilon(\overline{D})) \leq P_{out}^\rho(f, \overline{D_\rho}, N_\varepsilon(\overline{D})) \\ &\leq \sum_{j=1}^r \max\{0, d_j \rho_j\} + \frac{1}{T} \int_0^T f(\omega_0(s)) ds. \end{aligned}$$

428 The first two inequalities follow from $U_\rho \subset U_0$ and $\overline{D_\rho} \subset N_\varepsilon(\overline{D})$. Since
 429 $P_{out}(f, \overline{D}) = \lim_{\varepsilon \rightarrow 0} P_{out}(f, \overline{D}, N_\varepsilon(\overline{D}))$, the assertion follows. ■

430 *6.4. Example*

431 The following example illustrates Theorem 16. Consider the following
 432 linear control system in \mathbb{R}^d ,

$$\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \underbrace{\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}}_{=:A} \begin{bmatrix} x \\ y \end{bmatrix} + \underbrace{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}_{=:B} \omega(t)$$

433 and assume that $\omega(t) \in U := [-1, 1] + u_0$ for some $u_0 \in (-1, 1)$. In this case,
 434 $0 \in \text{int}U$, the pair (A, B) is controllable, and A is hyperbolic with eigenvalues
 435 given by $\lambda_{\pm} = 1 \pm i$. There exists a unique control set $D \subset \mathbb{R}^2$ such that
 436 $(0, 0) \in \text{int}D$, and \bar{D} is compact.

437 We may interpret the control functions $\omega(t)$ and also u_0 as external forces
 438 acting on the system. Take $f \in C(U, \mathbb{R})$ as $f(u) := |u - u_0|$, then $(S_{\tau}f)(\omega)$
 439 represents the impulse of $\omega - u_0$ until time τ . For a subset $K \subset D$ a (τ, K, D) -
 440 spanning set \mathcal{S} represents a set of external forces ω that cause the system to
 441 remain in D when it starts in K . By Theorem 16 we obtain for a compact
 442 subset $K \subset D$ with nonempty interior that

$$P_{inv}(f, K, Q) = 2 + \min_{u \in U} f(u) = 2 + \min_{u \in [-1, 1] + u_0} |u - u_0|.$$

443 Here $P_{inv}(f, K, Q)$ represent the exponential growth rate of the amount of
 444 total impulse required of the external forces $\omega - u_0$ acting on the system
 445 to remain in D as time tends to infinity. The minimum of f is attained in
 446 $u = u_0$, which does not correspond to an equilibrium if $u_0 \neq 0$. Hence [6,
 447 Corollary 21] (cf. Remark 18) could not be applied in this case.

448 **References**

- 449 [1] V. AYALA AND A. DA SILVA, *Controllability of linear systems on Lie*
 450 *groups*, SIAM J. Control Optim. 55 (2017), pp. 1332–1343.
- 451 [2] V. AYALA AND L.A.B. SAN MARTIN, *Controllability properties of a*
 452 *class of control systems on Lie groups*, in: Nonlinear Control in the Year
 453 2000 Vol. 1, A. Isidori, F. Lamnabhi-Lagarrigue, and W. Respondek,
 454 eds., Lecture Notes in Control and Inform. Sciences, Vol. 258, Springer
 455 2001, pp. 83–92.

- 456 [3] V. AYALA, A. DA SILVA AND G. ZSIGMOND, *Control sets of linear*
457 *systems on Lie groups*, NODEA Nonlinear Differ. Equ. and Appl. 24(8)
458 (2017), pp. 1-15.
- 459 [4] V. AYALA AND J. TIRAO, *Linear control systems on Lie groups and*
460 *controllability*, in: Differential Geometry and Control, Proc. Sympos.
461 Pure Math. 64, G. Ferreyra, R. Gardner, H. Hermes, and H. Sussmann,
462 eds., AMS, Providence, RI, 1999, pp. 47–64.
- 463 [5] F. COLONIUS, J.A.N. COSSICH AND A.J. SANTANA, *Invariance pres-*
464 *sure for control systems*, J. Dyn. Diff. Equations 31(1) (2019), pp 1–23.
- 465 [6] F. COLONIUS, J.A.N. COSSICH AND A.J. SANTANA, *Invariance pres-*
466 *sure of control sets*, SIAM J. Control Optim. 56(6) (2018), pp. 4130–
467 4147.
- 468 [7] F. COLONIUS, C. KAWAN AND G. NAIR, *A note on topological feedback*
469 *entropy and invariance entropy*, Systems & Control Letters 62(5), 377–
470 381 (2013)
- 471 [8] F. COLONIUS AND C. KAWAN, *Invariance entropy for control systems*,
472 SIAM J. Control Optim. 48 (2009), pp. 1701-1721.
- 473 [9] F. COLONIUS AND W. KLIEMANN, *The Dynamics of Control*.
474 Birkhäuser Boston, Boston, MA, 2000.
- 475 [10] A. DA SILVA, *Outer invariance entropy for linear systems on Lie groups*,
476 SIAM J. Control Optim. 52 (2014), pp. 3917–3934.
- 477 [11] A. DA SILVA, *Controllability of linear systems on solvable Lie groups*,
478 SIAM J. Control Optim., 54 (2016), pp. 372-390.
- 479 [12] A. DA SILVA AND C. KAWAN, *Invariance entropy of hyperbolic control*
480 *sets*, Disc. and Cont. Dyn. Systems. 36(1) (2016), pp. 97-136.
- 481 [13] A. DA SILVA AND C. KAWAN, *Robustness of critical bit rates for prac-*
482 *tical stabilization of networked control systems*, Automatica 93 (2018),
483 pp. 397-406.
- 484 [14] D. HINRICHSSEN AND A.J. PRITCHARD, *Mathematical Systems Theory*,
485 *Vol. 2*, in preparation, 2020.

- 486 [15] Y. HUANG AND X. ZHONG, *Carathéodory–Pesin structures associated*
487 *with control systems*, Systems and Control Letters 112 (2018), pp. 36-41.
- 488 [16] C. KAWAN, *Invariance entropy of control sets*, SIAM J. Control Optim.
489 49 (2011), pp. 732-751.
- 490 [17] C. KAWAN, *Invariance Entropy for Deterministic Control Systems. An*
491 *Introduction*. Lecture Notes in Math. Vol. 2089, Springer, Berlin, 2013.
- 492 [18] G. NAIR, R. J. EVANS, I. MAREELS, AND W. MORAN, *Topological*
493 *feedback entropy and nonlinear stabilization*, IEEE Trans. Aut. Control
494 49 (2004), pp. 1585–1597.
- 495 [19] X. ZHONG, Y. HUANG, *Invariance pressure dimensions for control sys-*
496 *tems*, J. Dyn. Diff. Equations, [https://doi.org/10.1007/s10884-018-9701-](https://doi.org/10.1007/s10884-018-9701-z)
497 [z](https://doi.org/10.1007/s10884-018-9701-z) (2018).