







Draft Genome Sequences of 12 *Leuconostoc carnosum* Strains Isolated from Cooked Ham Packaged in a Modified Atmosphere and from Fresh Sausages

Francesco Candelieri,^a  Stefano Raimondi,^a Gloria Spampinato,^a Moon Yue Feng Tay,^{c,d}  Alberto Amaretti,^{a,b}  Joergen Schlundt,^{c,d}  Maddalena Rossi^{a,b}

^aDepartment of Life Sciences, University of Modena and Reggio Emilia, Modena, Italy

^bBIOGEST—SITEIA, University of Modena and Reggio Emilia, Modena, Italy

^cNanyang Technological University Food Technology Centre (NAFTEC), Singapore

^dSchool of Chemical and Biomedical Engineering, Nanyang Technological University, Singapore

ABSTRACT *Leuconostoc carnosum* is a lactic acid bacterium that preferentially colonizes meat. In this work, we present the draft genome sequences of 12 *Leuconostoc carnosum* strains isolated from modified-atmosphere-packaged cooked ham and fresh sausages. Three strains harbor bacteriocin genes.

Leuconostoc carnosum is a heterofermentative, catalase-negative lactic acid bacterium (LAB) comprising the major component of the microbiota in modified-atmosphere-packaged (MAP) cooked ham at the end of the shelf life, where it reaches a concentration of 10^7 to 10^8 CFU/g (1). After a few weeks of shelf life, a high dose of alive *L. carnosum* may be ingested when consuming MAP cooked ham (1). *L. carnosum* easily colonizes several other meat-based food matrices, as suggested by the species name (2–6). It has been associated with ham spoilage (7, 8), but some strains have been proposed as bioprotective starters (4, 9) due to their production of bacteriocins against *Listeria monocytogenes* (5, 6). When we started this project, a sole completed genome sequence was available in GenBank (*L. carnosum* JB16, accession number [SAMN02603179](https://doi.org/10.1093/genbank/SAMN02603179)), obtained from a strain isolated from kimchi (10). Twelve *L. carnosum* genomes have been sequenced to characterize this species, belonging to 9 strains isolated from MAP cooked ham and 3 from fresh sausages, according to Raimondi et al. (1, 2) (Table 1).

For whole-genome sequencing, each strain was cultivated in brain heart infusion broth (Becton, Dickinson, USA) at 30°C for 48 h under microaerophilic conditions. Biomass was collected by centrifugation for 5 min at $5,000 \times g$. The genomic DNA was extracted with a QIAamp DNA minikit (Qiagen GmbH, Germany). The concentration was determined with a Qubit fluorometer (Invitrogen, USA). The DNA samples were submitted to the Singapore Centre for Environmental Life Sciences Engineering (SCELSE), where they were tagged with Illumina TruSeq high-throughput (HT) DNA dual barcodes for library preparation and sequenced on an Illumina HiSeq 2500 instrument (USA). For each sample, 250-bp paired-end reads were obtained. The raw and postprocessed reads were checked for quality with FastQC v0.11.7 (11). Cutadapt v1.16 was used to trim Illumina adapters, and reads with a quality score lower than 20 (–overlap = 15, –minimum-length = 30, –quality-cutoff = 20) were removed (12). The trimmed reads were assembled with SPAdes v3.12 (–careful –cov-cutoff auto –k auto) (13), and the quality of the assemblies was evaluated using QUAST v5.0.2 (14). Trimming, quality checking, and assembly were performed on the Galaxy platform (<https://usegalaxy.eu/>) (15). The completeness of the assemblies was determined with CheckM v1.0.8 (16). The taxonomy was confirmed with KmerFinder on the CGE server (<https://cge.cbs.dtu.dk/services/KmerFinder/>) (17). The assembled genomes were ordered with Mauve v2.4.0

Citation Candelieri F, Raimondi S, Spampinato G, Tay MYF, Amaretti A, Schlundt J, Rossi M. 2020. Draft genome sequences of 12 *Leuconostoc carnosum* strains isolated from cooked ham packaged in a modified atmosphere and from fresh sausages. Microbiol Resour Announc 9:e01247-19. <https://doi.org/10.1128/MRA.01247-19>.

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2020 Candelieri et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Maddalena Rossi, maddalena.rossi@unimore.it.

Received 3 October 2019

Accepted 9 December 2019

Published 9 January 2020

TABLE 1 Statistics of assembled genomes

GenBank accession no.	SRA accession no.	No. of raw reads	Strain	Source	Country of origin	Genome size (bp)	No. of contigs ^a	N ₅₀ (bp)	Coverage (x) ^b	G+C content (%) ^a	No. of CDS ^c	No. of rRNAs ^d	No. of tRNAs ^d	Completeness (%) ^e
VBXG01000000	SRR10389907	2,103,282	WC0318	Cooked ham	Italy	1,745,630	15	1,142,374	638	37.2	1,739	5	51	99.476
VBXF01000000	SRR10389906	1,718,930	WC0319	Cooked ham	Italy	1,700,071	11	1,137,580	521	37.1	1,679	5	49	99.476
VBXE01000000	SRR10389899	1,848,530	WC0320	Cooked ham	Poland	1,812,114	18	1,109,738	561	37.1	1,853	5	51	99.476
VBXD01000000	SRR10389898	1,753,950	WC0321	Cooked ham	France	1,804,293	40	1,106,035	532	37.1	1,854	5	52	99.476
VBXC01000000	SRR10389897	1,696,897	WC0322	Cooked ham	France	1,853,239	16	306,690	515	37	1,898	5	50	99.476
VBXB01000000	SRR10389896	1,847,670	WC0323	Cooked ham	Germany	1,773,698	23	256,026	560	37.2	1,802	5	52	99.476
VBXA01000000	SRR10389895	1,515,136	WC0324	Cooked ham	Germany	1,765,760	13	1,137,276	460	37.1	1,768	5	49	99.476
VBWY01000000	SRR10389893	1,654,963	WC0325	Cooked ham	Italy	1,830,248	27	229,291	502	37.2	1,881	5	52	99.476
VBWW01000000	SRR10389905	1,893,372	WC0326	Sausage	Italy	1,770,048	22	412,621	574	37.2	1,767	5	49	99.476
VBWV01000000	SRR10389904	1,872,273	WC0327	Sausage	Italy	1,769,683	20	412,621	568	37.2	1,768	5	49	99.476
VBWS01000000	SRR10389901	1,774,316	WC0328	Sausage	Italy	1,815,949	36	239,739	538	37.1	1,826	5	51	98.953
VBWR01000000	SRR10389900	1,498,696	WC0329	Cooked ham	France	1,650,966	14	256,123	455	37.2	1,644	5	52	99.476

^a Determined using QUAST v5.0.2.^b Coverage has been calculated as (2 × 250 × number of reads)/reference genome length.^c Determined using the RAST server.^d Determined using Prokka v1.13, with default parameters.^e Determined using CheckM v1.0.8.

(18), using the sequence of *L. carnosum* JB16 as a reference. The draft genomes were annotated on the RAST server (19) and with Prokka v1.13 (20). BAGEL4 (21) was used to identify bacteriocin genes. The sequence data were deposited in GenBank under the BioProject accession number PRJNA542256. Table 1 shows the GenBank accession number, number of raw reads, assembly statistics, and features of the genomes. The average G+C content (\pm the standard deviation) was $37.14\% \pm 0.07\%$. An average of 1,790 coding sequences (CDS) per genome were annotated, with genome sizes ranging between 1,650,966 and 1,853,239 bp and an average completeness of 99.43%. According to *in silico* analysis, 3 out of 12 strains were identified as potential bacteriocin producers.

Data availability. The sequence data were deposited in GenBank under BioProject accession number PRJNA542256. Table 1 lists, for each sample, the GenBank accession number and the number of raw reads.

ACKNOWLEDGMENTS

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

We acknowledge the Singapore Centre for Environmental Life Sciences Engineering (SCELSE) for sequencing the samples. We thank Kelyn L. G. Seow and Yalei Xu for the support in wet lab work.

REFERENCES

- Raimondi S, Luciani R, Sirangelo TM, Amaretti A, Leonardi A, Ulrici A, Foca G, D'Auria G, Moya A, Zuliani V, Seibert TM, Søtoft-Jensen J, Rossi M. 2019. Microbiota of sliced cooked ham packaged in modified atmosphere throughout the shelf life: microbiota of sliced cooked ham in MAP. *Int J Food Microbiol* 289:200–208. <https://doi.org/10.1016/j.ijfoodmicro.2018.09.017>.
- Raimondi S, Nappi MR, Sirangelo TM, Leonardi A, Amaretti A, Ulrici A, Magnani R, Montanari C, Tabanelli G, Gardini F, Rossi M. 2018. Bacterial community of industrial raw sausage packaged in modified atmosphere throughout the shelf life. *Int J Food Microbiol* 280:78–86. <https://doi.org/10.1016/j.ijfoodmicro.2018.04.041>.
- Geeraerts W, Pothakos V, De Vuyst L, Leroy F. 2018. Variability within the dominant microbiota of sliced cooked poultry products at expiration date in the Belgian retail. *Food Microbiol* 73:209–215. <https://doi.org/10.1016/j.fm.2018.01.019>.
- Budde BB, Hornbaek T, Jacobsen T, Barkholt V, Koch AG. 2003. *Leuconostoc carnosum* 4010 has the potential for use as a protective culture for vacuum-packed meats: culture isolation, bacteriocin identification, and meat application experiments. *Int J Food Microbiol* 83:171–184. [https://doi.org/10.1016/S0168-1605\(02\)00364-1](https://doi.org/10.1016/S0168-1605(02)00364-1).
- van Laack RLJM, Schillinger U, Holzapfel WH. 1992. Characterization and partial purification of a bacteriocin produced by *Leuconostoc carnosum* LA44A. *Int J Food Microbiol* 16:183–195. [https://doi.org/10.1016/0168-1605\(92\)90079-i](https://doi.org/10.1016/0168-1605(92)90079-i).
- Parente E, Moles M, Ricciardi A. 1996. Leucocin F10, a bacteriocin from *Leuconostoc carnosum*. *Int J Food Microbiol* 33:231–243. [https://doi.org/10.1016/0168-1605\(96\)01159-2](https://doi.org/10.1016/0168-1605(96)01159-2).
- Björkroth KJ, Vandamme P, Korkeala HJ. 1998. Identification and characterization of *Leuconostoc carnosum*, associated with production and spoilage of vacuum-packaged, sliced, cooked ham. *Appl Environ Microbiol* 64:3313–3319.
- Samelis J, Björkroth J, Kakouri A, Rementzis J. 2006. *Leuconostoc carnosum* associated with spoilage of refrigerated whole cooked hams in Greece. *J Food Prot* 69:2268–2273. <https://doi.org/10.4315/0362-028x-69.9.2268>.
- Jacobsen T, Budde BB, Koch AG. 2003. Application of *Leuconostoc carnosum* for biopreservation of cooked meat products. *J Appl Microbiol* 95:242–249. <https://doi.org/10.1046/j.1365-2672.2003.01970.x>.
- Jung JY, Lee SH, Jeon CO. 2012. Complete genome sequence of *Leuconostoc carnosum* strain JB16, isolated from kimchi. *J Bacteriol* 194:6672–6673. <https://doi.org/10.1128/JB.01805-12>.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17:10–12. <https://doi.org/10.14806/ej.17.1.200>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUASt: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Cech M, Chilton J, Clements D, Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltmann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D. 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res* 46:W537–W544. <https://doi.org/10.1093/nar/gky379>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Hasman H, Saputra D, Sicheritz-Ponten T, Lund O, Svendsen CA, Frimodt-Møller N, Aarestrup FM. 2014. Rapid whole-genome sequencing for detection and characterization of microorganisms directly from clinical samples. *J Clin Microbiol* 52:139–146. <https://doi.org/10.1128/JCM.02452-13>.
- Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <https://doi.org/10.1101/gr.2289704>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisano K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- van Heel AJ, de Jong A, Song C, Viel JH, Kok J, Kuipers OP. 2018. BAGEL4: a user-friendly Web server to thoroughly mine RiPPs and bacteriocins. *Nucleic Acids Res* 46:W278–W281. <https://doi.org/10.1093/nar/gky383>.