



UNIVERSITÀ
DEGLI STUDI
DI PADOVA

Università degli Studi di Padova

Padua Research Archive - Institutional Repository

Non-compact subsets of the Zariski space of an integral domain

Original Citation:

Availability:

This version is available at: 11577/3329231 since: 2020-03-04T15:14:00Z

Publisher:

University of Illinois

Published version:

DOI:

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

NON-COMPACT SUBSETS OF THE ZARISKI SPACE OF AN INTEGRAL DOMAIN

DARIO SPIRITO

ABSTRACT. Let V be a minimal valuation overring of an integral domain D and let $\text{Zar}(D)$ be the Zariski space of the valuation overrings of D . Starting from a result in the theory of semistar operations, we prove a criterion under which the set $\text{Zar}(D) \setminus \{V\}$ is not compact. We then use it to prove that, in many cases, $\text{Zar}(D)$ is not a Noetherian space, and apply it to the study of the spaces of Kronecker function rings and of Noetherian overrings.

1. INTRODUCTION

The *Zariski space* $\text{Zar}(K|D)$ of the valuation rings of a field K containing a domain D was introduced (under the name *abstract Riemann surface*) by O. Zariski, who used it to show that resolution of singularities holds for varieties of dimension 2 or 3 over fields of characteristic 0 [32, 33]. In particular, Zariski showed that $\text{Zar}(K|D)$, endowed with a natural topology, is always a compact space [34, Chapter VI, Theorem 40]; this result has been subsequently improved by showing that $\text{Zar}(K|D)$ is a spectral space (in the sense of Hochster [18]), first in the case where K is the quotient field of D [4, 5], and then in the general case [8, Corollary 3.6(3)]. The topological aspects of the Zariski space has subsequently been used, for example, in real and rigid algebraic geometry [19, 31] and in the study of representation of integral domains as intersections of valuation overrings [26, 27, 28]. In the latter context, i.e., when K is the quotient field of D , two important properties for subspaces of $\text{Zar}(K|D)$ to investigate are the properties of compactness and of Noetherianess.

In this paper, we concentrate on the case where K is the quotient field of D , studying subspaces of $\text{Zar}(K|D) = \text{Zar}(D)$ that are *not* compact. The starting point is a criterion based on semistar operations, proved in [8, Theorems 4.9 and 4.13] (see also [11, Proposition 4.5] for a slightly stronger version) and integrated, as in [9, Example 3.7], with

Date: November 24, 2017.

2010 Mathematics Subject Classification. Primary: 13F30; Secondary: 13A15, 13A18, 13B22, 54D30.

Key words and phrases. Zariski space; integral closure; valuation rings; semistar operations; spectral spaces; Kronecker function rings.

This work was partially supported by GNSAGA of *Istituto Nazionale di Alta Matematica*.

the use of the two-faced definition of the integral closure/ b -operation, either through valuation overrings or through equations of integral dependence (see e.g. [20, Chapter 6]). In particular, we analyze sets of the form $\text{Zar}(D) \setminus \{V\}$, where V is a minimal valuation overring of D : we show in Section 3 that such a space is compact only if V can be obtained from D in a very specific way (more precisely, as the integral closure of a localization of a finitely generated algebra over D), and we follow up in Sections 4 and 5 by showing that this condition implies a bound on the dimension of V in relation with the dimension of D (Proposition 4.3) and a quite strict condition on the intersection of sets of prime ideals of D (Theorem 5.1). Section 6 is dedicated to a brief application of these criteria to the study of Kronecker function rings (the definition will be recalled later).

In Section 7, we consider the set $\text{Over}(D)$ of overrings of D (which is known to be itself a spectral space [7, Proposition 3.5]). Using the result proved in the previous sections, we show that, when D is a Noetherian domain, some distinguished subspaces of $\text{Over}(D)$ (for example, the subspace of overrings of D that are Noetherian) are not spectral.

2. PRELIMINARIES AND NOTATION

2.1. Spectral spaces. A topological space X is a *spectral space* if there is a ring R such that X is homeomorphic to the prime spectrum $\text{Spec}(R)$, endowed with the Zariski topology. Spectral spaces can be characterized in a purely topological way as those spaces that are T_0 , compact, with a basis of open and compact subset that is closed by finite intersections and such that every irreducible closed subset has a generic point (i.e., it is the closure of a single point) [18, Proposition 4].

On a spectral space X it is possible to define two new topologies: the *inverse* and the *constructible* topology.

The *inverse topology* is the topology on X having, as a basis of closed sets, the family of open and compact subspaces of X . Endowed with the inverse topology, X is again a spectral space [18, Proposition 8]; moreover, a subspace $Y \subseteq X$ is closed in the inverse topology if and only if Y is compact (in the original topology) and $Y = Y^{\text{gen}}$ [8, Remark 2.2 and Proposition 2.6], where

$$\begin{aligned} Y^{\text{gen}} &:= \{z \in X \mid z \leq y \text{ for some } y \in Y\} = \\ &= \{z \in X \mid y \in \text{Cl}(z) \text{ for some } y \in Y\}, \end{aligned}$$

with $\text{Cl}(z)$ denoting the closure of the singleton $\{z\}$ (again, in the original topology) and \leq is the order induced by the original topology [17, d-1], which coincides on $\text{Spec}(R)$ with the set-theoretic inclusion.

The *constructible topology* on X (also called *patch topology*) is the coarsest topology such that the open and compact subsets of X are both open and closed. Endowed with the constructible topology, X

is a spectral space that is also Hausdorff (see [30, Propositions 3 and 5], [29] or [14, Proposition 5]), and the constructible topology is finer than both the original and the inverse topology. A subset of X closed in the constructible topology is said to be a *proconstructible subset* of X ; if Y is proconstructible, then it is a spectral space when endowed with the topology induced by the original spectral topology of X , and the constructible topology on Y is exactly the topology induced by the constructible topology on X (this follows from [3, 1.9.5(vi-vii)]).

2.2. Noetherian spaces. A topological space X is *Noetherian* if X verifies the ascending chain condition on the open subsets, or equivalently if every subspace of X is compact. Examples of Noetherian spaces are finite spaces and the prime spectra of Noetherian rings. If $\text{Spec}(R)$ is a Noetherian space, then every proper ideal of R has only finitely many minimal primes (see e.g. the proof of [2, Chapter 4, Corollary 3, p.102] or [1, Chapter 6, Exercises 5 and 7]).

2.3. Overrings and the Zariski space. Let $D \subseteq K$ be an extension of integral domains. We denote the set of all rings contained between D and K by $\text{Over}(K|D)$; if K is a field (not necessarily the quotient field of D), the set of all valuation rings containing D with quotient field K is denoted by $\text{Zar}(K|D)$, and it is called the *Zariski space* (or the *Zariski-Riemann space*) of D .

The *Zariski topology* on $\text{Over}(K|D)$ is the topology having, as a subbasis, the sets of the form

$$B(x_1, \dots, x_n) := \{T \in \text{Over}(K|D) \mid x_1, \dots, x_n \in T\},$$

as $\{x_1, \dots, x_n\}$ ranges among the finite subsets of K . Under this topology, both $\text{Over}(K|D)$ [7, Proposition 3.5] and its subspace $\text{Zar}(K|D)$ [5, 4] are spectral spaces, and the order induced by this topology is the inverse of the set-theoretic inclusion. In particular, every $Y \subseteq \text{Over}(K|D)$ with a minimum element is compact, and, if Z is an arbitrary subset of $\text{Over}(K|D)$, then $Z^{\text{gen}} = \{T \in \text{Over}(K|D) \mid T \supseteq A \text{ for some } A \in Z\}$.

We denote by $\text{Zar}_{\min}(D)$ the set of minimal elements of $\text{Zar}(D)$; since $\text{Zar}(D)$ is a spectral space, every $V \in \text{Zar}(D)$ contains an element $W \in \text{Zar}_{\min}(D)$.

If K is the quotient field of D , then we set $\text{Over}(K|D) =: \text{Over}(D)$ and $\text{Zar}(K|D) =: \text{Zar}(D)$. Elements of $\text{Over}(D)$ are called *overrings* of D , elements of $\text{Zar}(D)$ are the *valuation overrings* of D and elements of $\text{Zar}_{\min}(D)$ are the *minimal valuation overrings* of D .

The *center map* is the application

$$\begin{aligned} \gamma: \text{Zar}(K|D) &\longrightarrow \text{Spec}(D) \\ V &\longmapsto \mathfrak{m}_V \cap D, \end{aligned}$$

where \mathfrak{m}_V is the maximal ideal of V . When $\text{Zar}(K|D)$ and $\text{Spec}(D)$ are endowed with the respective Zariski topologies, the map γ is continuous ([34, Chapter VI, §17, Lemma 1] or [4, Lemma 2.1]), surjective (this follows, for example, from [1, Theorem 5.21] or [15, Theorem 19.6]) and closed [4, Theorem 2.5].

2.4. Semistar operations. Let D be a domain with quotient field K . Let $\mathbf{F}(D)$ be the set of D -submodules of K , $\mathcal{F}(D)$ be the set of fractional ideals of D , and $\mathcal{F}_f(D)$ be the set of finitely generated fractional ideals of D .

A *semistar operation* on D is a map $\star : \mathbf{F}(D) \rightarrow \mathbf{F}(D)$, $I \mapsto I^\star$, such that, for every $I, J \in \mathbf{F}(D)$ and every $x \in K$,

- (1) $I \subseteq I^\star$;
- (2) if $I \subseteq J$, then $I^\star \subseteq J^\star$;
- (3) $(I^\star)^\star = I^\star$;
- (4) $x \cdot I^\star = (xI)^\star$.

Given a semistar operation \star , the map \star_f is defined on every $E \in \mathbf{F}(D)$ by

$$E^{\star_f} = \bigcup \{F^\star \mid F \in \mathcal{F}_f(D), F \subseteq E\}.$$

The map \star_f is always a semistar operation; if $\star = \star_f$, then \star is said to be of *finite type*. Two semistar operations of finite type \star_1, \star_2 are equal if and only if $I^{\star_1} = I^{\star_2}$ for every $I \in \mathcal{F}_f(D)$. See [25] for general informations about semistar operations.

If $\Delta \subseteq \text{Zar}(D)$, then \wedge_Δ is defined as the semistar operation on D such that

$$I^{\wedge_\Delta} := \bigcap \{IV \mid V \in \Delta\}$$

for every D -submodule I of K ; a semistar operation of type \wedge_Δ is said to be a *valuative semistar operation*. By [11, Proposition 4.5], \wedge_Δ is of finite type if and only if Δ is compact (in the Zariski topology of $\text{Zar}(D)$). If $\Delta, \Lambda \subseteq \text{Zar}(D)$, then $\wedge_\Delta = \wedge_\Lambda$ if and only if $\Delta^{\text{gen}} = \Lambda^{\text{gen}}$ [10, Lemma 5.8(1)], while $(\wedge_\Delta)_f = (\wedge_\Lambda)_f$ if and only if Δ and Λ have the same closure with respect to the inverse topology [8, Theorem 4.9]. The semistar operation $\wedge_{\text{Zar}(D)}$ is usually denoted by b and called the *b-operation*.

3. THE USE OF MINIMAL VALUATION DOMAINS

The starting point of this paper is the following well-known result.

Proposition 3.1 (see e.g. [20, Proposition 6.8.2]). *Let I be an ideal of an integral domain D ; let $x \in D$. Then, $x \in IV$ for every $V \in \text{Zar}(D)$ if and only if there are $n \geq 1$ and $a_1, \dots, a_n \in D$ such that $a_i \in I^i$ and*

$$(1) \quad x^n + a_1x^{n-1} + \dots + a_{n-1}x + a_n = 0.$$

An inspection of the proof of the previous proposition given in [20] shows that this result does not really rely on the fact that I is an ideal of D , or on the fact that $x \in D$; indeed, it applies to every D -submodule I of the quotient field K , and to every $x \in K$. In the terminology of semistar operations, this means that, for each $I \in \mathbf{F}(D)$, $I^b = I^{\wedge_{\text{Zar}(D)}}$ is exactly the set of $x \in K$ that verifies an equation like (1), with $a_i \in I^i$. We are interested in generalizing that proof in a different way; we need the following definitions.

Definition 3.2. *Let D be an integral domain and let $\Delta, \Lambda \subseteq \text{Over}(D)$. We say that Λ dominates Δ if, for every $T \in \Delta$ and every $M \in \text{Max}(T)$, there is a $A \in \Lambda$ such that $T \subseteq A$ and $1 \notin MA$.*

For example, $\text{Zar}(D)$ dominates every subset of $\text{Over}(D)$, while the set of localizations of D dominates $\{D\}$.

Definition 3.3. *Let D be an integral domain. We denote by $D[\mathcal{F}_f]$ the set of finitely generated D -algebras of $\text{Over}(D)$, or equivalently*

$$D[\mathcal{F}_f] := \{D[I] : I \in \mathcal{F}_f(D)\}.$$

Even if the proof of the following result essentially repeats the proof of [20, Proposition 6.8.2], we replay it here for clarity.

Proposition 3.4. *Let D be an integral domain, and suppose that $\Delta \subseteq \text{Zar}(D)$ dominates $D[\mathcal{F}_f]$. Then, for every finitely generated ideal I of D , $I^{\wedge_{\Delta}} = I^b$.*

Proof. Clearly, $I^b \subseteq I^{\wedge_{\Delta}}$. Suppose thus that $x \in I^{\wedge_{\Delta}}$, $x \neq 0$, and let $I = (i_1, \dots, i_k)D$. Define $J := x^{-1}I \in \mathcal{F}_f(D)$, and let $A := D[J] = D[x^{-1}i_1, \dots, x^{-1}i_k]$; by definition, $J \subseteq A$.

If $JA \neq A$, then there is a maximal ideal M of A containing J , and thus, by domination, there is a valuation domain $V \in \Delta$ containing A whose maximal ideal \mathfrak{m}_V is such that $JV \subseteq \mathfrak{m}_V$, and thus $IV \subseteq x\mathfrak{m}_V$. However, $x \in I^b \subseteq IV$, which implies $x \in x\mathfrak{m}_V$, a contradiction.

Hence, $JA = A$, i.e., $1 = j_1a_1 + \dots + j_na_n$ for some $j_t \in J$, $a_t \in A$; expliciting the elements of A as elements of $D[J]$ and using $J = x^{-1}I$, we find that there must be an $N \in \mathbb{N}$ and elements $i_t \in I^t$ such that $x^N = i_1x^{N-1} + \dots + i_{N-1}x + i_N$, which gives an equation of integral dependence of x over I . Therefore, $x \in I^b$, as requested. \square

We can now use the properties of valuative semistar operations to study compactness.

Proposition 3.5. *Let D be an integral domain, and let $\Delta \subseteq \text{Zar}(D)$ be a set that dominates $D[\mathcal{F}_f]$. Then, Δ is compact if and only if it contains $\text{Zar}_{\min}(D)$.*

Proof. If Δ contains $\text{Zar}_{\min}(D)$, then \mathcal{U} is an open cover of Δ if and only if it is an open cover of $\text{Zar}(D)$; thus, Δ is compact since $\text{Zar}(D)$ is.

Conversely, suppose Δ is compact. By Proposition 3.4, $I^{\wedge\Delta} = I^b$ for every finitely generated ideal I ; hence, $(\wedge_{\Delta})_f = b_f = b$. By [10, Lemma 5.8(1)], it follows that the closure of Δ with respect to the inverse topology of $\text{Zar}(D)$ is the whole $\text{Zar}(D)$; however, since Δ is compact, its closure in the inverse topology is exactly $\Delta^{\text{gen}} = \Delta^{\dagger} = \{W \in \text{Zar}(D) \mid W \supseteq V \text{ for some } V \in \Delta\}$. Hence, Δ must contain $\text{Zar}_{\min}(D)$. \square

Thus, to find a subset of $\text{Zar}(D)$ that is not compact, it is enough to find a Δ that dominates $D[\mathcal{F}_f]$ but that does not contain $\text{Zar}_{\min}(D)$. The easiest case where this criterion can be applied is when $\Delta = \text{Zar}(D) \setminus \{V\}$ for some $V \in \text{Zar}_{\min}(D)$.

Theorem 3.6. *Let D be an integral domain and let $V \in \text{Zar}_{\min}(D)$. If $\text{Zar}(D) \setminus \{V\}$ is compact, then V is the integral closure of $D[x_1, \dots, x_n]_M$ for some $x_1, \dots, x_n \in K$ and some $M \in \text{Max}(D[x_1, \dots, x_n])$.*

Proof. If $\Delta := \text{Zar}(D) \setminus \{V\}$ is compact, then by Proposition 3.5 it cannot dominate $D[\mathcal{F}_f]$. Hence, there is a finitely generated fractional ideal I such that Δ does not dominate $A := D[I]$, and so a maximal ideal M of A such that $1 \in MW$ for every $W \in \Delta$. In particular, $A \neq K$ (otherwise M would be (0)).

However, there must be a valuation ring containing A_M whose center (on A_M) is MA_M , and the unique possibility for this valuation ring is V : it follows that V is the unique valuation ring centered on MA_M . However, the integral closure of A_M is the intersection of the valuation rings with center MA_M (since every valuation ring containing A_M contains a valuation ring centered on MA_M [15, Corollary 19.7]); thus, V is the integral closure of A_M . \square

4. THE DIMENSION OF V

Before embarking on using Theorem 3.6, we prove a simple yet general result.

Proposition 4.1. *Let D be an integral domain. If $\text{Zar}(D)$ is a Noetherian space, so is $\text{Spec}(D)$.*

Proof. The claim follows from the fact that $\text{Spec}(D)$ is the continuous image of $\text{Zar}(D)$ through the center map γ , and that the image of a Noetherian space is still Noetherian. \square

Note that the converse of this proposition is far from being true (this is, for example, a consequence of Proposition 5.4 or of Proposition 7.1).

The problem in using Theorem 3.6 is that it is usually difficult to control the behaviour of finitely generated algebras over D . We can, however, control the behaviour of the prime spectrum of D .

$$\begin{array}{ccccccc}
& & D[X_1, \dots, X_n] & \hookrightarrow & D[X_1, \dots, X_n]_{\mathfrak{a}} & & \\
& \nearrow & \downarrow & & \downarrow & & \\
D & \hookrightarrow & A = D[a_1, \dots, a_n] & \hookrightarrow & A_M \simeq \frac{D[X_1, \dots, X_n]_{\mathfrak{a}}}{\mathfrak{b}} & \hookrightarrow & V
\end{array}$$

FIGURE 1. Rings involved in the proof of Proposition 4.3.

Lemma 4.2. *Let D be an integral domain, and let $V \in \text{Zar}(D)$ be the integral closure of D_M , for some $M \in \text{Spec}(D)$. Then, the set of prime ideals of D contained in M is linearly ordered.*

Proof. Let P, Q be two prime ideals of D contained in M ; then, $PD_M, QD_M \in \text{Spec}(D_M)$. Since $D_M \subseteq V$ is an integral extension, $PD_M = P' \cap D_M$ and $QD_M = Q' \cap D_M$ for some $P', Q' \in \text{Spec}(V)$; however, V is a valuation domain, and thus (without loss of generality) $P' \subseteq Q'$. Hence, $PD_M \subseteq QD_M$ and $P \subseteq Q$, as requested. \square

Proposition 4.3. *Let D be an integral domain, let $V \in \text{Zar}_{\min}(D)$ and suppose that $\text{Zar}(D) \setminus \{V\}$ is compact. Let $\iota_V : \text{Spec}(V) \rightarrow \text{Spec}(D)$ be the canonical spectral map associated to the inclusion $D \hookrightarrow V$. For every $P \in \text{Spec}(D)$, $|\iota_V^{-1}(P)| \leq 2$; in particular, $\dim(V) \leq 2 \dim(D)$.*

Proof. Suppose $|\iota_V^{-1}(P)| > 2$: then, there are prime ideals $Q_1 \subsetneq Q_2 \subsetneq Q_3$ of V such that $\iota_V(Q_1) = \iota_V(Q_2) = \iota_V(Q_3) =: P$. If $\text{Zar}(D) \setminus \{V\}$ is compact, by Theorem 3.6 there is a finitely generated D -algebra $A := D[a_1, \dots, a_n]$ such that V is the integral closure of A_M , for some maximal ideal M of A . We can write A_M as a quotient $\frac{D[X_1, \dots, X_n]_{\mathfrak{a}}}{\mathfrak{b}}$, where X_1, \dots, X_n are independent indeterminates and $\mathfrak{a}, \mathfrak{b} \in \text{Spec}(D[X_1, \dots, X_n])$. Since $A_M \subseteq V$ is an integral extension, $Q_i \cap A \neq Q_j \cap A$ if $i \neq j$.

For $i \in \{1, 2, 3\}$, let \mathfrak{q}_i be the prime ideal of $D[X_1, \dots, X_n]$ whose image in A is Q_i ; then, $\mathfrak{q}_1, \mathfrak{q}_2$ and \mathfrak{q}_3 are distinct, $\mathfrak{q}_i \cap D = P$ for each i , and the set of ideals between \mathfrak{q}_1 and \mathfrak{q}_3 is linearly ordered (by Lemma 4.2). However, the prime ideals of $D[X_1, \dots, X_n]$ contracting to P are in a bijective and order-preserving correspondence with the prime ideals of $F[X_1, \dots, X_n]$, where F is the quotient field of D/P ; since $F[X_1, \dots, X_n]$ is a Noetherian ring, there are an infinite number of prime ideals between the ideals corresponding to \mathfrak{q}_1 and \mathfrak{q}_3 . This is a contradiction, and $|\iota_V^{-1}(P)| \leq 2$.

For the ‘‘in particular’’ statement, take a chain $(0) \subsetneq Q_1 \subsetneq \dots \subsetneq Q_k$ in $\text{Spec}(V)$. Then, the corresponding chain of the $P_i := Q_i \cap D$ has length at most $\dim(D)$, and moreover $\iota^{-1}((0)) = \{(0)\}$. Hence, $k+1 \leq 2 \dim(D) + 1$ and $\dim(V) \leq 2 \dim(D)$. \square

The *valuative dimension* of D , indicated by $\dim_v(D)$, is defined as the supremum of the dimensions of the valuation overrings of D ; we have always $\dim(D) \leq \dim_v(D)$, and $\dim_v(D)$ can be arbitrarily large

with respect to $\dim(D)$ [15, Section 30, Exercises 16 and 17]. In particular, with the notation of the previous proposition, the cardinality of $\iota_V^{-1}(P)$ can be arbitrarily large: for example, if (D, \mathfrak{m}) is local and one-dimensional, then $|\iota_V^{-1}(\mathfrak{m})| = \dim_v(D)$.

Corollary 4.4. *Let D be an integral domain such that $\text{Zar}(D)$ is Noetherian. Then, $\dim_v(D) \leq 2 \dim(D)$.*

Proof. If $\text{Zar}(D)$ is Noetherian, then in particular $\text{Zar}(D) \setminus \{V\}$ is compact for every $V \in \text{Zar}_{\min}(D)$. Hence, $\dim(V) \leq 2 \dim(D)$ for every $V \in \text{Zar}_{\min}(D)$, by Proposition 4.3; since, if $W \supseteq V$ are valuation domain, $\dim(W) \leq \dim(V)$, the claim follows. \square

Proposition 4.5. *Let D be an integral domain, and let $V \in \text{Zar}_{\min}(D)$ be such that $\text{Zar}(D) \setminus \{V\}$ is compact; let $(0) \subsetneq P_1 \subsetneq \dots \subsetneq P_k$ be the chain of prime ideals of V and let $Q_i := P_i \cap D$. Denote by $ht(P)$ the height of the prime ideal P . Then:*

(a) *for every $0 \leq t \leq \dim(D)$, we have*

$$\dim(V) \leq \dim_v(D_{Q_t}) + 2(\dim(D) - ht(Q_t));$$

(b) *if D_{Q_t} is a valuation domain, then*

$$\dim(V) \leq 2 \dim(D) - ht(Q_t).$$

Proof. (a) Let $(0) \subsetneq Q^{(1)} \subsetneq Q^{(2)} \subsetneq \dots \subsetneq Q^{(s)}$ be the chain $(0) \subseteq Q_1 \subseteq \dots \subseteq Q_k$ without the repetitions, and let a be the index such that $Q^{(a)} = Q_t$. For every $b > a$, by the proof of Proposition 4.3 there can be at most two prime ideals of V over $Q^{(b)}$; on the other hand, V_{P_t} is a valuation overring of D_{Q_t} , and thus $t = \dim(V_{P_t}) \leq \dim_v(D_{Q_t})$. Therefore,

$$\dim(V) \leq t + 2(s - a) \leq \dim_v(D_{Q_t}) + 2(\dim(D) - ht(Q_t))$$

since each ascending chain of prime ideals starting from Q_t has length at most $\dim(D) - ht(Q_t)$.

Point (b) follows, since $\dim(V) = \dim_v(V)$ for every valuation domain V . \square

Example 4.6. A class of integral domain whose Zariski space is Noetherian is constituted by the class of Prüfer domains with Noetherian spectrum. Indeed, if D is a Prüfer domain then the valuation overrings of D are exactly the localizations of D at prime ideals; thus, the center map γ establishes a homeomorphism between $\text{Zar}(D)$ and $\text{Spec}(D)$. Thus, if the latter is Noetherian also the former is Noetherian.

In this case, $\dim(D) = \dim_v(D)$.

Example 4.7. It is also possible to construct domains whose Zariski space is Noetherian but with $\dim(D) \neq \dim_v(D)$. For example, let L be a field, and consider the ring $A := L + YL(X)[[Y]]$, where X and Y are independent indeterminates. Then, the valuation overrings

of A different from $F := L(X)((Y))$ are the rings in the form $V + YL(X)[[Y]]$, as V ranges among the valuation rings containing L and having quotient field $L(X)$; that is, $\text{Zar}(A) \setminus \{F\} \simeq \text{Zar}(L(X)|L)$. By the following Corollary 5.5, $\text{Zar}(A)$ is a Noetherian space.

From this, we can construct analogous examples of arbitrarily large dimension. Indeed, if R is an integral domain with quotient field K , and $T := R + XK[[X]]$, then as above $\text{Zar}(T)$ is composed by $K((X))$ and by rings of the form $V + XK[[X]]$, as V ranges in $\text{Zar}(R)$; in particular, $\text{Zar}(T) = \{K((X))\} \cup \mathcal{X}$, where $\mathcal{X} \simeq \text{Zar}(R)$. Thus, $\text{Zar}(T)$ is Noetherian if $\text{Zar}(R)$ is. Moreover, $\dim(T) = \dim(R) + 1$ and $\dim_v(T) = \dim_v(R) + 1$.

Consider now the sequence of rings $R_1 := L + YL(X)[[Y]]$, $R_2 := R_1 + Y_2Q(R_1)[[Y_2]]$, \dots , $R_n := R_{n-1} + Y_nQ(R_{n-1})[[Y_n]]$, where $Q(R)$ indicates the quotient field of R and each Y_i is an indeterminate over $Q(R_{i-1})((Y_{i-1}))$. Recursively, we see that each $\text{Zar}(R_n)$ is Noetherian, while $\dim(R_n) = n \neq n + 1 = \dim_v(R_n)$.

5. INTERSECTIONS OF PRIME IDEALS

The results of the previous sections, while very general, are often difficult to apply, because it is usually not easy to determine the valuative dimension of a domain D . More applicable criteria, based on the prime spectrum of D , are the ones that we will prove next.

Theorem 5.1. *Let D be a local integral domain, and suppose there is a set $\Delta \subseteq \text{Spec}(D)$ and a prime ideal Q such that:*

- (1) $Q \notin \Delta$;
- (2) no two members of Δ are comparable;
- (3) $\bigcap \{P \mid P \in \Delta\} = Q$;
- (4) D_Q is a valuation domain.

Then, for any minimal valuation overring V of D contained in D_Q , $\text{Zar}(D) \setminus \{V\}$ is not compact; in particular, $\text{Zar}(D)$ is not Noetherian.

Proof. Note first that, since V is a minimal valuation overring, its center M on D must be the maximal ideal of D [15, Corollary 19.7]. Suppose that $\text{Zar}(D) \setminus \{V\}$ is compact: by Theorem 3.6, there is a finitely generated D -algebra $A := D[x_1, \dots, x_n]$ such that V is the integral closure of A_M for some $M \in \text{Max}(A)$.

Let $I := x_1^{-1}D \cap \dots \cap x_n^{-1}D \cap D = (D :_D x_1) \cap \dots \cap (D :_D x_n)$. If $I \subseteq Q$, then $(D :_D x) \subseteq Q$ for some $x_i := x$; then, since D_Q is flat over D ,

$$(D_Q :_{D_Q} x) = (D :_D x)D_Q \subseteq QD_Q,$$

and in particular $x \notin D_Q$. However, $V \subseteq D_Q$, and thus $x \notin V$, a contradiction. Hence, we must have $I \not\subseteq Q$.

In this case, there must be a prime ideal $P_1 \in \Delta$ not containing I . Moreover, $I \cap P_1 \not\subseteq Q$ too, and thus there is another prime $P_2 \in \Delta$,

$P_1 \neq P_2$, not containing I . By Lemma 4.2, the prime ideals of A inside M are linearly ordered; in particular, we can suppose without loss of generality that $\text{rad}(P_2A) \subseteq \text{rad}(P_1A)$.

Let now $t \in P_2 \setminus P_1$; then, $t \in \text{rad}(P_1A)$, and thus there are $p_1, \dots, p_k \in P_1$, $a_1, \dots, a_n \in A$ such that $t^e = p_1a_1 + \dots + p_ka_k$ for some positive integer e . For each i , $a_i = B_i(x_1, \dots, x_n)$, where B_i is a polynomial over D of total degree d_i ; let $d := \sup\{d_1, \dots, d_k\}$, and take an $r \in I \setminus P_1$ (recall that $I \not\subseteq P_1$). Then, $r^d B_i(x_1, \dots, x_n) \in D$ for each i ; therefore,

$$r^{dt^e} = p_1 r^d a_1 + \dots + p_k r^d a_k \in p_1 D + \dots + p_k D \subseteq P_1.$$

However, by construction, both r and t are out of P_1 ; since P_1 is prime, this is impossible. Hence, $\text{Zar}(D) \setminus \{V\}$ is not compact, and $\text{Zar}(D)$ is not Noetherian. \square

The first corollaries of this result can be obtained simply by putting $Q = (0)$. Recall that a *G-domain* (or *Goldman domain*) is an integral domain such that the intersection of all nonzero prime ideals is nonzero. They were introduced by Kaplansky for giving a new proof of Hilbert's Nullstellensatz (see for example [22, Section 1.3]).

Corollary 5.2. *Let D be a local domain of finite dimension, and suppose that D is not a G-domain. Then, $\text{Zar}(D) \setminus \{V\}$ is not compact for every $V \in \text{Zar}_{\min}(D)$.*

Proof. Since D is finite-dimensional, every prime ideal of D contains a prime ideal of height 1; since D is not a G-domain, it follows that the intersection of the set $\text{Spec}^1(D)$ of the height-1 prime ideals of D is (0) . The localization $D_{(0)}$ is the quotient field of D , and thus a valuation domain; therefore, we can apply Theorem 5.1 to $\Delta := \text{Spec}^1(D)$. \square

Corollary 5.3. *Let D be a local domain. If D has infinitely many height-1 primes, then $\text{Zar}(D)$ is not Noetherian.*

Proof. Let I be the intersection of all height-1 prime ideals. If $I \neq (0)$, every height-one prime of D would be minimal over I ; since there is an infinite number of them, $\text{Spec}(D)$ would not be Noetherian, and by Proposition 4.1 neither $\text{Zar}(D)$ would be Noetherian. Hence, $I = (0)$. But then we can apply Theorem 5.1 (for $Q = I$). \square

Note that the hypothesis that D is local is needed in Theorem 5.1 and in Corollary 5.3: for example, \mathbb{Z} has infinitely many height-1 primes, and $\bigcap\{P \mid P \in \text{Spec}^1(D)\} = (0)$, but $\text{Zar}(\mathbb{Z}) \simeq \text{Spec}(\mathbb{Z})$ is a Noetherian space.

Proposition 5.4. *Let D be an integral domain. If D is not a field, then $\text{Zar}(D[X])$ is not a Noetherian space.*

Proof. Since D is not a field, there exist a nonzero prime ideal P of D . For any $a \in P$, let \mathfrak{p}_a be the ideal of $D[X]$ generated by $X - a$;

then, each \mathfrak{p}_a is a prime ideal of height 1, $\mathfrak{p}_a \neq \mathfrak{p}_b$ if $a \neq b$, and $\bigcap\{\mathfrak{p}_a \mid a \in P\} = (0)$.

The prime ideal $\mathfrak{m} := PD[X] + XD[X]$ contains every \mathfrak{p}_a ; by Corollary 5.3, $\text{Zar}(D[X]_{\mathfrak{m}})$ is not Noetherian. Therefore, neither $\text{Zar}(D[X])$ is Noetherian. \square

Corollary 5.5. *Let $F \subseteq L$ be a transcendental field extension.*

- (a) *If $\text{trdeg}_F(L) = 1$ and L is finitely generated over F then $\text{Zar}(L|F)$ is Noetherian.*
- (b) *If $\text{trdeg}_F(L) > 1$ then $\text{Zar}(L|F)$ is not Noetherian.*

Proof. (a) Let $L = F(\alpha_1, \dots, \alpha_n)$; without loss of generality we can suppose that α_1 is transcendental over F . Then, the extension $F(\alpha_1) \subseteq L$ is algebraic and finitely generated, and thus finite.

Each $V \in \text{Zar}(L|F)$ must contain either α_1 or α_1^{-1} ; therefore, $\text{Zar}(L|F) = \text{Zar}(L|F[\alpha_1]) \cup \text{Zar}(L|F[\alpha_1^{-1}])$. However, $\text{Zar}(L|A) = \text{Zar}(A')$ for every domain A , where we denote by A' is the integral closure of A in L ; since $F[\alpha_1]$ (respectively, $F[\alpha_1^{-1}]$) is a principal ideal domain and $F(\alpha_1) \subseteq L$ is finite, the integral closure of $F[\alpha_1]$ (resp., $F[\alpha_1^{-1}]$) is a Dedekind domain, and thus $\text{Zar}(L|F[\alpha_1]) = \text{Zar}(F[\alpha_1]') \simeq \text{Spec}(F[\alpha_1]')$ is Noetherian. Being the union of two Noetherian spaces, $\text{Zar}(L|F)$ is itself Noetherian.

(b) Suppose $\text{trdeg}_F(L) > 1$. Then, there are $X, Y \in L$ such that $\{X, Y\}$ is an algebraically independent set over F ; in particular, we have a continuous surjective map $\text{Zar}(L|F) \longrightarrow \text{Zar}(F(X, Y)|F)$ given by $V \mapsto V \cap F(X, Y)$. However, $\text{Zar}(F(X, Y)|F)$ contains $\text{Zar}(F[X, Y])$; by Proposition 5.4, the latter is not Noetherian, since $F[X, Y]$ is the polynomial ring over $F[X]$, a domain of dimension 1. Thus, $\text{Zar}(L|F)$ is not Noetherian. \square

The condition that $\bigcap\{P \mid P \in \Delta\} = Q$ of Theorem 5.1 can be slightly generalized, requiring only that the intersection is contained in Q . However, doing so we can only prove that $\text{Zar}(D)$ is not Noetherian, without always finding a specific V such that $\text{Zar}(D) \setminus \{V\}$ is not compact.

Proposition 5.6. *Let D be a local integral domain, and suppose there is a set $\Delta \subseteq \text{Spec}(D)$ and a prime ideal Q such that:*

- (1) $Q \notin \Delta$;
- (2) *no two members of Δ are comparable;*
- (3) $\bigcap\{P \mid P \in \Delta\} \subseteq Q$;
- (4) D_Q is a valuation domain.

Then, $\text{Zar}(D)$ is not Noetherian.

Proof. If $\text{Spec}(D)$ is not Noetherian, by Proposition 4.1 neither is $\text{Zar}(D)$; suppose that $\text{Spec}(D)$ is Noetherian.

Let $I := \bigcap\{P \mid P \in \Delta\}$; since an overring of a valuation domain is still a valuation domain, we can suppose that Q is a minimal prime

of I . Since D has Noetherian spectrum, the radical ideal I has only a finite number of minimal primes, say $Q =: Q_1, Q_2, \dots, Q_n$; let $\Delta_i := \{\mathfrak{p} \in \Delta \mid Q_i \subseteq \mathfrak{p}\}$ and $I_i := \bigcap \{\mathfrak{p} \mid \mathfrak{p} \in \Delta_i\}$. By standard properties of minimal primes, $\Delta = \Delta_1 \cup \dots \cup \Delta_n$ and $I = I_1 \cap \dots \cap I_n$.

In particular, $I_1 \cap \dots \cap I_n \subseteq Q$; hence, $I_k \subseteq Q$ for some k . However, $Q_k \subseteq I_k$, and thus $Q_k \subseteq Q$; since different minimal primes of the same ideal are not comparable, $k = 1$ and $Q \subseteq I_1 \subseteq Q$, i.e., $I_1 = Q$. Then, Δ_1 is a family of primes satisfying the hypothesis of Theorem 5.1; in particular, $\text{Zar}(D)$ is not Noetherian. \square

An *essential prime* of a domain D is a $P \in \text{Spec}(D)$ such that D_P is a valuation domain. D is an *essential domain* if it is equal to the intersection of the localizations of D at the essential primes. If, moreover, the family of the essential primes is compact, then D can be called a *Prüfer v -multiplication domain* (*PvMD* for short) [12, Corollary 2.7]; note that the original definition of PvMDs was given through star operations (more precisely, D is a PvMD if and only if D_P is a valuation ring for every t -maximal ideal P [16, 21]).

Proposition 5.7. *Let D be an essential domain. Then, $\text{Zar}(D)$ is Noetherian if and only if D is a Prüfer domain with Noetherian spectrum.*

Proof. If D is a Prüfer domain with Noetherian spectrum, then $\text{Zar}(D) \simeq \text{Spec}(D)$ is Noetherian (see Example 4.6). Conversely, suppose $\text{Zar}(D)$ is Noetherian: by Proposition 4.1, $\text{Spec}(D)$ is Noetherian. Let \mathcal{E} be the set of essential prime ideals of D : since $\text{Spec}(D)$ is Noetherian, \mathcal{E} is compact, and thus D is a PvMD.

Suppose by contradiction that D is not a Prüfer domain. Then, there is a maximal ideal M of D such that D_M is not a valuation domain; since the localization of a PvMD is a PvMD [21, Theorem 3.11], and $\text{Zar}(D_M)$ is a subspace of $\text{Zar}(D)$, without loss of generality we can suppose $D = D_M$, i.e., we can suppose that D is local.

Since \mathcal{E} is compact, every $P \in \mathcal{E}$ is contained in a maximal element of \mathcal{E} ; let Δ be the set of such maximal elements. Clearly, $D = \bigcap \{D_P \mid P \in \Delta\}$. If Δ were finite, D would be an intersection of finitely many valuation domains, and thus it would be a Prüfer domain [15, Theorem 22.8]; hence, we can suppose that Δ is infinite. Let $I := \bigcap \{P \mid P \in \Delta\}$.

Each $P \in \Delta$ contains a minimal prime of I ; however, since $\text{Spec}(D)$ is Noetherian, I has only finitely many minimal primes. It follows that there is a minimal prime Q of I that is not contained in Δ ; in particular, $\bigcap \{P \mid P \in \Delta\} \subseteq Q$, and thus we can apply Proposition 5.6. Hence, $\text{Zar}(D)$ is not Noetherian, which is a contradiction. \square

Remark 5.8. The previous proof can be interpreted using the terminology of the theory of star operations. Indeed, any essential prime P is a t -ideal, i.e., $P = P^t$, where (for any ideal J of D) $J^t := \bigcup \{D \mid$

$(D : I) \mid I \subseteq J$ is finitely generated} [21, Lemma 3.17] and if D is a PvMD then the set Δ of the maximal elements of \mathcal{E} is exactly the set of *t-maximal ideals*, i.e., the set of the ideals I such that $I = I^t$ and $J \neq J^t$ for every proper ideal $I \subsetneq J$.

Corollary 5.9. *Let D be a Krull domain. Then, $\text{Zar}(D)$ is Noetherian if and only if $\dim(D) = 1$, i.e., if and only if D is a Dedekind domain.*

Proof. If $\dim(D) = 1$ then D is Noetherian and so is $\text{Zar}(D)$. If $\dim(D) > 1$, then D is not a Prüfer domain; since each Krull domain is a PvMD, we can apply Proposition 5.7. \square

Note that this corollary can also be proved directly from Corollary 5.3 since, if D is Krull, and $P \in \text{Spec}(D)$ has height 2 or more, then D_P has infinitely many height-1 primes.

6. AN APPLICATION: KRONECKER FUNCTION RINGS

Let D be an integrally closed integral domain with quotient field K . For every $V \in \text{Zar}(D)$, let $V(X) := V[X]_{\mathfrak{m}_V[X]} \subseteq K(X)$, where \mathfrak{m}_V is the maximal ideal of V . If $\Delta \subseteq \text{Zar}(D)$, the *Kronecker function ring* of D with respect to Δ is

$$\text{Kr}(D, \Delta) := \bigcap \{V(X) \mid V \in \Delta\};$$

equivalently,

$$\text{Kr}(D, \Delta) = \{f/g \mid f, g \in D[X], g \neq 0, \mathbf{c}(f) \subseteq (\mathbf{c}(g))^{\wedge \Delta}\},$$

where $\mathbf{c}(f)$ is the content of f and \wedge_{Δ} is the semistar operation defined in Section 2.4. See [15, Chapter 32] or [13] for general properties of Kronecker function rings.

The set of Kronecker function rings is exactly the set of overrings of the basic Kronecker function ring $\text{Kr}(D, \text{Zar}(D))$; this set is in bijective correspondence with the set of finite-type valuative semistar operations [15, Remark 32.9], or equivalently with the set of nonempty subsets of $\text{Zar}(D)$ that are closed in the inverse topology [8, Theorem 4.9].

Let $\mathcal{K}(D)$ be the set of Kronecker function rings T of D such that $T \cap K = D$. Then, $\mathcal{K}(D)$ is in bijective correspondence with the set of finite-type valuative *star* operations, or equivalently with the set of inverse-closed representation of D through valuation rings, i.e., the sets $\Delta \subseteq \text{Zar}(D)$ that are closed in the inverse topology and such that $\bigcap \{V \mid V \in \Delta\} = D$ [27, Proposition 5.10].

It has been conjectured [23] that $\mathcal{K}(D)$ is either a singleton (in which case D is said to be a *vacant domain*; see [6]) or infinite, and this has been proved to be the case for a wide class of pseudo-valuation domains [6, Theorem 4.10]. As a consequence of the following proposition, we will prove this conjecture for another class of domains.

Proposition 6.1. *Let D be an integrally closed integral domain such that $1 < |\mathcal{K}(D)| < \infty$. Then, there is a minimal valuation overring V of D such that $\text{Zar}(D) \setminus \{V\}$ is compact.*

Proof. Suppose $|\mathcal{K}(D)| > 1$. Then, there is an inverse-closed representation Δ of D different from $\text{Zar}(D)$; let $\Lambda := \text{Zar}(D) \setminus \Delta$. For each $W \in \Lambda$, let $\Delta(W) := \Delta \cup \{W\}^\uparrow$; then, every $\Delta(W)$ is an inverse-closed representation of D , and $\Delta(W) \neq \Delta(W')$ if $W \neq W'$ (since, without loss of generality, $W \not\subseteq W'$, and thus $W \notin \Delta(W')$). Hence, each $W \in \Lambda$ give rise to a different member of $\mathcal{K}(D)$; since $|\mathcal{K}(D)| < \infty$, it follows that Λ is finite.

If now V is minimal in Λ , then $\text{Zar}(D) \setminus \{V\} = \Delta \cup (\Lambda \setminus \{V\})$ is closed by generizations; since Λ is finite, it follows that $\text{Zar}(D) \setminus \{V\}$ is the union of two compact subspaces, and thus it is itself compact. \square

Corollary 6.2. *Let D be an integrally closed local integral domain, and suppose there exist a set $\Delta \subseteq \text{Spec}(D)$ of incomparable nonzero prime ideals such that $\bigcap \{P \mid P \in \Delta\} = (0)$. Then, $|\mathcal{K}(D)| \in \{1, \infty\}$.*

Proof. By Theorem 5.1, each $\text{Zar}(D) \setminus \{V\}$ is noncompact. The claim now follows from Proposition 6.1. \square

7. OVERRINGS OF NOETHERIAN DOMAINS

If D is a Noetherian domain, Theorem 3.6 admits a direct application, without using any of the results proved in Sections 4 and 5. Indeed, if D is Noetherian with quotient field K , then it is the same for any localization of $D[x_1, \dots, x_n]$, for arbitrary $x_1, \dots, x_n \in K$; thus, the integral closure of $D[x_1, \dots, x_n]_M$ is a Krull domain for each maximal ideal M of $D[x_1, \dots, x_n]$ ([24, (33.10)] or [20, Theorem 4.10.5]). Since a domain that is both Krull and a valuation ring must be a field or a discrete valuation ring, Theorem 3.6 implies that $\text{Zar}(D) \setminus \{V\}$ is not compact as soon as V is a minimal valuation overring of dimension 2 or more.

We can actually say more than this; the following is a proof through Proposition 3.5 of an observation already appeared in [9, Example 3.7].

Proposition 7.1. *Let D be a Noetherian domain with quotient field K , and let Δ be the set of valuation overrings of D that are Noetherian (i.e., Δ is the union of $\{K\}$ with the set of discrete valuation overrings of D). Then, Δ is compact if and only if $\dim(D) = 1$.*

Proof. If $\dim(D) = 1$, then $\Delta = \text{Zar}(D)$, and thus it is compact.

On the other hand, for every ideal I of D , $I^{\Delta} = I^b$ [20, Proposition 6.8.4]; however, if $\dim(D) > 1$, then $\text{Zar}(D)$ contains elements of dimension 2, and thus Δ cannot contain $\text{Zar}_{\min}(D)$. The claim now follows from Proposition 3.5. \square

Remark 7.2.

- (1) The equality $I^{\Delta} = I^b$ holds also if we restrict Δ to be the set of discrete valuation overrings of D whose center is a maximal ideal of D [20, Proposition 6.8.4]. For each prime ideal of height 2 or more, by passing to D_P , we can thus prove that the set of discrete valuation overrings of D with center P is not compact (and in particular it is infinite).
- (2) The previous proposition also allows a proof of the second part of Corollary 5.5 without using Theorem 5.1, since $F[X, Y]$ is a Noetherian domain of dimension 2.

By Proposition 7.1, in particular, the space Δ of Noetherian valuation overrings of D (where D is Noetherian and $\dim(D) \geq 2$) is not a spectral space, since it is not compact. Our next purpose is to see Δ as an intersection $X \cap \text{Zar}(D)$, for some subset X of $\text{Over}(D)$, and use this representation to prove facts about X . We start with using the inverse topology.

Proposition 7.3. *Let D be a Noetherian domain with quotient field K , and let:*

- X_1 be the set of all overrings of D that are Noetherian and of dimension at most 1;
- X_2 be the set of all overrings of D that are Dedekind domains (K included).

For $i \in \{1, 2\}$, the following are equivalent:

- (i) X_i is compact;
- (ii) X_i is spectral;
- (iii) X_i is proconstructible in $\text{Over}(D)$;
- (iv) $\dim(D) = 1$.

Proof. (i) \implies (iii). In both cases, $X = X^{\text{gen}}$: for X_1 see [22, Theorem 93], while for X_2 see e.g. [15, Theorem 40.1] (or use the previous result and [15, Corollary 36.3]). (iii) \implies (ii) \implies (i) always holds.

(iv) \implies (i). If $\dim(D) = 1$, then $X_1 = \text{Over}(D)$, while $X_2 = \text{Over}(D')$, where D' is the integral closure of D , and both are compact since they have a minimum.

(iii) \implies (iv). If X_i is proconstructible, so is $X_i \cap \text{Zar}(D)$ (since $\text{Zar}(D)$ is also proconstructible), and in particular $X_i \cap \text{Zar}(D)$ is compact. However, in both cases, $X_i \cap \text{Zar}(D)$ is exactly the set of Noetherian valuation overrings of D ; by Proposition 7.1, $\dim(D) = 1$. \square

Remark 7.4. The equivalence between the first three conditions of Proposition 7.3 holds for every subset $X \subseteq \text{Over}(D)$ such that $X = X^{\text{gen}}$ (and every domain D). In particular, it holds if X is the set of overrings of D that are principal ideal domains, and, with the same proof of the other cases, we can show that if D is Noetherian and these conditions hold, then $\dim(D) = 1$. However, it is not clear if, when D is Noetherian and $\dim(D) = 1$, this set is actually compact.

Another immediate consequence of Proposition 7.1 is that the set $\text{NoethOver}(D)$ of Noetherian overrings of D is not proconstructible as soon as D is Noetherian and $\dim(D) \geq 2$: indeed, if it were, then $\text{NoethOver}(D) \cap \text{Zar}(D) = \Delta$ would be proconstructible, against the fact that Δ is not compact. However, this is also a consequence of a more general result. We need a topological lemma.

Lemma 7.5. *Let $Y \subseteq X$ be spectral spaces. Suppose that there is a subbasis \mathcal{B} of X such that, for every $B \in \mathcal{B}$, both B and $B \cap Y$ are compact. Then, Y is a proconstructible subset of X .*

Proof. The hypothesis on \mathcal{B} implies that the inclusion map $Y \hookrightarrow X$ is a spectral map; by [3, 1.9.5(vii)], it follows that Y is a proconstructible subset of X . \square

Proposition 7.6. *Let D be an integral domain with quotient field K , and let $D[\mathcal{F}_f]$ be the set of finitely generated D -algebras contained in K .*

- (a) $D[\mathcal{F}_f]$ is dense in $\text{Over}(D)$, with respect to the constructible topology.
- (b) Let X such that $D[\mathcal{F}_f] \subseteq X \subseteq \text{Over}(D)$. Then, X is spectral in the Zariski topology if and only if $X = \text{Over}(D)$.

Proof. (a) A basis of the constructible topology is given by the sets of type $U \cap (X \setminus V)$, as U and V ranges in the open and compact subsets of $\text{Over}(D)$. Such an U can be written as $B_1 \cup \dots \cup B_n$, where each $B_i = B(x_1^{(i)}, \dots, x_n^{(i)})$ is a basic open set of $\text{Over}(D)$; thus, we can suppose that $U = B(x_1, \dots, x_n)$. Suppose $\Omega := U \cap (X \setminus V)$ is nonempty; we claim that $A := D[x_1, \dots, x_n] \in \Omega \cap D[\mathcal{F}_f]$. Clearly $A \in D[\mathcal{F}_f]$ and $A \in U$; let $T \in \Omega$. Then, $T \in U$, and thus $A \subseteq T$; therefore, A is in the closure $\text{Cl}(T)$ of T , with respect to the Zariski topology. But $X \setminus V$ is closed, and thus $\text{Cl}(T) \subseteq X \setminus V$; i.e., $A \in X \setminus V$. Hence, $A \in \Omega \cap D[\mathcal{F}_f]$, which in particular is nonempty, and $D[\mathcal{F}_f]$ is dense.

(b) Suppose X is spectral. For every x_1, \dots, x_n , the set $X \cap B(x_1, \dots, x_n)$ has a minimum (i.e., $D[x_1, \dots, x_n]$), so it is compact. Since the family of all $B(x_1, \dots, x_n)$ is a basis, by Lemma 7.5 it follows that X is proconstructible. By the previous point, we must have $X = \text{Over}(D)$. \square

Corollary 7.7. *Let D be a Noetherian domain. The spaces*

- $\text{NoethOver}(D) := \{T \in \text{Over}(D) \mid T \text{ is Noetherian}\}$, and
- $\text{KrullOver}(D) := \{T \in \text{Over}(D) \mid T \text{ is a Krull domain}\}$

are spectral if and only if $\dim(D) = 1$.

Proof. If $\dim(D) = 1$, then the claim follows by Proposition 7.3.

If $\dim(D) \geq 2$, then $\text{NoethOver}(D)$ is not spectral by Proposition 7.6(b) and the Hilbert Basis Theorem; the case of $\text{KrullOver}(D)$ follows in the same way, since $\text{KrullOver}(D) \cap B(x_1, \dots, x_n)$ has always a minimum (i.e., the integral closure of $D[x_1, \dots, x_n]$). \square

More generally, consider a property \mathcal{P} of Noetherian domains such that every field and every discrete valuation ring satisfies \mathcal{P} ; for example, \mathcal{P} may be the property of being regular, Gorenstein or Cohen-Macaulay. Let $X_{\mathcal{P}}(D)$ be the set of overrings of D satisfying \mathcal{P} ; then, $X_{\mathcal{P}}(D) \cap \text{Zar}(D)$ is not compact, and thus $X_{\mathcal{P}}(D)$ is not proconstructible. On the other hand, if $X_{\mathcal{P}}(T)$ is compact for every overring of D that is finitely generated as a D -algebra, then by Lemma 7.5 it follows that $X_{\mathcal{P}}(D)$ cannot be a spectral space. Thus, the assignment $D \mapsto X_{\mathcal{P}}(D)$ cannot be “too good”: either some $X_{\mathcal{P}}(T)$ is not compact, or $X_{\mathcal{P}}(D)$ is not spectral.

Question. Let \mathcal{P} be the property of being regular, the property of being Gorenstein or the property of being Cohen-Macaulay. Is it possible to characterize for which Noetherian domains D there is a $T \in \text{Over}(D)$ such that $X_{\mathcal{P}}(T)$ is not compact and for which $X_{\mathcal{P}}(D)$ is not spectral?

8. ACKNOWLEDGMENTS

Section 6 was inspired by a talk given by Daniel McGregor at the congress “Recent Advances in Commutative ring and Module Theory” in Bressanone (June 13-17, 2017). I also thank the referee for his numerous suggestions, which improved the paper.

REFERENCES

- [1] M. F. Atiyah and I. G. Macdonald. *Introduction to commutative algebra*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
- [2] Nicolas Bourbaki. *Commutative algebra. Chapters 1–7*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1989. Translated from the French, Reprint of the 1972 edition.
- [3] Jean Dieudonné and Alexander Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. I. *Inst. Hautes Études Sci. Publ. Math.*, (20):259, 1964.
- [4] David E. Dobbs, Richard Fedder, and Marco Fontana. Abstract Riemann surfaces of integral domains and spectral spaces. *Ann. Mat. Pura Appl.* (4), 148:101–115, 1987.
- [5] David E. Dobbs and Marco Fontana. Kronecker function rings and abstract Riemann surfaces. *J. Algebra*, 99(1):263–274, 1986.
- [6] Alice Fabbri. Integral domains having a unique Kronecker function ring. *J. Pure Appl. Algebra*, 215(5):1069–1084, 2011.
- [7] Carmelo A. Finocchiaro. Spectral spaces and ultrafilters. *Comm. Algebra*, 42(4):1496–1508, 2014.

- [8] Carmelo A. Finocchiaro, Marco Fontana, and K. Alan Loper. The constructible topology on spaces of valuation domains. *Trans. Amer. Math. Soc.*, 365(12):6199–6216, 2013.
- [9] Carmelo A. Finocchiaro, Marco Fontana, and Dario Spirito. New distinguished classes of spectral spaces: a survey. In *Multiplicative Ideal Theory and Factorization Theory: Commutative and Non-Commutative Perspectives*. Springer Verlag, 2016.
- [10] Carmelo A. Finocchiaro, Marco Fontana, and Dario Spirito. Spectral spaces of semistar operations. *J. Pure Appl. Algebra*, 220(8):2897–2913, 2016.
- [11] Carmelo A. Finocchiaro and Dario Spirito. Some topological considerations on semistar operations. *J. Algebra*, 409:199–218, 2014.
- [12] Carmelo A. Finocchiaro and Francesca Tartarone. On a topological characterization of Prüfer v -multiplication domains among essential domains. *J. Commut. Algebra*, to appear.
- [13] Marco Fontana and K. Alan Loper. An historical overview of Kronecker function rings, Nagata rings, and related star and semistar operations. In *Multiplicative ideal theory in commutative algebra*, pages 169–187. Springer, New York, 2006.
- [14] Marco Fontana and K. Alan Loper. The patch topology and the ultrafilter topology on the prime spectrum of a commutative ring. *Comm. Algebra*, 36(8):2917–2922, 2008.
- [15] Robert Gilmer. *Multiplicative ideal theory*. Marcel Dekker Inc., New York, 1972. Pure and Applied Mathematics, No. 12.
- [16] Malcolm Griffin. Some results on v -multiplication rings. *Canad. J. Math.*, 19:710–722, 1967.
- [17] Klaas Pieter Hart, Jun-iti Nagata, and Jerry E. Vaughan, editors. *Encyclopedia of general topology*. Elsevier Science Publishers, B.V., Amsterdam, 2004.
- [18] Melvin Hochster. Prime ideal structure in commutative rings. *Trans. Amer. Math. Soc.*, 142:43–60, 1969.
- [19] Roland Huber and Manfred Knebusch. On valuation spectra. In *Recent advances in real algebraic geometry and quadratic forms (Berkeley, CA, 1990/1991; San Francisco, CA, 1991)*, volume 155 of *Contemp. Math.*, pages 167–206. Amer. Math. Soc., Providence, RI, 1994.
- [20] Craig Huneke and Irena Swanson. *Integral closure of ideals, rings, and modules*, volume 336 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2006.
- [21] B. G. Kang. Prüfer v -multiplication domains and the ring $R[X]_{N_v}$. *J. Algebra*, 123(1):151–170, 1989.
- [22] Irving Kaplansky. *Commutative rings*. The University of Chicago Press, Chicago, Ill.-London, revised edition, 1974.
- [23] Daniel J. McGregor. Personal communication.
- [24] Masayoshi Nagata. *Local rings*. Interscience Tracts in Pure and Applied Mathematics, No. 13. Interscience Publishers a division of John Wiley & Sons New York-London, 1962.
- [25] Akira Okabe and Ryūki Matsuda. Semistar-operations on integral domains. *Math. J. Toyama Univ.*, 17:1–21, 1994.
- [26] Bruce Olberding. Noetherian spaces of integrally closed rings with an application to intersections of valuation rings. *Comm. Algebra*, 38(9):3318–3332, 2010.
- [27] Bruce Olberding. Affine schemes and topological closures in the Zariski-Riemann space of valuation rings. *J. Pure Appl. Algebra*, 219(5):1720–1741, 2015.

- [28] Bruce Olberding. Topological aspects of irredundant intersections of ideals and valuation rings. In *Multiplicative Ideal Theory and Factorization Theory: Commutative and Non-Commutative Perspectives*. Springer Verlag, 2016.
- [29] Jean-Pierre Olivier. Anneaux absolument plats universels et épimorphismes à buts réduits. In *Séminaire Samuel. Algèbre commutative, 2: Les épimorphismes d'anneaux. 1967–1968*.
- [30] Jean-Pierre Olivier. Anneaux absolument plats universels et épimorphismes d'anneaux. *C. R. Acad. Sci. Paris Sér. A-B*, 266:A317–A318, 1968.
- [31] N. Schwartz. Compactification of varieties. *Ark. Mat.*, 28(2):333–370, 1990.
- [32] Oscar Zariski. The reduction of the singularities of an algebraic surface. *Ann. of Math. (2)*, 40:639–689, 1939.
- [33] Oscar Zariski. The compactness of the Riemann manifold of an abstract field of algebraic functions. *Bull. Amer. Math. Soc.*, 50:683–691, 1944.
- [34] Oscar Zariski and Pierre Samuel. *Commutative algebra. Vol. II*. Springer-Verlag, New York, 1975. Reprint of the 1960 edition, Graduate Texts in Mathematics, Vol. 29.

E-mail address: spirito@mat.uniroma3.it

DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DEGLI STUDI “ROMA TRE”, ROMA, ITALY