

LE MATEMATICHE

Vol. LXXV (2020) – Issue I, pp. 57–66

doi: 10.4418/2020.75.1.3

## BREAKING THROUGH BORDERS WITH $\sigma$ -HARMONIC MAPPINGS

GIOVANNI ALESSANDRINI - VINCENZO NESI

We consider mappings  $U = (u^1, u^2)$ , whose components solve an arbitrary elliptic equation in divergence form in dimension two, and whose respective Dirichlet data  $\varphi^1, \varphi^2$  constitute the parametrization of a simple closed curve  $\gamma$ . We prove that, if the interior of the curve  $\gamma$  is not convex, then we can find a parametrization  $\Phi = (\varphi^1, \varphi^2)$  such that the mapping  $U$  is not invertible.

*Dedicato a chi sconfinava frontiere geografiche o ideologiche,  
a chi travalica stereotipi e va oltre i pregiudizi.*

### 1. Introduction

Let  $B = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < 1\}$  denote the unit disk. We denote by  $\sigma = \sigma(x)$ ,  $x \in B$ , a possibly non-symmetric matrix having measurable entries and satisfying the ellipticity conditions

$$\begin{aligned} \sigma(x)\xi \cdot \xi &\geq K^{-1}|\xi|^2, \text{ for every } \xi \in \mathbb{R}^2, x \in B, \\ \sigma^{-1}(x)\xi \cdot \xi &\geq K^{-1}|\xi|^2, \text{ for every } \xi \in \mathbb{R}^2, x \in B, \end{aligned} \tag{1.1}$$

---

Received on June 28, 2019

*AMS 2010 Subject Classification:* 30C62, 35J55

*Keywords:* Elliptic equations, Beltrami operators, quasiconformal mappings.

G. A. has been supported by PRIN 2017 “Direct and inverse problems for partial differential equations: theoretical aspects and applications”.

V. N. has been supported by Sapienza - Ateneo 2018 Project “Stationary and Evolutive Problems in Mathematical Physics and Materials Science”.

for a given constant  $K \geq 1$ .

Given a homeomorphism  $\Phi = (\varphi^1, \varphi^2)$  from the unit circle  $\partial B$  onto a simple closed curve  $\gamma \subset \mathbb{R}^2$ , we denote by  $D$  the bounded domain such that  $\partial D = \gamma$ .

Consider the mapping  $U = (u^1, u^2) \in W_{loc}^{1,2}(B; \mathbb{R}^2) \cap C(\bar{B}; \mathbb{R}^2)$  whose components are the solutions to the following Dirichlet problems

$$\begin{cases} \operatorname{div}(\sigma \nabla u^i) = 0, & \text{in } B, \\ u^i = \varphi^i, & \text{on } \partial B, i = 1, 2. \end{cases} \quad (1.2)$$

We call such a  $U$  a  $\sigma$ -harmonic mapping.

In the last two decades, it has been investigated, by the present authors and others, under which conditions can one assure that  $U$  is an invertible mapping between  $B$  and  $D$ .

The classical starting point for this issue is the celebrated Radò–Kneser–Choquet Theorem [10, 11, 13, 16] which asserts that assuming  $\sigma = I$ , the identity matrix, (that is:  $u^1, u^2$  are harmonic) if  $D$  is convex then  $U$  is a homeomorphism. Generalizations to equations with variable coefficients have been obtained in [2, 7] and to certain nonlinear systems in [6, 8, 14]. Counterexamples [3, 10] show that if  $D$  is not convex then the invertibility of  $U$  may fail. In fact Choquet [10] proved that, whenever  $D$  is not convex, there exists a homeomorphism  $\Phi : \partial B \rightarrow \gamma$  such that the corresponding harmonic ( $\sigma = I$ ) mapping  $U$  is not invertible. The proof is crucially based on the classical mean value property of harmonic functions. Also the counterexample in [3] is limited to the purely harmonic case.

In [3, 5] the present authors investigated which additional conditions are needed for invertibility in the case of a possibly non-convex target  $D$ . Let us recall the main result in that direction.

**Theorem 1.1.** *Let  $\Phi$  and  $U$  be as above stated. Assume that the entries of  $\sigma$  satisfy  $\sigma_{ij} \in C^\alpha(\bar{B})$  for some  $\alpha \in (0, 1)$  and for every  $i, j = 1, 2$ . Assume also that  $U \in C^1(\bar{B}; \mathbb{R}^2)$ . The mapping  $U$  is a diffeomorphism of  $\bar{B}$  onto  $\bar{D}$  if and only if*

$$\det DU > 0 \quad \text{everywhere on } \partial B. \quad (1.3)$$

The object of the present note is to extend the construction by Choquet to  $\sigma$ -harmonic mappings with arbitrary coefficient matrix  $\sigma$ . The main result will be as follows.

**Theorem 1.2.** *Given a homeomorphism  $\Psi : \partial B \rightarrow \gamma \subset \mathbb{R}^2$ , let  $D$  be the bounded domain such that  $\partial D = \gamma$ . Assume that  $D$  is not convex. For every  $\sigma = \sigma(x)$ , satisfying (1.1), there exists a  $C^\infty$  diffeomorphism  $\Xi : \partial B \rightarrow \partial B$  such that, posing  $\Phi = \Psi \circ \Xi$ , the  $\sigma$ -harmonic mapping  $U$  solving (1.2) is not invertible.*

Note that the parametrization  $\Phi$  of the curve  $\gamma$  is as much smooth as the original one  $\Psi$ . In particular, if  $\Psi$  is  $C^{1,\alpha}$  so is  $\Phi$ . Hence under the hypothesis of Hölder continuity of  $\sigma$ , it turns out that  $U$  is  $C^{1,\alpha}$  up to the boundary. As a consequence, we obtain that the hypothesis (1.3) in Theorem 1.1 is indeed non-trivial.

Let us illustrate what should be the features of a candidate counterexample: first we recall that Kneser [13] noticed that, in the purely harmonic case, if it is a-priori known that  $U(B) \subset D$ , then indeed  $U$  is invertible, whether or not  $D$  is convex. The observation by Kneser, is merely of topological nature, see also Duren [11, p. 31], and hence it actually extends to the  $\sigma$ -harmonic case, for any  $\sigma$ . That is, in order to violate invertibility in general, we must provide a mapping  $U$  whose image exceeds  $D$ .

Viceversa, again by elementary topological arguments, if  $U$  is one-to-one on all of  $\bar{B}$ , then it is an open mapping, hence a homeomorphism. Therefore it maps  $\partial B$  onto  $\gamma$  and  $B$  onto  $D$ . In other terms, if  $U$  maps some point of  $B$  outside of  $\bar{D}$ , then it cannot be one-to-one.

In conclusion, in order to construct an example of a non-invertible  $\sigma$ -harmonic mapping  $U$ , whose boundary data  $\Phi : \partial B \rightarrow \gamma$  is invertible, it is necessary and sufficient that  $U$  *trespasses* the boundary  $\gamma$ , or in other words, that  $U$  maps some interior point of  $B$  outside of  $\bar{D}$ . This will be indeed the crux of our argument below.

## 2. $\sigma$ -harmonic measure

Given  $\sigma$  as in (1.1), and  $\varphi \in C(\partial B)$ , consider the scalar Dirichlet problem

$$\begin{cases} \operatorname{div}(\sigma \nabla u) = 0, & \text{in } B, \\ u = \varphi, & \text{on } \partial B, \end{cases} \quad (2.1)$$

the, by now, classical theory of divergence structure elliptic equation tells us that there exists a unique weak solution  $u \in W_{loc}^{1,2}(B) \cap C(\bar{B})$ , see for instance [12, Theorem 8.30]. In particular the functional

$$C(\partial B) \ni \varphi \rightarrow u(0) \in \mathbb{R}$$

is bounded and linear. Hence there exists a Radon measure  $\omega_\sigma$  on  $\partial B$  such that

$$u(0) = \int_{\partial B} \varphi d\omega_\sigma .$$

We call  $\omega_\sigma$  the  $\sigma$ -harmonic measure. Note that, being  $u \equiv 1$  the solution to (2.1) when  $\varphi \equiv 1$ , we trivially have  $\omega_\sigma(\partial B) = 1$ .

From examples due to Modica and Mortola and to Caffarelli, Fabes and Kenig [9, 15], it is known that the  $\sigma$ -harmonic measure may not be absolutely continuous with the arclength measure. Still, some kind of continuity holds. For every  $P \in \partial B$  and for every  $r > 0$  let us denote

$$\Delta_r(P) = \partial B \cap B_r(P).$$

We prove the following.

**Lemma 2.1.** *For every  $P \in \partial B$  we have*

$$\lim_{r \rightarrow 0^+} \omega_\sigma(\Delta_r(P)) = 0. \quad (2.2)$$

*Proof.* Let  $h_r$  be the Perron solution to the Dirichlet problem

$$\begin{cases} \operatorname{div}(\sigma \nabla h_r) = 0, & \text{in } B, \\ h_r = \chi_{\Delta_r(P)}, & \text{on } \partial B, \end{cases} \quad (2.3)$$

our aim is to prove that

$$\lim_{r \rightarrow 0^+} h_r(0) = 0.$$

We start considering the selfadjoint case, that is when  $\sigma = \sigma^T$ . We extend  $\sigma = I$  outside of  $B$ .

Let  $D_r$  be the annulus  $B_2(P) \setminus \overline{B_r(P)}$ , and let  $c_r$  be the solution of the following Dirichlet problem

$$\begin{cases} \operatorname{div}(\sigma \nabla c_r) = 0, & \text{in } D_r, \\ c_r = 0, & \text{on } \partial B_2(P), \\ c_r = 1, & \text{on } \partial B_r(P). \end{cases} \quad (2.4)$$

By the maximum principle, we have

$$0 \leq h_r \leq c_r, \text{ on } B \setminus \overline{B_r(P)}.$$

Because of selfadjointness, we have

$$\begin{aligned} & \int_{D_r} \sigma \nabla c_r \cdot \nabla c_r = \\ & = \min \left\{ \int_{D_r} \sigma \nabla v \cdot \nabla v \mid v \in W^{1,2}(D_r), v = 0 \text{ on } \partial B_2(P), v = 1 \text{ on } \partial B_r(P) \right\}. \end{aligned}$$

Choosing

$$v(x) = \frac{\log \frac{2}{|x-P|}}{\log \frac{2}{r}},$$

we compute

$$\begin{aligned} \int_{D_r} \sigma \nabla c_r \cdot \nabla c_r &\leq K \int_{D_r} |\nabla v|^2 = \\ &= 2\pi K \frac{1}{\log \frac{2}{r}} \rightarrow 0 \text{ as } r \rightarrow 0. \end{aligned}$$

Next we invoke a more or less standard form of Poincaré inequality, the emphasis being on the uniformity of the inequality with respect to the small radius  $r$ . A proof is outlined in Section 4 below.

**Lemma 2.2.** *For every  $w \in W^{1,2}(D_r)$ , having zero trace on  $\partial B_2(P)$ , we have*

$$\int_{D_r} w^2 \leq 16 \int_{D_r} |\nabla w|^2.$$

Consequently we obtain  $\|c_r\|_{W^{1,2}(D_r)} \rightarrow 0$  as  $r \rightarrow 0$ , and by an interior boundedness estimate [12, Theorem 8.17],  $c_r(0) \rightarrow 0$ , and the thesis follows.

Now we remove the symmetry assumption on  $\sigma$ .

It is well-known that there exists  $k_r \in W^{1,2}(B)$ , called the *stream function* of  $h_r$  such that

$$\nabla k_r = J \sigma \nabla h_r, \quad (2.5)$$

where the matrix  $J$  denotes the counterclockwise  $90^\circ$  rotation

$$J = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad (2.6)$$

see, for instance, [1]. Denoting

$$f = h_r + ik_r, \quad (2.7)$$

it is well-known that  $f$  solves the Beltrami type equation

$$f_{\bar{z}} = \mu f_z + \nu \overline{f_z} \quad \text{in } B, \quad (2.8)$$

where, the so called complex dilatations  $\mu, \nu$  are given by

$$\mu = \frac{\sigma_{22} - \sigma_{11} - i(\sigma_{12} + \sigma_{21})}{1 + \text{Tr} \sigma + \det \sigma}, \quad \nu = \frac{1 - \det \sigma + i(\sigma_{12} - \sigma_{21})}{1 + \text{Tr} \sigma + \det \sigma}, \quad (2.9)$$

and satisfy the following ellipticity condition

$$|\mu| + |\nu| \leq k < 1, \quad (2.10)$$

where the constant  $k$  only depends on  $K$ , see [4, Proposition 1.8] and the notation  $\text{Tr} A$  is used for the trace of a square matrix  $A$ . We can also write

$$f_{\bar{z}} = \tilde{\mu} f_z \text{ in } B,$$

where  $\tilde{\mu}$  is defined almost everywhere by

$$\tilde{\mu} = \mu + \frac{\overline{f_z}}{f_z} \nu,$$

and consequently we obtain

$$\operatorname{div}(\tilde{\sigma} \nabla h_r) = 0, \text{ in } B$$

where  $\tilde{\sigma}$  is given by

$$\tilde{\sigma} = \begin{bmatrix} \frac{|1 - \tilde{\mu}|^2}{1 - |\tilde{\mu}|^2} & -\frac{2\Im m(\tilde{\mu})}{1 - |\tilde{\mu}|^2} \\ -\frac{2\Im m(\tilde{\mu})}{1 - |\tilde{\mu}|^2} & \frac{|1 + \tilde{\mu}|^2}{1 - |\tilde{\mu}|^2} \end{bmatrix},$$

which satisfies uniform ellipticity conditions of the form (1.1) with a new constant  $\tilde{K}$  only dependent on  $K$ , see, for instance, [4], but in addition is symmetric. Hence we may proceed as before, just replacing  $\sigma$  with  $\tilde{\sigma}$  in (2.3) and obtain again

$$\lim_{r \rightarrow 0^+} h_r(0) = 0. \quad \square$$

The above Lemma can be seen as a continuity result for the cumulative distribution function associated to  $\omega_\sigma$ .

Given two points  $P, Q \in \partial B$  we denote by  $\widehat{PQ}$  the arc of the unit circle  $\partial B$  which connects  $P$  to  $Q$ , moving in the counterclockwise direction. The above Lemma, along with Harnack's inequality, implies the following straightforward consequence.

**Corollary 2.3.** *For every  $P \in \partial B$ , the function*

$$\partial B \ni Q \rightarrow \omega_\sigma(\widehat{PQ}) \in [0, 1]$$

*is a strictly increasing, onto and continuous function, as  $Q$  performs a full counterclockwise rotation on  $\partial B$  starting from  $P$  and ending on  $P$  itself. Moreover, for every  $P \in \partial B$ , there exists exactly one point  $Q \in \partial B$  such that*

$$\omega_\sigma(\widehat{PQ}) = \omega_\sigma(\widehat{QP}) = \frac{1}{2}.$$

### 3. Assembling a parametrization

Let us consider a given homeomorphism  $\Psi : \partial B \rightarrow \gamma \subset \mathbb{R}^2$ , let us fix two distinct points  $a, b \in \gamma$ . For any  $\varepsilon > 0$  let  $\alpha, \beta$  two disjoint simple open arcs in  $\gamma$  such that

$$a \in \alpha \subset B_\varepsilon(a), b \in \beta \subset B_\varepsilon(b).$$

Denote

$$A = \Psi^{-1}(a), B = \Psi^{-1}(b),$$

and

$$\widehat{A^-A^+} = \Psi^{-1}(\alpha), \widehat{B^-B^+} = \Psi^{-1}(\beta).$$

Having fixed points  $P, Q \in \partial B$  such that

$$\omega_\sigma(\widehat{PQ}) = \omega_\sigma(\widehat{QP}) = \frac{1}{2}$$

for any  $r, 0 < r < 1$  we select a  $C^\infty$  diffeomorphism  $\Xi_r : \partial B \rightarrow \partial B$  such that

$$\Xi_r(\Delta_r(P)) = \widehat{A^+B^-}, \Xi_r(\Delta_r(Q)) = \widehat{B^+A^-}.$$

In other words, setting  $\widehat{P^-P^+} = \Delta_r(P)$ ,  $\widehat{Q^-Q^+} = \Delta_r(Q)$ , we need to construct a diffeomorphism  $\Xi_r$  which maps the points  $P^-, P^+, Q^-, Q^+$  to the points  $A^+, B^-, B^+, A^-$  in their respective order. More generally, we can prove the following Lemma, whose proof is deferred to the next Section 4.

**Lemma 3.1.** *Let  $N \geq 2$  and let  $P_1, \dots, P_N$  be distinct, cyclically ordered points on  $\partial B$  and let  $Q_1, \dots, Q_N$  be another  $N$ -tuple of distinct, cyclically ordered points on  $\partial B$ . There exists a  $C^\infty$  diffeomorphism  $\Xi : \partial B \rightarrow \partial B$  such that  $\Xi(P_n) = Q_n$  for every  $n = 1, \dots, N$ .*

*Proof of Theorem 1.2.* We let  $\Phi_r = \Psi \circ \Xi_r$  and consider  $U = U_r$  as the solution to (1.2) when  $\Phi = \Phi_r$ . If  $D$  is not convex, we may find two points  $a, b \in \gamma$  such that the open segment with endpoints  $a, b$  lies outside  $\bar{D}$ . In particular

$$\frac{1}{2}(a+b) \notin \bar{D}.$$

We have

$$U_r(0) = \int_{\partial B} \Phi_r d\omega_\sigma$$

and we may split  $\partial B$  into the four arcs  $\widehat{P^-P^+}, \widehat{P^+Q^-}, \widehat{Q^-Q^+}, \widehat{Q^+P^-}$ . Let  $M > 0$  be such that  $\gamma \subset B_M(0)$ , then we evaluate

$$\left| \int_{\widehat{P^-P^+}} \Phi_r d\omega_\sigma \right| \leq M \omega_\sigma(\Delta_r(P)) \rightarrow 0$$

as  $r \rightarrow 0$  and, analogously,

$$\left| \int_{\widehat{Q^-Q^+}} \Phi_r d\omega_\sigma \right| \leq M\omega_\sigma(\Delta_r(Q)) \rightarrow 0.$$

Conversely,  $\Phi_r(\widehat{P^+Q^-}) \subset \beta \subset B_\varepsilon(b)$  and  $\Phi_r(\widehat{Q^+P^-}) \subset \alpha \subset B_\varepsilon(a)$ , that is

$$|\Phi_r - b| < \varepsilon \text{ on } \widehat{P^+Q^-}, \quad |\Phi_r - a| < \varepsilon \text{ on } \widehat{Q^+P^-}.$$

Note also that

$$\lim_{r \rightarrow 0^+} \omega_\sigma(\widehat{P^+Q^-}) = \lim_{r \rightarrow 0^+} \omega_\sigma(\widehat{Q^+P^-}) = \frac{1}{2}.$$

Hence we may find  $r > 0$  small enough and a constant  $C > 0$  such that

$$|U_r(0) - \frac{1}{2}(a+b)| \leq C\varepsilon$$

and, in conclusion, with  $r, \varepsilon$  small enough,  $U = U_r$  is such that

$$U(0) \notin \overline{D}. \quad \square$$

#### 4. Auxiliary proofs

*Proof of Lemma 2.2.* As is customary in this context, it suffices to consider  $w \in C^1(\overline{D_r})$ ,  $w(P + 2e^{i\vartheta}) = 0$  for all  $\vartheta$ . Hence, for every  $\rho \in (r, 2)$  we have

$$w^2(P + \rho e^{i\vartheta}) = - \int_\rho^2 \frac{\partial}{\partial s} w^2(P + se^{i\vartheta}) ds,$$

hence

$$w^2(P + \rho e^{i\vartheta}) \leq 2 \int_\rho^2 |w| |\nabla w|(P + se^{i\vartheta}) ds.$$

Consequently

$$\int_{D_r} w^2 \leq 2 \int_0^{2\pi} d\vartheta \int_r^2 \rho d\rho \int_\rho^2 |w| |\nabla w|(P + se^{i\vartheta}) ds,$$

and, using the inequalities  $0 < r \leq \rho \leq s$ ,

$$\begin{aligned} \int_{D_r} w^2 &\leq 2 \int_0^{2\pi} d\vartheta \int_r^2 d\rho \int_\rho^2 |w| |\nabla w|(P + se^{i\vartheta}) s ds \leq \\ &\leq 2 \int_0^{2\pi} d\vartheta \int_0^2 d\rho \int_r^2 |w| |\nabla w|(P + se^{i\vartheta}) s ds, \end{aligned}$$

that is

$$\int_{D_r} w^2 \leq 4 \int_{D_r} |w| |\nabla w|,$$

and by Schwarz inequality the thesis follows.  $\square$



*Proof of Lemma 3.1.* Up to rotations, we may assume  $P_n = e^{i\vartheta_n}, Q_n = e^{i\varphi_n}, n = 1, \dots, N$  where

$$0 = \vartheta_1 < \dots < \vartheta_N < 2\pi, 0 = \varphi_1 < \dots < \varphi_N < 2\pi.$$

We may construct a continuous, strictly increasing, piecewise linear function  $f$  mapping the interval  $[0, 2\pi]$  onto itself, such that

$$f(\vartheta_n) = \varphi_n \text{ for every } n = 1, \dots, N,$$

we may consider to extend  $f$  to  $\mathbb{R}$  in such a way that  $f(\vartheta) - \vartheta$  is  $2\pi$ -periodic. We may also require that its corner points  $\xi_1, \dots, \xi_J \in [0, 2\pi]$  are distinct from the points

$$0 = \vartheta_1, \dots, \vartheta_N, \vartheta_{N+1} = 2\pi.$$

Let  $\delta = \min \{|\vartheta_n - \xi_j| \mid n = 1, \dots, N + 1, j = 1, \dots, J\}$ . Let  $\chi_\varepsilon$  be a family of  $C^\infty$ , mollifying kernels, supported in  $[-\varepsilon, \varepsilon]$ , even symmetric with respect to 0. Fixing  $\varepsilon < \delta$  and denoting

$$g = \chi_\varepsilon * f,$$

we compute  $g(\vartheta_n) = f(\vartheta_n)$  for all  $n$ , we obtain that  $g$  is  $C^\infty$  with positive derivative everywhere and we conclude that

$$\Xi(e^{i\vartheta}) = e^{ig(\vartheta)}$$

fulfils the thesis. □

### REFERENCES

- [1] Alessandrini, G. & Magnanini, R.: Elliptic equations in divergence form, geometric critical points of solutions, and Stekloff eigenfunctions. *SIAM J. Math. Anal.*, **25** 5, 1259–1268 (1994).
- [2] Alessandrini, G. & Nesi, V.: Univalent  $\sigma$ -harmonic mappings. *Arch. Ration. Mech. Anal.*, **158** 2, 155–171 (2001).
- [3] Alessandrini, G. & Nesi, V.: Invertible harmonic mappings, beyond Kneser. *Ann. Scuola Norm. Sup. Pisa, Cl. Sci.* **V**, **8** 5, 451–468 (2009). Errata Corrige. *Ann. Scuola Norm. Sup. Pisa, Cl. Sci.* **V**, **17** 2, 815–818 (2017).
- [4] Alessandrini, G. & Nesi, V.: Beltrami operators, non-symmetric elliptic equations and quantitative Jacobian bounds. *Ann. Acad. Sci. Fenn. Math.*, **34** 1, 47–67 (2009).

- [5] Alessandrini, G. & Nesi, V.: Globally diffeomorphic  $\sigma$ -harmonic mappings, arXiv:1906.00902, submitted.
- [6] Alessandrini, G. & Sigalotti, M.: Geometric properties of solutions to the anisotropic p-Laplace equation in dimension two, *Ann. Acad. Sci. Fenn. Math.* **21**, 249–266 (2001).
- [7] Bauman, P., Marini, A. & Nesi, V.: Univalent solutions of an elliptic system of partial differential equations arising in homogenization. *Indiana Univ. Math. J.*, **50** 2, 747–757 (2001).
- [8] Bauman, P. & Phillips, D.: Univalent minimizers of polyconvex functionals in two dimensions. *Arch. Rational Mech. Anal.*, **126** 2, 161–181 (1994).
- [9] Caffarelli, L., Fabes, E. & Kenig, C.: Completely Singular Elliptic-Harmonic Measures. *Indiana University Mathematics Journal*, 30(6), 917-924, (1981).
- [10] Choquet, G.: Sur un type de transformation analytique généralisant la représentation conforme et définie au moyen de fonctions harmoniques, *Bull. Sci. Math. (2)* **69** 156–165, (1945).
- [11] Duren, P.: Harmonic mappings in the plane. Cambridge Tracts in Mathematics, vol. 156, Cambridge University Press, Cambridge (2004).
- [12] Gilbarg, D. and Trudinger, N. S.: Elliptic Partial Differential Equations of Second Order. Grundlehren der Mathematischen Wissenschaften, vol 224, 2nd edn, Springer, Berlin (1983).
- [13] Kneser, H.: Lösung der Aufgabe 41. *Jber. Deutsch. Math.-Verein.*, **35**, 123–124 (1926).
- [14] Iwaniec, T., Koski, A. & Onninen, J.: Isotropic p-harmonic systems in 2D Jacobian estimates and univalent solutions. *Rev. Mat. Iberoam.* **32**, 57–77 (2016).
- [15] Modica, L. & Mortola, S.: Construction of a singular elliptic-harmonic measure, *Manuscripta Math.* **33** 1, 81–98 (1980) .
- [16] Radó, T.: Aufgabe 41, *Jber. Deutsch. Math.-Verein.* **35**, 49 (1926).

*GIOVANNI ALESSANDRINI*

*formerly Dipartimento di Matematica e Geoscienze*

*Università di Trieste*

*Via Valerio 12/b, 34100 Trieste, Italia*

*e-mail: 55gioale@gmail.com*

*VINCENZO NESI*

*Dipartimento di Matematica “G. Castelnuovo”*

*Sapienza, Università di Roma*

*Piazzale A. Moro 2, 00185 Roma, Italia*

*e-mail: nesi@mat.uniroma1.it*