

Geological reactive systems from the mantle to the abyssal sub-seafloor: Preface

Godard Marguerite^{1,*}, Fumagalli Patrizia², Jamtveit Bjørn³, Ménez Bénédicte⁴

¹ Géosciences Montpellier, CNRS, Université de Montpellier, 34095 Montpellier, FRANCE

² Dipartimento di Scienze della Terra "Ardito Desio", Università degli Studi di Milano, ITALY

³ The Njord Centre, Department of Geosciences, University of Oslo, 0316 Oslo, NORWAY

⁴ Institut de Physique du Globe de Paris, Sorbonne Paris Cité, Université Paris Diderot, CNRS, 75005 Paris, FRANCE

Contact :

Godard Marguerite
Géosciences Montpellier
Université de Montpellier
Campus Triolet cc60
Place Eugène Bataillon
34095 Montpellier FRANCE
Email : Marguerite.Godard@umontpellier.fr

1 **Geological reactive systems from the mantle to the abyssal sub-seafloor: Preface**

2 Godard Marguerite, Fumagalli Patrizia, Jamtveit Bjørn, Ménez Bénédicte

3

4 The formation and alteration of the oceanic lithosphere represent one of the main processes for
5 energy and chemical exchanges between the deep Earth and its outer envelopes. However, the steep
6 thermal gradients characterizing these environments, especially at the main thermal and lithological
7 interfaces along mid-ocean accretion zones (Figure 1), mean that the physical and chemical
8 mechanisms controlling these exchanges remain poorly understood. Yet, these interfaces are the
9 main transitions for the physical and rheological properties of rocks, such as permeability and
10 viscosity, that control melt focussing and transport from the partially molten mantle to the surface,
11 as well as deformation mechanisms and the influx of seawater into the cooling oceanic lithosphere.
12 These processes also give rise to hydrothermal systems that produce economically valuable ore-
13 deposits and play a major role in the global chemical budget. Some hydrothermal reactions produce
14 hydrogen and abiotic hydrocarbon, hence these extreme environments sustain life and they are
15 potentially implicated in its origin. Finally, these processes determine the architecture and
16 composition of the lithosphere plunging into the deep Earth along subduction zones, and contribute
17 to a broad range of mechanisms driving arc magmatism and localization of earthquakes in these
18 regions.

19 Traditionally the formation of oceanic lithosphere has been envisioned as a suite of mantle and
20 crustal magmatic processes, followed by high- to low-temperature hydrothermal processes, which in
21 turn supported the development of diverse ecosystems; in this scheme, each step was considered as
22 occurring independently and in different domains, whilst the interactions and feedbacks controlling
23 mass and energy transfers at their boundaries had been mostly overlooked. However, over the
24 recent years, the role of the physical and (bio-)chemical processes occurring at the interfaces
25 bounding the Earth's envelopes has been progressively recognized. These boundary layers
26 characterize the architecture of Earth's oceanic lithosphere and they are determined by two
27 competitive processes at mid-ocean ridges: conduction and advection of heat from depth, which is
28 dominantly controlled by mantle upwelling (spreading rate) and melt transport from the mantle to
29 the surface, and cooling by conduction and hydrothermal circulation. These boundary layers
30 correspond to isotherms marking transitions in rocks properties and therefore, in turn, they depend
31 on the composition of the lithosphere and are expected to vary greatly from fast-spreading layered
32 oceanic lithosphere (Penrose Model) to slow-spreading heterogeneous oceanic lithosphere (Figure
33 1). Understanding such complex highly reactive geological systems requires a shift of approach and
34 the development of new scientific tools to comprehend and integrate the role of transport of
35 magmatic melts and hydrothermal fluids through the oceanic lithosphere, and their feed-back on its

36 physical and (bio-)chemical properties. This volume brings together a series of articles addressing
37 these challenges with natural observations and laboratory experiments.

38

39 *From mantle lithosphere to magmatic crust*

40 Basch et al. (2018), Ferrando et al. (2018) and Borghini et al. (2018) investigate the nature of the
41 transition from mantle to gabbroic lower crust with the aim to characterize and identify the
42 magmatic processes controlling this transition. Basch et al. (2018) document a transition from mantle
43 peridotites to olivine-rich troctolites, the most primitive end-member of the lower gabbroic crust,
44 preserved in the Mt. Maggiore Ophiolite (Corsica, France). Ferrando et al. (2018) focus on the study
45 of olivine-rich troctolites interlayered within a drilled gabbroic section of heterogeneous oceanic
46 lithosphere (Integrated Ocean Drilling Program (IODP) Expeditions 304/305 Site U1309 Atlantis
47 Massif 30°N, Mid-Atlantic Ridge). Basch et al. (2018) and Ferrando et al. (2018) combine petro-
48 structural and geochemical analyses and numerical modelling of major and trace element variations
49 to determine the respective roles of fractional crystallization and reactive porous melt flow during
50 the incipient stages of the formation of gabbroic oceanic crust. This approach provides criteria to
51 identify and quantify melt/rock reactions and to estimate the contribution of mantle rocks to the
52 formation of the lower crust. These studies give evidences for a likely progressive formation of the
53 lower oceanic crust at the expense of the shallow mantle, and shed light on the control that such
54 processes can exert on melt evolution in the oceanic crust. To better constrain the chemical and
55 physical parameters driving these processes, Borghini et al. (2018) investigate the origin of olivine-
56 rich troctolites and studied melt-olivine reaction using specifically designed experiments. They
57 observe textural development comparable with disequilibrium features observed in natural olivine-
58 rich troctolites (e.g., Basch et al. (2018), Ferrando et al. (2018)) and demonstrate the control of
59 starting melt composition and melt-olivine ratio on modal composition and mineral chemistry of
60 olivine-rich troctolites.

61

62 *High temperature hydrothermalism and cooling: Impact on architecture and composition of newly
63 formed crust*

64 Koepke et al. (2018), Grant and Harlov (2018), Currin et al. (2018a, b) and Zihlmann et al. (2018)
65 investigate the role of fluids and of hydrothermalism on differentiation processes during cooling of
66 the gabbroic mafic lower crust and their impact on the architecture and composition of the newly
67 formed crust. Koepke et al. (2018) present the first phase-equilibria study of the late stages of MORB
68 differentiation and explore the role of water activity in these systems. On this basis, they propose a
69 two-step differentiation model for the formation of oxide gabbros in slow spread magmatic crust and
70 at the transition from lower to upper layered crust as well as for the formation of highly evolved

71 lavas at fast- and intermediate spreading mid-ocean ridges. This model emphasizes the prevalence of
72 oxygen fugacity on oxide differentiation suites including late formation of minerals such as apatite
73 and amphibole during cooling down to temperatures of 800-900°C. Grant and Harlov (2018) and
74 Currin et al. (2018a) use experimental approaches to study the behaviour of these systems in the
75 presence of NaCl-brines in order to understand fluid-rock interactions associated with high
76 temperature hydrothermalism (900° to 500°C) not only in oceanic gabbros but also in the deep
77 continental mafic crust. Grant and Harlov (2018) show a correlation between the reactivity of olivine-
78 plagioclase assemblages, water activity and fluid NaCl concentrations. They identify the chemical and
79 temperature conditions the most favourable for the formation of Cl-rich amphiboles in these
80 systems. Currin et al. (2018a) further investigate the processes leading to the formation of
81 amphiboles, and in particular of Cl-rich amphiboles, at the pressure and temperature conditions
82 expected during hydrothermal reactions between seawater-derived fluids and the gabbroic oceanic
83 lower crust. They show that the composition of amphibole is highly variable in hydrothermal systems
84 and that their Cl content is affected not only by the composition of hydrothermal fluids but also by
85 the extent of fluid/rock interactions. Zihlmann et al. (2018) and Currin et al. (2018b) provide new
86 constraints on the role of deep hydrothermal systems at mid-oceanic ridges through detailed field,
87 petrologic and geochemical studies of lower gabbroic sections of layered oceanic crust preserved by
88 the Oman Ophiolite. Zihlmann et al. (2018) identify fault zones as the main flow paths for high
89 temperature hydrothermalism at mid-ocean ridges. They establish that such focussed flow zones
90 contribute to the fast cooling of the gabbroic lower oceanic crust and to the global hydrothermal
91 geochemical fluxes. Currin et al. (2018b) focus on the formation of Cl-rich amphiboles and show that
92 fluid-rock interactions leading to their formation occur in rock-dominated environment.

93

94 *Serpentinization, sulphides, carbon and geo-resources*

95 The linkages between magmatic, hydrothermal and (bio-)geochemical processes becomes even
96 more prominent when investigating the mechanisms building the shallow oceanic lithosphere and
97 their impact on global geochemical cycles, deep sea natural resources and the development of life as
98 illustrated by Früh-Green et al. (2018). These authors present an overview of the results of the first
99 oceanic drilling expedition dedicated to investigate the interplay between magmatism,
100 serpentinization processes and microbial activity in the shallow subsurface (IODP Expedition 357,
101 Atlantis Massif). This expedition provided a unique sampling of the suite of altered and deformed
102 ultramafic and mafic rocks forming one of the major detachment fault zone along the Mid-Atlantic
103 Ridge as well as first biogeochemical and microbiological characterization of the shallow ultramafic
104 subseafloor.

105 Rouméjon et al. (2018), Escario et al. (2018) and Pastore et al. (2018) investigate the
106 thermodynamic and hydrodynamic conditions and the localization of serpentinization reactions, the
107 dominant hydration process in the shallow mantle lithosphere. Rouméjon et al. (2018) document the
108 development of serpentine minerals along detachment faults at slow- and ultraslow-spreading ridges
109 using samples drilled at Atlantis Massif (IODP Expedition 357) and dredged along the easternmost
110 Southwest Indian Ridge (SWIR, 62–65°E). The studied serpentine minerals have similar textures in
111 these environments, yet they have variable compositions when associated to gabbros, which suggest
112 cross-contamination by hydrothermal fluids. They are also isotopically heterogeneous for oxygen
113 isotopes down to scales of $\sim 100 \mu\text{m}$, and these variations depend on water/rock ratio. Pastore et al.
114 (2018) developed an innovative scanning magnetic microscopy technique allowing mapping, at the
115 millimeter to micrometer scale in serpentinized peridotites, the distribution of magnetite, one of the
116 mineral products of serpentinization reactions. This technique sheds new light on the strong
117 heterogeneity and variability in the direction of the magnetization with respect to the pristine
118 sample in serpentinized samples. Escario et al. (2018) investigate the effects of solute transport on
119 reaction paths during incipient serpentinization of olivine cores using a reactive-percolation
120 experimental approach. They show that, for the same initial fluid and mineral compositions,
121 serpentinization reaction paths vary depending on local flow distribution (at the microscale) and that
122 kinetics- and transport-controlled reaction paths can coexist at the sample scale. These mechanisms
123 favour the development of mineralogical and compositional heterogeneities. These results suggest a
124 contribution of flow rate on the development of the different serpentinization reactions paths
125 observed in the basement and in fault zones in serpentinized oceanic mantle lithosphere, where they
126 are commonly ascribed to changes in the composition of hydrothermal fluids resulting from
127 interactions with different lithologies along flow paths (e.g., Rouméjon et al, this volume).

128 Los et al. (2018) studied experimentally the formation of sulphides in relation to low temperature
129 hydrothermalism at mid-ocean ridges and more particularly the role of the composition of basement
130 rocks on sulfidation efficiency. Basalt, troctolite, dunite and serpentinite were reacted with H_2S -rich
131 hydrothermal fluids. Sulphides could be observed only in the basalt experiment indicating that
132 olivine-rich seafloor lithologies are not favourable to sulphidation reactions. These results challenge
133 the commonly proposed anhydrite formation models and open new research paths.

134 Noel et al. (2018) and Menzel et al. (2018) studied in natural samples the formation of carbonates
135 at the expense of ferromagnesian minerals, a process commonly associated to the alteration of
136 mantle peridotite. They carried out a detailed petro-structural, geochemical and carbon and oxygen
137 isotope study of variously altered peridotites from the Oman and the Newfoundland ophiolites. Noel
138 et al. (2018) document successive episodes of carbonate-forming reactions from the Oman Ophiolite,
139 first associated to serpentinization close to a mid-ocean ridge, then to the different stages of the

140 emplacement of the ophiolite. They demonstrate structural and chemical linkages between
141 serpentinization and carbonate-forming reactions during the cooling of the oceanic mantle
142 lithosphere in the presence of CO₂-bearing fluids, and evidence the control of inherited mantle fabric
143 on the field scale orientation of late carbonate veins. Menzel et al. (2018) document a complete
144 sequence of carbonate-forming and redox reactions preserved at the interface in the Advocate
145 Ophiolite Complex (Newfoundland, Canada). They show that this reaction sequence was triggered by
146 an influx of CO₂-rich fluids and that the devolatilization of neighbouring meta-sediments during
147 subduction is the most likely source of these fluids. They suggest that carbonate-bearing peridotites
148 can act as a carbon flux pathway beyond sub-arc depths.

149 Carbon occurs also as abiotic organic compounds as well as organic metabolic byproducts or
150 remnants of microbial ecosystems in the shallow mantle lithosphere. Ménez et al. (2018) show that
151 organic carbon can influence secondary mineral formation as well as the speciation and mobility of
152 transition metals during low temperature serpentinization reactions (< 200 °C). This mechanism has
153 possible implications for understanding ore formation during late active serpentinization of ophiolitic
154 massifs and/or for subsurface carbon dioxide storage in ultramafic rocks.

155

156

157 **Acknowledgements**

158 We thank Marco Scambelluri for supportive interactions throughout the edition of this volume. We
159 express our sincere thanks to all the reviewers who have participated in evaluating the articles
160 published in this special issue. The editors and many authors of this volume were supported by
161 funding from the People Programme (Marie Curie Actions) of the European Union's Seventh
162 Framework Programme FP7/2007-2013 under the REA grant agreement no. 608001 to develop their
163 research on the complex reactive systems driving the formation and evolution of the oceanic
164 lithosphere.

165

166

167 Figure Caption

168

169 Figure 1: Boundary layers marking the transition of mantle asthenosphere to the shallow sub-
170 seafloor at mid-ocean ridges for heterogeneous (left-hand side) or layered (right-hand side) oceanic
171 lithosphere. They correspond mainly to isotherms : (1) Asthenosphere-lithosphere boundary layer: (i)
172 mechanical boundary (**LithM**): transition between a visco-plastic asthenospheric mantle and a rigid
173 lithosphere (ca. 1000 °C; note that it depends also on the deformation and composition of the
174 mantle); (ii) chemical/petrological boundary (**LithT**): marks the crystallization of clinopyroxene and /
175 or plagioclase (ca 1180°C), this process locally blocks magma flow and induces a change in
176 permeability (and viscosity?) of the partially molten mantle; (2) **Moho**: seismic interface interpreted
177 as marking (i) a change in lithology in layered lithosphere, from mantle peridotite to magmatic crust
178 (max 1180°C at mid-ocean ridges), and (ii) the transition from serpentinite to peridotite in
179 heterogeneous mantle lithosphere (350-500 °C); (3) Lithosphere / hydrosphere interface i.e. limit of
180 penetration of hydrothermal fluids in the lithosphere (**Hy**): (i) In a layered lithosphere, this limit is
181 generally assumed to be the brittle-ductile transition in the oceanic crust (ca. 700°C-750 °C);
182 however, recent works suggest that very high temperature (up to 975°C, **HyHT**) hydrothermalism
183 could develop in lower gabbros; (ii) the dominant mechanisms driving the hydration of the
184 heterogeneous lithosphere remain debated (permeability and role of large faults vs. diffuse
185 penetration); assuming that the lithosphere / hydrosphere limit corresponds to the Moho implies
186 that hydrothermal fluids can interact with the lithosphere down to several kilometre deep.

187

188 **References**

- 189 Basch, V., Rampone, E., Crispini, L., Ferrando, C., Ildefonse, B., Godard, M., 2018. From mantle
190 peridotites to hybrid troctolites: textural and chemical evolution during melt-rock interaction
191 history (Mt.Maggiore, Corsica, France). *Lithos*, Special Volume "ABYSS", doi :
192 10.1016/j.lithos.2018.02.025
- 193 Borghini, G., Francomme, J., Fumagalli, P., 2018. Melt-dunite interactions at 0.5 and 0.7 GPa:
194 experimental constraints on the origin of olivine-rich troctolites, *Lithos*, Special Volume "ABYSS",
195 doi:10.1016/j.lithos.2018.09.022
- 196 Currin, A., Koepke, J., Almeev, R., Beerman, O., 2018. Interaction of highly saline fluid and olivine
197 gabbro: experimental simulation of deep hydrothermal processes involving amphibole at the base
198 of the oceanic crust, *Lithos*, Special Volume "ABYSS", doi:10.1016/j.lithos.2018.09.017
- 199 Currin, A., Wolff, P.E., Koepke, J., Almeev, R., Zhang, C., Ildefonse, B., Teagle, D., Zihlmann, B., 2018.
200 Chlorine-rich amphibole in deep layered gabbros as evidence for brine/rock interaction in the lower
201 oceanic crust: a case study from the Wadi Wariyah, Samail Ophiolite, Sultanate of Oman, *Lithos*,
202 Special Volume "ABYSS", doi:10.1016/j.lithos.2018.09.015
- 203 Escario, S., Godard, M., Gouze, P., Leprovost, R. 2018. Experimental study of the effects of solute
204 transport on reaction paths during incipient serpentinization. *Lithos*, Special Volume "ABYSS", doi :
205 10.1016/j.lithos.2018.09.020
- 206 Ferrando, C., Godard, M., Ildefonse, B., Rampone, E., 2018. Melt transport and mantle assimilation at
207 Atlantis Massif (IODP Site U1309): constraints from geochemical modeling. *Lithos*, Special Volume
208 "ABYSS" doi: 10.1016/j.lithos.2018.01.012
- 209 Früh-Green, G.L., Orcutt, B.N., Rouméjon, S., Lilley, M.D., Morono, Y., Cotterill, C., Green, S., Escartin,
210 J., John, B.E., McCaig, A.M., Cannat, M., Ménez, B., Schwarzenbach, E.M., Williams, M.J., Morgan,
211 S., Lang, S.Q., Schrenk, M.O., Brazelton, W.J., Akizawa, N., Boschi, C., Dunkel, K.G., Quéméneur, M.,
212 Whattam, S.A., Mayhew, L., Harris, M., Bayrakci, G., Behrmann, J.-H., Herrero-Bervera, E., Hesse, K.,
213 Liu, H.-Q., Ratnayake, A.S., Twing, K., Weis, D., Zhao, R., Bilenker, L., 2018. Magmatism,
214 serpentinization and life: Insights through drilling the Atlantis Massif (IODP Expedition 357). *Lithos*,
215 Special Volume "ABYSS", doi:10.1016/j.lithos.2018.09.012.
- 216 Koepke, J., Botcharnikov, R.E., Natland, J.H., 2018. Crystallization of late-stage MORB under varying
217 water activities and redox conditions: Implications for the formation of highly evolved lavas and
218 oxide gabbro in the ocean crust, *Lithos*, Special Volume "ABYSS", doi : 10.1016/j.lithos.2018.10.001
- 219 Los, C., Bach, W., 2018. Sulfidation of major rock types of the oceanic lithosphere; an experimental
220 study at 250°C and 400 bars. *Lithos*, Special Volume "ABYSS", doi:10.1016/j.lithos.2018.02.006

221 Ménez, B., Pasini, V., Guyot, F., Benzerara, K., Bernard, S., Brunelli, D., 2018. Mineralizations and
222 transition metal mobility driven by organic carbon during low-temperature serpentinization. *Lithos*,
223 Special Volume "ABYSS". 10.1016/j.lithos.2018.1007.1022

224 Menzel, M.D., Garrido, C.J., López Sánchez-Vizcaíno, V., Marchesi, C., Hidas, K., Escayola, M.P.,
225 Delgado Huertas, A., 2018. Carbonation of mantle peridotite by CO₂-rich fluids: the formation of
226 listvenites in the Advocate ophiolite complex (Newfoundland, Canada). *Lithos Special Volume*
227 "ABYSS", doi: 10.1016/j.lithos.2018.06.001.

228 Noël, J., Godard, M., Oliot, E., Martinez, I., Williams, M., Boudier, F., Rodriguez, O., Chaduteau, C.,
229 Escario, S., Gouze, P., 2018 Evidence of polygenetic carbon trapping in the Oman Ophiolite: petro-
230 structural, geochemical, and carbon and oxygen isotope study of the Wadi Dima harzburgite-hosted
231 carbonates (Wadi Tayin massif). *Lithos Special Volume "ABYSS"*, 10.1016/j.lithos.2018.08.020

232 Pastore, Z., Ter Maat, G., Church, N., McEnroe, S.A, Oda, H., Fumagalli, P., 2018. Millimeter to
233 Micrometer Scale Mapping of serpentinized dunite by High Resolution Magnetic Microscopy. *Lithos*
234 Special Volume "ABYSS", doi : 10.1016/j.lithos.2018.09.018

235 Rouméjon, S., Williams, M.J., Früh-Green, G.L., 2018. In-situ oxygen isotope analyses in serpentine
236 minerals: Constraints on serpentinization during tectonic exhumation at slow- and ultraslow-
237 spreading ridges. *Lithos Special Volume "ABYSS"* , doi: 10.1016/j.lithos.2018.09.021.

238 Zihlmann, B., Mueller, S., Koepke, J., Strauss, H., Teagle, D.A.H., 2018. Hydrothermal fault zones in
239 the lower oceanic crust: An example from Wadi Gideah, Samail ophiolite, Oman. *Lithos, Special*
240 *Volume "ABYSS"*, 10.1016/j.lithos.2018.09.008