



## Planta Medica

### Natural Products and Medicinal Plant Research

#### Editor

Prof. Dr. Adolf Nahrstedt  
Institut für Pharmazeutische Biologie  
und Phytochemie der Universität  
Hittorfstraße 56  
0-48149 Munster  
Germany  
Fax: +4925183-38341  
E-mail: nahrste@uni-muenster.de

#### Co-Editors

Prof. Dr. W. Sarz, Munster  
Prof. Dr. R. Bauer, Graz  
Prof. Dr. G. Franz, Regensburg  
Prof. Dr. M. Hamburger, Basel  
Prof. Dr. I. Khan, Oxford, MS

#### Editorial Advisory Board

Y. Asakawa, Tokushima  
L. Bohlin, Uppsala  
I. Calis, Ankara  
S. Canigual, Barcelona  
A. M. Clark, Oxford, MS  
O. Guo, Peking  
R. Maffei Facino, Milano  
S. Habtemariam, London  
G. Harnischfeger, Salzgitter  
W. Herz, Tallahassee  
M. Hesse, Zurich  
K. Hostettmann, Lausanne  
P. J. Houghton, London  
P. Joseph-Nathan, Mexico City  
G. Kahl, Frankfurt/M  
W. Kreis, Erlangen  
U. Matern, Marburg  
I. Merfort, Freiburg  
S. Nyiredy, Budakalasz  
K.-M. Oksman-Caldentey, Espoo  
P. Potier, Gif-sur-Yvette  
V. Schulz, Berlin  
J. F. Verbist, Nantes  
E. J. Verspohl, Munster  
A. Vlietinck, Antwerp  
A. Vollmar, Munchen  
H. Vuorela, Helsinki  
H. Yamada, Tokio

Georg Thieme Verlag KG  
Stuttgart · New York  
Rüdigerstraße 14  
0-70469 Stuttgart  
Postfach 3011 20  
0-70451 Stuttgart

#### Reprint

© Georg Thieme Verlag KG  
Stuttgart · New York

Thieme New York  
333 Seventh Avenue  
New York, NY 10001

# Tucumanin, a $\beta$ -Hydroxy- $\gamma$ -lactone Bistetrahydrofuranic Acetogenin from *Annona cherimolia*, is a Potent Inhibitor of Mitochondrial Complex I

Isabel Barrachina<sup>1</sup>, Adriana Neske<sup>2</sup>, Susana Granell<sup>1</sup>,  
Almudena Bermejo<sup>1</sup>, Nadia Chahboune I, Nouredine El Aouad<sup>1</sup>,  
Olga Alvarez<sup>2</sup>, Alicia Bardon<sup>2</sup>, M. Carmen Zafra-Polo<sup>1</sup>

## Abstract

A new {3-hydroxy- $\gamma$ -methyl- $\gamma$ -lactone bistetrahydrofuranic acetogenin, tucumanin, with the infrequent symmetrical *threo/trans/threo/trans/threo* relative configuration at the tetrahydrofuran rings was isolated from *Annona cherimolia* (Annonaceae) seeds. The inhibitory potency on the mitochondrial complex I of acetogenins with this relative configuration (tucumanin and asimicin) was compared with that shown by the corresponding pairs with an asymmetrical *threo/trans/threo/trans/erythro* relative configuration (laherradurin/rolliniastatin-2, and itrabin/molvizarin). All these compounds act as selective inhibitors of mitochondrial complex I in the 0.18 - 1.55 nM range.

"Cherimoya" is an edible fruit from a tropical tree (*Annona cherimolia* Mill., Annonaceae) native to Peru that is cultivated worldwide. In previous chemical studies on the seeds of *A. cherimolia* cultivated in a small pseudo-tropical area in the south of Spain, on the coast of Granada, four monotetrahydrofuranic (mono-THF) and eleven bistetrahydrofuranic (bis-THF) acetogenins were isolated [1], [2]. Recently, three new cytotoxic mono-THF acetogenins, annocherin, annomolin, and annocherimolin, were isolated from seeds of "cherimoya" grown in plantations of Southern California [31,14].

The Annonaceous acetogenins are claimed as potent inhibitors of NADH:ubiquinone oxidoreductase (complex I) in the mitochondrial electron transport chain [5]. Among them, the {3-hydroxy- $\gamma$ -methyl- $\gamma$ -lactone acetogenins, a type of acetogenin only isolated from the seeds of *A. cherimolia*, are more potent complex I inhibitors than any other natural acetogenin [6]. The high potency of this class of acetogenins led us to further investigate the composition of seeds collected in Tucuman (Argentina).

Affiliation: <sup>1</sup> Departamento de Farmacología. Facultad de Farmacia. Universidad de Valencia, Burjassot. Spain. <sup>2</sup> Departamento de Química Orgánica. Facultad de Química. Universidad Nacional de Tucumán. Tucumán. Argentina

Correspondence: Prof. Dr. M. Carmen Zafra-Polo. Departamento de Farmacología. Facultad de Farmacia. Avda. Vicent Andrés Estelles s/n. 46100 Burjassot. Valencia. Spain. Phone: +34-96-354-4944. Fax: +34-96-354-4943. E-mail: carmen.zafra@uv.es

Received: January 29, 2004. Accepted: June 4, 2004

Bibliography: *Planta Med* 2004; 70: 866-868. © Georg Thieme Verlag KG Stuttgart. New York. DOI:10.1055/s-2004-827237. ISSN 0032-0943

Fractionation of the methanolic extract furnished 11 known acetogenins (almunequin, asimicin, cherimolin-1, itrabin, laherradurin, molvizarin, motrilin, neoannonin, rolliniastatin-2, squamocin, squamocin-B) and a new acetogenin, tucumanin (1). The molecular weight of 1 was established by the  $[M+Na]^+$  ion observed at  $m/z = 647.45183$  (calcd. 647.44988 for  $C_{37}H_{65}O_7Na$ ) in the HR-ESI-MS. Absorptions at 204 nm and at 1760  $cm^{-1}$  in the UV and IR spectra, respectively, are in accordance with the presence of a saturated  $\gamma$ -lactone ring in 1 [1]. The  $^1H$ -NMR spectrum and the 2D heteronuclear correlation ( $^1H$ - $^1H$  COSY 45) of 1, prove the existence of an ABX spin system, characteristic of the three lactone ring protons of a (3-hydroxy- $\gamma$ -methyl- $\gamma$ -lactone acetogenin [7], [8]: H-2 ( $\alpha$ ) at  $\delta = 2.58$ , H-35 ( $\beta$ ) at  $\delta = 4.19$ , and H-36 ( $\gamma$ ) at  $\delta = 4.49$ . These protons are correlated in 2D heteronuclear NMR experiments (HMQC and HMBC) with three methine carbons at  $\delta = 43.7$ ,  $\delta 73.7$ , and  $\delta 82.4$ , respectively (Fig.1). The relative configuration of the lactone moiety deduced from the coupling constants,  $J_{\alpha-\beta} = 5.5$  Hz (*cis*) and  $J_{\beta-\gamma} = 1.0$  Hz (*trans*), was in agreement with that reported by Sibi et al. for the synthetic laherradurin [9].

The presence of an  $\alpha, \alpha'$ -dihydroxylated bis-THF system with the infrequent symmetrical *threo/trans/threo/trans/threo* relative configuration was deduced from  $^1H$ - $^{13}C$  NMR heteronuclear correlations between the signals at  $\delta = 3.38$  and  $\delta = 74.0$  (two methine alcohol), and between the signals at  $\delta = 3.81$  and  $\delta = 83.1$  (two methine tetrahydrofuran) [10]. This system is placed between C-15 and C-24 based on the fragment ions observed in the EI-MS (Fig.1). Compound 1 is the first {3-hydroxy- $\gamma$ -methyl- $\gamma$ -lactone adjacent bis-THF acetogenin with a *threo/trans/threo/trans/threo* relative configuration.

In previous studies we have shown that the inhibitory potency of acetogenins is modulated by several structural factors. Therefore, it is relevant to compare the inhibitory potency of acetogenins with this type of configuration (tucumanin, 1 and asimicin, 2) with that of the corresponding pairs with an asymmetrical *threo/trans/threo/trans/erythro* relative configuration (laherradurin, 3/rolliniastatin-2, 5, and itrabin, 4/molvizarin, 6). As potency criterion we used the  $IC_{50}$  against the integrated NADH oxidase activity of beef heart open submitochondrial particles [5], [6], [12], which evaluates the complex I activity in a more physiological environment. Results are shown in Table 1.

The {3-hydroxyacetogenins with a *threo/trans/threo/trans/erythro* relative configuration, laherradurin (3) and itrabin (4) ( $IC_{50} = 0.18$  and  $0.21$  nM, respectively) are more potent inhibitors of the mitochondrial complex I than the corresponding  $\alpha, \beta$ -unsaturated  $\gamma$ -lactone acetogenins, rolliniastatin-2 (5) and molvizarin (6) ( $IC_{50} = 0.51$  and  $1.55$  nM, respectively). However, tucumanin (1) ( $IC_{50} = 0.57$  nM) shows an inverted tendency when compared to asimicin (2) ( $IC_{50} = 0.33$  nM), both with a *threo/trans/threo/trans/threo* relative configuration. Therefore, our data show that, indeed, the terminal  $\gamma$ -methyl- $\gamma$ -lactone ring and the relative configuration at the THF system are both relevant structural factors for a potent inhibition of the mitochondrial complex I, because of a double binding to the active site of the enzyme to which both moieties probably contribute.






Fig. 1  $^1H$ - $^{13}C$  NMR heteronuclear correlations ( $^{13}C$  in parentheses) spectral data, and significant EIMS fragment ions of tuconanin (1).

Table 1 Inhibitory potency of acetogenins 1- 6

	Compound						rotenone [6]
	2	3	4	5	6		
NADH oxidase $_{1(50)}$ (nM)	0.57 $\pm$ 0.07*	0.33 $\pm$ 0.03'	0.18 $\pm$ 0.02*	0.21 $\pm$ 0.03'	0.51 $\pm$ 0.03'	1.551'0.17'	5.10 $\pm$ 0.09

Significant difference compared with rotenone; \*  $P < 0.001$ .

## Material and Methods

Optical rotation (Perkin-Elmer 241 polarimeter), IR spectra (Satellite FTIR-Mattson 980514 spectrometer), MS (VG Auto Spec Fisons spectrometer),  $^1H$ - and  $^{13}C$ -NMR, DEPT, HMBC, HMQC, tH-tH-COSY (Varian Unity 400 MHz spectrometer) using the solvent signal as reference ( $CDCl_3$ ) at  $\delta = 7.26$  and 77.0). Chromatograph-

ic separations were carried out by column chromatography on silica gel 60 H(5 - 40  $\mu m$ , 7336 Merck). Semi-preparative HPLC was carried out on a LiChroCart<sup>R</sup> 100 RP-18 column (25 x 1 em i. d., 10  $\mu m$  particle size), flow rate 4.5 mL/min. using MeOH-H<sub>2</sub>O-THF. Analytical TLC was performed on Merck precoated silica gel 60 F<sub>254</sub> plates.

The fruits of *Annona cherimolia* Mill. ("cherimoya") were collected at Yerba Buena, Tucuman, Argentina, in February 2000. A voucher sample (No. 515092) was deposited at the Herbarium of Instituto Lillo of Tucuman.

The dried and powdered seeds of cherimoya (350 g) were percolated with methanol (2 x 3 L). Evaporation of the solvent yielded a crude MeOH extract (15 g) which was further partitioned between CH<sub>2</sub>Cl<sub>2</sub> (3 x 100 mL) and H<sub>2</sub>O (100 mL). The CH<sub>2</sub>Cl<sub>2</sub> layer was washed with water, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated in vacuum to give 9 g of a solid extract with an acetogenin mixture, which was chromatographed on a silica gel 60 H (315 g) column, using gradients of hexane/EtOAc (90 : 10; 80 : 20; 70 : 30; 60 : 40; 50 : 50; 40 : 60; 20 : 80 and 10 : 90) giving eight fractions (F-I to F-VIII, 250 mL each). Fraction F-III (hexane/EtOAc 70 : 30, 660 mg) was again chromatographed on silica gel 60 H (23 g) and eluted with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (95 : 5) to give five subfractions (50 mL each), one of them (F-III<sub>1</sub>) was applied to semipreparative HPLC with MeOH-H<sub>2</sub>O-THF (87 : 13 : 5) affording neoannonin {12 mg, R<sub>f</sub> = 10.8 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +16.2° (CHCl<sub>3</sub>, e 0.8)}; F-IV (hexane/EtOAc, 60 : 40, 2 g) was chromatographed on silica gel 60 H (70 g) and eluted with CH<sub>2</sub>Cl<sub>2</sub>/EtOAc/MeOH (50 : 45 : 5) to yield eight subfractions (100 mL each). F-IV<sub>5</sub> and F-IV<sub>6</sub> by semipreparative HPLC with MeOH-H<sub>2</sub>O-THF 80 : 20 : 5 afforded squamocin {90 mg, R<sub>f</sub> = 16.6 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +14.0° (CHCl<sub>3</sub>, e 0.13), squamocin-B (16 mg, R<sub>f</sub> = 8.7 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +27.6° (MeOH, e 0.2) and motrilin (5 mg, R<sub>f</sub> = 14.8 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +10.8° (MeOH, c 0.13)), and asimicin {2, 5 mg, R<sub>f</sub> = 29.4 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +11.0° (CHCl<sub>3</sub>, c 0.22), molvizarin (6, 10 mg, R<sub>f</sub> = 13.9 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +10.0° (CHCl<sub>3</sub>, e 0.2) and rolliniastatin-2 (5, 5 mg, R<sub>f</sub> = 11.3 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +12.8° (CHCl<sub>3</sub>, e 0.26)), respectively. F-VI (hexane/EtOAc, 40 : 60, 500 mg) was chromatographed on silica gel (18 g) and eluted with CH<sub>2</sub>Cl<sub>2</sub>/EtOAc/MeOH (60 : 13 : 10) to give four subfractions (75 mL each). One of them (F-VI<sub>2</sub>) was applied to semipreparative HPLC with MeOH-H<sub>2</sub>O-THF (80 : 20 : 5) affording cherimolin-1 {15 mg, R<sub>f</sub> = 8.9 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +64.0° (MeOH, e 0.3) and almunequin (5 mg, R<sub>f</sub> = 5.8 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +13.5° (CHCl<sub>3</sub>, c 0.14)}. F-VIII (hexane/HOAc, 10 : 90, 250 mg) was chromatographed on silica gel (9 g) and eluted with CH<sub>2</sub>Cl<sub>2</sub>/HOAc/MeOH (10 : 85 : 5) to give three subfractions (50 mL each). F-VIII<sub>2</sub> was applied to semipreparative HPLC with MeOH-H<sub>2</sub>O-THF (80 : 20 : 5) affording itrabin {4, 40 mg, R<sub>f</sub> = 2.1 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +21.0° (MeOH, c 0.2), laherradurin (3, 60 mg, R<sub>f</sub> = 3.5 min, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +21.0° (MeOH, e 0.09) and tucumanin (1, 11 mg, R<sub>f</sub> = 4.2 min)}.

**Tucumanin** (1): waxy compound, [ $\alpha$ ]<sub>D</sub><sup>25</sup>: +1.5° (EtOH, e 0.8); UV (EtOH):  $\lambda$ <sub>max</sub> (log  $\epsilon$ ) = 204 nm (3.8). IR (film):  $\nu$ <sub>max</sub> = 3360, 2923, 2853, 1760, 1465, 1052 cm<sup>-1</sup>. HR-LSI-MS:  $m/z$  = 647.45183 [M + Na]<sup>+</sup> (calcd. for C<sub>37</sub>H<sub>68</sub>O<sub>7</sub>Na: 647.44988). EI-MS, 1H-NMR (400 MHz), and 13C-NMR (100 MHz), see Fig.1.

The inhibitory potency of acetogenins as complex I inhibitors was assayed using submitochondrial particles from beef heart as previously reported [5], [11], [12]. NADH oxidase activity was measured as the aerobic oxidation of 75  $\mu$ M NADH in the absence of external quinone substrates and other inhibitors of the respiratory chain. Rotenone (Sigma-Aldrich St. Louis, MO, USA) was used as the positive control. Data from four titrations were used to determine the statistic means and the standard deviation.

Significance of differences was evaluated by the Student's *t* test, with *P* < 0.001 being regarded as significant.

## Acknowledgements

This research was supported by the Spanish Direccin General de Investigacin under grant SAF 2001 -3142. and by the Secretaria del Plan Valenciano de Investigacin Cientfica, Desarrollo Tecnolgico e Innovacin de la Generalitat Valenciana under grant GVOI-282.

## References

- 1 Cavé A, Figadère B, Laurens A, Cortes D. Acetogenins from Annonaceae. In: Herz W, Kirby GW, Moore RE, Steglich W, Tamm Ch, editors. Progress in the chemistry of Organic Natural Products: Vol.70 Wien: Springer-Verlag, 1997: p 81-288
- 2 Zaft'a-Polo MC, Figadere B, Gallardo T, Tormo JR, Cortes D. Natural acetogenins from Annonaceae, synthesis and mechanisms of action. Phytochemistry 1998; 48: 1087-1117
- 3 Woo MH, Kim DH, Fotopoulos SS, McLaughlin JL. Annocherin and (2,4)-cis- and trans-annocherinones, monotetrahydrofuran annonaceous acetogenins with a C-7 carbonyl group from *Annona cherimolia* seeds. J Nat Prod 1999; 62: 1250-5
- 4 Kim DH, Ma ES, Suk KD, SonJK, LeeJS, Woo M-H, 2001. Annonolin and annocherimolin, new cytotoxic annonaceous acetogenins from *Annona cherimolia* seeds. J Nat Prod 1996; 64: 502-506
- 5 Degli Esposti M, Ghelli A, Ratta M, Cortes D, Estornell E. Natural substances (acetogenins) from the family Annonaceae are powerful inhibitors of mitochondrial NADH dehydrogenase (complex I). Biochem J 1994; 301: 161-7
- 6 Tormo JR, Estornell E, Gallardo T, Gonzalez MC, Cavé Ad, Granell S, Cortes D, Zafra-Polo Me.  $\gamma$ -Lactone functionalized antitumoral acetogenins are the most potent inhibitors of mitochondrial complex I. Bioorg Med Chem Lett 2001; 11: 681-4
- 7 Cortes D, Myint SH, Leboeuf M, Cavé A. A new type of cytotoxic acetogenins: the tetrahydrofuranic  $\beta$ -hydroxy methyl  $\gamma$ -lactones. Tetrahedron Lett 1991; 32: 6133-4
- 8 Cortes D, Myint SH, Dupont B, Davoust D. Bioactive acetogenins from seeds of *Annona cherimolia*. Phytochemistry 1993; 32: 1475-82
- 9 Sibi M, Lu J, Talbacka CL. Enantiospecific synthesis of trisubstituted butyrolactone natural products and their analogs. J Org Chem 1996; 61:7848-55
- 10 Cortes D, Figadere B, Cavé A. Bis-tetrahydrofuran acetogenins from Annonaceae. Phytochemistry 1993; 32: 1467-73
- 11 Fato R, Estornell E, Di Bernardo S, Pallotti F, Parenti-Castelli G, Lenaz G. Steady-state kinetics of the reduction of coenzyme Q analogs by complex I (NADH:ubiquinone oxidoreductase) in bovine heart mitochondria and submitochondrial particles. Biochemistry 1996; 35: 2705-16
- 12 Gallardo T, Zafra-Polo MC, Tormo JR, Gonzalez MC, Franck X, Estornell E, Cortes D. Semisynthesis of antitumoral acetogenins: SAR of functionalized alkyl-chain bis-tetrahydrofuranic acetogenins, specific inhibitors of mitochondrial complex I. J Med Chem 2000; 43: 4793-800