

University of Massachusetts Medical School

eScholarship@UMMS

---

Open Access Articles

Open Access Publications by UMMS Authors

---

2020-01-15

## Glycan profiles of gp120 protein vaccines from four major HIV-1 subtypes produced from different host cell lines under non-GMP or GMP conditions

Shixia Wang

*University of Massachusetts Medical School*

*Et al.*

Let us know how access to this document benefits you.

Follow this and additional works at: <https://escholarship.umassmed.edu/oapubs>



Part of the [Amino Acids, Peptides, and Proteins Commons](#), [Immunology of Infectious Disease Commons](#), [Immunoprophylaxis and Therapy Commons](#), [Virology Commons](#), [Virus Diseases Commons](#), and the [Viruses Commons](#)

---

### Repository Citation

Wang S, Voronin Y, Zhao P, Ishihara M, Mehta N, Porterfield M, Chen Y, Bartley C, Hu G, Han D, Wells L, Tiemeyer M, Lu S. (2020). Glycan profiles of gp120 protein vaccines from four major HIV-1 subtypes produced from different host cell lines under non-GMP or GMP conditions. Open Access Articles. <https://doi.org/10.1128/JVI.01968-19>. Retrieved from <https://escholarship.umassmed.edu/oapubs/4127>

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](#).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

1       **Glycan profiles of gp120 protein vaccines from four major HIV-1 subtypes**  
2       **produced from different host cell lines under non-GMP or GMP conditions**

3

4       Shixia Wang<sup>a,b</sup>, Yegor Voronin<sup>b</sup>, Peng Zhao<sup>c</sup>, Mayumi Ishihara<sup>c</sup>, Nickita Mehta<sup>c</sup>, Mindy  
5       Porterfield<sup>c</sup>, Yuxin Chen<sup>a</sup>, Christopher Bartley<sup>d</sup>, Guangnan Hu<sup>a</sup>, Dong Han<sup>a</sup>, Lance  
6       Wells<sup>c</sup>, Michael Tiemeyer<sup>c</sup>, Shan Lu<sup>a#</sup>

7

8       <sup>a</sup>Department of Medicine, University of Massachusetts Medical School, Worcester, MA;

9       <sup>b</sup>Worcester HIV Vaccine, Inc., Worcester, MA;

10      <sup>c</sup>Complex Carbohydrate Research Center, University of Georgia, Athens, GA;

11      <sup>d</sup>Waisman Biomanufacturing, Madison, WI;

12

13

14

15      **Running Title:** HIV-1 gp120 protein glycan profiles

16      **Key Words:** HIV-1, gp120, Env protein, glycosylation, vaccine

17      **Abstract length:** 234 words

18      **Manuscript length:** 5277 words

19

20      **#Corresponding Author:**

21             Shan Lu

22             Department of Medicine

23             University of Massachusetts Medical School

24             55 Lake Avenue North, Worcester, MA 01532

25             Phone: 508-856-6791

26             Email: [shan.lu@umassmed.edu](mailto:shan.lu@umassmed.edu)

27

28 **Abstract**

29 Envelope glycoprotein (Env) of human immunodeficiency virus type 1 (HIV-1) is an  
30 important target for the development of an HIV vaccine. Extensive glycosylation of Env  
31 is an important feature that both protects the virus from antibody responses and serves  
32 as a target for some highly potent broadly neutralizing antibodies. Therefore, analysis  
33 of glycans on recombinant Env proteins is highly significant. Here we present  
34 glycosylation profiles of recombinant gp120 proteins from four major clades of HIV-1 (A,  
35 B, C, and AE) produced either as research-grade material in 293 and CHO cells or as  
36 two independent lots of clinical material under GMP conditions. Almost all potential N-  
37 linked glycosylation sites were at least partially occupied in all proteins. The occupancy  
38 rates were largely consistent among proteins produced under different conditions,  
39 although a few sites showed substantial variability even between two GMP lots. Our  
40 data confirmed previous studies in the field showing high abundance of oligomannose  
41 on Env protein, with 40-50% of glycans having Man<sub>5</sub>-Man<sub>9</sub> on all four proteins under all  
42 production conditions. Overall the differences in occupancy and glycan forms among  
43 Env from different subtypes produced under different conditions were less dramatic than  
44 anticipated and antigenicity analysis with a panel of six monoclonal antibodies showed  
45 that all four gp120s maintained their antibody-binding profiles, including antibodies that  
46 recognize glycan forms. Such findings have major implications to the final production of  
47 a clinical HIV vaccine including Env glycoprotein components.

48

49

50 **Importance**

51 HIV-1 Env protein is a major target for the development of an HIV-1 vaccine. Env is  
52 covered with a large number of sugar-based glycan forms – about 50% of the Env  
53 molecular weight is composed of glycans. Glycan analysis of recombinant Env proteins  
54 is important to understand its roles in viral pathogenesis and immune responses. The  
55 current report presents the first extensive comparison of glycosylation patterns of  
56 recombinant gp120 proteins from four major clades of HIV-1 produced in two different  
57 cell lines, grown at either laboratory condition or at 50L GMP scale across different lots.  
58 Information learned in this study is valuable for the further design and production of HIV-  
59 1 Env proteins as the critical components of HIV-1 vaccine formulations.

60

61 **Introduction**

62 The envelope glycoprotein (Env) of human immunodeficiency virus type 1 (HIV-1) plays  
63 the key role in viral entry and serves as a major target for a preventive HIV-1 vaccine.

64 Env is heavily glycosylated with N-glycans comprising about half of its molecular weight.

65 In the course of HIV-1 evolution, either within an individual patient or on the population  
66 level, viral gene mutations may lead to the disappearance of certain glycosylation sites  
67 and to the appearance of new ones. This shifting glycan shield protects the Env

68 proteins from the engagement of antibodies elicited during the course of viral infection  
69 and contributing to the growth of escaping viral mutants in chronic infection (1).

70 Nevertheless, some glycan features are highly conserved even across clades such that

71 glycans contribute to several key antigenic domains recognized by broadly neutralizing  
72 antibodies. For example, the highly conserved N332 glycan is important for the binding

73 of monoclonal antibodies (mAbs) PGT128 and 10-1074, while N160 glycan is

74 recognized by mAbs PG9 and PG16 (2, 3). Another conserved feature of HIV glycan

75 shield is high abundance of unusual oligomannose forms, which normally serve as

76 intermediates in mammalian glycan synthesis (4–6).

77

78 Glycosylation pattern can be expected to vary depending on multiple factors impacting

79 glycoprotein synthesis, including the viral strain, the form in which Env is expressed

80 (gp120, gp140, gp160), the cell type, the protein expression levels, and even the

81 metabolic state of the cell (7). In past studies, the exact proportion of oligomannose on

82 Env varied from 17% to 98% with levels of 40-75% being common for both monomeric

83 gp120 and native trimers (5, 8, 9). However, analysis of two batches of membrane-

84 anchored Env showed remarkable consistency of forms found at each glycosylation site  
85 (9, 10), indicating that glycosylation patterns are generally preserved when the same  
86 Env protein is produced under identical conditions and that the differences in  
87 oligomannose content reflect either virus- or host cell specific factors.

88

89 In previous studies the viral clade-specific differences in the abundance of  
90 oligomannose have been attributed to the differences in the total number and regional  
91 density of the glycosylation sites, with higher glycan densities correlating with higher  
92 oligomannose content (4). On the other hand, comparisons of glycosylation of the  
93 same Env proteins expressed in 293 and CHO cells revealed mostly similar  
94 oligomannose content, similar occupancy, and similar glycan profiles, but with some  
95 notable exceptions (11, 12). It was observed that more complex glycans were present  
96 on CHO-derived clade C gp120 as compared to 293-derived protein, particularly at two  
97 sites (N386 and N392).

98

99 The increasing understanding of the impact of glycans on HIV Env immunogenicity and  
100 the increased focus on recombinant Env proteins after the RV144 trial led to the  
101 growing appreciation of the importance of these features to the design of Env-based  
102 protein vaccines against HIV-1(13). In particular, characterization of glycan profiles of  
103 recombinant Env proteins will be important to interpret the resulting antibody responses.

104

105 Various approaches of producing and purifying recombinant Envs for laboratory  
106 research and for clinical studies can be employed, but there's limited information based

107 on well-controlled studies for how different approaches may impact Env glycosylation.  
108 While transiently-transfected 293 cells are often used to produce research-grade  
109 proteins, the proteins for clinical use are usually produced in stably transfected CHO  
110 cells. Clinical material is usually produced in bioreactors that have larger volumes and  
111 reach higher cell density than those used in research-grade protein production. Diverse  
112 purification processes are used both in the lab and during clinical-grade protein  
113 purification, such as antibody-based affinity columns, size-exclusion chromatography,  
114 lectin-based columns, or the industry preferred ion-exchange columns. Recently, a few  
115 recombinant GMP-grade Env-based vaccines have been characterized by analysis of  
116 glycans (14, 15) but the number of Env proteins included in those studies was limited  
117 and they did not provide direct comparison between different cell lines or between GMP  
118 and non-GMP for the production of the same Env proteins.  
119  
120 Here we expanded the study to characterize glycosylation profiles of four recombinant  
121 gp120 proteins from four major clades of HIV-1 (subtypes A, B, C, and AE) produced  
122 under GMP condition for a Phase I human clinical trial (HVTN124). We analyze two  
123 separate GMP lots of the same four gp120 proteins comparing them to the same four  
124 gp120 proteins produced under non-GMP condition in CHO and 293F cells. Our results  
125 provide much needed information on the Env glycan patterns among different viral  
126 clades and between different preparations of the same protein. Such information is not  
127 only valuable for the better understanding on the variation of Env glycan patterns but  
128 also critical for the establishment of quality control standards for the production of  
129 clinical grade Env-based HIV-1 vaccines.

130 **Results**

131 **HIV-1 gp120 Env proteins from four clades and produced under different**

132 **conditions**

133 The four gp120 glycoproteins included in the current study were selected based on the  
134 immunogenicity analysis of a large panel of HIV-1 Env variants (16) and were included  
135 in a polyvalent DNA prime-protein boost HIV vaccine formulation currently going  
136 through a phase I clinical study at HIV Vaccine Trials Network (HVTN, protocol 124).  
137 They represent three primary isolates from clades A, B, and C, as well as a consensus  
138 variant from the AE clade. Their amino acid sequences have a low homology to each  
139 other in the range of 75-80 % (Fig.1A). They have 23-26 potential N-linked  
140 glycosylation sites (PNGSs), which are distributed throughout the sequence in a similar,  
141 but distinct manner (Fig.1B).

142

143 Research-grade gp120 proteins were produced by transient transfection of 293F cells  
144 and from stably-transfected CHO cells in a laboratory setting and the proteins were  
145 purified using lectin-based columns. For the GMP manufacturing process, stably  
146 transfected CHO cells expressing each of the proteins were grown in 50L bioreactors  
147 and the purification process involved ion-exchange columns. Two separate GMP  
148 manufacturing runs have been done under identical conditions, which allowed us the  
149 opportunity to compare the consistency of glycosylation profiles from one GMP lot to  
150 another.

151

152 **Glycan analysis of research-grade proteins**



153 Glycan heterogeneity for gp120 proteins produced as research-grade reagents in CHO  
154 and 293F cell lines were first analyzed. Digestion with PNGase F was used to release  
155 glycans from the gp120 proteins and the released glycans were permethylated and  
156 analyzed by NSI-MSn to characterize glycan structural features. Representative  
157 profiles for clade B gp120 are shown on Fig. 2A. The types of glycan forms found on  
158 proteins produced in CHO and 293F cells were generally very similar and only a few  
159 types comprised more than 10% of the total glycans (Fig. 2B). A large proportion of  
160 oligomannose forms (Man<sub>7</sub>-Man<sub>9</sub>) was detected in both preparations. A diverse group  
161 of complex glycans were also present, as well as some hybrid forms. The results for  
162 proteins from three other clades were similar (data not shown).

163

164 To study occupancy rate at each PNGS, proteins were digested with several proteases  
165 to produce peptides for LC/MS analysis and then consecutively digested with EndoH  
166 and PNGase F to allow detection of occupancy by different types of glycans at each  
167 PNGS. Digestion with EndoH cleaves N-linked glycans between the two N-  
168 acetylglucosamine (GlcNAc) residues in the core region of the glycan chain on high-  
169 mannose and hybrid, but not complex, glycans, leaving one GlcNAc still bound to the  
170 protein. Treatment with PNGase F removes all glycans that have not been impacted by  
171 EndoH treatment and leaves an aspartic acid residue at the site of N-linked  
172 glycosylation, which can be distinguished from the original asparagine by mass  
173 spectrometry of the peptides. Therefore, consecutive digestion with EndoH and  
174 PNGaseF allowed us to distinguish between oligomannose and complex glycans at

175 each site. The presence of the original asparagine in the peptide indicates that the  
176 PNG site has not been glycosylated.

177

178 Analysis of PNGS occupancy of research grade 293F- and CHO-produced proteins  
179 showed that most of PNGSs were at least partially occupied in both cases (Fig. 3A).  
180 N141, N186 and N339 in clade B protein, N186 and N397 in clade C protein, and N465  
181 in clade AE protein were the only sites that showed less than 20% occupancy in our  
182 analysis. A large proportion of oligomannose glycans was observed for all four gp120  
183 proteins, with higher proportions in glycans described as innate mannose patch.

184

185 Glycosylation profiles of proteins produced in 293F and CHO cells were remarkably  
186 similar (Fig. 3B). Most of the variation in occupancy was under 30 percentage points.  
187 Clade B proteins showed the largest variation with CHO-produced protein being more  
188 glycosylated than 293-produced protein. In some cases the changes were not in the  
189 total occupancy, but in the relative abundance of oligomannose and complex glycans.  
190 For example, N262 in clade B proteins was almost 100% occupied, but 293-produced  
191 protein carried an equal mixture of oligomannose and complex glycans, and CHO-  
192 produced proteins had almost exclusively oligomannose glycans at this site.

193

#### 194 **Glycan profile analysis of four GMP-grade gp120 proteins**

195 The four GMP-grade gp120 proteins have been produced in stably expressing CHO cell  
196 lines at 50L scale and purified using a multi-step chromatography. Two separate lots  
197 have been manufactured using the same master cell bank CHO cells and the same

198 fermentation and downstream purification process, which allowed us to investigate the  
199 lot-to-lot variability of GMP grade gp120 protein preparations. The glycan forms  
200 identified from GMP grade gp120 proteins are shown in Fig.4, and the relative amounts  
201 of these glycan forms between two GMP lots are show in Fig.5. For most gp120  
202 proteins, oligomannose (Man<sub>5</sub>-Man<sub>9</sub>) constituted 40-50% of the total glycans, while  
203 complex glycans were about 35-45%, and the rest were hybrid glycans or  
204 paucimannose (Man<sub>3</sub>-Man<sub>4</sub>). The least-processed Man<sub>9</sub> and Man<sub>8</sub> forms predominated  
205 on clade A and clade B gp120 proteins, while clade C and clade AE proteins showed  
206 higher percentages of Man<sub>5</sub>. Percentage of paucimannose showed the largest variation  
207 both between the clades and especially between the two lots of clade C gp120 protein,  
208 where in one of the lots paucimannose represented 23% of all glycans. Other proteins  
209 showed a more consistent glycan composition for two independent lots. Complex  
210 glycans were predominantly (63-87%) sialylated and >85% were core fucosylated  
211 (Fig.6).

212

213 Thus all variants of GMP-grade gp120 proteins exhibited high proportion of  
214 oligomannose glycans. The comparison of the two GMP lots showed mostly  
215 comparable glycan composition for three tested variants of gp120 proteins (clades A, B  
216 and AE) and some variation in the amount of paucimannose for clade C variant.

217

#### 218 **PNGSs occupancy analysis of GMP-grade gp120 proteins**

219 Next, PNGS occupancy was mapped for proteins from one of the GMP lots and  
220 compared to the CHO-produced research-grade proteins. The overall occupancy profile

221 of the GMP-grade gp120 proteins was generally similar to that of research-grade  
222 proteins, although the GMP proteins tended to have higher proportion of complex  
223 glycans (Fig.7). Analysis of the protein from the second lot did not reveal any major  
224 differences with the first lot (data not shown).

225

226 The oligomannose glycans were not equally distributed among the PNGSs. N262,  
227 N289/N295, N332, and N363 were enriched in oligomannose, corresponding to the  
228 intrinsic mannose patch that has been noted for HIV Env previously. While present on  
229 all four proteins, the patch was more pronounced for clade A, B, and C proteins, while  
230 oligomannose was more evenly distributed on clade AE protein. Clade C had an  
231 unusual enrichment in oligomannose at the C-terminus (N406, N442, N448, N463) that  
232 was absent in the other three proteins.

233

#### 234 **Antigenicity analysis among 4 gp120 proteins**

235 Finally, we sought to test whether any observed lot-to-lot variation in PNGS occupancy  
236 and glycan composition, no matter how minor they may be, had an effect on key  
237 antigenic features of these proteins. Using the affinity-measuring Octet Qke system and  
238 a panel of probing reagents we tested the preservation of key epitopes on these four  
239 gp120 protein: CD4 binding site (IgG-CD4 and mAb VRC01), V2 loop (mAb 2158), V3  
240 region (mAb R16), gp120 bridging sheet and loop F overlapping with CD4bs (mAb  
241 R53), and glycan forms (mAb 2G12 and mAb PGT128). The epitope for the 2G12  
242 antibody is believed to include mannose-rich glycans at positions 295, 332, 392, 386,  
243 and 448 (17). The PGT128 epitopes includes glycans at positions 332 and 301, as well  
244 as the C-terminal end of the V3 loop (18). While we observed differences in affinities of

245 these reagents, the differences between two lots of each clade were minimal (Table 1).  
246 This includes similar affinity of glycan-dependent 2G12 and PGT128 to the lot 2 of clade  
247 C protein, which exhibited the unusually high proportion of paucimannose compared to  
248 lot 1. Thus, our results demonstrate that observed differences in glycan profiles did not  
249 result in changes in affinity of antibodies targeting key epitopes of four gp120 proteins,  
250 including glycan-binding antibodies, indicating that proteins produced under different  
251 conditions mostly preserved their antigenic structure.  
252

253 **Discussion**

254 This report presents the first extensive comparison of glycosylation patterns of  
255 recombinant gp120 proteins from four clades of HIV-1 across two different cell lines,  
256 grown either at laboratory scale or at 50L GMP conditions, purified using different  
257 methods, and across two GMP lots under identical conditions. Our results show that in  
258 all four gp120 proteins included in the current study, the majority of PNG sites were  
259 occupied by glycans with occupancy rates usually above 50%. We also found that  
260 among glycans found on these proteins, oligomannose forms represented 40-50% and  
261 were concentrated in the previously described intrinsic mannose patch. At the same  
262 time, glycosylation profiles were basically very similar, with a low level of variability  
263 under different conditions. These differences did not affect binding by a panel of  
264 antibodies targeting key immunologic epitopes of gp120 proteins, indicating that the  
265 observed low level glycan differences should not have a major impact on protein  
266 immunogenicity.

267

268 The most prominent feature of HIV Env glycans distinguishing them from glycans on  
269 host proteins is the presence of oligomannose. In agreement with previous reports, we  
270 observed high proportion of oligomannose glycans (Man<sub>5</sub>-Man<sub>9</sub>) on all four proteins  
271 produced under all conditions confirming that this is a characteristic feature of HIV  
272 envelope. Even though previous studies reported a rather wide range of the percentage  
273 of oligomannose, we consistently observed that in all cases it constituted 40-50% of  
274 glycans. At least one recent study reported that gp120 derived from infectious virions  
275 contained 50% oligomannose glycans (19), which suggests that glycosylation patterns

276 of native Env are generally preserved on the recombinant gp120 proteins. This  
277 preservation is important for HIV vaccine development aiming to elicit antibodies that  
278 bind and neutralize HIV virions.

279

280 The research-grade gp120 protein was produced in 293F and CHO cells and purified  
281 using lectin columns, while the GMP-grade protein was produced in CHO cells grown at  
282 50L scale and purified using ion exchange columns. Nevertheless, we found only minor  
283 differences in glycan occupancy and glycan content between proteins produced under  
284 different conditions, indicating that these features are primarily determined by viral  
285 sequence and not by the producing cells or purification process.

286

287 Our comparison of two independent GMP-grade lots of gp120 proteins showed a  
288 consistent glycosylation pattern between them, but also did show some differences,  
289 including a significant increase in paucimannose content in one of the lots of clade C  
290 protein. It should be noted that this particular lot differed from all other GMP lots in  
291 having a significantly higher yield of the protein. We do not have enough data to  
292 establish the causal relationship between these two observations, but hypothesize that  
293 high levels of protein production overwhelmed medial and trans-Golgi glycan-  
294 processing machinery in producing cells, resulting in secretion of proteins with glycans  
295 that were fully trimmed by cis-Golgi mannosidases but incompletely branched,  
296 extended, and capped. Further work is needed to test the factors impacting variability  
297 of glycosylation during GMP manufacturing and acceptable variability levels should be

298 established. Special attention should be paid to optimizations that boost protein  
299 production in cells as that may have an impact on cellular glycosylation machinery.  
300



301 **Materials and Methods**

302 **Production of Non-GMP gp120 proteins**

303 The four gp120 glycoproteins used in this report are from HIV-1 clade A isolate  
304 92UG037.8, clade B isolate JRFL, clade C isolate 93MW965.26 and clade AE  
305 consensus, respectively (16). The non-GMP research grade HIV-1 gp120 proteins were  
306 produced using two protein expression systems: transiently transfected 293F cell  
307 expression and stably transfected CHO cell expression system. The codon optimized  
308 gp120-coding DNA inserts cloned in the vector pJW4303 were used in both 293F and  
309 CHO cells. To produce gp120 proteins from transiently transfect 293F cells, the serum-  
310 free 293F cell supernatant was collected at 72 hours after transfection. To express  
311 gp120 proteins from CHO cells, the serum-free culture supernatant of stable transfected  
312 CHO cells were collected. The harvested gp120 proteins from both 293F and CHO  
313 cells in research grade were purified using a Lectin column and verified by SDS-PAGE  
314 and Western blot analysis as previously described (20–22).

315

316 **Production of GMP grade gp120 proteins**

317 The codon optimized gp120 gene inserts from the same four clades (A, B, C and AE) as  
318 described above for research grade proteins were transfected into CHO DG44 cells  
319 (Invitrogen, CA), respectively, and used to establish the master cell banks. Each of  
320 these four gp120 expressing stable CHO GD44 cells were grown in 50L bioreactors and  
321 the cell culture supernatants were collected after 8-10 days of fermentation and purified  
322 through a down stream purification process including the anion-exchange, cation-  
323 exchange and size-exclusion steps under the GMP conditions. The purity of each

324 gp120 proteins is at the range of 96-98% based on their release certificates. The same  
325 purified gp120 proteins are currently being tested in a Phase I clinical trial under  
326 HVTN124 protocol at six major US medical centers.

327

### 328 **Detection of occupancy of N-linked glycosylation sites on gp120 proteins**

329 An aliquot of each gp120 protein was buffered to alkaline pH, reduced, alkylated, and  
330 digested with a combination of proteases including Lys-C (Promega), Arg-C (Promega),  
331 Glu-C (Promega), and trypsin (Promega). Following digestion, the proteins were  
332 deglycosylated by Endo-H (Promega) followed by PNGaseF (Glyko®, Prozyme)  
333 treatment in the presence of  $^{18}\text{O}$ -water. The resulting peptides were separated on an  
334 Acclaim PepMap RSLC C18 column (75  $\mu\text{m}$  x 15 cm) and eluted into the nano-  
335 electrospray ion source of an Orbitrap Fusion™ Lumos™ Tribrid™ mass spectrometer  
336 (Thermo Fisher Scientific) with a 240-min linear gradient consisting of 0.8-80%  
337 acetonitrile over 180 min at a flow rate of 200 nL/min. Full MS analysis was conducted  
338 in the Orbitrap and automated MS/MS analysis using collision-induced dissociation was  
339 conducted in the ion trap. Resulting data was analyzed using a combination of  
340 Proteome Discover/Sequest and ProteoIQ/Provalt to generate a 1% false discovery rate  
341 for protein assignments. Site occupancy was calculated using spectral counts assigned  
342 to the  $^{18}\text{O}$ -Asp-containing (PNGaseF-cleaved) and/or HexNAc-modified (EndoH-  
343 cleaved) peptides and their unmodified counterparts. The positivity cut-off for spectral  
344 counts was set at 10% of the spectral count for the most abundant peptide in each  
345 LC/MS run. Peptides with spectral counts below the positivity cut-off were not included  
346 in the analysis.

347

348 **N-linked glycan profiling analysis for gp120 proteins**

349 A 20 µg aliquot of each gp120 sample was denatured by boiling in SDS. Upon cooling,  
350 the SDS was removed by precipitation as its potassium salt. Denatured proteins were  
351 buffered, recombinant peptide-N-glycanase F (PNGaseF) was added, and the mixture  
352 was incubated overnight to release N-linked glycans. The released N-linked glycans  
353 were freed from residual enzyme, deglycosylated protein, and other contaminants by  
354 passage over C18 Sep-pak. The released, purified glycans were permethylated using  
355 methyl iodide (CH<sub>3</sub>I) under basic conditions in an aprotic solvent (DMSO), followed by  
356 recovery through organic extraction. For mass spectrometric analysis, one-half of the  
357 total permethylated glycans released from 20 µg of protein (10 µg equivalent of protein)  
358 was supplemented by the addition of 10 pmol of an exogenous glycan standard  
359 (maltotetraose) that had previously been permethylated with isotopically heavy  
360 methyl iodide (<sup>13</sup>CH<sub>3</sub>I). The sample glycans spiked with standard were directly infused  
361 into an LTQ/Orbitrap mass spectrometer fitted with a nanospray ionization interface  
362 (NSI-MSn; Orbitrap Discovery, Thermo-Fisher). Glycans were detected in full MS mode  
363 and by total ion mapping (TIM), in which automated CID is performed on small,  
364 overlapping m/z windows. TIM allows the unbiased detection of ions that give  
365 fragmentation patterns consistent with glycan structural topologies (23). Glycan signal  
366 intensities were recovered from extracted full MS spectra (Xtract, Thermo-Fisher) and  
367 glycan identities were assigned based on exact mass and CID fragmentation.  
368 Graphical representations of monosaccharide residues are presented in accordance  
369 with the broadly accepted Symbolic Nomenclature For Glycans, SNFG, and glycan

370 analysis was performed in keeping with the MIRAGE guidelines for glycomic studies  
371 (24, 25).

372

### 373 **IgG-CD4 protein and gp120-specific monoclonal antibodies (mAbs)**

374 The IgG-CD4 protein used in the Octet Qke assays was a fusion protein of human CD4  
375 domains 1 and 2 with human IgG1 Fc at the C-terminus produced by transient  
376 transfection of 293F cells and His-tagged purification. The gp120 CD4 binding site  
377 (CD4bs) monoclonal antibody VRC01 (26) was produced from transiently transfected  
378 293F cells using the molecular clones coding for VRC01 heavy and light chains  
379 obtained from NIH AIDS Reagent Program and purified with a Protein A column. The  
380 gp120 glycan-specific human mAb 2G12 (27) was purchased from Polymun. The  
381 gp120 glycan specific mAb PGT128 (18) was provided by Wayne Koff from International  
382 AIDS Vaccine Initiative (IAVI). The gp120-V2 specific human mAb 2158 (28) was  
383 purchased from Dr. Susan Zolla-Pazner's lab at Mount Sinai School of Medicine. The  
384 gp120 V3 and C4 specific rabbit mAbs R16 and R53 as reported (29, 30) were  
385 produced from transiently transfected 293F cells using paired heavy and light chain  
386 molecular clones and purified using a Protein A column. The IgG-CD4 protein and  
387 monoclonal antibodies produced by this study were verified before use.

388

### 389 **Antigenicity analysis of gp120 proteins with Octet Qke**

390 The antigenicity of gp120 proteins was tested using IgG-CD4 and gp120-specific  
391 monoclonal antibodies by Octet Qke (ForteBio) based on biolayer interferometry. The  
392 IgG-CD4 or each gp120-specific human mAb was individually loaded on to Protein G  
393 sensors at 20 µg/mL and the individual gp120-specific rabbit mAb was loaded to Protein

394 A sensors at 10  $\mu\text{g/mL}$  diluted in ForteBio kinetics buffer. After capture, tips were  
395 washed in kinetics buffer and a baseline measurement recorded. The tips were then  
396 incubated in wells containing serial dilutions of individual gp120 protein (600nM-0.4nM)  
397 to measure the association rate ( $K_{on}$ ) and dissociation rate ( $K_{off}$ ). The antibody binding  
398 kinetics and  $K_D$  values ( $K_{off}/K_{on}$ ) were collected by the ForteBio Data Analysis  
399 software package v7.1 using a 1:1 fitting model for IgG-CD4 and mAbs VRC01,  
400 PGT128, 2G12, 2158 and R16, and using a 2:1 fitting model for mAb R53.

401

402

403

404 **Acknowledgements**

405 This study was supported in part by NIH grants U19AI082676, P01AI082274,  
406 R01AI065250, R21/R33AI087191, U19AI09646, and R01AI39290, as well as by the  
407 grant OPP1033112 from the Bill and Melinda Gates Foundation. The funders had no  
408 role in study design, data collection and interpretation, or the decision to submit the  
409 work for publication.

410 **References**

411

- 412 1. Reitter JN, Means RE, Desrosiers RC. 1998. A role for carbohydrates in immune  
413 evasion in AIDS. *Nat Med* 4:679–684.
- 414 2. Mouquet H, Scharf L, Euler Z, Liu Y, Eden C, Scheid JF, Halper-Stromberg A,  
415 Gnanapragasam PNP, Spencer DIR, Seaman MS, Schuitemaker H, Feizi T,  
416 Nussenzweig MC, Bjorkman PJ. 2012. Complex-type N-glycan recognition by  
417 potent broadly neutralizing HIV antibodies. *Proc Natl Acad Sci U S A* 109:E3268–  
418 E3277.
- 419 3. Walker LM, Phogat SK, Chan-Hui P-Y, Wagner D, Phung P, Goss JL, Wrin T,  
420 Simek MD, Fling S, Mitcham JL, Lehrman JK, Priddy FH, Olsen OA, Frey SM,  
421 Hammond PW, Protocol G Principal Investigators, Kaminsky S, Zamb T, Moyle M,  
422 Koff WC, Poignard P, Burton DR. 2009. Broad and potent neutralizing antibodies  
423 from an African donor reveal a new HIV-1 vaccine target. *Science* 326:285–289.
- 424 4. Bonomelli C, Doores KJ, Dunlop DC, Thaney V, Dwek RA, Burton DR, Crispin M,  
425 Scanlan CN. 2011. The glycan shield of HIV is predominantly oligomannose  
426 independently of production system or viral clade. *PLoS One* 6:e23521.
- 427 5. Doores KJ, Bonomelli C, Harvey DJ, Vasiljevic S, Dwek RA, Burton DR, Crispin M,  
428 Scanlan CN. 2010. Envelope glycans of immunodeficiency virions are almost  
429 entirely oligomannose antigens. *Proc Natl Acad Sci U S A* 107:13800–13805.

- 430 6. Pritchard LK, Spencer DIR, Royle L, Bonomelli C, Seabright GE, Behrens A-J, Kulp  
431 DW, Menis S, Krumm SA, Dunlop DC, Crispin DJ, Bowden TA, Scanlan CN, Ward  
432 AB, Schief WR, Doores KJ, Crispin M. 2015. Glycan clustering stabilizes the  
433 mannose patch of HIV-1 and preserves vulnerability to broadly neutralizing  
434 antibodies. *Nat Commun* 6:7479.
- 435 7. Stanley P, Taniguchi N, Aebi M. 2015. N-Glycans, p. . *In* Varki, A, Cummings, RD,  
436 Esko, JD, Stanley, P, Hart, GW, Aebi, M, Darvill, AG, Kinoshita, T, Packer, NH,  
437 Prestegard, JH, Schnaar, RL, Seeberger, PH (eds.), *Essentials of Glycobiology*,  
438 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor (NY).
- 439 8. Go EP, Irungu J, Zhang Y, Dalpathado DS, Liao H-X, Sutherland LL, Alam SM,  
440 Haynes BF, Desaire H. 2008. Glycosylation site-specific analysis of HIV envelope  
441 proteins (JR-FL and CON-S) reveals major differences in glycosylation site  
442 occupancy, glycoform profiles, and antigenic epitopes' accessibility. *J Proteome*  
443 *Res* 7:1660–1674.
- 444 9. Go EP, Chang Q, Liao H-X, Sutherland LL, Alam SM, Haynes BF, Desaire H.  
445 2009. Glycosylation site-specific analysis of clade C HIV-1 envelope proteins. *J*  
446 *Proteome Res* 8:4231–4242.
- 447 10. Go EP, Herschhorn A, Gu C, Castillo-Menendez L, Zhang S, Mao Y, Chen H, Ding  
448 H, Wakefield JK, Hua D, Liao H-X, Kappes JC, Sodroski J, Desaire H. 2015.  
449 Comparative Analysis of the Glycosylation Profiles of Membrane-Anchored HIV-1  
450 Envelope Glycoprotein Trimers and Soluble gp140. *J Virol* 89:8245–8257.



- 451 11. Go EP, Liao H-X, Alam SM, Hua D, Haynes BF, Desaire H. 2013. Characterization  
452 of host-cell line specific glycosylation profiles of early transmitted/founder HIV-1  
453 gp120 envelope proteins. *J Proteome Res* 12:1223–1234.
- 454 12. Pritchard LK, Vasiljevic S, Ozorowski G, Seabright GE, Cupo A, Ringe R, Kim HJ,  
455 Sanders RW, Doores KJ, Burton DR, Wilson IA, Ward AB, Moore JP, Crispin M.  
456 2015. Structural Constraints Determine the Glycosylation of HIV-1 Envelope  
457 Trimers. *Cell Rep* 11:1604–1613.
- 458 13. Rerks-Ngarm S, Pitisuttithum P, Nitayaphan S, Kaewkungwal J, Chiu J, Paris R,  
459 Prensri N, Namwat C, de Souza M, Adams E, Benenson M, Gurunathan S,  
460 Tartaglia J, McNeil JG, Francis DP, Stablein D, Birx DL, Chunsuttiwat S,  
461 Khamboonruang C, Thongcharoen P, Robb ML, Michael NL, Kunasol P, Kim JH.  
462 2009. Vaccination with ALVAC and AIDSVAX to Prevent HIV-1 Infection in  
463 Thailand. *N Engl J Med* 361:2209–20.
- 464 14. Wen Y, Trinh HV, Linton CE, Tani C, Norais N, Martinez-Guzman D, Ramesh P,  
465 Sun Y, Situ F, Karaca-Griffin S, Hamlin C, Onkar S, Tian S, Hilt S, Malyala P,  
466 Lodaya R, Li N, Otten G, Palladino G, Friedrich K, Aggarwal Y, LaBranche C, Duffy  
467 R, Shen X, Tomaras GD, Montefiori DC, Fulp W, Gottardo R, Burke B, Ulmer JB,  
468 Zolla-Pazner S, Liao H-X, Haynes BF, Michael NL, Kim JH, Rao M, O'Connell RJ,  
469 Carfi A, Barnett SW. 2018. Generation and characterization of a bivalent protein  
470 boost for future clinical trials: HIV-1 subtypes CR01\_AE and B gp120 antigens with  
471 a potent adjuvant. *PloS One* 13:e0194266.

- 472 15. Zambonelli C, Dey AK, Hilt S, Stephenson S, Go EP, Clark DF, Winger M,  
473 Labranche C, Montefiori D, Liao H-X, Swanstrom RI, Desaire H, Haynes BF, Carfi  
474 A, Barnett SW. 2016. Generation and Characterization of a Bivalent HIV-1 Subtype  
475 C gp120 Protein Boost for Proof-of-Concept HIV Vaccine Efficacy Trials in  
476 Southern Africa. *PloS One* 11:e0157391.
- 477 16. Wang S, Chou T-H, Hackett A, Efros V, Wang Y, Han D, Wallace A, Chen Y, Hu G,  
478 Liu S, Clapham P, Arthos J, Montefiori D, Lu S. 2017. Screening of primary gp120  
479 immunogens to formulate the next generation polyvalent DNA prime-protein boost  
480 HIV-1 vaccines. *Hum Vaccines Immunother* 13:2996–3009.
- 481 17. Sanders RW, Venturi M, Schiffner L, Kalyanaraman R, Katinger H, Lloyd KO,  
482 Kwong PD, Moore JP. 2002. The Mannose-Dependent Epitope for Neutralizing  
483 Antibody 2G12 on Human Immunodeficiency Virus Type 1 Glycoprotein gp120. *J*  
484 *Virol* 76:7293–7305.
- 485 18. Pejchal R, Doores KJ, Walker LM, Khayat R, Huang P-S, Wang S-K, Stanfield RL,  
486 Julien J-P, Ramos A, Crispin M, Depetris R, Katpally U, Marozsan A, Cupo A,  
487 Maloveste S, Liu Y, McBride R, Ito Y, Sanders RW, Ogohara C, Paulson JC, Feizi  
488 T, Scanlan CN, Wong C-H, Moore JP, Olson WC, Ward AB, Pognard P, Schief  
489 WR, Burton DR, Wilson IA. 2011. A potent and broad neutralizing antibody  
490 recognizes and penetrates the HIV glycan shield. *Science* 334:1097–1103.
- 491 19. Struwe WB, Chertova E, Allen JD, Seabright GE, Watanabe Y, Harvey DJ, Medina-  
492 Ramirez M, Roser JD, Smith R, Westcott D, Keele BF, Bess JW, Sanders RW,  
493 Lifson JD, Moore JP, Crispin M. 2018. Site-Specific Glycosylation of Virion-Derived

- 494 HIV-1 Env Is Mimicked by a Soluble Trimeric Immunogen. *Cell Rep* 24:1958-  
495 1966.e5.
- 496 20. Vaine M, Wang S, Hackett A, Arthos J, Lu S. 2010. Antibody responses elicited  
497 through homologous or heterologous prime-boost DNA and protein vaccinations  
498 differ in functional activity and avidity. *Vaccine* 28:2999–3007.
- 499 21. Wang S, Pal R, Mascola JR, Chou T-HW, Mboudjeka I, Shen S, Liu Q, Whitney S,  
500 Keen T, Nair BC, Kalyanaraman VS, Markham P, Lu S. 2006. Polyvalent HIV-1  
501 Env vaccine formulations delivered by the DNA priming plus protein boosting  
502 approach are effective in generating neutralizing antibodies against primary human  
503 immunodeficiency virus type 1 isolates from subtypes A, B, C, D and E. *Virology*  
504 350:34–47.
- 505 22. Wang S, Kishko M, Wan S, Wang Y, Brewster F, Gray GE, Violari A, Sullivan JL,  
506 Somasundaran M, Luzuriaga K, Lu S. 2012. Pilot study on the immunogenicity of  
507 paired Env immunogens from mother-to-child transmitted HIV-1 isolates. *Hum*  
508 *Vaccines Immunother* 8:1638–1647.
- 509 23. Aoki K, Perlman M, Lim J-M, Cantu R, Wells L, Tiemeyer M. 2007. Dynamic  
510 developmental elaboration of N-linked glycan complexity in the *Drosophila*  
511 *melanogaster* embryo. *J Biol Chem* 282:9127–9142.
- 512 24. Varki A, Cummings RD, Aebi M, Packer NH, Seeberger PH, Esko JD, Stanley P,  
513 Hart G, Darvill A, Kinoshita T, Prestegard JJ, Schnaar RL, Freeze HH, Marth JD,  
514 Bertozzi CR, Etzler ME, Frank M, Vliegenthart JF, Lütke T, Perez S, Bolton E,

- 515 Rudd P, Paulson J, Kanehisa M, Toukach P, Aoki-Kinoshita KF, Dell A, Narimatsu  
516 H, York W, Taniguchi N, Kornfeld S. 2015. Symbol Nomenclature for Graphical  
517 Representations of Glycans. *Glycobiology* 25:1323–1324.
- 518 25. York WS, Agravat S, Aoki-Kinoshita KF, McBride R, Campbell MP, Costello CE,  
519 Dell A, Feizi T, Haslam SM, Karlsson N, Khoo K-H, Kolarich D, Liu Y, Novotny M,  
520 Packer NH, Paulson JC, Rapp E, Ranzinger R, Rudd PM, Smith DF, Struwe WB,  
521 Tiemeyer M, Wells L, Zaia J, Kettner C. 2014. MIRAGE: the minimum information  
522 required for a glycomics experiment. *Glycobiology* 24:402–406.
- 523 26. Wu X, Yang Z-Y, Li Y, Hogerkorp C-M, Schief WR, Seaman MS, Zhou T, Schmidt  
524 SD, Wu L, Xu L, Longo NS, McKee K, O'Dell S, Louder MK, Wycuff DL, Feng Y,  
525 Nason M, Doria-Rose N, Connors M, Kwong PD, Roederer M, Wyatt RT, Nabel  
526 GJ, Mascola JR. 2010. Rational design of envelope identifies broadly neutralizing  
527 human monoclonal antibodies to HIV-1. *Science* 329:856–861.
- 528 27. Trkola A, Purtscher M, Muster T, Ballaun C, Buchacher A, Sullivan N, Srinivasan K,  
529 Sodroski J, Moore JP, Katinger H. 1996. Human monoclonal antibody 2G12  
530 defines a distinctive neutralization epitope on the gp120 glycoprotein of human  
531 immunodeficiency virus type 1. *J Virol* 70:1100–1108.
- 532 28. Pinter A, Honnen WJ, He Y, Gorny MK, Zolla-Pazner S, Kayman SC. 2004. The  
533 V1/V2 Domain of gp120 Is a Global Regulator of the Sensitivity of Primary Human  
534 Immunodeficiency Virus Type 1 Isolates to Neutralization by Antibodies Commonly  
535 Induced upon Infection. *J Virol* 78:5205–5215.

- 536 29. Chen Y, Vaine M, Wallace A, Han D, Wan S, Seaman MS, Montefiori D, Wang S,  
537 Lu S. 2013. A novel rabbit monoclonal antibody platform to dissect the diverse  
538 repertoire of antibody epitopes for HIV-1 Env immunogen design. *J Virol* 87:10232–  
539 10243.
- 540 30. Pan R, Chen Y, Vaine M, Hu G, Wang S, Lu S, Kong X-P. 2015. Structural analysis  
541 of a novel rabbit monoclonal antibody R53 targeting an epitope in HIV-1 gp120 C4  
542 region critical for receptor and co-receptor binding. *Emerg Microbes Infect* 4:e44.
- 543
- 544

545

546

**Figure legends**

547

548 Figure 1. A. Sequences of the gp120 proteins belonging to clade A (92UG037.8), B  
549 (JRFL), C (93MW965.26), and AE (consensus) aligned to the reference strain HXB2.

550 Identical amino acids are shown as dots, gaps are indicated with dashes, numbers

551 correspond to the HXB2 sequence. Variable regions V1-V5 of gp120 are indicated

552 above the sequences. PNGSs predicted based on the consensus glycosylation

553 sequence are shown in red font and marked with stars above the sequences. B.

554 Summary of glycosylation site distribution among the four gp120 proteins. N indicates

555 the presence of the PNGS in the sequence.

556

557 Figure 2. A. Representative MS spectra of glycans identified on research grade gp120

558 proteins produced in CHO cells (top) and 293F cells (bottom); The gp120 protein from

559 clade B (strain JR-FL) protein is shown here as a representative example. Glycan forms

560 corresponding to the most abundant peaks are shown using the standard glycan icons.

561 Mass peaks corresponding to oligomannose and complex glycan forms are indicated.

562 B. Quantitative comparison of relative proportions of various glycan forms (in %) on

563 CHO- and 293F-derived gp120-B proteins. Glycan structures found at proportions

564 greater than 10% of total glycans are highlighted in grey.

565

566 Figure 3. A. PNGS occupancy analysis of gp120 proteins expressed in 293F cells (top

567 row) and CHO cells (bottom row). The percent of glycan occupancy at each PNGS are

568 shown for each of the gp120 proteins (clades A, B, C and AE) as indicated on the top of  
569 each panel. Green and purple bars indicate the percent of oligomannose glycans and  
570 complex glycans, respectively, and grey bars means the site was not occupied by a  
571 glycan. ND indicates peptides not detected. B. Differences (%) in glycan occupancy  
572 between the 293F- and CHO-produced proteins at each PNGS for each of the gp120  
573 proteins. Percentages for oligomannose and complex glycans were compared at each  
574 site and the difference was plotted based on which protein had larger amount of that  
575 glycan. For example, at position N187 in clade A protein, 293F-derived protein had 25  
576 percentage points more complex glycans than CHO-derived protein, while CHO-derived  
577 protein had 18 percentage points more oligomannose glycans than 293F-derived  
578 protein.

579

580 Figure 4. MS spectra of glycans released from four gp120 proteins (A, B, C and AE)  
581 produced under GMP conditions. Glycan forms corresponding to the most abundant  
582 peaks are shown using the standard glycan icons. Spectra corresponding to  
583 oligomannose and complex glycan forms are highlighted below.

584

585 Figure 5. Quantitation of the types of glycan forms released from two independent lots  
586 (Lot 1 & Lot 2) of gp120 proteins (A, B, C and AE) produced under GMP conditions.  
587 Numbers indicate the percentages of the corresponding glycan forms or structural  
588 classes as color coded for Man5 (M5), Man6 (M6), Man7 (M7), Man8 (M8), Man9 (M9),  
589 Pauci (Man 3 and Man4), Hybrid, and Complex.

590

591 Figure 6. The analysis of sialylation (left panel) and core fucosylation (right panel)  
592 modifications of complex glycans of each gp120 protein (A, B, C or AE) produced from  
593 2 different GMP lots. Only complex glycans were included in the analysis. “1” and “2”  
594 under each bar of the graph indicate lot 1 and lot 2 GMP-grade gp120 proteins,  
595 respectively. “A”, “B”, “C” and “AE” at the bottom of the graph indicate gp120 protein  
596 from clades A, B, C and AE, respectively.

597

598 Figure 7. A. PNGS occupancy analysis of GMP-grade gp120 proteins.

599 The percent of glycan occupancy at each PNGS is shown for each of the gp120  
600 proteins (clades A, B, C and AE) as indicated on the top of each panel. Green and  
601 purple bars indicate the oligomannose glycans and complex glycans, respectively, and  
602 grey bars means the site was not occupied by a glycan. ND indicates peptides not  
603 detected. B. Differences (%) in glycan occupancy in gp120 proteins produced in CHO  
604 cells under research-grade conditions and under GMP conditions at each PNGS. The  
605 plots were generated as described in Figure 3B.

606

607

608



609

**Table 1.** Binding affinity of two GMP lots of gp120 proteins,  $K_d$  (M)

GMP gp120 proteins	gp120 specific mAb						IgG-CD4
	CD4bs	Glycan		V2	V3	C4	
	VRC01	2G12	PGT128	2158	R16	R53	
Clade A Lot 1	6.04E-09	3.11E-10	7.68E-10	6.85E-09	5.59E-10	6.76E-08	5.48E-09
Clade A Lot 2	5.96E-09	1.80E-10	7.62E-10	6.46E-09	2.81E-10	2.76E-08	3.62E-09
Clade B Lot 1	3.11E-08	<1.0E-12	2.46E-09	4.55E-09	3.49E-10	1.16E-08	7.44E-09
Clade B Lot 2	2.83E-08	<1.0E-12	2.10E-09	3.26E-09	4.56E-10	1.32E-08	4.87E-09
Clade C Lot 1	4.53E-08	5.54E-09	3.25E-08	2.61E-08	3.80E-10	5.63E-09	1.57E-07
Clade C Lot 2	5.21E-08	2.24E-09	2.03E-08	3.31E-08	4.27E-10	4.98E-08	1.15E-07
Clade AE Lot 1	5.25E-08	8.02E-09	3.71E-08	3.05E-08	5.87E-10	8.89E-09	2.46E-08
Clade AE Lot 2	3.53E-08	8.62E-09	2.62E-08	3.32E-08	3.22E-10	5.63E-09	1.84E-08

610



Fig 2. Glycoform analysis of clade B gp120 proteins expressed in CHO and 293F cells.




Fig 3. A. PNGS occupancy analysis of gp120 proteins expressed in CHO and 293F cells. B. Differences in glycosylation between 293- and CHO-produced proteins.




Fig 4. Glycan profiles of gp120 proteins produced by GMP manufacturing.




Fig 5. Glycan profiles of gp120 proteins produced by GMP manufacturing.






Fig 6. Sialylation and core fucosylation of GMP grade gp120s.




Fig 7. Glycan occupancy analysis of GMP-grade proteins.

