



Directed topological complexity

Eric Goubault, Aurélien Sagnier, Michael Färber

► To cite this version:

Eric Goubault, Aurélien Sagnier, Michael Färber. Directed topological complexity. Journal of Applied and Computational Topology, Springer, 2019. hal-02434377

HAL Id: hal-02434377

<https://hal.archives-ouvertes.fr/hal-02434377>

Submitted on 10 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Directed topological complexity

Eric Goubault*, Aurélien Sagnier†, Michael Farber‡

December 27, 2018

Abstract

It has been observed that the very important motion planning problem of robotics mathematically speaking boils down to the problem of finding a section to the path-space fibration, raising the notion of topological complexity, as introduced by M. Farber. The above notion fits the motion planning problem of robotics when there are no constraints on the actual control that can be applied to the physical apparatus. In many applications, however, a physical apparatus may have constrained controls, leading to constraints on its potential future dynamics. In this paper we adapt the notion of topological complexity to the case of directed topological spaces, which encompass such controlled systems, and also systems which appear in concurrency theory. We study its first properties, make calculations for some interesting classes of spaces, and show applications to a form of directed homotopy equivalence.

Keywords Directed topology, topological complexity, controlled systems, homotopy theory.

1 Introduction

In this paper we adapt the notion of topological complexity [Farber, 2003], [Farber, 2008], to the case of directed topological spaces. Let us briefly motivate the interest in a notion of “directed” topological complexity. It has been observed that the very important motion planning problem of robotics mathematically speaking boils down to the problem of finding a section to the path-space fibration

$$\chi : X^I \rightarrow X \times X \quad (1)$$

where $\chi(p) = (p(0), p(1))$; here X^I denotes the space of all continuous paths $p : I = [0, 1] \rightarrow X$. If this section can be continuous, then the complexity $\text{TC}(X)$ is the lowest possible (equals to one), otherwise, $\text{TC}(X)$ is defined as the minimal number of “discontinuities” that would encode such a section. The notion of topological complexity is understandable both algorithmically, and topologically, e.g. $\text{TC}(X) = 1$ is equivalent for X to be contractible.

*LIX, Ecole Polytechnique, CNRS, Université Paris-Saclay, 91128 Palaiseau, France, goubault@lix.polytechnique.fr

†LIX & CMAP, Ecole Polytechnique, CNRS, Université Paris-Saclay, 91128 Palaiseau, France, aurelien.sagnier@polytechnique.edu

‡School of Mathematical Sciences, Queen Mary University of London, United Kingdom, m.farber@qmul.ac.uk

Generally speaking, the topological complexity $\mathrm{TC}(X)$ is defined as the Schwartz genus of the path space fibration.

The above definition fits the motion planning problem of robotics when there are no constraints on the actual control that can be applied to the physical apparatus that is supposed to be moved from the state a to the state b . In many applications, however, a physical apparatus may have dynamics that can be described as an ordinary differential equation in the state variables $x \in \mathbb{R}^n$ in time t , and parameterised by the control parameters $u \in \mathbb{R}^p$,

$$\dot{x}(t) = f(t, x(t), u(t)). \quad (2)$$

The control parameters $u(t)$ are usually restricted to lie within a set $u \in U$. Equivalently, as is well-known, one may describe the variety of trajectories of the control system (2) is by using the language of differential inclusions,

$$\dot{x}(t) \in F(t, x(t)), \quad (3)$$

where $F(t, x(t))$ is the set of all $f(t, x(t), u)$ with $u \in U$. Under some well-investigated conditions this differential inclusion can be proven to have solutions, at least locally. Under these conditions, the set of solutions of the differential inclusion (3) naturally forms a *directed space*, compare [Grandis, 2009], see also section 2 below. We observe in this paper that the motion planning problem of robotics in the presence of control constraints equates to finding sections to the analogue of the path space fibration (1), i.e. the map taking a d-path to the pair of its end points¹. This material is developed in the following sections where we work in the generality of directed spaces. In particular we introduce the notion of a directed homotopy equivalence which has precisely, and in a certain non technical sense, minimally, the right properties with respect to the directed version of topological complexity.

2 Definitions

The context of a d-space was introduced in [Grandis, 2009]; we will restrict ourselves later to a more convenient category of d-spaces, that ought to be thought of as some kind of cofibrant replacement of more general (but sometimes pathological) d-spaces.

Definition 1 ([Grandis, 2009]). *A directed topological space, or a d-space $X = (X, PX)$ is a topological space equipped with a set PX of continuous maps $p : I \rightarrow X$ (where $I = [0, 1]$ is the unit segment with the usual topology inherited from \mathbb{R}), called directed paths or d-paths, satisfying three axioms :*

- every constant map $I \rightarrow X$ is directed;
- PX is closed under composition with continuous non-decreasing maps from $I \rightarrow I$;
- PX is closed under concatenation.

Note that for a d-space X , the paths space PX is a topological space, equipped with the compact-open topology.

A map $f : X \rightarrow Y$ between d-spaces is a *d-map* if it is continuous and for any directed path $p \in PX$ the path $f \circ p : I \rightarrow Y$ belongs to PY . In other words we require that f preserves directed paths.

¹That map would most likely not qualify for being called a fibration in the directed setting.

Remark. Given a topological space X equipped with a set D of paths $p : I \rightarrow X$, closed under concatenation and such that the union of the images $p(I)$, for $p \in D$ is X , we call saturation \overline{D} of D the smallest set of paths containing D that forms a d-structure on X . The saturation of D is just made of all composites of path of D with continuous and non-decreasing maps from I to I .

d-spaces in control theory. Consider a differential inclusion

$$\dot{x} \in F(x) \tag{4}$$

where F is a map from \mathbb{R}^n to $\wp(\mathbb{R}^n)$, the set of all subsets of \mathbb{R}^n . A function $x : [0, \infty) \rightarrow \mathbb{R}^n$ is a *solution* of inclusion (4) if x is absolutely continuous and for almost all $t \in \mathbb{R}$ one has $\dot{x}(t) \in F(x(t))$, see [Aubin and Cellina., 1984]. In general, there can be many solutions to a differential inclusion.

Lemma 1. [Aubin and Cellina., 1984] *Suppose a set-valued map $F : \mathbb{R}^n \rightsquigarrow \mathbb{R}^n$ is an upper semicontinuous function of x and such that the set $F(x)$ is closed and convex for all x . Then there exists a solution to Equation (4) defined on an open interval of time.*

Consider a smooth manifold X and an upper semicontinuous set-valued mapping $x \mapsto F(x)$ where for $x \in X$ the image $F(x)$ is a convex cone contained in the tangent space to X at point x , i.e. $F(x) \subset T_x X$. Let PX denote the saturation of the set of all solutions to the differential inclusion $\dot{x} \in F(x)$. Then the pair (X, PX) is a d-space.




Figure 1: The semantics of $Pa.Va.Pb.Vb|Pa.Va.Pb.Vb$.

d-spaces in concurrency and distributed systems theory. The semantics of concurrent and distributed systems can be given in terms of d-spaces, more specifically in terms of geometric realizations [Fajstrup, Goubault, Haucourt, Mimram and Raussen, 2016] of certain pre-cubical sets. As an example, consider the following concurrent program, made of two processes T_1, T_2 , and two binary semaphores a, b , i.e. resources, that can only be accessed locked by one of the two processes [Fajstrup, Goubault, Haucourt, Mimram and Raussen, 2016] at a time : $T_1 = Pa.Va.Pb.Vb, T_2 = Pa.Va.Pb.Vb$, in the notations. This means that process T_1 is locking a (Pa), then relinquishing the lock on a (Va), then locking b (Pb), and finally relinquishing the lock of b (Vb). Process T_2 does the same sequence of actions. The semantics of this concurrent program is depicted in Figure 1 : it is a partially ordered space X , i.e. a topological space with a global order \leq , closed in $X \times X$. Its d-space structure is given by

choosing dipaths to be paths $p : I \rightarrow X$ such that p is non-decreasing. A number of such dipaths are depicted in Figure 1.

The d-paths map. In what follows, we will be particularly concerned with the following map :

Definition 2. Let (X, PX) be a d -space. Define the d -paths map

$$\chi : PX \rightarrow X \times X$$

by $\chi(p) = (p(0), p(1))$ where $p \in PX$.

This map is analogous to the classical path-space fibration (1); the essential distinction is that in the directed setting χ , as defined above, is not necessary a fibration.

Since PX contains only directed paths, the image of χ is a subset of $X \times X$, denoted

$$\Gamma_X = \{(x, y) \in X \times X \mid \exists p \in PX, p(0) = x, p(1) = y\}.$$

In the classical case, one do not need to force the restriction to the image of the path space fibration since the notions of contractibility and path-connectedness are simple enough to be defined separately. In the directed setting, d -contractibility, and “ d -connectedness” are not simple notions and will be defined here through the study of the d -path space map.

Notations: For $a, b \in X$, the symbol $PX(a, b)$ will denote the subspace of PX consisting of all d -paths from the point $a \in X$ to the point $b \in X$. We denote by $*$ the concatenation map

$$PX(a, b) \times PX(b, c) \rightarrow PX(a, c).$$

Note that $PX(a, b)$ is non-empty if and only if $(a, b) \in \Gamma_X$.

Any d -map $f : X \rightarrow Y$ induces continuous maps $\Gamma f : \Gamma_X \rightarrow \Gamma_Y$ and $Pf : PX \rightarrow PY$, such that the diagram

$$\begin{array}{ccc} PX & \xrightarrow{Pf} & PY \\ \downarrow \chi_X & & \downarrow \chi_Y \\ \Gamma_X & \xrightarrow{\Gamma f} & \Gamma_Y \end{array}$$

commutes.

3 Directed topological complexity

Let (X, PX) be a d -space such that X is an Euclidean Neighbourhood Retract (ENR).

Definition 3. The directed topological complexity $\overrightarrow{\mathcal{TC}}(X, PX)$ of a d -space (X, PX) is the minimum number n (or ∞ if no such n exists) such that there exists a map $s : \Gamma_X \rightarrow PX$ (not necessarily continuous) and Γ_X can be partitioned into n ENRs

$$\Gamma_X = F_1 \cup F_2 \cup \dots \cup F_n, \quad F_i \cap F_j = \emptyset, \quad i \neq j,$$

such that

- $\chi \circ s = Id$, i.e. s is a (non-necessarily continuous) section of χ ;
- $s|_{F_i} : F_i \rightarrow PX$ is continuous.

A collection of such ENRs, F_1, \dots, F_n , with n equal to the directed topological complexity of X is called a patchwork.

Example in control theory. As in [Farber, 2008], a motion planner, for the dynamics described by the differential inclusion (4) is a section of the d-paths map produced by the differential inclusion. A section $s : \Gamma_X \rightarrow PX$ associates to any pair of points $(x, y) \in \Gamma_X$ an “admissible” path $s(x, y) = \gamma \in PX$ with $\gamma(0) = x$ and $\gamma(1) = y$.

Example in concurrency and distributed systems theory. Examine again Figure 1; a section of χ is just a scheduler for the actions of the processes T_1 and T_2 .

In the theory of usual (i.e. undirected) topological complexity [Farber, 2003], [Farber, 2008], there are several other equivalent definitions, for example the topological complexity $\text{TC}(X)$ is also the minimal cardinality of the covering of $X \times X$ by open (resp. closed) sets admitting continuous sections; moreover, the book [Farber, 2008] contains four different definitions of $\text{TC}(X)$ leading to the equivalent notions of $\text{TC}(X)$. In the directed case, however, the definitions with open or closed covers lead to notions which can be distinct between themselves as well as distinct from the notion with the ENR partitions given above.

Example 1. Consider the interval $I = [0, 1]$ with the d-structure given by the set of all non-decreasing paths, i.e. $p : [0, 1] \rightarrow [0, 1]$ such that $p(t) \leq p(t')$ for any $t \leq t'$. The space Γ_I is $\{(x, y); x \leq y\}$ and the map $\chi : PI \rightarrow \Gamma_I$ admits a continuous section

$$s(x, y)(t) = (1 - t)x + ty$$

where $t \in [0, 1]$. Hence $\overrightarrow{\text{TC}}(I) = 1$.

Note that in this example the space Γ_I is contractible and the map χ is a fibration with a contractible fibre.

Example 2. Let us consider the directed circle $\overrightarrow{\mathbb{S}^1}$ shown on the figure below:



It is a directed graph homeomorphic to the circle S^1 which is the union of two directed intervals $I_+ \cup I_-$; the d-paths of $\overrightarrow{\mathbb{S}^1}$ are the d-paths lying in one of the intervals I_{\pm} . We see that $P(\overrightarrow{\mathbb{S}^1}) = P(I_+) \cup P(I_-)$ and $P(I_+) \cap P(I_-)$ is a 2-point set containing the two constant paths $p_b(t) \equiv b$ and $p_e(t) \equiv e$. Similarly, one has $\Gamma_{\overrightarrow{\mathbb{S}^1}} = \Gamma_{I_+} \cup \Gamma_{I_-}$ and the intersection $\Gamma_{I_+} \cap \Gamma_{I_-}$ is a 3 point set $\{(b, b), (b, e), (e, e)\}$. Since each of the sets $\Gamma_{I_{\pm}}$ is contractible we obtain that $\Gamma_{\overrightarrow{\mathbb{S}^1}}$ is homotopy equivalent to the wedge $S^1 \vee S^1$.

Next we observe that the map $\chi : P\overrightarrow{\mathbb{S}^1} \rightarrow \Gamma_{\overrightarrow{\mathbb{S}^1}}$ admits no continuous section over any neighbourhood U of the point $(b, e) \in \Gamma_{\overrightarrow{\mathbb{S}^1}}$. To show this one notes that the preimage $\chi^{-1}(b, e)$ has two connected components, one of which consists of the d-paths lying in I_+ and the other of the d-paths lying in I_- . Any open set $U \subset \Gamma_{\overrightarrow{\mathbb{S}^1}}$ containing (b, e) must contain a pair $(x^+, y^+) \in \Gamma_{I_+}$ and a pair $(x^-, y^-) \in \Gamma_{I_-}$, arbitrarily close to (b, e) . Moreover, we may find two sequences $(x_n^{\pm}, y_n^{\pm}) \in \Gamma_{I_{\pm}}$ of points converging to (b, a) and the limits of any section over U along these sequences would land in different connected component of $\chi^{-1}(b, e)$. Hence, we obtain $\overrightarrow{\text{TC}}(\overrightarrow{\mathbb{S}^1}) \geq 2$. On the other hand, we may represent $\Gamma_{\overrightarrow{\mathbb{S}^1}}$ as the union

$$\Gamma_{\overrightarrow{\mathbb{S}^1}} = F_1 \cup F_2$$

where $F_1 = \Gamma_{I_+}$ and $F_2 = \Gamma_{I_-} - \{(b, e), (b, b), (e, e)\}$ and using the previous example we see that over each of the sets F_1, F_2 there exists a continuous section of χ . Hence we obtain

$$\overrightarrow{\text{TC}}(\overrightarrow{\mathbb{S}^1}) = 2. \quad (5)$$

4 Regular d-spaces

Definition 4. A d-space (X, PX) will be called regular if one can find a partition

$$\Gamma_X = F_1 \cup F_2 \cup \cdots \cup F_n, \quad n = \overrightarrow{\text{TC}}(X)$$

onto ENRs such that the map χ admits a continuous section over each F_i and, additionally, the sets $\bigcup_{i=1}^r F_i$ are closed for any $r = 1, \dots, n$.

Note the following property of the sets which appear in Definition 4:

$$\overline{F}_i \cap F_{i'} = \emptyset \quad \text{for } i < i'. \quad (6)$$

In the “undirected” theory of $\text{TC}(X)$ this property is automatically satisfied, see Proposition 4.12 of [Farber, 2008].

All examples of d-spaces which appear in this paper are regular. At present we know of no examples of d-spaces which are not regular; we plan to address this question in more detail elsewhere.

Example 3. The directed circle $\overrightarrow{\mathbb{S}^1}$ is regular as follows from the construction of Example 2.

The Cartesian product of d-spaces (X, PX) and (Y, PY) has a natural d-space structure. Any path $\gamma : [0, 1] \rightarrow X \times Y$ has the form $\gamma(t) = (\gamma_X(t), \gamma_Y(t))$ and we declare γ to be directed if its both coordinates are directed, i.e. $\gamma_X \in PX$ and $\gamma_Y \in PY$. Note that $\Gamma_{X \times Y} = \Gamma_X \times \Gamma_Y$.

Proposition 1. If the d-spaces (X_i, PX_i) are regular, where $i = 1, 2, \dots, k$, then

$$\overrightarrow{\text{TC}}(X_1 \times X_2 \times \cdots \times X_k) - 1 \leq \sum_{i=1}^k \left[\overrightarrow{\text{TC}}(X_i) - 1 \right]. \quad (7)$$

Proof. Denote $\overrightarrow{\text{TC}}(X_i) = n_i + 1$ and let

$$\Gamma_{X_i} = F_0^i \cup F_1^i \cup \cdots \cup F_{n_i}^i$$

be a partition as in the Definition 4, i.e. each set F_j^i is an ENR, the map χ admits a continuous section over F_j^i and each union $F_0^i \cup \cdots \cup F_r^i$ is closed, $r = 0, \dots, n_i$. Denoting $X = \prod_{i=1}^k X_i$ and identifying the space Γ_X with the product $\prod_{i=1}^k \Gamma_{X_i}$, we see that the sets

$$F_{j_1}^1 \times F_{j_2}^2 \times \cdots \times F_{j_k}^k$$

form a ENR partition of Γ_X , where each index j_s runs through $0, 1, \dots, n_s$. The continuous sections $F_{j_s}^s \rightarrow PX_s$, where $s = 1, \dots, k$, obviously produce continuous sections

$$\sigma_{j_1 j_2 \dots j_s} : F_{j_1}^1 \times F_{j_2}^2 \times \cdots \times F_{j_k}^k \rightarrow PX.$$

Consider the sets

$$\bigcup_{j_1+\dots+j_k=j} F_{j_1}^1 \times F_{j_2}^2 \times \dots \times F_{j_k}^k = G_j \subset \Gamma_X, \quad (8)$$

with $j = 0, 1, \dots, N$, where $N = n_1 + n_2 + \dots + n_k$. We observe that the terms of the union (8) are pairwise disjoint and open in G_j (due to (6)) and hence the collection of continuous maps $\sigma_{j_1 j_2 \dots j_s}$ defines a continuous section $G_j \rightarrow PX$. This proves that $\overrightarrow{\text{TC}}(X) \leq N + 1$ as claimed. \square

Corollary 1. *The directed torus $(\overrightarrow{\mathbb{S}^1})^n$ satisfies $\overrightarrow{\text{TC}}((\overrightarrow{\mathbb{S}^1})^n) \leq n + 1$.*

Proof. This follows from Proposition 1 and 3. \square

Definition 5. *We say that a d-space X is strongly connected if $\Gamma_X = X \times X$.*

In other words, in a strongly connected d-space X for any pair (x, y) in $X \times X$ there exists a directed path $\gamma \in PX$ with $\gamma(0) = x$, $\gamma(1) = y$.

Proposition 2. *For any strongly connected d-space X one has $\text{TC}(X) \leq \overrightarrow{\text{TC}}(X)$.*

Proof. Let X be strongly connected and let $\Gamma_X = X \times X = F_1 \cup F_2 \cup \dots \cup F_n$ be a partition into the ENRs as in Definition 3 with $n = \overrightarrow{\text{TC}}(X)$. Then the same partition can serve for the path space fibration $X^I \rightarrow X \times X$ which implies our result. \square

Example 4. Consider the directed loop \mathbb{O}^1 which can be defined as the unit circle

$$S^1 = \{z \in \mathbb{C}; |z| = 1\} \subset \mathbb{C}$$

with the d-structure described below. Any continuous path $\gamma : [0, 1] \rightarrow S^1$ can be presented in the form $\gamma(t) = \exp(i\phi(t))$ where the function $\phi : [0, 1] \rightarrow \mathbb{R}$ is defined uniquely up to adding an integer multiple of $\pm 2\pi$. We declare a path γ to be positive if the function $\phi(t)$ is nondecreasing. It is obvious that the obtained d-space is strongly connected. Hence, using Proposition 2, we obtain $\overrightarrow{\text{TC}}(\mathbb{O}^1) \geq \text{TC}(S^1) = 2$. On the other hand, we can partition $S^1 \times S^1 = F_1 \cup F_2$ where $F_1 = \{(z_1, z_2) \in S^1 \times S^1; z_1 = z_2\}$ and $F_2 = \{(z_1, z_2) \in S^1 \times S^1; z_1 \neq z_2\}$. It is clear that we obtain a section of χ over F_1 by assigning the constant path at z for any pair $(z, z) \in F_1$. A continuous section of χ over F_2 can be defined as follows by moving z_1 along the circle in the positive direction towards z_2 with constant velocity. We conclude that

$$\overrightarrow{\text{TC}}(\mathbb{O}^1) = 2. \quad (9)$$

Besides, we see that the directed loop \mathbb{O}^1 is regular.

Corollary 2. *One has,*

$$\overrightarrow{\text{TC}}((\mathbb{O}^1)^n) = n + 1,$$

i.e. the directed topological complexity of the directed n -dimensional torus $(\mathbb{O}^1)^n$ equals $n + 1$.

Proof. First we apply (9) and Proposition 1 to obtain the inequality $\overrightarrow{\text{TC}}((\mathbb{O}^1)^n) \leq n + 1$. Next we observe that $(\mathbb{O}^1)^n$ is strongly connected and, by Proposition 2, $\overrightarrow{\text{TC}}((\mathbb{O}^1)^n) \geq \text{TC}((S^1)^n) = n + 1$. \square

5 Directed graphs

Let G be a directed connected graph, i.e. each edge of G has a specified orientation. One naturally defines a d-structure on G as follows. Each edge of G can be identified either with the directed interval I (see Example 1) or with the loop \mathbb{O}^1 (see Example 4) and “small directed paths”, i.e. the paths lying on an edge, are the directed paths specified in Example 1 and Example 4. In general, the directed paths of G are concatenations of small directed paths.

For a directed graph G the set Γ_G has the following property: if a pair (x, y) belongs to Γ_G where x is an internal point of an edge e and $y \notin e$ then all pairs (x', y) also belong to Γ_G where $x' \in \text{Int}(e)$.

Proposition 3. $\overline{\text{TC}}(G) \leq 3$.

Proof. Consider the following partition $\Gamma_G = F_1 \cup F_2 \cup F_3$ where

- F_1 is the set of pairs of vertices (α_i, α_j) of G which are in Γ_G ;
- F_2 is the set of pairs $(x, y) \in \Gamma_G$ made of a vertex, and the interior of an arc;
- F_3 is the set of pairs $(x, y) \in \Gamma_G$ with x and y lying in the interiors of arcs.

For each pair of vertices $(\alpha_i, \alpha_j) \in \Gamma_G$ fix a directed path γ_{ij} from α_i to α_j . This defines a section of χ over F_1 . Note that all pairs (α_i, α_i) belong to Γ_G and the path γ_{ii} can be chosen to be constant.

Consider now an oriented edge e and a vertex α_j such that $(x, \alpha_j) \in \Gamma_G$ for an internal point $x \in \text{Int}(e)$. Let α_i be the end point of e and let γ_{x, α_i} denote the constant velocity path along e from x to α_i . A continuous section of χ over $\text{Int}(e) \times \alpha_j$ can be defined as $(x, \alpha_j) \mapsto \gamma_{x, \alpha_i} * \gamma_{ij}$ where $*$ stands for concatenation. A continuous section over $\alpha_j \times \text{Int}(e)$ can be defined similarly, and hence we have a continuous section of χ over F_2 .

Finally we describe a continuous section of χ over F_3 . Consider two oriented edges e and e' where we shall first assume that $e \neq e'$. Let α denote the end point of e and β denote the initial point of e' . We define a section of χ by

$$(x, y) \mapsto \gamma_{x, \alpha} * \gamma_{\alpha \beta} * \gamma_{\beta y}$$

for $x \in \text{Int}(e)$ and $y \in \text{Int}(e')$. Here $\gamma_{x\alpha}$ denotes a constant velocity directed path along e connecting x to α ; the path $\gamma_{\beta y}$ is defined similarly and $\gamma_{\alpha\beta}$ is a positive path from α to β .

Finally we consider the case when $e = e'$. For a pair $(x, y) \in \Gamma_G$ with $x, y \in \text{Int}(e)$ we define the section by $(x, y) \mapsto \gamma_{xy}$ where γ_{xy} is a constant velocity path along e from x to y .

All the partial sections described above over various parts of F_3 obviously combine into a continuous section over F_3 . □

The following example shows that the directed topological complexity can be smaller than the usual complexity.

Example 5. Consider the following graph :



A patchwork for $\Gamma_G : F_1 = \{(b, e)\}$ and $F_2 = \Gamma_G \setminus F_1$. We thus have $\overrightarrow{\text{TC}}(G) = 2$ (here again, it is easy to see that there is no global section). But $\text{TC}(G) = 3$.

However in the special case of strongly connected graphs, the directed and classical topological complexity coincide:

Proposition 4. *Let G be a strongly connected directed graph. Then*

$$\overrightarrow{\text{TC}}(G) = \text{TC}(G) = \min(b_1(G), 2) + 1.$$

Proof. By [Farber, 2008], we know that $\text{TC}(G) = \min(b_1(G), 2) + 1$. As G is strongly connected, we have $\overrightarrow{\text{TC}}(G) \geq \text{TC}(G) = \min(b_1(G), 2) + 1$, see Proposition 2. To prove that we have in fact an equality consider the following cases:

- $b_1(G) = 0$. Since G is contractible and strongly connected, G must be a single point. Then $\overrightarrow{\text{TC}}(G) = 1$ and the result follows.
- $b_1(G) = 1$. It is easy to see that in this case G must be a cycle, i.e. G has n vertices v_1, v_2, \dots, v_n and n oriented edges e_1, e_2, \dots, e_n where e_i connects v_i with v_{i+1} for $i = 1, \dots, n-1$ and e_n connects v_n and v_1 . We see that $\overrightarrow{\text{TC}}(G) = 2$ similarly to Example 4. As we have seen already, $\overrightarrow{\text{TC}}(\overrightarrow{\mathbb{S}^1}) = 2$.
- $b_1(G) \geq 2$. Then $\text{TC}(G) = 3$ (see above) and hence $\overrightarrow{\text{TC}}(G) \geq 3$. On the other hand, $\overrightarrow{\text{TC}}(G) \leq 3$ by Proposition 3. Thus $\overrightarrow{\text{TC}}(G) = 3$.

□

6 Higher-dimensional directed spaces

We begin by recalling the definition of “geometric” precubical sets [Fajstrup, 2005]. The interest [Fajstrup, Goubault, Haucourt, Mimram and Raussen, 2016] of such precubical sets is that the precubical semantics of most programs is a geometric precubical set. Also they are sufficiently tractable for us to compute, in some cases, their directed topological complexity, or more precisely, the directed topological complexity of their directed geometric realization, that we call, cubical complexes (see Definition 7).

Definition 6. *A precubical set C is geometric when it satisfies the following conditions:*

1. no self-intersection: *two distinct iterated faces of a cube in C are distinct*
2. maximal common faces: *two cubes admitting a common face admit a maximal common face.*

Definition 7. *A cubical complex is K is a topological space of the form*

$$K = \left(\bigsqcup_{\lambda \in \Lambda} I^{n_\lambda} \right) / \approx$$

where Λ is a set, $(n_\lambda)_{\lambda \in \Lambda}$ is a family of integers, and \approx is an equivalence relation, such that, writing $p_\lambda : I^{n_\lambda} \rightarrow K$ for the restriction of the quotient map $\bigsqcup_{\lambda \in \Lambda} I^{n_\lambda} \rightarrow K$, we have

1. for every $\lambda \in \Lambda$, the map p_λ is injective,
2. given $\lambda, \mu \in \Lambda$, if $p_\lambda(I^{n_\lambda}) \cap p_\mu(I^{n_\mu}) \neq \emptyset$ then there is an isometry from a face J_λ of I^{n_λ} to a face J_μ of I^{n_μ} such that $p_\lambda(x) = p_\mu(y)$ if and only if $y = h_{\lambda, \mu}(x)$.

As shown in [Goubault and Mimram, 2016] :

Proposition 5. *The realization of a geometric precubical set is a cubical complex.*

Generalising Proposition 3 one may show that

$$\overrightarrow{\text{TC}}(X) \leq 2 \dim(X) + 1$$

for nice cubical complexes X . We shall address this question elsewhere.

6.1 The directed spheres

Let \square^n be the cartesian product of n copies of the unit segment with the d-structure generated by the standard ordering on $[0, 1]$. Its d-space structure is generated by a partially-ordered space [Fajstrup, Goubault, Haucourt, Mimram and Raussen, 2016].

Definition 8. *The directed sphere $\overrightarrow{\mathbb{S}}^n$ of dimension n is defined as the boundary $\partial \square^{n+1}$ of the hypercube \square^{n+1} . Its d-structure is inherited from the one of \square^{n+1} .*

Proposition 6. $\overrightarrow{\text{TC}}(\overrightarrow{\mathbb{S}}^n) = 2$ for any $n \geq 1$.

The case $n = 1$ is covered by Example 2; see [Borat and Grant, 2019] for the general case.

7 Directed homotopy equivalence and topological complexity

As for now, there is no uniquely well-established notion of directed homotopy equivalence between directed spaces, although there has been numeral proposals, among which one linked to our present problem [Goubault, 2017].

We take the view here that directed homotopy equivalences should at least induce equivalent trace categories, viewed with enough structure. We will show in the following sections that directed topological complexity is an invariant of simple equivalences that should be implied by any “reasonable” directed equivalences.

7.1 A simple dihomotopy equivalence, and dcontractibility

In [Goubault, 2017], one of the authors introduced a notion of dihomotopy equivalence. The most important ingredient are that two equivalent d-spaces should be homotopy equivalent in some naive way, and their trace spaces should be homotopy equivalent as well². First, we need to define continuous gradings :

Definition 9. *Let $v, v' \in V$, $q : U \rightarrow V$ be a continuous map, and $W \subseteq U \times U$ be the inverse image by $q \times q$ of (v, v') . Suppose we have a map*

$$h : PV(v, v') \times W \rightarrow PU$$

²In [Goubault, 2017], an extra “bisimulation relation” was added to the definition, that we do not use here.

which is continuous and is such that for all $(u, u') \in W$, $h(p, u, u') \in PU(u, u')$.

In this case, we say that h is continuously graded over W , and by abuse of notation, we write this graded map as a $h : PV \multimap PU$ given by grading $h_{u,u'} : PV(q(u), q(u')) \rightarrow PU(u, u')$, varying continuously for $(u, u') \in W$ in $PU^{PV(v,v')}$, with the compact-open topology.

Any reasonable dihomotopy equivalence should be in particular a d-map inducing a (classical) homotopy equivalence that also induced (classical) homotopy equivalences on the corresponding path spaces. We call this minimum requirement, a simple dihomotopy equivalence :

Definition 10. *Let X and Y be two d-spaces. A simple dihomotopy equivalence between X and Y is a d-map $f : X \rightarrow Y$ such that :*

- *f is a d-homotopy equivalence between X and Y , i.e. a homotopy equivalence with homotopy inverse a d-map $g : Y \rightarrow X$.*
- *There exists a map $F : PY \multimap PX$, continuously graded by $F_{x,x'} : PY(f(x), f(x')) \rightarrow PX(x, x')$ for $(x, x') \in \Gamma_X$, such that $(Pf_{x,x'}, F_{x,x'})$ is a homotopy equivalence³ between $PX(x, x')$ and $PY(f(x), f(x'))$*
- *There exists a map $G : PX \multimap PY$, continuously graded by $G_{y,y'} : PX(g(y), g(y')) \rightarrow PY(y, y')$ for $(y, y') \in \Gamma_Y$ such that $(Pg_{y,y'}, G_{y,y'})$ is a homotopy equivalence between $PY(y, y')$ and $PX(g(y), g(y'))$.*

We sometimes write (f, g, F, G) for the full data associated to the simple dihomotopy equivalence $f : X \rightarrow Y$.

Remark : This definition clearly bears a lot of similarities with Dwyer-Kan weak equivalences in simplicial categories (see e.g. [Bergner, 2004]). The main ingredient of Dwyer-Kan weak equivalences being exactly that Pf induces a homotopy equivalence. But our definition adds continuity and directedness requirements which are instrumental to our theorems and to the classification of the underlying directed geometry.

Example 6. • Let X, Y be two directed spaces. Suppose X and Y are isomorphic as d-spaces i.e. that there exists $f : X \rightarrow Y$ a dmap, which has an inverse, also a dmap. Then X and Y are simply directed homotopy equivalent. The proof goes as follows. Take $f = u$, $g = u^{-1}$, $Pg = F$ the pointwise application of u^{-1} on paths in Y and $Pf = G$ the pointwise application of u on paths in X . This data obviously forms a directed homotopy equivalence.

- The directed unit segment \vec{I} is simply dihomotopically equivalent to a point. Consider the unique map $f : \vec{I} \rightarrow \{*\}$, and $g : \{*\} \rightarrow \vec{I}$ (the inclusion of the point as 0 in \vec{I}). Define $F : P\{*\} \rightarrow P\vec{I}$ by $F(*)$ being the constant map on 0 and $G : P\vec{I} \rightarrow P\{*\}$ to be the unique possible map (since $P\{*\}$ is a singleton $\{*\}$).

As expected, directed topological complexity is an invariant of simple dihomotopy equivalence :

³ Pf is the map on paths which is the natural pointwise extension of f , i.e. $Pf(u)$ is the path $t \rightarrow f(u(t))$ when u is a path in X .

Proposition 7. *Let X and Y be two simply dihomotopically equivalent d -spaces. Then $\overrightarrow{\text{TC}}(X) = \overrightarrow{\text{TC}}(Y)$.*

Proof. As X and Y are dihomotopy equivalent, we have $f : X \rightarrow Y$ and $g : Y \rightarrow X$ dmaps, which form a homotopy equivalence between X and Y . We also get G a continuously graded map from PX to PY , which can be restricted to $G_{y,y'} : PX(g(y), g(y')) \rightarrow PY(y, y')$, inverse modulo homotopy to $Pg_{y,y'}$; and F a continuously graded map from PY to PX such that its restrictions to $PX(x, x')$, for $(x, x') \in \Gamma_X$, $F_{x,x'} : PX(x, x') \rightarrow PY(f(x), f(x'))$ is inverse modulo homotopy to $Pf_{x,x'}$.

Suppose first $k = \overrightarrow{\text{TC}}(X)$. Thus we can write $\Gamma_X = F_1^X \cup \dots \cup F_k^X$ such that we have a map $s : \Gamma_X \rightarrow PX$ with $\chi \circ s = Id$ and $s|_{F_i^X}$ is continuous.

Define $F_i^Y = \{u \in \Gamma_Y \mid g(u) \in F_i^X\}$ (which is either empty or an ENR as F_i^X is ENR and g is continuous) and define $t|_{F_i^Y}(u) = G_u \circ s|_{F_i^X} \circ g(u) \in PY(u)$ for all $u \in F_i^Y \subseteq \Gamma_Y$. This is a continuous map in u since $s|_{F_i^X}$ is continuous, g is continuous, and G is continuous and graded. Therefore $\overrightarrow{\text{TC}}(Y) \leq \overrightarrow{\text{TC}}(X)$.

Conversely, suppose $l : \overrightarrow{\text{TC}}(Y)$, $\Gamma_Y = F_1^Y \cup \dots \cup F_l^Y$ such that we have a map $t : \Gamma_Y \rightarrow PY$ with $\chi \circ t = Id$ and $t|_{F_i^Y}$ is continuous. Now define $F_i^X = \{u \in \Gamma_X \mid f(u) \in F_i^Y\}$ (which is either empty or an ENR as F_i^Y is ENR and f is continuous) and define $s|_{F_i^X}(u) = F_u \circ t|_{F_i^Y} \circ f(u) \in PX(u)$ for all $u \in F_i^X \subseteq \Gamma_X$. This is a continuous map in u since $t|_{F_i^Y}$ is continuous, f is continuous, and F is continuous and graded. Therefore $\overrightarrow{\text{TC}}(X) \leq \overrightarrow{\text{TC}}(Y)$. Hence we conclude that $\overrightarrow{\text{TC}}(X) = \overrightarrow{\text{TC}}(Y)$ and directed topological complexity is an invariant of dihomotopy equivalence. \square

A very simple application is that some spaces must have directed topological complexity of 1 :

Definition 11. *A d -space X is dcontractible if it is dihomotopically equivalent to a point.*

By applying Proposition 7, as the directed topological complexity of a point is 1, all dcontractible spaces have complexity 1, as in the undirected case. Similarly to the undirected case again, although with extra conditions, the converse is also true :

Theorem 1. *Suppose X is a contractible d -space. Then, the dipath space map has a continuous section if and only if X is dcontractible.*

Proof. As X is contractible, we have $f : X \rightarrow \{a_0\}$ (the constant map) and $g : \{a_0\} \rightarrow X$ (the inclusion) which form a (classical) homotopy equivalence. Trivially, f and g are dmaps, and form a d-homotopy equivalence.

Suppose that we have a continuous section s of χ . There is an obvious inclusion map $i : \{s(a, b)\} \rightarrow PX(a, b)$, which is graded in a and b . Define R to be this map. Now the constant map $r : PX(a, b) \rightarrow \{s(a, b)\}$ is a retraction map for i .

We define

$$H : PX \times [0, 1] \rightarrow PX$$

$$(u, t) \rightarrow v \text{ s.t. } \begin{cases} v(x) = u(x) & \text{if } 0 \leq x \leq \frac{t}{2} \\ v(x) = s\left(u\left(\frac{t}{2}\right), u\left(1 - \frac{t}{2}\right)\right) \left(\frac{x - \frac{t}{2}}{1 - t}\right) & \text{if } \frac{t}{2} \leq x \leq 1 - \frac{t}{2} \\ v(x) = u(x) & \text{if } 1 - \frac{t}{2} \leq x \leq 1 \end{cases}$$

($H(u, t)$ is extended by continuity for $t = 1$ as being equal to u)

As concatenation and evaluation are continuous and as s is continuous in both arguments H is continuous in $u \in PX$ and in t . H induces families $H_{a,b} : PX(a, b) \times [0, 1] \rightarrow PX(a, b)$, and because H is continuous in u in the compact-open topology, this family $H_{a,b}$ is continuous in a and b in X .

Finally, we note that $H(u, 1) = u$ and $H(u, 0) = s(u(0), u(1)) = i \circ r(u)$. Hence r is a deformation retraction and $PX(a, b)$ is homotopy equivalent to $\{s(a, b)\}$ and has the homotopy type we expect (is contractible for all a and b), meaning that R is a (graded) homotopy equivalence.

Conversely, suppose X is dcontractible. We have in particular a continuous map $R : \{*\} \rightarrow PX$, which is graded in $(a, b) \in \Gamma_X$. Define $s(a, b) = R_{a,b}(*)$, this is a continuous section of χ . \square

Remark : Sometimes, we do not know right away, in the theorem above, that X is contractible. But instead, there is an initial state in X , i.e. a state a_0 from which every point of X is reachable. Suppose then that, as in the Theorem above, χ has a continuous section $s : \Gamma_X \rightarrow PX$. Consider $s'(a, b) = s^{-1}(a_0, a) * s(a_0, b)$ the concatenation of the inverse dipath, going from a to a_0 , with the dipath going from a_0 to b : this is a continuous path from a to b for all a, b in X . Now, s' is obviously continuous since concatenation, and s , are. By a classical theorem [Farber, 2008], this implies that X is contractible and the rest of the theorem holds.

Example 7. Direct applications of Proposition 7 show that :

- Directed n -tori \mathbb{O}^{1^n} and \mathbb{O}^{1^m} cannot be simply dihomotopically equivalent when $n \neq m$.
- Directed n -tori \mathbb{O}^{1^n} cannot be dihomotopically equivalent to any directed graph for $n \geq 3$.

7.2 Natural homology, and dcontractibility

We now come to make a first connection between some invariants that have been introduced in directed topology (see e.g. [Dubut, Goubault and Goubault-Larrecq, 2015]), like natural homology, [Dubut, Goubault and Goubault-Larrecq, 2016].

We first recap the construction of such invariants.

A *monotonic reparametrization* r is a monotonic continuous surjection from $[0, 1]$ to $[0, 1]$.

Let X be a pospace, i.e. a topological space together with a closed order $\leq \subseteq X \times X$. X is then a particular d-space with the directed paths being the continuous and increasing maps from the unit segment, with the standard ordering, to X .

Let now p and q two dipaths from a to b in X . We say that p is *reparametrized in* q if there exists a monotonic reparametrization γ such that $p \circ \gamma = q$. The *trace* of p , written $\langle p \rangle$ is the equivalence class modulo monotonic reparametrization.

Now we can put together all dipaths from point a to point b , modulo monotonic reparametrization in a topological space:

Let X be a pospace and a and $b \in X$. We topologize the set of traces of dipaths from a to b , with the compact-open topology. Its quotient $\overline{\mathfrak{T}}(X)(a, b)$ by reparametrization, with the quotient topology is called the *trace space in X from a to b* (see [Raussen, 2009]).

Definition 12. We define \mathcal{T}_X to be the category whose:

- objects are traces of X
- morphisms (also called extensions) from $\langle p \rangle$ to $\langle q \rangle$ with p , a dipath from x to y and q , one from x' to y' are pairs of traces $(\langle \alpha \rangle, \langle \beta \rangle)$ such that $\langle q \rangle = \langle \alpha \star p \star \beta \rangle$

We then define $\overrightarrow{T}_*(X) : \mathcal{T}_X \rightarrow \mathbf{Top}_*$ which maps:


- every trace $\langle p \rangle$ with p from x to y to the pointed space $(\overrightarrow{\mathcal{T}}(X)(x, y), \langle p \rangle)$
- every extension $(\langle \alpha \rangle, \langle \beta \rangle)$ with α dipath from x' to x and β dipath from y to y' to the continuous map $\langle \alpha \star _ \star \beta \rangle : \overrightarrow{\mathcal{T}}(X)(x, y) \rightarrow \overrightarrow{\mathcal{T}}(X)(x', y')$ which maps $\langle p \rangle$ to $\langle \alpha \star p \star \beta \rangle$.

We can now define the natural homology functors :

Definition 13 (Natural homology). We define for $n \geq 1$, $\overrightarrow{H}_n(X) : \mathcal{T}_X \rightarrow \mathcal{M}$ (where \mathcal{M} is **Ab**) composing $\overrightarrow{T}_*(X)$ with the $(n-1)^{th}$ homology group functor H_{n-1} .

Remark. \mathcal{T}_X is actually the category of factorization (or twisted arrow category) of the category whose objects are points of X and morphisms are traces and this makes $\overrightarrow{T}_*(X)$ into a natural system in the sense of [Baues and Wirsching, 1985].


Example 8. (taken from [Dubut, Goubault and Goubault-Larrecq, 2016]) We consider the pospace $\overrightarrow{\mathbb{S}^1}$ again, which is made up of two directed segments a and b where their initial points are identified, and their final points are identified too. In the following picture, we distinguish two particular points x and y on a , with $x < y$ (respectively x' and y' on b , with $x' < y'$), which we will use to describe the category of factorization \mathcal{T}_{a+b} as well as the natural homology $\overrightarrow{H}_n(a+b)$.




The description of \mathcal{T}_{a+b} is now as follows. Objects of \mathcal{T}_{a+b} are dipaths, which can be either:

- constant dipaths, $0, x, y, x', y', 1$, for all points x, y, x', y' that we chose to distinguish in the picture of $a+b$.
- non constant and non maximal dipaths of the form $[0, x], [x, y], [y, 1]$ etc.
- maximal dipaths a and b

We chose below to draw a picture of a subcategory of \mathcal{T}_{a+b} , where x, y, x' and y' are any distinguished points of a and b as discussed before. The extension morphisms in \mathcal{T}_{a+b} are pictured below as arrows ; for instance, there is an extension morphism from dipath $[x, y]$ to $[0, y]$ and to $[x, 1]$, among other extension morphisms:



Now, we can picture a subdiagram of $\vec{H}_1(a+b)$, by applying the homology functor on the trace spaces from the starting point to the end point of the dipaths, objects of \mathcal{T}_{a+b} . For instance, the trace space $\vec{\mathfrak{X}}(a+b)(x,y)$ (respectively $\vec{\mathfrak{X}}(a+b)(0,y)$) corresponding to dipath $[x,y]$ (respectively $[0,y]$) in the diagram above, is just a point, hence has zeroth homology group equal to \mathbb{Z} (respectively \mathbb{Z}). All other zeroth homology groups are trivial with the exception of the ones corresponding to the two maximal dipaths (up to reparametrization) a and b , going from 0 to 1. In that case, $\vec{\mathfrak{X}}(a+b)(0,1)$ is composed of two points, that we can identify with a and b , and has \mathbb{Z}^2 (or $\mathbb{Z}[a,b]$ with the identification we just made) as zeroth homology. Now the extension morphism from $[0,y]$ to a induces a map in homology which maps the only generator of $H_0(\vec{\mathfrak{X}}(a+b)(0,y))$ to generator a in $\mathbb{Z}[a,b]$ as indicated in the picture below:



We now define bisimulation as in [Dubut, Goubault and Goubault-Larrecq, 2015]. A bisimulation between functor categories into Abelian groups $P : F \rightarrow Ab$, and $Q : G \rightarrow Ab$ is a “relation” labelled with such isomorphisms of Abelian groups, i.e. is a set of triples

$$(\sigma, \eta, \tau)$$

which is hereditary in the following sense :

- for all $\langle \alpha, \beta \rangle \in F$ from x to x' , if $(x, \eta, y) \in R$, there exists $\langle \gamma, \delta \rangle \in G$ from y to y' such that $(x', \eta', y') \in R$ and such that the following diagram commutes :

$$\begin{array}{ccc}
 P(x) & \xrightarrow{\eta} & Q(y) \\
 \langle \alpha, \beta \rangle \downarrow & & \downarrow \langle \gamma, \delta \rangle \\
 P(x') & \xrightarrow{\eta} & Q(y')
 \end{array}$$

- for all $\langle \gamma, \delta \rangle \in G$ from y to y' , if $(x, \eta, y) \in R$, there exists $\langle \alpha, \beta \rangle \in F$ from x to x' such that $(x', \eta', y') \in R$ and such that the following diagram, as above, commutes up to homotopy

$$\begin{array}{ccc}
P(x) & \xrightarrow{\eta} & Q(y) \\
\langle \alpha, \beta \rangle \downarrow & & \downarrow \langle \gamma, \delta \rangle \\
P(x') & \xrightarrow{\eta'} & Q(y')
\end{array}$$

The main connection with directed topological complexity is as follows :

Proposition 8. *Let X be a d -space. X has directed topological complexity of one (i.e. is dicontractible) implies that its natural homologies $\vec{H}_n(X)$ are all bisimulation equivalent to either, $1_{\mathbb{Z}} : \mathbf{1} \rightarrow \mathbb{Z}$ for $n = 1$, or to $1_0 : \mathbf{1} \rightarrow 0$ for $n > 1$, defined as :*

- $\mathbf{1}$ is the terminal category, with one object 1 and one morphism (the identity on 1)
- $1_{\mathbb{Z}}(1) = \mathbb{Z}$, $1_0(1) = 0$.

Proof. Suppose that X has directed topological complexity of 1. Then by Theorem 1, all trace spaces $\vec{\mathfrak{T}}(X)(x, y)$ are contractible, for all $(x, y) \in \Gamma_X$, hence $\vec{H}_1(X)(x, y) = \mathbb{Z}$ and $\vec{H}_n(X)(x, y) = 0$ for $n > 1$. Therefore the natural homology functors are all constant, either with value \mathbb{Z} or with value 0, and it is a simple exercise to see that the relation between \mathcal{T}_X and $\mathbf{1}$ which relates all objects of \mathcal{T}_X to the only object 1 of $\mathbf{1}$ is hereditary, hence is a bisimulation equivalence. \square

Example 9. We get back to example $\vec{\mathbb{S}}^1$. Its first homology functor was calculated in Example 8 and is not a constant functor (it contains \mathbb{Z}^2 and \mathbb{Z} in its image). Therefore $\vec{\mathbb{S}}^1$ cannot have directed topological complexity of 1. It is also easy to see that the first natural homology functor of \mathbb{O}^1 is $\mathbb{Z}^{\mathbb{N}}$ between two equal points and hence cannot have directed topological complexity of 1.

References

- [Aubin and Cellina., 1984] J.-P. Aubin and A. Cellina. *Differential Inclusions, Set-Valued Maps And Viability Theory*. Number 264 in Grundle. der Math. Wiss. Springer, 1984.
- [Baues and Wirsching, 1985] Hans-Joachim Baues and Gunther Wirsching, Cohomology of small categories, *Journal of Pure and Applied Algebra*, 38(2-3), 187-211 (1985)
- [Bergner, 2004] J. Bergner. A model category structure on the category of simplicial categories. *Trans. Amer. Math. Soc.*, 2004.
- [Borat and Grant, 2019] Ayse Borat and Mark Grant. Directed Topological Complexity of Spheres. To appear in the same volume, 2019.
- [Dubut, Goubault and Goubault-Larrecq, 2015] Jérémy Dubut, Eric Goubault, and Jean Goubault-Larrecq. Natural homology. In *Automata, Languages, and Programming - 42nd International Colloquium, ICALP 2015, Kyoto, Japan, July 6-10, 2015, Proceedings, Part II*, pages 171–183, 2015.

- [Dubut, Goubault and Goubault-Larrecq, 2016] Jérémy Dubut, Éric Goubault, and Jean Goubault-Larrecq. Directed homology theories and Eilenberg-Steenrod axioms. *Applied Categorical Structures*, pages 1–33, 2016.
- [Fajstrup, 2005] Lisbeth Fajstrup. Dipaths and Dihomotopies in a Cubical Complex. *Adv. Appl. Math.*, 35(2), aug, 2005.
- [Farber, 2003] M. Farber. *Topological complexity of motion planning*. *Discrete Comput. Geom.* 29 (2003), no. 2, 211221.
- [Farber, 2008] M. Farber. *Invitation to Topological Robotics*. Zurich lectures in advanced mathematics. European Mathematical Society, 2008.
- [Goubault, 2017] Eric Goubault. On directed homotopy equivalences and a notion of directed topological complexity. CoRR abs/1709.05702, <http://arxiv.org/abs/1709.05702>, 2017.
- [Fajstrup, Goubault, Haucourt, Mimram and Raussen, 2016] Lisbeth Fajstrup, Eric Goubault, Emmanuel Haucourt, Samuel Mimram, and Martin Raussen. *Directed Algebraic Topology and Concurrency*. Springer Publishing Company, 2016.
- [Goubault and Mimram, 2016] Eric Goubault and Samuel Mimram. Directed homotopy in non-positively curved spaces, 2016.
- [Grandis, 2009] Marco Grandis. *Directed Algebraic Topology, Models of non-reversible worlds*. Cambridge University Press, 2009.
- [Raussen, 2009] Martin Raussen, Trace spaces in a pre-cubical complex. *Topology and its Applications*, 156(9), 1718-1728 (2009)