

Antifungal activity and evaluation of phenolics contents of dill *Anethum graveolens* L. extracts original from Algeria

Tifouri Benzina Zeyneb^{1*}, Toumi Benali Fawzia¹, Benyahia Mohamed¹ and Moon Joon-kwan²

¹Laboratoire Ecodéveloppement des espaces, Département des Sciences de l'Environnement, Faculté des Sciences de la Nature et de la vie, Université Djillali Liabès, Sidi Bel Abbès 22000, Algérie

²Laboratory of Pesticide Chemistry and Toxicology, Department of Plant Life and Environmental Sciences, Hankyong National University, Anseong, Republic of Korea

Received 03 October 2018; Revised 14 November 2019

The present study aims to evaluate the antifungal activity of dill *Anethum graveolens* L. against two fungus pathogens of cereals in Algeria, where quantification of the secondary metabolite was investigated in different plant part (whole plant, seeds) and two soil conditions, one is non-salty and second salty soil. The polyphenols, flavonoids, and tannins ranged from 6.25±0.74 mg/g dried weight for the tannins to 167±9.47 mg/g dried weight for polyphenols. The antifungal activity of extracts was tested against *Rhynchosporium secalis* and *Pyrenophora tritici-repentis*. The mycelial growth of the tested fungi was also determined to evaluate the antifungal activity. The ideal concentration showed a significant reduction of the two tested fungi was found to be 10 mg/mL, and the mycelial growth inhibition was determined after 7 days of incubation. Observations on the mycelial growth after extracts treatment decreased growth diameters. The effect of the hexanoic extract on inhibition of *P. tritici-repentis* was 91.17% and the methanolic extract inhibited *R. secalis* at 93.42% as the highest inhibition rate. Thus, the extract of dill could be used to control fungus pathogens as a potential source of eco-friendly fungicide.

Keywords: *Anethum graveolens* L., Antifungal activity, *Pyrenophora tritici-repentis*, *Rhynchosporium secalis*, Secondary metabolite.

IPC Code; Int. cl. (2015.01)-A61K 36/00, A61K 36/23, A61K 125/00, A61K 127/00, A61K 135/00, A01P 3/00

Introduction

Wheat and barley are staple food in Algeria and many other countries. In Algeria, they are susceptible to attack by numerous diseases such as leaf scald and tan spot caused by *Rhynchosporium secalis* in barley¹ and *Pyrenophora tritici-repentis* in wheat² respectively. The losses outcome of these two fungi attack is economically significant³. The use of fungicides to control the fungal attack of the crop has been practised for very long time ago but in recent years the fungicides are used with a small amount reported with a negative impact on the non-target organisms and consumers⁴. However, several environmental and health problems were investigated due to harmful effects (carcinogenicity), severe toxicity, and persistence of the synthetic molecules⁵ where the use of ecofriendly pesticides became a critical board for the world leaders.

Plant extracts aid as antifungal agent⁶ against a wide range of fungi⁷. *Anethum graveolens* L., an

aromatic herb⁸ belonging to the family Apiaceae ex Umbelliferae⁹ is known for the bioactive compounds it contain¹⁰. The quinones seem to be the key substance to impede some fungi¹¹.

Dill has been reported to be used in traditional medicine and its antibacterial activity has been reported in earlier study¹². The antihyperlipidemic and antihypercholesterolemic properties have also been studied earlier¹³. From the beneficial virtues of this plant, heightening of milk production in nursing women and countersigns menstruation in females have been observed¹⁴. Studies have been conducted to study the antifungal activity of dill seed oil against spoilage fungi¹⁵⁻¹⁸. In addition, studies have reported high antifungal activity of the dill essential oil¹⁹⁻²¹.

Due to the lack of studies about the antifungal activity of the crude extract of *A. graveolens*, the present research was undertaken to evaluate the antifungal activity of different dill plant parts which were grown in two soil conditions (salty soil, normal soil) against two fungi strain pathogens that infect wheat and barley.

*Correspondent author

Email: lune03@hotmail.fr, zinebbenzina7@gmail.com

Material and Methods

Plant material

The plant (stem, leaf, and root) at flowering stage and seeds of *A. graveolens* were harvested from the Technical Institute of Vegetable and Industrial Crops (ITCMI) in two different soil conditions, at Sidi Bel Abbès where the soil is normal (non-saline) and Mohammadia with saline soil. Samples were washed, dried at room temperature, ground and stored for future use.

Extraction procedure

The extraction was performed following two methods, by fractionation for the quantification of secondary metabolites and by Soxhlet for the antifungal activity.

Fractionation

The crude methanolic extract was obtained by the method of Benhammou *et al.*²², after evaporation, the residue of each sample was weighed and stored for next step. The methanolic crude extracts were dissolved in boiled water, the flavonoids were extracted by ethyl acetate and n-butanol following the technique of Bekkara *et al.*²³. The tannins extracted by the process described by Zhang *et al.*²⁴, the ethyl acetate extract is used after evaporation.

Soxhlet extraction

The direct Soxhlet extraction was performed by the method of Harboune²⁵, with some modifications, 10 g of each sample was extracted by methanol, ethanol and hexane separately for 3 hours, the extracts were evaporated under vacuum using Büchi R-200 and the residue pondered and stored for further use. The aqueous extracts were obtained by mixing 10 g of powder with distillate water under reflux for 2 hours²⁶.

Secondary metabolites quantification

Total phenols

The total phenolic contents of tannins ethyl acetate extracts and flavonoids n-butanol, ethyl acetate extracts of dill plant (non-salt, salt) and seeds (non-salt, salt) were determined using the Folin-Ciocalteu according to the method described by Singleton and Rossi²⁷.

Total flavonoids

The total flavonoids of the n-butanol and ethyl acetate extracts were determined by the Aluminum chloride assay using a method described by Zhishen *et al.*²⁸.

Total condensed tannins

Vanillin assay was used²⁹ to determine the tannins contents in tannin ethyl acetate extracts of dill plant of Mohammadia and Sidi Bel Abbès as well as seeds.

Antifungal activity

Fungal strain

P. tritici-repentis and *R. secalis* were isolated from the host crop and purified through repeated planting method in Potato Dextrose Agar (PDA) media, the strains were identified according to the morphological characteristics of the colony in the National Institute of Agronomic Research of Algeria (INRAA). The fungal strain cultures were maintained in PDA media at 4 °C.

Antifungal assay principal

Exactly 20 mL of Potato Dextrose Agar (PDA) medium was placed in each Petri Dish, to which a necessary quantity of the *A. graveolens* extract was added to obtain 1.25, 2.5, 5, and 10 mg/mL concentrations of the ethanolic, methanolic, hexanoic and aqueous extract. The effect on the mycelial growth of the tested fungi was determined following the method of Brewer³⁰ and Leach³¹. The mycelial growth was determined with the following equation:

$$L = D - d/2 \quad \dots \text{Eq (1)}$$

where L mycelian growth, D: colony diameter and d the explant diameter.

Sample without any treatment was considered as control. The Petri dishes were then incubated for 7 days at 28 °C. The mycelial growth was calculated regularly from the first day after incubation until the last day of incubation.

The inhibition rate of the fungus colony is calculated according to the method of Kordali *et al.*³².

$$\text{IR}\% = [(D_t - D) / D_t] \times 100 \quad \dots \text{Eq (2)}$$

where IR inhibition rate of the mycelial growth, D_t diameter of mycelial growth of the control and D diameter of mycelial growth of the extract.

Statistical analysis

All the measurements were replicated in a triple for each treatment, and the data reported as mean ± standard deviations. Significant differences between the mean values were determined at ($P < 0.01$), following MANOVA. The statistical analysis was performed using SPSS V20.

Results and Discussion

Total polyphenols, flavonoids, and tannins

The phenolic contents are represented in Table 1, the results were ranked from 62.79±5.98 mg gallic acid equivalent (GAE)/g dry weight to 212.3±4.2 mg/g dry weight for the polyphenols, from 6.11±0.29 to 23.99±0.86 mg catechin equivalent (CE)/g dry weight for the flavonoids and from 6.25±0.74 to 6.73±0.75 mg gallic acid equivalent (GAE)/g dry weight for tannins. Phenols comprise a great variety of compounds, such as polyphenols, flavonoids, condensed tannins³³. The result showed that the ethyl acetate obtained from fractionation of flavonoids contains the highest quantity of phenolics compound followed by the ethyl acetate of tannins, whereas the n-butanol had the lowest value which may be referred to the polarity of the solvent.

Comparing the contents on polyphenols and flavonoids of the plant part revealed that the seeds are significantly richer than the whole plant (stems, flower, and root), however, the study of Yung-Shin Shyu³⁴, showed that dill flower contains more polyphenols and flavonoids than the leaf, seeds and root³⁵ respectively, the flower contents is 196.65 mg/g dried extract, 67.10 mg/g dried extract polyphenols and flavonoids respectively. The fresh herb comprises more flavonoids than the dry samples used in this study, where the HPLC analysis showed that the flavonoid content up to

100 mg/g fresh weight³⁶. The use of bio-fertilizers on dill affect the flavonoids contents³⁷ which confirm that the stress induces more production of bioactive nevertheless the plant and seed coming from salty soil are significantly containing more polyphenols and flavonoids than the plant and seed in normal soil condition and is important to note that the result obtained in this study showed that *A. graveolens* original from Algeria contains more polyphenols and flavonoids than wild dill³⁸. For the tannins, there is no significance between the samples, where the plant, seeds, non-salt and salt conditions contain almost the same quantity of tannins.

Antifungal activity

Mycelial growth

One day after the incubation of the fungus in PDA without extracts, *P. tritici-repentis* was grown 4 mm, whereas, the growth of *R. secalis* was relatively slow by 2 mm, the growth of two fungi in different extracts was shown in the Figs. (1–4).

The results indicated that; the mycelial growth was considerably reduced with increasing concentration of *A. graveolens* extracts while their growth increased with incubation time. From the Figs. (1,2) the mycelial growth of *P. tritici-repentis* by the plant extract was reduced at 5 mm for 10 mg/mL of hexanoic non-salt plant extract, followed by aqueous extract (7 mm), methanolic extract (9 mm) and




Fig. 1 — Mycelial growth of *Pyrenophora tritici-repentis* in dill, non-salt plant(a) and non-salt seed extracts (b).

Table 1 — Phenolics contents in dill plant and seed extracts, and various soluble fraction (mg/g)

Extracts	Polyphenols ^{T,F,E}			Flavonoids ^{T,F}		Tannins ^E
	T	F	E	T	F	E
Non-salt plant	66.66 ± 5.98	121.04 ± 4.31	78.40 ± 5.94	12.33 ± 0.41	8.25 ± 0.60	6.68 ± 0.70
Non-salt seed	142.19 ± 2.5	153.12 ± 6.43	62.79 ± 5.98	7.23 ± 1.17	6.11 ± 0.29	6.58 ± 0.38
Salt plant	76.01 ± 9.63	96.54 ± 9.63	87.79 ± 7.57	7.85 ± 0.12	23.99 ± 0.86	6.25 ± 0.74
Salt seed	212.30 ± 4.20	167.70 ± 9.47	90.56 ± 9.44	6.88 ± 0.18	12.63 ± 0.43	6.73 ± 0.75

T- tannins ethylacetate extract, F-flavonoids n-butanol extract, and E-ethylacetate extract

ethanolic extract (13 mm) respectively. For the salt plant extract the hexanoic extract reduced the mycelial growth at 3 mm by the concentration 10 mg/mL and by the same concentration the aqueous extract stopped the growth at 4 mm, the highest mycelial growth showed by the methanolic extract by 7 mm. the non-salt seed extracts showed the best mycelial reduction by 10 mg/mL of the hexanoic extract followed by the same concentration of aqueous extract, however, the salt seeds extract

reduced the mycelial growth as well as the non-salt seeds extract for the hexanoic extract but reduced more for the ethanolic and methanolic extract (5 and 4.5mm) respectively.

The drop of mycelial growth of *R. secalis* seen in Figs. 3,4. The growth stopped at 3.5 mm for the ethanolic extract and 4.5 mm for the hexanoic and aqueous extract of non-salt plant while it was reduced at 2.5 mm for the methanolic salt plant extract with the same concentration of 10 mg/mL, the non-salt




Fig. 2 — Mycelial growth of *Pyrenophora tritici-repentis* in dill, salt plant (a) and salt seed extracts (b).




Fig. 3 — Mycelial growth of *Rhynchosporium secalis* in dill, non-salt plant (a) and non-salt seed extracts (b).




Fig. 4 — Mycelial growth of *Rhynchosporium secalis* in dill, salt plant (a) and salt seed extracts (b).




Fig. 5 — Inhibition rate of the fungi in different dill extracts of salt and non-salt plant and seeds, *Pyrenophora tritici-repentis* (a) and *Rhynchosporium secalis* (b).

seeds extract showed equal mycelial growth reduction at 10 mg/mL of methanolic, hexanoic and aqueous extract (6.5 mm) while the salt seeds extracts at the same concentration reduce the growth at 2.5 mm for the aqueous extract and at 3 mm for the ethanolic extract.

Inhibition rate

The inhibition rate of *P. tritici-repentis* (Fig. 5a) ranged between 61.76 to 91.17% for the plant samples of different extracts and between 70.58 to 89.70% for seed extracts. Fig. 5b shows the inhibition rate of dill extract on *R. secalis* which fluctuated between 84.21 to 93.10% for the Plant extracts; also, it was recorded between 81.57 to 93.42% for seed extracts.

Recently, world demand for effective, safe, and natural products without synthetic chemicals to control food has been high. Aromatic volatile products of plant secondary metabolism have formed the basis of many applications in food flavouring and preservation industries^{18,39}. In our study, the different extract of *A. graveolens* showed pronounced antifungal efficacy against all the tested fungi. Mycelium growth was recorded to decrease with increasing concentrations of the extract. However, at 10 mg/mL, the fungal colony of *P. tritici-repentis* was significantly reduced by the hexanoic extract of all the samples which confirm the sensibility of the fungus to the essential oil of dill¹⁸. This study reported that; dill seed oil is found to be highly effective for controlling the growth of *A. niger* and *A. flavus* at doses of 6 mL by poison food techniques. On the other hand, *R. secalis* was more affected by the ethanolic extract. The plant and seeds obtained from salty soil condition showed significant antifungal activity on both fungi than the samples from non-salt conditions.

In vitro studies on *A. graveolens* illustrate its potential as an ideal antifungal agent against fungi.

Therefore, the present results clearly demonstrate that the different extract of *A. graveolens* significantly reduces the development of two fungus pathogens for wheat and barley which are a very important crop worldwide. The *in vivo* experiment shows that contact of produce to the vapours may persuade inferior confrontation to fungal challenge⁴⁰. Nevertheless, higher plant extract concentrations reported necessary in the field than in laboratory⁴¹, which may be available for favourable conditions to fungus development.

Conclusion

Wheat and barley are the main food crop in Algeria and many other countries. These crops are susceptible to attack by leaf scald and tan spot caused by *Rhynchosporium secalis* in barley and *Pyrenophora tritici-repentis* in wheat, respectively. The present study showed that dill original from Algeria has antifungal activity against cereals fungus pathogens, however, the saline stress has significantly increased this activity. *Anethum graveolens*, is a rich aromatic plant with the phenolics contents, where the seeds are advised to be more explored, whereas the soil salinity increase significantly the phenolic contents in the plant and seeds. Dill could be one of the best eco-friendly fungicides. Furthermore, a study of the impact of different level of salinity on the production of a secondary metabolite is highly needed.

Conflict of interest

The authors have no conflict to declare.

References

- Williams K, Donnellan S, Smyl C, Scott L and Wallwork H, Molecular variation in *Rhynchosporium secalis* isolates obtained from hotspots, *Australas Plant Pathol*, 2003, **32**(2), 257-262.

- 2 Ameen G, Kariyawasam G, Shi G, Friesen T L, Faris J D, *et al.*, Molecular manipulation of the mating-type system and development of a new approach for characterizing pathogen virulence in *Pyrenophora tritici-repentis*, *Fungal Genet Biol*, 2017, **109**, 16-25.
- 3 Singh D P, *Management of wheat and barley diseases*, Apple academic press, 2018, 5-6.
- 4 Keith J B and Derek W H, Fungicide resistance in crop pathogens: how can it be managed?, 2nd edition, (Fungicide Resistance Action Committee), 2007, 6-7.
- 5 Lingk W, Health risk evaluation of pesticide contamination in drinking water, *Gesunde Pflanz*, 1991, **43**, 21-25.
- 6 Ali-Shtayeh M S and Ghdeib S I A, Antifungal activity of plant extracts against dermatophytes, *Mycoses*, 1999, **42**, 665-672.
- 7 Wilson C L, Solar J M, El Ghaouth A and Wisniewski M E, Rapid evaluation of plant extracts and essential oils for antifungal activity against *Botrytis cinerea*, *Plant Disease*, 1997, **81**(2), 204-210.
- 8 Hekmatzadeh S F, Bazarganipour F, Malekzadeh J, Goodarzi F and Aramesh S, A randomized clinical trial of the efficacy of applying a simple protocol of boiled *Anethumgraveolens* seeds on pain intensity and duration of labor stages, *Complement Ther Med*, 2014, **22**(6), 970-976.
- 9 Karklelienė R, Dambrauskienė E, Juškevičienė D, Radzevičius A, Rubinskienė M, *et al.*, Productivity and nutritional value of dill and parsley, *Hort Sci*, 2014, **41**(3), 131-137.
- 10 Kmiecik W, Lisiewska Z and Slupski J, Relationship between the yield and quality of green dill and the height of plants, *Folia Hort*, 2005, **17**(2), 37-52.
- 11 Maruzzella J C and Liguori L, The *in vitro* antifungal activity of essential oils, *J Am Pharm Assoc*, 1958, **47**(4), 250-254.
- 12 Raffi F and Shahverdi A R, Comparison of essential oils from three plants for enhancement of antimicrobial activity of nitrofurantoin against enterobacteria, *Chemotherapy*, 2007, **53**, 21-25.
- 13 Yazdanparast R and Alavi M, Antihyperlipidaemic and antihypercholesterolaemic effects of *Anethum graveolens* leaves after the removal of furocoumarins, *Cytobios*, 2001, **105**, 185-191.
- 14 Monsefi M, Ghasemi M and Bahaoddini A, The effects of *Anethum graveolens* L. on female reproductive system, *Phytother Res*, 2006, **20**(10), 865-868.
- 15 Elgayyar M, Draughon F A, Golden D A and Mount J R, Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms, *J Food Prot*, 2001, **64**(7), 1019-1024.
- 16 Fatope M O, Marwah R G, Onifade A K, Ochei J E and Al Mahroqi Y K, ¹³C NMR analysis and antifungal and insecticidal activities of Oman dill herb oil, *Pharm Biol*, 2006, **44**(1), 44-49.
- 17 Lopez P, Sanchez C, Batlle R and Nerin C, Solid- and vapor-phase antimicrobial activities of six essential oils: susceptibility of selected foodborne bacterial and fungal strains, *J Agric Food Chem*, 2005, **53**, 6939-6946.
- 18 Singh G, Maurya S, De Lampasona M P and Catalan C, Chemical constituents, antimicrobial investigations, and antioxidative potentials of *Anethum graveolens* L. essential oil and acetone extract, *J Food Sci*, 2005, **70**(4), 208-215.
- 19 Amiri A, Dugas R, Pichot A L and Bompeix G, *In vitro* and *in vivo* activity of eugenol oil (*Eugenia caryophyllata*) against four important postharvest apple pathogens, *Int J Food Microbiol*, 2008, **126**, 13-19.
- 20 Anthony S, Abeywickrama K, Dayananda R, Wijeratnam S W and Arambewela L, Fungal pathogens associated with banana fruit in Sri Lanka, and their treatment with essential oils, *Mycopathologia*, 2004, **157**(1), 91-97.
- 21 Dikbas N, Kotan R, Dadasoglu F and Sahin, F, Control of *Aspergillus flavus* with essential oil and methanol extract of *Satureja hortensis*, *Int J Food Microbiol*, 2008, **124**(2), 179-182.
- 22 Benhammou N, Bekkara F A and Panovska T K, Antioxidant activity of methanolic extracts and some bioactive compounds of *Atriplex halimus*, *CR Chim*, 2009, **12**(12), 1259-1266.
- 23 Bekkara F, Jay M, Viricel M R and Rome S, Distribution of phenolic compounds within seed and seedlings of two *Vicia faba* cvs differing in their seed tannin content, and study of their seed and root phenolic exudations, *Plant Soil*, 1998, **203**(1), 27-36.
- 24 Zhang S Y, Zheng C G, Yan X Y and Tian W X, Low concentration of condensed tannins from catechu significantly inhibits fatty acid synthase and growth of MCF-7 cells, *Biochem Biophys Res Commun*, 2008, **371**(4), 654-658.
- 25 Harborne J B, *Phytochemical methods: A guide to modern techniques of plant analysis*, 5th Edition, Chapman & Hall, London, 1998, 21-72.
- 26 Applebaum S W, Marco S and Birk Y, Saponins as possible factors of resistance of legume seeds to the attack of insects, *J Agric Food Chem*, 1969, **17**(3), 618-622.
- 27 Singleton V L and Rossi J A, Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents, *Am J Enol Viticult*, 1965, **16**(3), 144-158.
- 28 Zhishen J, Mengcheng T and Jianming W, The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals, *Food Chem*, 1999, **64**(4), 555-559.

- 29 Julkunen-Tiitto R, Phenolic constituents in the leaves of northern willows: Methods for the analysis of certain phenolics, *J Agric Food Chem*, 1985, **33**(2), 213-217.
- 30 Brewer D, Studies in *Asochytapiri*, *Can J Bot*, 1960, **38**, 705-717.
- 31 Leach C M, The quantitative and qualitative relationship of ultraviolet and visible radiation to the induction of reproduction in *Asochytapisi*, *Can J Bot*, 1962, **40**(12), 1577-1602.
- 32 Kordali S, Cakir A, Zengin H and Duru M E, Antifungal activities of the leaves of three *Pistacia* species grown in Turkey, *Fitoterapia*, 2003, **74**, 164-167.
- 33 Shahidi F and Naczki M, Phenolics in food and nutraceuticals, edited by CRC Press, 2004, 421-426.
- 34 Yung-Shin S, Jau-Tien L, Yuan-Tsung C, Chia-Jung C and Deng-Jye Y, Evaluation of antioxidant ability of ethanolic extract from dill (*Anethum graveolens* L.) flower, *Food Chem*, 2009, **115**(2), 515-521.
- 35 Saloua K, Lella O A, Ouafae E Y, Amal S, Bahia B, *et al.*, Quantitative study of root the plant (*Anethumgraveolens*) and evaluation of their antioxidant activity, *J Pharm Phytochem*, 2017, **6**(5), 2735-2740.
- 36 Justesen U and Knuthsen P, Composition of flavonoids in fresh herbs and calculation of flavonoids intake by use of herbs in traditional Danish dishes, *Food Chem*, 2001, **73**(2), 245-250.
- 37 Said-Al Ahl H A, Sarhan A M, Dahab A D M A, Zeid E S N A, Ali M S, *et al.*, Flavonoids, essential oil and its constituents of *Anethum graveolens* L. herb affected by nitrogen and Bio-Fertilizers, *Agric Biol Sci J*, 2015, **1**(3), 105-109.
- 38 Ksouri A, Dob T, Belkebir A, Lamari L, Krimat S, *et al.*, Total phenolic, antioxidant, antimicrobial activities and cytotoxicity study of wild *Anethum graveolens* L., *Int J Pharm Phytochem Res*, 2015, **7**(6), 1025-1032.
- 39 Rahman A and Kang S C, *In vitro* control of food-borne and food spoilage bacteria by essential oil and ethanol extracts of *Lonicera japonica* Thunb, *Food Chem*, 2009, **116**(3), 670-675.
- 40 Tzortzakis N G, Impact of cinnamon oil-enrichment on microbial spoilage of fresh produce, *Innov Food Sci Emerg Technol*, 2009, **10**(1), 97-102.
- 41 Farbood M I, MacNeil J H and Ostovar K, Effect of rosemary spice extractive on growth of microorganisms in meats, *J Milk Food Technol*, 1976, **39**(10), 675-679.