

ONLINE FIRST

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 0015-5659

e-ISSN: 1644-3284

Histomorphometrical evaluation of tibial subchondral bone after moderate running in male and female Wistar rats

Authors: M. N. Nazem, M. Sahebozamani, R. Mosavian Naeini, S. M. Sajjadian

DOI: 10.5603/FM.a2020.0030

Article type: ORIGINAL ARTICLES

Submitted: 2019-05-07

Accepted: 2019-11-30

Published online: 2020-03-11

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Folia Morphologica" are listed in PubMed.

Histomorphometrical evaluation of tibial subchondral bone after moderate running in male and female Wistar rats

Running head: Histomorphometrical evaluation of tibial subchondral bone after moderate running

M.N. Nazem, M. Sahebozamani, R. Mosavian Naeini, S.M. Sajjadian

Shahid Bahonar University of Kerman, Kerman, Iran

Address for correspondence: Mohammad Naser, Nazem, Associate Professor of Anatomy, Histology and Embryology, Department of Basic Sciences, School of Veterinary Medicine, Shahid Bahonar University of Kerman, Pazhuhesh Square, Kerman, Iran. P. O. Box: 76169133, tel: +983431322954, fax: +983433257447, e-mail: nnazem@uk.ac.ir; nasernazem@yahoo.com

Abstract

Exercise has been shown to be beneficial to the skeleton, in both humans and animals. This study was done to test the sex-related difference in the risk of OA (osteoarthritis) of the knee joint and also on the subchondral bone after moderate running exercise.

Forty male and female Wistar rats were randomly assigned to four equal groups (2 male and 2 female groups) in a same condition. Ten of each sex were selected as control groups, while running exercises were performed in remained 20 male and female rats using a motor treadmill to motivate rats to run daily distances of 1 km at 5 days/week within six weeks.

On day 43, all animals were sacrificed and the knee articular cartilage and also histomorphometric parameters of subchondral bone and mid shaft of tibia were evaluated. Results showed mild OA in both male and female runner groups. Results in male runner rats were significantly lesser than that in female runners. On the other hand, the difference

in female runner group showed significant changes in comparison with other groups in histomorphometric parameters in tibia.

Obtained results showed that the development of knee OA and subchondral bone changes may be related to the sex differences. Although there was no synovitis in male runners but female runner group showed mild hyperemia dropsy with a moderate synovitis in this region.

Key words: histomorphometry, running, subchondral bone, tibia, rat

INTRODUCTION

According to human (Howe et al., 2015) and animal (Jarvinen et al., 2001) studies, exercise and mechanical loading are beneficial to the skeleton. Despite the obvious effects of high mechanical loading on the skeleton, the capacity of skeleton to maintain the exercise-induced bone gain remains unknown (Seeman, 2002).

Based on multiple studies with experimental models, increased mechanical loading or physical activity triggers bone formation and growth and has positive effects on the skeleton (Tromp et al., 2006; Joo et al., 2003; Iwamoto et al., 2011). Studies show that running exercise increases cortical and cancellous tibial bone mass in aged female rats (Chen et al., 1994).

The articular cartilage health is dependent on the mechanical features of its bony subchondral bed. It has been shown that bone changes are associated with osteoarthritis (OA) (Radin and Rose, 1986). The architecture of subchondral metaphyseal bone is not uniform. The organization of trabecular sheets is in a way that they can transfer stress from the overlying layers of articular cartilage, subchondral plate, and calcified cartilage to metaphyseal cortices (Lee et al., 2011; Kenneth et al., 2005).

In this study, we aimed to evaluate the effects of moderate running exercise on the subchondral bone of tibia in the knee joint, histological assessment and grading of the synovial membrane changes and to conduct a histomorphometric assessment of the mid-shaft of tibia.

MATERIALS AND METHODS

All procedures involving the experimental use of animals were approved by the Animal Ethics Committee, a branch of the Research Council of the Veterinary School in Shahid Bahonar University, Kerman Province, Iran.

Forty adult male and female Wistar rats (9-10 weeks of age) with no significant difference in their body weight (251.59 ± 12.31 g) were randomly divided into four equal groups. All rats were allowed a 5-day adaptation period in a room with controlled conditions (temperature 22–25 °C and humidity 60–70 %) before starting the experiment (Seyyedini and Nazem, 2017). Ten of each sex were selected as male and female control groups while running exercises were performed in remained 20 male and female rats for 6 weeks on a motor-driven rodent treadmill (Model T510, DRI Co., Taiyuan, Taiwan), speed of 20 m/min with inclination of 18° for 60 minutes each day on 5 days of every week.

During the exercise period, all animals were fed a standard diet (Javaneh Khorasan Co., Iran).

On day 43, all control and runner rats were sacrificed by overdose injection of pentobarbital sodium (100mg/100 g body weight, intraperitoneal injection). The body weight was recorded. Then whole of both knee joints were taken and fixed in 10% buffered formaldehyde for 1 day and then was placed in a new solution of 10% buffered formaldehyde for 10 days. Decalcification in 10% formic acid solution was followed by embedding of the complete knee joints in this solution for 50 days. 5µm thickness serial sections of frontal whole left knee joint were prepared in order to obtain the thickest cartilage section from the lateral condyle of the tibia. Obtained thickest sections of each tibia were stained with hematoxylin-eosin method. Sections were evaluated using the Mankin's scheme (Mankin et al., 1971). We only differed safranin-O staining with hematoxylin-eosin method. The histological evaluation system for OA (osteoarthritis) was classified into four categories: Mankin score of 0, no OA; Mankin scores of 1-5, mild OA; scores of 6-10, moderate OA; and scores of 11-14, severe OA (Pap et al., 1998). In order to histological grading of synovial layer changes, Krenn et al's method was used. Based on

their results, 0 to 1 corresponds to no synovitis, 2 to 4 to a slight synovitis, 5 to 6 to a moderate synovitis, and 7 to 9 to a strong synovitis (Krenn et al., 2002).

Histomorphometric analyses were taken on sections using method described by Renner et al. 2006. Thickness from subchondral bone to articular surface was measured at the middle of lateral condyle. In each section, chondrocyte cells were counted within a $120,000 \mu\text{m}^2$ area including both calcified layer and articular surface (Renner et al., 2006).

All right knee joints were opened and examined for gross knee articular surface evaluation. On the other hand, the right tibia was dissected out. The tibia was cut into two equal parts. Thus, the proximal half of tibia was collected. The samples were proceeding similar to the left ones. Serial longitudinal sections were cut with a rotary microtome and mounted on saline-coated slides glass after preparation of histological specimens. Before staining, the sections were deparaffinized in 100% xylene and rehydrated in graded ethanol (Chang et al., 2010). The longitudinal sections of the proximal tibia, between medial and lateral intercondyloid spines, and transvers sections of the mid shaft of tibia were cut at 5- μm thickness with a microtome.

Histomorphometric measurements of the cancellous bone of the proximal tibia and the cortical bone of the tibial shaft were examined using an Olympus light microscope and photomicrographs were taken by an attached eyepiece camera (Dino-eye, AM-7023, 5Mp, Taiwan) on 40 \times and 100 \times magnification. Ten random fields were selected in each slide. Histomorphometric parameters were described according to the American Society for Bone and Mineral Research (ASBMR) nomenclature committee (Vidal et al., 2012)

The histomorphometric parameters evaluated for cancellous bone were bone volume (BV: the percentage of occupied area by calcified bone in relation to the total area.), ratio of BV to TV which included trabecular bone volume (BV/TV), trabecular number (TbN: number of trabeculae that a line through a trabecular compartment would hit per millimetre of its length), trabecular thickness (TbTh: the thickness of 40 complete trabecular bone packet on straight, rod-like trabecular structures in each section), osteoblast surface (ObS/BS: percent of bone surface occupied by osteoblasts and osteoclast surface (OcS/BS: percent of bone surface occupied by osteoclasts) (Kulak and Dempster, 2010, Vidal et al., 2012).

The histomorphometric parameters evaluated for tibial shaft were total tissue area (Tt Ar), marrow cavity area (Mc Ar) and cortical bone area (Cb Ar). These parameters were chosen based on a previous study (Chang et al., 2010). The results were obtained by using an Olympus light microscope and an attached eyepiece digital lens (Dino-eye, AM-7023, 5Mp, Taiwan).

Results were expressed as Means±SE using the software SPSS 16 (version 16, Chicago, USA). Statistical analysis was carried out using an independent T-test and one-way ANOVA. Also the Tukey's test was used for post hoc analysis with significance set at $P<0.05$.

RESULTS

Weight

In the female exercise group, body weight decreased markedly, compared to the control group at the end of the intervention (Table 1).

Bone histomorphometry and histopathology

Knee joint: Gross findings

The surface of the femoral cartilage was found to be smooth in both female and male control groups, whereas in four male runner rats, the articular surface was irregular after the sixth week. In the female exercise group, superficial cartilage irregularities were detected in the trochlea and femoral condyles.

Histological finding

Our findings were indicative of mild OA in both female and male exercise groups. The results in male runner rats were less significant than those of female runners (Table 2). Based on the findings, the synovial layer was normal in both female and male control groups. Most male runner rats in the control group had a normal condition.

There was no synovitis in male runners. However, the synovial joint seemed almost coarse and thick in the female runner group, with mild hyperemia dropsy. There was also a moderate synovitis in this group (Table 2).

In the female exercise group, OcS/BS% was found to be significantly lower than that of the female control group (Table 3).

In the female runner group, ObS/BS% showed a significant increase compared to the female controls (Table 3).

The trabecular number (TbN) increased significantly in both female and male exercise groups, compared to the controls. On the other hand, the trabecular thickness (Tt Ar) difference between female runner and control rats was significant (Table 3).

The BV/TV ratio was higher in male and female exercise groups, compared to the corresponding control groups. The difference between female runner rats was significant (Table 3).

A greater total tissue area (Tt Ar) and a smaller marrow cavity area (Mc Ar) were found in both exercise groups, compared to the controls. The difference between the control and runner female groups was significant (Table 3). On the other hand, the cortical bone area (Cb Ar) was significantly greater in the female exercise group in comparison with the corresponding control group (Table 3).

DISCUSSION

In the present study, moderate-intensity running was found to damage the articular cartilage; therefore, it is considered a strenuous exercise, especially for females. Based on our findings, in moderate-intensity exercise, women experience more cartilage changes, compared to men with a similar body weight. According to previous research, the cartilage volume of normal men is 33% to 42% larger than that of normal women, as indicated in radiographic analyses (Ding et al., 2003).

Moreover, Jones et al. (2000) showed that the knee cartilage of men is significantly more than that of females. It was concluded that sex differences in cartilage development could explain the variations in OA of the knee in later stages of life. In addition, Faber et al.

(2001) reported that after adjustments for height and body weight, gender differences did not influence cartilage volume. Nevertheless, it was reported that body and bone size had significant effects on this difference, suggesting a direct relationship between cartilage volume and bone size (Eckstein et al., 2001; Jones et al., 2000).

Evidence suggests the thicker cartilage of men, compared to women (Ni et al., 2011). In contrast, no gender-related effects have been reported after adjustments for body weight and tibial head diameter (Eckstein et al., 2006). Some significant changes were detected in rats with matching age and body weight; therefore, other parameters might explain gender differences

Previous studies show that running exercise decreases fat mass (Morseth et al., 2011; Howe et al., 2015). Our obtained weight results were similar to this study

The main bone function is to bear gravity-induced mechanical forces and muscle contraction and to facilitate efficient body locomotion (Pajamaki et al., 2003). In a study by Iwamoto et al. (2000), to preserve the positive effects of exercise (for eight weeks), continuous exercise was found to be necessary. To explain the mechanism through which exercise leads to increased bone mass on histomorphometry, Yeh et al. (1993) showed that suppressed bone resorption and improved bone formation might be responsible for the positive effects of exercise on the cancellous bones of tibia in aged female rats. Moreover, Chen et al. (1994) showed that exercise leads to cortical bone mass accumulation in the tibia of aged female rats. Our data demonstrated that moderate running exercise resulted in significant increases in both cancellous and cortical bone mass (BV/TV and Ct Ar, respectively) in a weight-bearing long bone, the tibia, in the female rats. The results of OcS/BS and ObS/BS percentage in female rats indicated that the response of cancellous bone to exercise differs according to the magnitude of mechanical loading to the bone. There is a threshold in mechanical loading to the bones when cancellous bone mass increases, with bone formation exceeding bone resorption (Table 3). These findings are consistent with the researchers' hypothesis (Iwamoto et al., 1998, 2000).

Joint tissues can sustain minimal damage without progressive change. It is known that cumulative micro-damage can lead to bony remodeling, stiffening, and cartilage lesion progression. Peak dynamic forces applied at a high rate cause subchondral bony

remodeling and stiffening (Hudelmaier et al., 2006). In this study, cortical and cancellous bone areas increased significantly in the tibia due to exercise. It should be noted that exercise duration and intensity were probably ideal for increasing cancellous bone mass in our study. Our results also demonstrated that the effect of exercise in females was more significant than males.

In a previous study, it was reported that moderate running exercise significantly increased cortical and cancellous bone mass in tibia (Iwamoto et al., 2011); our findings are in consistence with this study. This finding might be attributed to the position of tibia and mechanical loading on the tibia during treadmill running (Iwamoto et al., 2011; Mosavian 2017). Moreover, Chang et al. (2010) showed that the cancellous bone response to exercise changes relative to the bone mechanical loading. This finding is consistent with a previous study, which indicated the positive site-specific effect of treadmill exercise on BMD, with predominant weight-bearing sites (Iwamoto et al. 2011).

CONCLUSIONS

Our results showed the moderate exercises are more harmful in females than males. Also subchondral bone changes in females may be more visible than males.

Acknowledgements

This research was financially supported by the research council of Shahid Bahonar University of Kerman (No: 1.7.1395). The authors would like to thank Mr. Mazhab Jaafari for providing histological laboratory methods and histology slides. The authors declared no conflict of interests.

Table 1. The changes (Mean±SE) of weight (g) at the first day and end day of the exercise period.

Groups		First day of exercise	End day of exercise
Control	Male	254.14±18.54	256.2±16.82
	Female	248.32±12.32	254.14±14.81
Runner	Male	253.32±21.08	241.37±17.03
	Female	249.23±16.01 ^a	230.08±17.01 ^b

^{a, b} in the same row with no common superscript show significant difference (P<0.05).

Table 2. Mankin's histopathology and synovial gradings (Mean±SE) of distal extremity of lateral condyle of tibia.

	Male		Female	
	Control	Runner	Control	Runner
Mankin*	0 ^a	1.02±0.38 ^a	0 ^a	5.12±0.89 ^b
Synovial Layer**	0 ^a	0.8±0.36 ^a	0 ^a	5.21±0.32 ^b

^{a, b} in the same row with no common superscript show significant difference (P <0.05).

*0: No OA (Osteoarthritis); 1-5: Mild OA; 6-10: Moderate OA; 11-14: Severe OA.

** 0 to 1 corresponds to no synovitis, 2 to 4 to a slight synovitis, 5 to 6 to a moderate synovitis, and 7 to 9 to a strong synovitis.

Table 3. Bone histomorphometry (Mean±SE) of cancellous and cortical parts of tibia*.

	Male		Female	
	Control	Runner	Control	Runner
<i>Cancellous part</i>				
TbN	6.75±0.75 ^a	10.43±0.75 ^b	7.25±0.63 ^a	19.25±0.75 ^c
TbTh (µm)	43.32±5.47 ^{ab}	46.54±4.7 ^{ab}	39.41±4.01 ^a	54.93±4.86 ^b
BV/TV (%)	58.32±5.71 ^a	60.3±5.32 ^a	55.43±4.91 ^a	72.83±6.32 ^b
OcS/BS (%)	8.42±0.6 ^a	8.12±0.52 ^{ab}	8.51±0.61 ^a	7.82±0.42 ^b
ObS/BS (%)	53.41±5.12 ^a	56.02±4.01 ^{ab}	51.07±4.92 ^a	62.41±3.12 ^b
<i>Cortical part</i>				
Tt Ar (mm²)	541.48±27.32 ^{ab}	558.92±31.27 ^{ab}	524.61±19.27 ^a	561.58±23.97 ^b
Mc Ar (mm²)	172.11±10.27 ^{ac}	161.92±9.81 ^a	180.01±14.31 ^{bc}	161.14±9.97 ^a
Cb Ar (mm²)	328.31±24.03 ^{ab}	333.12±19.81 ^{ab}	312.42±16.81 ^a	348.91±23.14 ^b

*The abbreviations are mentioned in material and methods.

^{a,b,c} in the same row with no common superscript show significant difference (P<0.05).

References

Chang TK, Huang CH, Huang CH, Chen HC, Cheng CK (2010) The influence of long-term treadmill exercise on bone mass and articular cartilage in ovariectomized rats. *BMC Musculoskeletal Disorders*, 11:185.

Chen MM, Yeh JK, Aloia JF, Tierney JM, Sprintz S (1994) Effect of treadmill exercise on tibial cortical bone in aged female rats: A histomorphometry and dual energy X-ray absorptiometry study. *Bone*, 15: 313–9.

Ding C, Cicuttini F, Scott F, Glisson M, Jones G (2003) Sex differences in knee cartilage volume in adults: role of body and bone size, age and physical activity. *Rheumatol*, 42:1317–1323.

Eckstein F, Cicuttini F, Raynauld JP, Waterton JC, Peterfy C (2006) Magnetic resonance imaging (MRI) of articular cartilage in knee osteoarthritis (OA): morphological assessment. *Osteoarthritis Cartilage*, 14: 46-75.

Eckstein F, Winzheimer M, Hohe J, Englmeier KH, Reiser M (2001) Interindividual variability and correlation among morphological parameters of knee joint cartilage plates: analysis with three-dimensional MR imaging. *Osteoarthritis Cartilage*, 9: 101–11.

Faber SC, Eckstein F, Lukasz S, Mühlbauer R, Hohe J, Englmeier KH, Reiser M (2001) Sex differences in knee joint cartilage thickness, volume and articular surface areas: assessment with quantitative three-dimensional MR imaging. *Skeletal Radiol*, 30:144–50.

Howe T, Shea B, Dawson LJ, Downie F, Murray A, Ross C, Harbour RT, Caldwell LM, Creed G (2015) Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Data Sys Rev*, 7:1-7. DOI: 10.1002/14651858.

Hudelmaier M, Glaser C, Hausschild A, Burgkart R, Eckstein F (2006) Effects of joint unloading and reloading on human cartilage morphology and function, muscle cross-sectional areas, and bone density - a quantitative case report. *J Musculoskelet Neuronal Interact*, 6(3):284-90.

Iwamoto J, Sato Y, Takeda T, Matsumoto H (2011) Effectiveness of exercise for osteoarthritis of the knee: A review of the literature. *World J Orthop*, 18:2(5):37-42. doi: 10.5312/wjo.v2.i5.37.

Iwamoto J, Takeda T, Ichimura S (1998) Effect of exercise on tibial and lumbar vertebral bone mass in mature osteopenic rats: Bone histomorphometry study. *J Orthop Sci*, 3(5):257-63.

Iwamoto J, Yeh JK, Aloia JF (2000) Effect of deconditioning on cortical and cancellous bone growth in the exercise trained young rats. *J Bone Miner Res*, 15:1842–1849.

- Jarvinen TLN, Kannus P, Sievanen H, Jozsa L, Heinonen OJ, Vieno T, Järvinen M (2001) Effects of remobilisation on rat femur are dose-dependent. *Scand J Med Sci Sports*, 11:292–298.
- Jones G, Glisson M, Hynes K, Cicuttini F (2000) Sex and site differences in cartilage development: a possible explanation for variations in knee osteoarthritis in later life. *Arthritis Rheumatol*, 43:2543–8.
- Joo YI, Sone T, Fukunaga M, Lim SG, Onodera S (2003) Effects of endurance exercise on three-dimensional trabecular bone microarchitecture in young growing rats. *Bone*, 33(4):485-493.
- Kenneth D, Brandt M (2005) Osteoarthritis. In: Kasper, Braunwald, Fauci, Hauser, Longo, Jameson, Editors: *Harrison's principles of internal medicine*. 15th ed. USA. BMC Graw – Hill Incl, 2036 – 45.
- Krenn V, Morawietz L, Haupl T, Neidel J, Petersen I, König A (2002) Grading of chronic synovitis—A histopathological grading system for molecular and diagnostic pathology. *Path Res Pract*, 198(5):317-325.
- Kulak CAM, Dempster DW (2010) Bone histomorphometry: a concise review for endocrinologists and clinicians. *Arch Bras Endocrinol Metab*, 54(2):87-98.
- Lee YJ, Park JA, Yang SH, Kim KY, Kim BK, Lee EY, Lee EB, Seo JW, Echtermeyer F, Pap T, Song YW (2011) Evaluation of osteoarthritis induced by treadmill-running exercise using the modified Mankin and the new OARSI assessment system. *Rheumat Inter*, 31(12): 1571-1576.
- Mankin HJ, Dorfman H, Lippiello L, Zarins A (1971) Biochemical and metabolic abnormalities in articular cartilage from osteoarthritic human hips II. Correlation of morphology with biochemical and metabolic data. *J Bone Joint Surg Am*, 53:523–537.
- Morseth B, Emaus N, Jørgensen L (2011) Physical activity and bone: The importance of the various mechanical stimuli for bone mineral density. A review. *Norsk Epidemiologi*, 20 (2): 173-178.
- Mosavian Naeini R, Sahebozamani M, Nazem MN (2017) Effect of moderate treadmill exercise on hip osteoarthritis in male and female Wistar rats. *Novel Biomed*, 4, 177-84.
- Ni GX, Zhan LQ, Gao MQ, Lei L, Zhou YZ, Pan YX (2011) Matrix metalloproteinase-3 inhibitor retards treadmill running-induced cartilage degradation in rats. *Arthritis Res Ther*, 13:1-11.
- Pajamaki I, Kannus P, Vuohelainen T, Sievänen H, Tuukkanen J, Järvinen M, Järvinen TL (2003) The bone gain induced by exercise in puberty is not preserved through a virtually life-long deconditioning: a randomized controlled experimental study in male rats. *J Bone Miner Res*, 18(3):544-52.

Pap G, Eberhardt R, Sturmer I, Machner A, Schwarzberg H, Roessner A, Neumann W (1998) Development of osteoarthritis in the knee joints of Wistar rats after strenuous running exercise in a running wheel by intracranial self-stimulation. *Path Res Prac*, 194: 41-47.

Radin EL, Rose RM (1986) Role of subchondral bone in the initiation and progression of cartilage damage. *Clin Orthop Relat Res*, 213: 34-40.

Renner AF, Carvalho E, Soares E, Mattiello-Rosa S (2006) The effect of a passive muscle stretching protocol on the articular cartilage. *Osteoarthritis Cartilage*, 14(2):196–202.

Seeman E (2002) An exercise in Geometry. *J Bone Miner Res*, 17:373–380.

Seyyedini S, Nazem MN (2017) Histomorphometric study of the effect of methionine on small intestine parameters in rat: an applied histologic study. *Folia Morphol*, 76(4):620–629. DOI: 10.5603/FM.a2017.0044.

Tromp AM, Bravenboer N, Tanck E, Oostlander A, Holzmann PJ, Kostense PJ, Roos JC, Burger EH, Huiskes R, Lips P (2006) Additional weight bearing during exercise and estrogen in the rat: the effect on bone mass, turnover, and structure. *Calcif Tissue Int*, 79(6):404-415.

Vidal B, Pinto A, Galvao MJ, Santos AR, Rodrigues A, Cascao R, Abdulghani S, Caetano-Lopes J, Ferreira A, Fonseca JE, Canhao H (2012) Bone histomorphometry revisited. *Acta Reumatol Port*, 37:294-300.

Yeh JK, Aloia JF, Chen MM, Tierney JM, Sprintz S (1993) Influence of exercise on cancellous bone of the aged female rat. *J Bone Miner Res*, 8:1117–25.