

Effect of Calcium Carbonate Residues from Cement Industries on the Phenolic Composition and Yield of Shiraz Grapes

I.O. Maya-Meraz¹, R. Pérez-Leal^{1*}, J.J. Ornelas-Paz², J.L. Jacobo-Cuellar¹, M.J. Rodríguez-Roque¹, R.M. Yañez-Muñoz¹, A. Cabello-Pasini^{3*}

(1) Universidad Autónoma de Chihuahua, Facultad de Ciencias Agrotecnológicas, Área de Fisiología Vegetal y Frutales de Clima Templado, Chih., Chihuahua, México

(2) Centro de Investigación en Alimentación y Desarrollo A.C., Laboratorio de Fitoquímicos y Nutrientes, Cuauhtémoc, Chih., México

(3) Universidad Autónoma de Baja California, Instituto de Investigaciones Oceanológicas, Ensenada, Baja California, México

Submitted publication: April 2019

Accepted for publication: October 2019

Key words: abiotic stress, biotic stress, calcium, grapevines, polyphenols, ripening

Phenolic compounds are secondary metabolites synthesised in response to biotic or abiotic stress in plants. This stress-induced increase in phenolic compound concentrations is generally activated by internal levels of abscisic acid (ABA). The exogenous application of ABA or calcium chloride on grapevines is also known to increase grape yield and alter the phenolic composition of grapes. Residues of cement industries such as calcium carbonates (CaCO₃) are a safe environmental source of calcium that could be used to induce the synthesis of phenolic compounds and act as a yield promoter in grapes and other crops. Consequently, the objective of this study was to evaluate the effect of cement industries' CaCO₃ residues (CaCO₃R) on the yield and concentration of phenolic compounds in Shiraz grapes. Thirteen phenolic compounds were identified and quantified by HPLC-DAD. Malvidin-3-*O*-glucoside was the major anthocyanin found in Shiraz grapes, and its concentration increased by more than 200% in CaCO₃R-treated vines. Similarly, the concentration of cinnamic acid, the main precursor of phenolic compounds, increased by more than 900% in grapes treated with CaCO₃ residues at harvest time. Finally, catechin, epicatechin and procyanidin B1 and B2 increased significantly at harvest time in CaCO₃R-treated grapes relative to the controls. In general, it was found that foliar application of CaCO₃ residues from the cement industry at véraison induced an increase in yield, and in the concentration and composition of phenolic compounds in grapes.

INTRODUCTION

Calcium (Ca) is a secondary messenger that triggers environmental adaptive mechanisms in plants as a response to biotic and abiotic stress (Ranty *et al.*, 2016). Biotic stress, resulting from the presence of insects or pathogens in grapevines, for example, triggers jasmonic acid synthesis in the tendrils. Similarly, abiotic stress due to exposure to extreme cold, high UV-B radiation, high levels of metal in the soil, high salinity or drought activates abscisic acid (ABA) synthesis in the leaves and results in stomatal closure (Taurino *et al.*, 2015; Vishwakamara *et al.*, 2017).

It has been demonstrated that Ca regulates at least three groups of Ca-dependent proteins in *Arabidopsis* and other plants. Proteins such as calmodulins (CMs), calcineurin B-like proteins (CBLs) and calcium-dependent protein kinases (CDPKs) can play a critical role in cellular signalling cascades. These Ca-dependent proteins operate as sensors and signal transmitters in response to salinity or drought stress, and can induce gene expression that activates ABA

synthesis and stomatal closure (Yang *et al.*, 2012; Zou *et al.*, 2015). In vineyards, regulated deficit irrigation is a common practice to induce drought stress in the plant at the onset of ripening or véraison in order to stimulate ABA synthesis and enhance pigment accumulation. However, this irrigation practice commonly reduces grape yield at harvest time (De-la-Hera-Orts *et al.*, 2005; Kyraleou *et al.*, 2017). Recent studies show that foliar ABA application enhances pigment concentration in tomatoes, strawberries, litchis and grapes (Singh *et al.*, 2014; Yamamoto *et al.*, 2015; Jia *et al.*, 2016). It was demonstrated that these ABA applications induce the expression of structural genes and transcription factor genes that activate the phenylpropanoid pathway and anthocyanin synthesis through their effect on the activity of phenylalanine ammonia-lyase (PAL) and chalcone synthase (CHS) enzymes (Ferrandino & Lovisolo, 2014; Koyama *et al.*, 2014; Villalobos-González *et al.*, 2016). However, exogenous applications in commercial farming

*Corresponding author: E-mail address: acabello@uabc.edu.mx; rleal@uach.mx

Acknowledgements: We would like to thank the Laboratorio de Fitoquímicos y Nutrientes of the Centro de Investigación en Alimentación y Desarrollo A.C of Cd. Cuauhtémoc, Mexico. The CaCO₃ residues were supplied by Grupo Cementos Chihuahua. This research was partially supported by CONACyT, Mexico

are not economically feasible due to the high cost of ABA. Consequently, other studies have explored the use of calcium chloride (CaCl₂) to increase the synthesis of total phenolic compounds and anthocyanins in plants (Al-Qurashi & Awad, 2013; Xu *et al.*, 2014; Martins *et al.*, 2018). It has been shown that the use of foliar CaCO₃ applications as a source of Ca increases the total content of phenolic compounds in olives (Squeo *et al.*, 2016). Moreover, CaCO₃ application also increase grape yield in commercial vineyards (Sabir *et al.*, 2014).

Overall, phenolic compounds from grapes play a key role in determining wine organoleptic characteristics and quality. Anthocyanins and stilbenes, for example, provide the red or purple coloration in wines, while phenolic acids and flavan-3-ols are responsible for astringency and bitterness. In addition to polyphenols, the overall characteristics and balance of the wine are provided by the chemical characteristics of the grapes at harvest. These include the concentration of total soluble solids (TSS), pH, total titratable acidity (TA), organic acid composition, and positive and negative aromas, among others (Nogales-Bueno *et al.*, 2013; Olivares *et al.*, 2017). Thus, the addition of various compounds to improve yield in commercial vineyards must maintain the balance among chemical characteristics and phenolic compounds. It is hypothesised in this study that foliar applications of CaCO₃ in grapevines during véraison will increase yield and enhance the accumulation of phenolic compounds in the grapes, without affecting chemical parameters.

Climate change has been associated with high anthropogenic carbon dioxide (CO₂) emissions into the environment. To mitigate this environmental impact, the cement industries use different processes to capture CO₂ emissions and generate residues as CaCO₃ (Anbu *et al.*, 2016). These residues could be used as a source of Ca in commercial farming operations. Particularly, CaCO₃ from cement industry residues (CaCO₃R) could provide an inexpensive source of Ca to increase grape yield and polyphenol concentrations in viticulture operations. Consequently, the objective of this study was to evaluate the effect of foliar CaCO₃R applications on the yield of, and on the total and individual concentrations of phenolic compounds in, Shiraz grapes.

MATERIAL AND METHODS

Chemicals and solvents

Malvidin-3-*O*-glucoside chloride, pelargonidin-3-*O*-glucoside chloride, cyanidin 3-*O*-glucoside chloride, cyanidin-3-*O*-galactoside, gallic acid, *trans*-cinnamic acid, *trans*-caftaric acid, protocatechuic acid, *trans*-resveratrol, (+)-catechin, (-)-epicatechin, and procyanidin dimers B1 and B2 standards were used for molecule identification and quantification. All reagents and solvents used were of analytical or HPLC grade, and were purchased from Sigma-Aldrich (St. Louis, Mo, USA), J.T. Baker (Baker-Mallinckrodt Inc, Mexico), and Merck (Merck, Germany).

Study site and treatments

Experiments were conducted in 2017 at an experimental vineyard (28° 25' N and 106° 51' W, 2 010 m above sea level) owned by the Universidad Autónoma de Chihuahua,

Chihuahua, Mexico. The vineyard consisted of eight-year-old Shiraz vines (1103P rootstock) planted at a 3 m distance between rows and 1 m between grapevines. Vines were trained using a Royat system, with rows oriented north-south on sandy loam soils. Drip irrigation was applied from bud break to véraison, and a humidity of 15.6% was maintained throughout the experimental period. Climatic conditions were monitored at a meteorological station (Quintas Lupitas of Unifrut). Rainfall is frequent during the ripening period at the study site, and the accumulation from the time of CaCO₃R application until harvest time was 201.4 mm. The average maximum and minimum temperatures were 23.9°C and 13.6°C respectively.

Calcium carbonate residues (Table 1) from a local cement industry were used as a source of Ca in the experimental treatments. A CaCO₃R (1% w/w) aqueous solution was prepared with tween-20 (0.05 mL L⁻¹) as adherent agent. The concentration of the solution was selected after conducting preliminary field tests. The CaCO₃R solution was applied to the grapevine leaves when clusters reached > 90% véraison using a backpack piston sprayer at a rate of 1 L plant⁻¹ (approx. 3.9 g Ca plant⁻¹, Table 2). Grape sampling started fifteen days after the CaCO₃R foliar application. Approximately 200 berries were randomly collected from each treatment once a week from véraison to harvest time.

Foliar elemental analysis

Forty grapevine leaves were randomly collected once a week before harvest. Leaves were rinsed with deionised water and oven-dried for 72 h in an oven at 70°C to constant weight. The concentration of calcium (Ca), potassium (K), phosphorus (P), magnesium (Mg), manganese (Mn), iron (Fe), copper (Cu) and zinc (Zn) in the leaves was determined using flame atomic absorption spectrometry with oxygen-acetylene flow (PinAAcle 900H, Perkin Elmer, Shelton, CT, USA) (Oliveira *et al.*, 2009). The flame emission measurements were determined at 422.7 nm (Ca), 766.5 nm (K), 285.2 nm (Mg), 327 nm (Cu), 372 nm (Fe), 403.1 nm (Mn) and 213.9 nm (Zn). Standards (0 to 20 ppm) were used to calibrate the instrument, and four replicates were analysed per sample.

Grape analysis

Chemical analysis

Fresh berries (50 per sample) were randomly collected (four replicates) from each treatment and control to conduct chemical and phenolic analyses. Samples were used to determine tristimulus colour (lightness value L*, colour channels a*

TABLE 1

Characterisation of CaCO₃ residues from cement industries in Chihuahua, Mexico.

Molecule	Concentration (% w/w)
Calcium (Ca)	20.4
Magnesium (Mg)	0.32
Calcium carbonate (CaCO ₃)	45.50
Silicon dioxide (SiO ₂)	33.77

TABLE 2
Phenological stages from dormancy to leaf fall and CaCO₃R application on Shiraz grapes.

Phenological stage	Gregorian day	Degree days (°C)
Dormancy	60	108.02
Bud swelling	73	152.44
Bud burst	78	178.80
Beginning of foliar development	83	205.37
Foliar development (> 3 leaves unfolded)	88	223.83
Inflorescence clearly visible	95	245.38
Inflorescence fully developed	99	268.74
Flower separating	110	339.03
Flowering	121	410.16
Fruit set	132	473.99
Beginning of véraison	185	990.15
* Véraison (> 90%)	198	1096.63
Berries ripe for harvest	242	1445.10
Leaves falling	293	1784.27

*Timing of foliar application of CaCO₃R.

and b*) using a tabletop colorimeter (CR-300 model, Minolta Co. Ltd, Osaka, Japan). Fruit firmness was determined using a TA-XT2i texture analyser (Texture Technology Corp., Scarsdale, N.Y., USA. and Stable Micro System Ltd, Godalming, UK). The texture analyser was equipped with a 7 mm diameter stainless steel striker pin that punctured the grape skin at a rate of 10 mm s⁻¹. The maximum puncturing force (in Newtons) was determined for 20 berries per treatment (four replicates). Berries were crushed and macerated for 1 h to evaluate pH (HANNA Instruments Inc., Woonsocket, USA), TA, TSS and phenolic compounds according to international methods for grapes and wine (OIV, 2017). Total soluble solids in the must were determined using a digital refractometer (ATAGO Co. Ltd., Osaka, Japan), and TA was determined by titration of the sample with 0.1 N NaOH.

Analysis of polyphenols

Polyphenols were extracted from the grapes in a dark room with a red safety light. Crushed berries (~3 g) were mixed with 40 mL of acetone/water/trifluoroacetic acid (AWTA, 70:29.9:0.1 v/v) and homogenised using an Ultra Turrax (IKA T18) at 1 000 rpm for 1 min. The sample was vortexed for 5 min, sonicated on ice for 5 min, and then filtered through a Whatman filter paper No. 4. Material on the filter was further rinsed with 40 mL AWTA until the tissue was decolorised. The acetone phase was eliminated in a rotavapor (39°C) and the solution was brought up to 50 mL with 0.01% TA. Finally, the extracts were filtered through a 0.45 µm membrane filter and stored in borosilicate amber vials for subsequent polyphenol analysis.

Total phenolic compound concentration was determined spectrophotometrically after the Folin-Ciocalteu reactions, and was quantified with gallic acid calibration

curves. Individual phenol analysis was performed in an Agilent 1200 series high-performance liquid chromatography (HPLC) system (Agilent, Palo Alto, Ca, USA) with a diode array detector (Ymc Inc. Miliford, Ma, USA) according to standard protocols (Ornelas-Paz *et al.*, 2017). The separation of phenolic compounds was performed with an XDB-C18 column (Agilent, Zorbax eclipse 4.6 x 150 mm 5 µm). The column was operated at 30°C with mobile phases consisting of 2% (v/v) acetic acid (A) and acetonitrile (B). The flow rate was maintained at 1.0 mL min⁻¹, with the following gradient: 100% A/0% B at 0 min; 93% A/7% B at 12 min; 89% A/11% B at 20 min; 86% A/14% B at 35 min; 84% A/16% B at 36 min; 82% A/18% B at 41 min; 76% A/24% B at 48 min; 70% A/30% B at 54 min; 65% A/35% B at 59 min; 50% A/50% B at 65 min; 35% A/65% B at 70 min; 25% A/85% B at 75 min; 15% A/85% B at 80 min; 5% A/95% B at 85-90 min; and 100% A at 95 min. Polyphenols were monitored with UV-visible spectra for anthocyanins (λ = 520 nm), stilbenes (λ = 320 nm), phenolic acids (λ = 280 nm) and flavan-3-ol groups (λ = 280 nm, except catechin, which was monitored at λ = 320 nm) (Wang *et al.*, 2016). Individual polyphenols were identified and quantified using standards and calibration curves. Compounds were identified using retention time of the peak, the shape of the chromatographic peak and the spectra of the extracted anthocyanins. The retention time and shape of the chromatographic peak was compared to that of specific standards. In addition, internal standards were injected with the samples to verify the compound.

Statistical analysis

Results were statistically evaluated based on a split-plot in time design. Analysis of variance and a least square means test were used to detect significant differences among

treatments, and were conducted using SAS System for Windows 9.0 (SAS Institute, Inc. Cary, N.C., USA, 2002) after testing for normality and homoscedasticity of the data. The significance level was set at 0.05 and the experiments were conducted using four replicates.

RESULTS

Effect of grapevine leaf CaCO₃R application on yield and chemical characteristics of grapes

Several macro- and micronutrient concentrations in grapevine leaves varied significantly as a result of the CaCO₃R application relative to the controls (Fig. 1). Thus, the relative concentration of Ca increased ~25% in CaCO₃R-treated grapevine leaves compared to the controls (Fig. 1A). In contrast, absolute or relative concentrations of K, Mg, Fe and Mn in CaCO₃R-treated grapevine leaves decreased significantly, but only by up to approximately 10% relative to the controls (Fig. 1A and 1B). Finally, the concentration of P, Cu and Zn in the leaves did not change significantly in the CaCO₃R-treated vines relative to the controls.

The grape yield and cluster weight of the Shiraz vines also varied as a result of CaCO₃R application (Fig. 2). While the number of clusters per vine was kept similar between treatments (Fig. 2A), the cluster weight and yield increased significantly, by more than 50%, in the CaCO₃R-treated grapevines relative to the controls (Fig. 2B and 2C).

In general, there were no significant changes in the chemical composition of the grapes as a result of CaCO₃R application (Table 3). The percentage of total soluble solids, %TA and pH of the grapes did not vary significantly as a result of CaCO₃R application relative to the control at harvest time. Similarly, the firmness and colour of the grapes were statistically similar between the control and the treated vines at harvest time.

Influence of pre-harvest CaCO₃R on the concentration of phenolic compounds in grapes

Total phenolic compounds, stilbenes and anthocyanins

Total phenolic concentration, as well as that of specific phenolic compounds in the berries, varied significantly

between the treated and control vines (Table 4). Thus, the total phenolic concentration was approximately threefold greater in the CaCO₃R-treated vines relative to the controls at harvest time, whereas *trans*-resveratrol did not differ significantly at harvest time between CaCO₃R-treated vines and the controls. The most abundant anthocyanin found in Shiraz grapes was malvidin-3-*O*-glucoside, which increased nearly threefold at harvest time in the CaCO₃R-treated vines relative to the controls. Similar to malvidin-3-*O*-glucoside, the mean concentrations of cyanidin-3-*O*-galactoside, cyanidin-3-*O*-glucoside and pelargonidin-3-*O*-glucoside increased two- to threefold in CaCO₃R-treated vines relative to the controls.

Phenolic acids and flavan 3-ols

At harvest time, the concentration of most phenolic acids and flavan-3-ols increased significantly in the grapes of CaCO₃R-treated vines relative to the controls (Table 5). The concentration of gallic acid in the grapes was 10- to 70-fold greater than that of the rest of the phenolic acids. In most cases, the concentration of phenolic acids and flavan-3-ols increased sharply 36 days after CaCO₃R application. The concentration of the phenolic acid *trans*-cinnamic acid significantly increased by 12-fold, while the concentration of caftaric acid increased twofold at harvest time in the berries of CaCO₃R-treated vines relative to those in the controls. In contrast, the concentration of gallic acid remained the same in the treatment and controls at harvest time. The sharp decrease in total and specific phenolic compounds 29 days after CaCO₃R application coincided with rainfall in the area.

Catechin, epicatechin, and procyanidin B1 and B2 were identified in Shiraz grapes (Table 5). Epicatechin was the major flavan-3-ol in berries, and at harvest time its concentration was twice as high in the CaCO₃R-treated vines than in the controls. Similar to epicatechin, the concentration of catechin and procyanidin B1 was two- to threefold greater at harvest time in the CaCO₃R-treated vines compared to the controls, whereas that of procyanidin B2 was only 35% greater in the CaCO₃R-treated vines than in the controls.

Overall, there was a significant increase in the relative




FIGURE 1

Effect of foliar CaCO₃R application on the relative concentration (%) of potassium (K), calcium (Ca), magnesium (Mg) and phosphorous (P), and the concentrations (in ppm) of copper (Cu), iron (Fe), manganese (Mg) and zinc (Zn) in the leaves of Shiraz grapes. Different letters above bars indicate statistically significant differences.




FIGURE 2

Effect of foliar applications of CaCO₃R on the average number of clusters per grapevine (\pm SE), cluster weight and grape yield at harvest time in Shiraz grapes. Different letters above bars indicate statistically significant differences.

concentration of polyphenolic compounds in Shiraz grapes relative to the controls at harvest time. The relative concentration of anthocyanins and flavan-3-ols increased ~twofold at harvest time in the berries of CaCO₃R-treated vines compared to that of the controls. Finally, the relative concentration of stilbene in the berries increased significantly, by ~30%, in the CaCO₃R-treated vines relative to in the controls. In contrast to these polyphenols, there was no detectable variation in the concentration of phenolic acids between treatments and controls at harvest time.

DISCUSSION

The results of the present study show that foliar CaCO₃R applications affect macro- and micronutrient concentrations, grape yield, and total and specific polyphenolic concentrations in vine and grape tissue. These results are consistent with the wide impact of calcium on plant physiology found in other studies. Increasing concentration of Ca, for example, increased the biomass of leaves but decreased root dry weight in tobacco plants (López-Lefebvre *et al.*, 2001). In contrast, respiration rates and ascorbic acid content in

cherries decreased with increasing Ca concentration (Wang *et al.*, 2014). Finally, foliar applications of CaCl₂ increased total phenolic content, especially that of anthocyanin, in strawberries (Xu *et al.*, 2014). Clearly, Ca plays a key role in the assimilation of macro- and micronutrients, the activation/deactivation of specific enzymes, and the expression of genes controlling key metabolic pathways. Similarly, the results of this study suggest that foliar applications of Ca from CaCO₃ residues produced by cement industries impact the overall physiology of wine-producing grapes.

The decrease in tissue K, Fe, Mg and Mn concentrations in CaCO₃R-treated grape leaves observed in this study is consistent with the results of prior studies. The decrease in tissue Mg was observed even though there was a small concentration (0.32%) of this element in the CaCO₃-residues used here. The concentration of P, K and Mg in leaves and fruits of cherry trees, for example, decreased as a result of foliar applications of Ca (Mikiciuk *et al.*, 2015). In addition, the concentration of Mg, Na and P decreased in the leaves and roots of tobacco plants as a result of increasing concentrations of Ca in the culture media (López-Lefebvre

TABLE 3
Effect of foliar CaCO₃R application on chemical parameters of Shiraz grapes during ripening. TSS = total soluble solids, TA = titratable acidity. Values indicate the means of four samples. CIELab = colour as defined by the International Commission on Illumination (CIE), L* = lightness value, a* and b* = colour channels.

Days after application (DDA)	TA (g L ⁻¹ tartaric acid equivalents)													
	TSS (°Brix)		pH		Firmness (N)		Colour (CIELab)		CaCO ₃ R		Control			
	Control	CaCO ₃ R	Control	CaCO ₃ R	Control	CaCO ₃ R	Control	CaCO ₃ R	L*	a*	b*	L*	a*	b*
15	16.07	16.14	11.13	11.42	2.89	2.88	2.02 ^b	2.32 ^a	32.58	3.11	-1.65 ^a	30.10	3.16	-0.09 ^b
22	19.17 ^a	17.24 ^b	11.00	11.00	3.08	3.06	2.07	2.13	30.39	3.35	-1.35 ^a	28.58	3.56	-0.83 ^b
29	19.36	19.30	10.46	10.19	2.90 ^b	2.99 ^a	1.23	1.31	29.39	3.44	-0.84 ^b	28.54	3.48	-1.28 ^a
36	20.39 ^a	19.51 ^b	8.21	8.35	3.23	3.26	1.56	1.61	30.39	3.02	-0.85	31.46	2.88	-0.76
44 (HD)	20.24	20.15	7.57	7.35	3.34	3.30	1.89	1.89	29.81	2.87	-0.47	31.12	2.77	-0.46
S.E.	0.159	0.190	0.0241	0.0765	0.0701	0.145	0.275	0.701	0.145	0.701	0.275	0.701	0.145	0.275

Harvest date (HD). Different letters indicate statistically significant differences ($p < 0.05$) between treatment and control. Standard error = S.E.

et al., 2001). Changes in the absorption of these ions can affect stomatal aperture (Ruiz *et al.*, 1993; Andrés *et al.*, 2014), the concentration of phenolic compounds, cell wall organisation (Martins *et al.*, 2018), the accumulation of sugars and organic acids in fruit (Prasad *et al.*, 2015), and also overall plant physiology. For example, it has been observed that a disruption in the accumulation of K in guard cells reduces cell turgidity and the opening of the stoma (López-Lefebvre *et al.*, 2001). Similar to K, the increase in Ca concentration in tissues was shown to reduce stomatal aperture in a number of plants (Ruiz *et al.*, 1993). The Ca-induced suppression of stomatal aperture recovers after Ca concentrations in the xylem decrease. Thus, increasing Ca concentration via foliar applications might reduce the K concentration in the leaves and reduce stomatal aperture. Furthermore, the results of the present study suggest that applications of foliar CaCO₃R could have a similar effect on stomatal closure as that observed in grapevines under restricted irrigation. Moreover, the low concentrations of Fe, Mg and Mn found in CaCO₃R-treated grapevine leaves could be related to other physiological processes, given that these elements function as cofactors in a number of key metabolic reactions such as photosynthesis.

The results of the effects of foliar ABA applications on the maturation of grapes are equivocal. On the one hand, some studies have found that the use of ABA and ethephon to enhance phenolic compounds in grapes does not increase grape yield and cluster weight (Peppi *et al.*, 2006; Coelho de Souza-Leaño *et al.*, 2014). However, other studies indicate that foliar ABA application can reduce grape yield and berry size (Alonso *et al.*, 2016). The results of the present study suggest that foliar applications of CaCO₃R positively affect the yield and cluster weight of Shiraz grapes. This is consistent with the findings in Narince grapes treated with fertilisers containing CaCO₃ (Sabir *et al.*, 2014). In contrast, the increase in yield and cluster weight in the present study could be attributed to the fact that treatment with CaCO₃R reduces the stomatal opening and prevents vine and grape dehydration.

External applications of chemical compounds can affect plant cell physiology and chemical fruit composition. Thus, in some cases, foliar application of ABA can increase the concentration of TSS and anthocyanins in grapes and litchis (Singh *et al.*, 2014; Yamamoto *et al.*, 2015). Other studies, however, have found that the same ABA application showed no differences in TSS, TA, pH or tristimulus colour relative to controls (Peppi *et al.*, 2006; Alonso *et al.*, 2016; Coelho de Souza-Leaño *et al.*, 2014). The results on the use of exogenous ABA applications are thus equivocal. These chemical parameters are of great interest for winemaking, since they determine the technological maturation of the grapes, potential alcohol content, and the balance between some organoleptic parameters of phenolic compounds in wines (Nogales-Bueno *et al.*, 2013). The increase in TSS and grape colour, however, is likely an artefact of grape dehydration rather than an increase in the synthesis of sugars or anthocyanins (Santiago *et al.*, 2013). The use of CaCO₃R in the experiment reported in the present study did not affect TSS, TA or pH in Shiraz grapes. This suggests that the use of CaCO₃R does not interact with the plant's

TABLE 4

Mean concentrations of total phenolic compounds, stilbene and anthocyanin in CaCO₃R-treated Shiraz grapes during ripening (mg g⁻¹ dry weight).

Compound and treatment	Days after application (DDA)				
	15	22	29	36	44 (HD)
Total phenolic compounds					
+ CaCO ₃ R	5.673	5.710 ^{a*}	1.855	3.340 ^{a*}	4.560 ^{a*}
Control	5.050	3.343 ^b	2.295	1.540 ^b	1.452 ^b
S.E.	0.635	0.635	0.635	0.550	0.078
<i>Stilbene</i>					
Trans-resveratrol					
+ CaCO ₃ R	0.086	0.105	0.102	0.148 ^{a*}	0.076
Control	0.072	0.090	0.079	0.063 ^b	0.052
S.E.	0.017	0.017	0.017	0.017	0.015
<i>Anthocyanins</i>					
Malvidin-3- <i>O</i> -glucoside					
+ CaCO ₃ R	1.690	3.152	2.291	5.787 ^{a**}	6.093 ^{a**}
Control	1.761	3.331	2.038	1.505 ^b	2.212 ^b
S.E.	0.361	0.312	0.361	0.361	0.442
Cyanidin-3- <i>O</i> -galactoside					
+ CaCO ₃ R	0.518	0.629	0.519	1.160 ^{a*}	1.108 ^{a*}
Control	0.442	0.618	0.582	0.435 ^b	0.494 ^b
S.E.	0.121	0.121	0.148	0.121	0.121
Cyanidin-3- <i>O</i> -glucoside					
+ CaCO ₃ R	0.321	0.412	0.320	0.540 ^{a*}	0.621 ^{a*}
Control	0.340	0.428	0.363	0.206 ^b	0.283 ^b
S.E.	0.050	0.061	0.050	0.050	0.050
Pelargonidin-3- <i>O</i> -glucoside					
+ CaCO ₃ R	0.204	0.299	0.157	0.477 ^{a*}	0.443 ^{a*}
Control	0.181	0.229	0.249	0.172 ^b	0.257 ^b
S.E.	0.040	0.032	0.032	0.032	0.032

Harvest date (HD). Different letters indicate significant differences between treatments (* = at $p < 0.05$ and ** = $p < 0.001$). S.E. = standard error

maturation process, and that the latter is regulated by climatic conditions. However, it is important to highlight that other characteristics, such as the absolute and relative concentrations of phenolic compounds, were positively affected in grapes treated with CaCO₃R.

Exogenous Ca application is used in the fruit industry to increase fruit firmness and shelf life. While the use of CaCl₂ was found to increase these characteristics in El-Bayadi table grapes (Al-Qurashi & Awad, 2013), leaf ABA application decreased the firmness of Flame Seedless grapes. The use of CaCO₃R in the present study did not increase Shiraz grape firmness at harvest. Further studies are thus necessary to determine the effect of these chemicals on the firmness and shelf life of different types of grapes. However, since Shiraz

grapes are used in the production of wines, it is likely that the application of CaCO₃R does not play a critical role in the production of wine-producing grapes.

Abcisic acid plays a key role in the biosynthesis of phenols in plants, including grapevines. This acid triggers the biosynthesis of phenolic compounds that filter UV radiation in grapevine leaves (Berli *et al.*, 2011). For example, total phenolic compounds were shown to increase as a result of foliar applications of ABA (Coelho de Souza-Leaño *et al.*, 2014; Yamamoto *et al.*, 2015; Alonso *et al.*, 2016; Wang *et al.*, 2016). Similarly, foliar applications of 50 mM CaCl₂ led to an increase in the concentration of total phenolic compounds in grapes and strawberries (Al-Qurashi & Awad, 2013; Wang *et al.*, 2013; Xu *et al.*, 2014). Furthermore, the use

TABLE 5
Phenolic acids and flavan-3-ols identified and quantified in CaCO₃R-treated Shiraz grapes and controls during ripening (mg g⁻¹ dry weight).

Compound and treatment	Days after application (DDA)				
	15	22	29	36	44 (HD)
<i>Phenolic acids</i>					
Trans-cinnamic acid					
+ CaCO ₃ R	0.021	0.030	0.041	0.155 ^{a**}	0.169 ^{a**}
Control	0.003	0.041	0.027	0.004 ^b	0.014 ^b
S.E.	0.006	0.006	0.006	0.006	0.006
Gallic acid					
+ CaCO ₃ R	30.558 ^{a*}	23.470 ^{a*}	26.420 ^{a*}	22.145 ^{a*}	4.299
Control	13.644 ^b	15.439 ^b	13.723 ^b	5.346 ^b	4.541
S.E.	3.885	3.885	3.884	2.800	3.205
Caftaric acid					
+ CaCO ₃ R	0.519	0.698 ^{a*}	0.317	0.549 ^{a*}	0.473 ^{a*}
Control	0.631	0.383 ^b	0.459	0.319 ^b	0.240 ^b
S.E.	0.073	0.074	0.074	0.074	0.064
Protocatechuic acid					
+ CaCO ₃ R	0.048	0.056	0.050	0.069 ^{a*}	0.061 ^{a*}
Control	0.044	0.044	0.039	0.034 ^b	0.036 ^b
S.E.	0.006	0.006	0.007	0.006	0.006
<i>Flavan-3-ols</i>					
Catechin					
+ CaCO ₃ R	0.253	0.214	0.378	0.403 ^{a*}	0.619 ^{a*}
Control	0.187	0.188	0.387	0.145 ^b	0.219 ^b
S.E.	0.039	0.039	0.045	0.045	0.039
Epicatechin					
+ CaCO ₃ R	1.583	1.916	1.195 ^{b*}	3.998	2.639 ^{a*}
Control	2.234	1.838	2.342 ^a	2.905	1.270 ^b
S.E.	0.275	0.275	0.318	0.389	0.318
Procyanidin B1					
+ CaCO ₃ R	0.904	0.356	0.732	0.424	1.106 ^{a*}
Control	0.908	0.228	0.886	0.454	0.421 ^b
S.E.	0.092	0.076	0.092	0.076	0.076
Procyanidin B2					
+ CaCO ₃ R	0.070	0.334 ^{a*}	0.366	0.636 ^{a*}	0.779 ^{a*}
Control	0.032	0.033 ^b	0.284	0.379 ^b	0.501 ^b
S.E.	0.082	0.099	0.099	0.099	0.082

Harvest date (HD). Different letters indicate significant differences between treatments (* = $p < 0.05$ and ** = $p < 0.001$). S.E. = standard error.

of CaCO₃ also increased the concentration of total phenolic compounds in olives (Squeo *et al.*, 2016). In agreement with these studies, the foliar application of approximately 40 mM

CaCO₃ here also increased phenolic compound concentration in the skins of Syrah grapes. Therefore, the results of the present study agree with the findings of previous studies

that showed an increase in total phenolic compounds as a result of the foliar application of Ca (CaCO₃ or CaCl₂). Calcium is a secondary messenger that regulates the groups of Ca-dependent proteins related to the synthesis of ABA and stomatal closure (Yang *et al.*, 2012; Zou *et al.*, 2015; Ranty *et al.*, 2016). Consequently, our results suggest that the use of CaCO₃R as a source of Ca could be more economical than the use of ABA or other exogenous compounds to regulate the phenolic concentration in grapes. Additionally, the application of Ca could be used to regulate the concentration of specific stilbenes, anthocyanins, phenolic acids and flavan-3-ols.

Stilbenes are phytoalexins synthesised as a defence mechanism against biotic stress in plants. The stilbene most commonly studied in *Vitis* species is *trans*-resveratrol. The application of a number of chemical compounds is known to affect the accumulation of stilbenes in vine tissue, including in the grapes. Exogenous applications of ABA to vines, for example, increase the concentration of *trans*-resveratrol in Beihong grapes. This increase was only observed 20 days after ABA application and rapidly decreased thereafter (Wang *et al.*, 2016). Furthermore, applications of CaCl₂ to Beihong and Hongbaladuo vines increased *trans*-resveratrol concentrations in the leaves and berry skins relative to the controls (Wang *et al.*, 2013). In agreement with these studies, foliar applications of CaCO₃R in the present study resulted in an increase in the concentration of *trans*-resveratrol as the grapes matured. However, the concentration decreased to levels similar as those measured in the controls at harvest time. The initial increase in *trans*-resveratrol as a result of CaCO₃R application might be a direct response to increased Ca availability. Calcium-dependent proteins, such as CDPKs, CMs and CBLs, trigger the synthesis of ABA (Boss *et al.*, 1996; Yang *et al.*, 2012; Zou *et al.*, 2015; Vandelle *et al.*, 2018). Thus, it is likely that foliar applications of CaCO₃ activate some Ca-dependent proteins. In turn, the decrease in *trans*-resveratrol at harvest time might be the result of substrate competition between stilbene and chalcone synthases, as both use the same substrate. Therefore, the competition for substrate could explain the relationship between the higher concentrations of anthocyanins and lower *trans*-resveratrol concentrations in CaCO₃R-treated grapes observed at harvest time in the present study.

Anthocyanins in grapes are extracted during maceration and play a key role in the production of red wines. Thus, in the last decade, exogenous substance applications to plants have been used to increase the synthesis of anthocyanins in fruits. Applications of ABA, for example, increased the anthocyanin content in litchis and grapes by 50% (Singh *et al.*, 2014; Koyama *et al.*, 2014). Similarly, exogenous applications of ABA (approximately 2 mM) increased the concentration of malvidin-3-*O*-glucoside and cyanidin in Crimson seedless and Malbec grapes (Coelho de Souza-Leaño *et al.*, 2014; Alonso *et al.*, 2016). External application of CaCl₂ was also shown to increase the concentration of cyanidin in strawberries and cherries (Wang *et al.*, 2014; Xu *et al.*, 2014). The presence of pelargonidin-3-*O*-glucoside and cyanidin-3-*O*-galactoside is not often reported in the skins of *Vitis vinifera* grapes. Lately, however, the presence of these molecules has been clearly established in Cabernet

sauvignon and Pinot noir grapes grown in high-altitude vineyards (approximately 2 500 m; He *et al.*, 2010). It is also likely that the presence of pelargonidin-3-*O*-glucoside and cyanidin-3-*O*-galactoside is related to the high elevation of our study site. The present study demonstrates for the first time that foliar CaCO₃R application increases the concentration of anthocyanins. Thus, this study suggests that exogenous application of 1% CaCO₃R is as effective as ABA or CaCl₂ in increasing the synthesis of anthocyanins. This further suggests that there are a number of factors involved in the ripening of grapes, including climatic conditions and the effects of calcium on leaves. Grapevines have different and complex responses to abiotic or biotic stress. Thus, our results suggest that CaCO₃R might be a source of Ca that acts as the secondary messenger and activates the response to abiotic stress, and possibly triggers ABA synthesis or expression in the leaves.

Phenolic acids are the first phenolic compounds synthesised mainly in the tissues of flowers and fruits. The phenylalanine ammonia-lyase enzyme catalyses the conversion of the amino acid phenylalanine to *trans*-cinnamic acid, the first phenolic acid (Abdel-Salam & Hassan, 2015). Foliar application of ABA and CaCl₂ to vines has been shown to change the concentration of phenolic acids in fruits. Thus, the application of ABA led to a decrease in the concentration of gallic acid in the grapes (Alonso *et al.*, 2016). It was also demonstrated that CaCl₂ application increased the concentration of *trans*-cinnamic acid, yet other phenolic acids such as *trans*-caftaric acid and protocatechuic acid were not affected (Kiselev *et al.*, 2013; Yamamoto *et al.*, 2015; Martins *et al.*, 2018). In agreement with findings obtained with CaCl₂ applications, the results of the present study show that foliar applications of CaCO₃R increase the concentration of *trans*-cinnamic acid, *trans*-caftaric acid and protocatechuic acid in Shiraz grapes at harvest time. In contrast, gallic acid increases only during early grape maturity. These results suggest that foliar application of CaCO₃R induces the synthesis of the main phenolic compound precursors.

Flavan-3-ols are responsible for the bitterness and astringency of wines (Kyrleou *et al.*, 2017). Monomeric and oligomeric forms in grapes are known to be affected by ABA application, e.g. the latter resulted in high concentrations of catechin and epicatechin in Yan and Cabernet Sauvignon grapes (Luan *et al.*, 2014). These studies demonstrate that the application of ABA to Cabernet Sauvignon grapes increases the activity of the leucoanthocyanidin reductase, the enzyme responsible for the synthesis of catechins (Lacampagne *et al.*, 2010). Furthermore, foliar application of CaCl₂ increased the concentration of epicatechin and procyanidin B1, while leading to a decrease in catechin and procyanidin B2 in Tempranillo and Graciano grapes (Abdel-Salam & Hassan, 2015). In the present study, the concentration of catechin, epicatechin and procyanidin B1 and B2 increased in grapes treated with CaCO₃R. This suggests that the application of CaCO₃R increases ABA expression in leaves and triggers leucoanthocyanidin reductase activity, resulting in an increase in monomeric and oligomeric forms of flavan-3-ols. In general, our results suggest that CaCO₃R treatment strongly affects the production of all phenolic compounds,

including anthocyanins, stilbenes, phenolic acids and flavan-3-ols. This further suggests that CaCO₃R could be used to enhance phenolic compounds such as anthocyanins and flavan-3-ols in grapes and others fruits.

CONCLUSIONS

The present study evaluated the effect of foliar applications of CaCO₃ residues from the cement industry on the concentration of phenolic compounds in Shiraz grapes. The results demonstrate that foliar applications of CaCO₃R increase grape yield and modify the concentration of ions that regulate stomatal closure in the treated vines. Furthermore, the increase in phenolic compound concentration in grapes from treated vines suggests that CaCO₃R activates the expression of genes involved in the synthesis of ABA or jasmonic acid. Finally, the use of CaCO₃ residues from the cement industry offers an effective and economical alternative to increase total phenolic compounds, including anthocyanins, in wine grapes.

LITERATURE CITED

- Abdel-Salam, M.M. & Hassan, M.A.M., 2015. Effect of CaCl₂ sprays on anthocyanins, pectin and phenolic acids fractionation in berry skin of two red grape (*Vitis vinifera* L.) cultivars during ripening. *Int. J. Acad. Res.* 7(1), 159-167.
- Alonso, R., Berli, F.J., Fontana, A., Piccoli, P. & Bottini, R., 2016. Malbec grape (*Vitis vinifera* L.) responses to the environment: Berry phenolics as influenced by solar UV-B, water deficit and sprayed abscisic acid. *Plant Physiol. Biochem.* 109, 84-90.
- Al-Qurashi, A.D. & Awad, A., 2013. Effect of pre-harvest calcium chloride and ethanol spray on quality of 'El-Bayadi' table grapes during storage. *Vitis* 52(2), 61-67.
- Anbu, P., Kang, C.H., Shin, Y.J. & So, J.S., 2016. Formation of calcium carbonate mineral by bacteria and its multiple applications. *Springer Plus* 5(1), 1-26.
- Andrés, Z., Pérez-Hormanche, J., Leidi, E.O., Schlücking, K., Steinhorst, L., McLachlan, D.H., Schumacher, K., Hetherington, A.M., Kudla, J., Cubero, B. & Pardo, J.M., 2014. Control of vacuolar dynamics and regulation of stomatal aperture by tonoplast potassium uptake. *Proc. Natl Acad. Sci. U. S. A.* 111(17), E1806-E1814.
- Berli, F.J., Fanzone, M., Piccoli, P. & Bottini, R., 2011. Solar UV-B and ABA are involved in phenol metabolism of *Vitis vinifera* L. increasing biosynthesis of berry skin polyphenols. *J. Agr. Food Chem.* 59(9), 4874-4884.
- Boss, P.K., Davies, C. & Robinson, S.P., 1996. Analysis of the expression of anthocyanin pathway genes in developing *Vitis vinifera* L. cv 'Shiraz' grape berries and the implications for pathway regulation. *Plant Physiol.* 111(4), 1059-1066.
- Coelho de Souza-Leaño, P., Coelho-Lima, M.A., Dias-Costa, J.P. & Gomes-da Trindade, D.C., 2014. Abscisic acid and ethephon for improving red color and quality of crimson seedless grapes grown in a tropical region. *Am. J. Enol. Viticult.* 66(1), 37-45.
- De-la-Hera-Orts, M.L., Martínez-Cutillas, A., López-Roca, J.M. & Gómez-Plaza, E., 2005. Effect on moderate irrigation on grape composition during ripening. *Span. J. Agric. Res.* 3(3), 352-361.
- Ferrandino, A. & Lovisololo, C., 2014. Abiotic stress effects on grapevine (*Vitis vinifera* L.): Focus on abscisic acid-mediated consequences on secondary metabolism and berry quality. *Environ. Exp. Bot.* 103, 138-147.
- He, F., He, J.J., Pan, Q.H. & Duan, C.Q., 2010. Mass-spectrometry evidence confirming the presence of pelargonidin-3-O-glucoside in the berry skins of Cabernet Sauvignon and Pinot Noir (*Vitis vinifera* L.). *Aust. J. Grape Wine Res.* 16, 464-468.
- Jia, H., Jiu, S., Zhang, Ch., Wang, Ch., Tariq, P., Liu, Z., Wang, B., Cui, L. & Fang, J., 2016. Abscisic acid and sucrose regulate tomato and strawberry fruit ripening through the abscisic acid-stress-ripening transcription factor. *Plant Biotechnol. J.* 14(10), 2045-2065.
- Kiselev, K.V., Dubrovina, A.S., Shumakova, O.A., Karetin, Y.A. & Manyakhin, A.Y., 2013. Structure and expression profiling of a novel calcium-dependent protein kinase gene, CDPK3a, in leaves, stems, grapes, and cell cultures of wild-growing grapevine *Vitis amurensis* Rupr. *Plant Cell Rep.* 32, 431-442.
- Koyama, R., Marinho de Assis, A., Yamamoto, L.Y., Borges, W.F. & Borges, R.S., 2014. Exogenous abscisic acid increases the anthocyanin concentration of berry and juice from 'Isabel' grapes (*Vitis labrusca* L.). *HortScience* 49(4), 460-464.
- Kyraleou, M., Kallithraka, S., Theodorou, N., Teissedre, P.L., Kotseridis, Y. & Kondouras, S., 2017. Changes in tannin composition of Syrah grape skins and seeds during fruit ripening under contrasting water conditions. *Molecules* 22, 1453.
- Lacampagne, S., Gagné, S. & Gény, L., 2010. Involvement of abscisic acid in controlling the proanthocyanidin biosynthesis pathway in grape skin: New elements regarding the regulation of tannin composition and leucoanthocyanidin reductase (LAR) and anthocyanidin reductase (ANR) activities and expression. *J. Plant Growth Regul.* 29(1), 81-90.
- López-Lefebvre, L.R., Rivero, R.M., García, P.C., Sánchez, E., Ruiz, J.M. & Romero, L., 2001. Effect of calcium on mineral nutrient uptake and growth of tobacco. *J. Sci. Food Agric.* 81(4), 1334-1338.
- Luan, L.Y., Zhang, Z.W., Xi, Z.M., Huo, S.S. & Ma, L.N., 2014. Comparing the effects of exogenous abscisic acid on the phenolic composition of 'Yan' and 'Cabernet Sauvignon' (*Vitis vinifera* L.) wines. *Eur. Food Res Technol.* 239(2), 203-213.
- Martins, V., Garcia, A., Costa, C., Sottomayor, M. & Gerós, H., 2018. Calcium- and hormone-driven regulation of secondary metabolism and cell wall enzymes in grape berry cells. *J. Plant Physiol.* 231, 57-67.
- Mikiciuk, G., Mikiciuk, M., Moździerz, E., Statkiewicz, M. & Chylewska, U., 2015. The effects of foliar nutrition with UnCa fertilizers on the chemical composition of leaves and fruits of sweet cherries. *J. Ecol. Eng.* 16(2), 116-119.
- Nogales-Bueno, J., Hernández-Hierro, J.M., Rodríguez-Pulido, F.J. & Heredia, F.J., 2013. Determination of technological maturity of grapes and total phenolic compounds of grape skin in red and white cultivars during ripening by near infrared hyperspectral image: A preliminary approach. *Food Chem.* 221, 1685-1690.
- OIV (International Organisation of Vine and Wine). 2017. Compendium of international methods of wine and musts analysis. Rue d'Aguesseau-75008 Paris. Online: <http://www.oiv.int/en/technical-standards-and-documents/methods-of-analysis/compendium-of-international-methods-of-analysis-of-wines-and-musts-2-vol> [accessed 15 July 2017].
- Olivares, D., Contreras, C., Muñoz, V., Rivera, S., González-Agüero, M., Retamales, J. & Defilippi, B.G., 2017. Relationships among color development, anthocyanins and pigment-related gene expression in 'Crimson seedless' grapes with abscisic acid and sucrose. *Plant Physiol. Biochem.* 115, 286-297.
- Oliveira, S.R., Gomes-Neto, J.A., Nóbrega, J.A. & Jones, B.T., 2009. Determination of macro-and micronutrients in plant leaves by high resolution continuum source flame atomic absorption spectrometry combining instrumental and ample preparation strategies. *Spectrochim. Acta B* 65(4), 316-320.

- Ornelas-Paz, J.J., Meza, M.B., Obenland, D., Rodríguez-Friscia, K., Jain, A., Thornton, S. & Prakash, A., 2017. Effect of phytosanitary irradiation on the postharvest quality of seedless kishu mandarins (*Citrus kinokuni mukaku kishu*). *Food Chem.* 230, 712-720.
- Peppi, M.C., Fidelibus, M.W. & Dokoozlian, N., 2006. Abscisic acid applications timing and concentration affect firmness, pigmentation, and color of 'Flame Seedless' grapes. *HortScience* 41(6), 1440-1445.
- Prasad, B., Dimri, D.C. & Bora, L., 2015. Effect of pre-harvest foliar spray of calcium and potassium on fruit quality of pear cv. Pathernakh. *Sci. Res. Essays* 10(11), 376-380.
- Ranty, B., Aldon, D., Cotellet, V., Galaud, J.P., Thuleau, P. & Mazars, C., 2016. Calcium sensors as key hub in plant responses to biotic and abiotic stresses. *Front. Plant Sci.* 7, 1-7.
- Ruiz, L.P., Atkinson, C.J. & Mansfield, T.A., 1993. Calcium in the xylem and its influence on the behavior of stoma. *Phil. Trans. R. Soc. B* 341, 67-74.
- Sabir, A., Yazar, K., Sabir, F., Kara, Z., Yazici, M.A. & Goksu, N., 2014. Vine growth, yield, berry quality attributes and leaf nutrient content of grapevines as influenced by seaweed extract (*Ascophyllum nodosum*) and nanosize fertilizer pulverizations. *Sci. Hortic.* 175, 1-8.
- Santiago, W.E., Dos-Reis-Tinini, R.C., Augustos-De-Oliveira, R. & Teruel, B.J., 2013. Partial dehydration of 'Niagara Rosada' grapes (*Vitis labrusca* L.) targeting increased concentration of phenolic compounds and soluble solids. *Afr. J. Biotechnol.* 12, 6474-6479.
- Singh, S.P., Saini, M.K., Singh, J., Pongener, A. & Sidhu, G.S., 2014. Preharvest applications of abscisic acid promotes anthocyanins accumulation in pericarp of litchi fruit without adversely affecting postharvest quality. *Postharvest Biol. Tech.* 96, 14-22.
- Squeo, G., Silletti, R., Summo, C., Paradiso, V.M., Pasqualone, A. & Caponio, F., 2016. Influence of calcium carbonate on extraction yield and quality of extra virgin oil from olive (*Olea europaea* L. cv. Coratina). *Food Chem.* 209, 65-71.
- Taurino, M., Ingresso, I., D'amico, L., De Domenico, S., Nicoletti, I., Corradini, D., Santino, A. & Giovino, G., 2015. Jasmonates elicit different sets of stilbenes in *Vitis vinifera* cv. Negramaro cell cultures. *SpringerPlus* 4(49), 2-11.
- Vandelle, E., Vannozzi, A., Wong, D., Danzi, D., Digby, A.M., Dal Santo, S. & Astegno, A., 2018. Identification, characterization, and expression analysis of calmodulin and calmodulin-like genes in grapevine (*Vitis vinifera*) reveal likely roles in stress responses. *Plant Physiol. Biochem.* 129, 221-237.
- Villalobos-González, L., Peña-Neira, A., Ibáñez, F. & Pastenes, C., 2016. Long-term effects of abscisic acid (ABA) on the grape berry phenylpropanoid pathway: Gene expression and metabolite content. *Plant Physiol. Biochem.* 105, 213-223.
- Vishwakarma, K., Upadhyay, N., Kumar, N., Yadav, G., Singh, J., Mishra, R.K., Kumar, V., Verma, R., Upadhyay, R.G., Pandey, M. & Sharma, S., 2017. Abscisic acid signaling and abiotic stress tolerance in plants: A review on current knowledge and future prospects. *Front. Plant Sci.* 8(161), 1-12.
- Wang, J., Wang, S., Liu, G., Edwards, E. J., Duan, W., Li, S. & Wang, L., 2016. The synthesis and accumulation of resveratrol are associated with veraison and abscisic acid concentration in Beihong (*Vitis vinifera* × *Vitis amurensis*) berry skin. *Front. Plant Sci.* 7, 1-11.
- Wang, L., Ma, L., Xi, H., Duan, W., Wang, J. & Li, Sh., 2013. Individual and combined effects of CaCl₂ and UV-C on the biosynthesis of resveratrol in grape leaves and berry skins. *J. Agric. Food Chem.* 61(29), 7135-7141.
- Wang, Y., Xie, X. & Long, L.E., 2014. The effect of postharvest calcium application in hydro-cooling water on tissue calcium content, biochemical changes, and quality attributes of sweet cherry fruit. *Food Chem.* 160, 23-30.
- Xu, W., Peng, H., Yang, T., Whitaker, B., Huang, L., Sun, J. & Chen, P., 2014. Effect of calcium on strawberry fruit flavonoid pathway gene expression and anthocyanin accumulation. *Plant Physiol. Biochem.* 82, 289-298.
- Yamamoto, L.Y., Marinho de Asis, A., Roberto, S.R., Bovolenta, Y.R., Nixdorf, S.N., García-Romero, E., Gómez-Alonso, S. & Hermosín-Gutiérrez, I., 2015. Application of abscisic acid (S-ABA) to cv. Isabel grapes (*Vitis vinifera* × *Vitis labrusca*) for color improvement: Effects on color, phenolic composition and antioxidant capacity of their grape juice. *Food Res. Int.* 77(3), 572-583.
- Yang, T., Peng, H., Whitaker, B.D. & Conway, W.S., 2012. Characterization of calcium/calmodulin-regulated SR/CAMTA gene family during tomato fruit development and ripening. *Plant Biol.* 12(19), 2-13.
- Zou, J.J., Li, X.D., Ratnasekera, D., Wang, C., Liu, W.X., Song, L.F., Zhang, W.Z. & Wu, W.H., 2015. Arabidopsis calcium-dependent protein kinases8 and catalase3 function in abscisic acid-mediated signaling and H₂O₂ homeostasis in stomatal guard cells under drought stress. *Plant Cell* 27(5), 1445-1460.