

# Effects of Cortisol and Dexamethasone on Insulin Signalling Pathways in Skeletal Muscle of the Ovine Fetus during Late Gestation

Juanita K. Jellyman<sup>1</sup>, Malgorzata S. Martin-Gronert<sup>2</sup>, Roselle L. Cripps<sup>2</sup>, Dino A. Giussani<sup>1</sup>, Susan E. Ozanne<sup>2</sup>, Qingwu W. Shen<sup>3</sup>, Min Du<sup>3</sup>, Abigail L. Fowden<sup>1</sup>, Alison J. Forhead<sup>1\*</sup>

**1** Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, United Kingdom, **2** Metabolic Research Laboratories, Institute of Metabolic Science, University of Cambridge, Cambridge, United Kingdom, **3** Department of Animal Science, University of Wyoming, Laramie, United States of America

## Abstract

Before birth, glucocorticoids retard growth, although the extent to which this is mediated by changes in insulin signalling pathways in the skeletal muscle of the fetus is unknown. The current study determined the effects of endogenous and synthetic glucocorticoid exposure on insulin signalling proteins in skeletal muscle of fetal sheep during late gestation. Experimental manipulation of fetal plasma glucocorticoid concentration was achieved by fetal cortisol infusion and maternal dexamethasone treatment. Cortisol infusion significantly increased muscle protein levels of Akt2 and phosphorylated Akt at Ser473, and decreased protein levels of phosphorylated forms of mTOR at Ser2448 and S6K at Thr389. Muscle GLUT4 protein expression was significantly higher in fetuses whose mothers were treated with dexamethasone compared to those treated with saline. There were no significant effects of glucocorticoid exposure on muscle protein abundance of IR- $\beta$ , IGF-1R, PKC $\zeta$ , Akt1, calpastatin or muscle glycogen content. The present study demonstrated that components of the insulin signalling pathway in skeletal muscle of the ovine fetus are influenced differentially by naturally occurring and synthetic glucocorticoids. These findings may provide a mechanism by which elevated concentrations of endogenous glucocorticoids retard fetal growth.

**Citation:** Jellyman JK, Martin-Gronert MS, Cripps RL, Giussani DA, Ozanne SE, et al. (2012) Effects of Cortisol and Dexamethasone on Insulin Signalling Pathways in Skeletal Muscle of the Ovine Fetus during Late Gestation. *PLoS ONE* 7(12): e52363. doi:10.1371/journal.pone.0052363

**Editor:** Josep V. Planas, Universitat de Barcelona, Spain

**Received:** November 7, 2011; **Accepted:** November 16, 2012; **Published:** December 26, 2012

**Copyright:** © 2012 Jellyman et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This study was funded by the Biotechnology and Biological Sciences Research Council and Tommy's the Baby Charity, UK. SEO is a British Heart Foundation Senior Fellow. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of this manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: [ajf1005@cam.ac.uk](mailto:ajf1005@cam.ac.uk)

## Introduction

In all species studied to date, there is an increase in the circulating glucocorticoid concentration in the fetus near term [1]. Glucocorticoids such as cortisol promote differentiation of specific cell types in fetal tissues at the expense of cell proliferation and growth. They are thereby responsible for both the normal decrease in fetal growth rate that occurs near term and the prepartum maturation of tissues essential for the successful transition from the intrauterine to the extrauterine environment at birth, including insulin-sensitive tissues such as skeletal muscle [1,2]. The maturational properties of glucocorticoids are beneficial in clinical practice where synthetic glucocorticoids such as dexamethasone are routinely administered to pregnant women at risk of preterm delivery in order to accelerate fetal maturation and improve survival and health outcomes of the premature neonate [3]. However, exposure of the fetus to antenatal glucocorticoid treatment, and endogenous glucocorticoids elevated in response to suboptimal intrauterine conditions, may impair fetal growth and alter insulin sensitivity and glucose metabolism in the adult animal [4,5,6].

The insulin receptor (IR) and insulin-like growth factor type 1 receptor (IGF-1R) are members of the family of receptor tyrosine kinases. When activated by insulin or IGFs, these kinases

phosphorylate insulin receptor substrates (IRS) which, in turn, lead to the activation of a number of cascades, including the phosphatidylinositol 3-kinase (PI3K) signalling pathway. This pathway can induce Akt phosphorylation which induces tissue-specific metabolic effects, such as translocation of the glucose transporter-4 (GLUT4) and consequently stimulation of glucose uptake in skeletal muscle and adipose tissue, activation of the peroxisome proliferator activated receptors involved in fat metabolism and storage, and stimulation of glycogen synthase activity and glycogen deposition in skeletal muscle and liver [7,8]. Activation of the PI3K pathway also leads to phosphorylation of mammalian target of rapamycin (mTOR) which controls the phosphorylation rate of proteins such as ribosomal protein S6 kinase (S6K) that in turn promotes protein translation [7,8]. Tissue mTOR expression thereby has an important role in the coordination of nutrient status, protein synthesis and tissue growth [9,10].

In fetal life, insulin has both glucoregulatory and anabolic actions from at least mid-gestation and developmental changes in the expression of insulin receptors and intracellular signalling proteins have been observed in fetal tissues, including skeletal muscle [11–14]. For example, the expression of GLUT4 in skeletal muscle of rats and baboons is lower in fetal compared to adult life and increases with gestational age [13,14]. However, little is

known about the regulation of insulin signalling pathways by glucocorticoids before birth. Therefore, the aim of the present study was to determine the effect of endogenous and synthetic glucocorticoids on the expression of insulin signalling proteins in skeletal muscle of fetal sheep. Glucocorticoid exposure *in utero* was hypothesised to decrease the abundance of proteins associated with growth, such as mTOR and S6K, and increase the proteins associated with glucose transport, such GLUT4. These changes may be responsible for the decline in fetal growth towards term and the preparation for greater dependence on insulin-stimulated glucose disposal after birth, respectively. Experimental manipulations of fetal plasma concentrations of glucocorticoid were achieved by fetal cortisol infusion and maternal dexamethasone treatment. These two experimental models enabled the impact of glucocorticoid exposure on insulin signalling proteins to be assessed.

## Methods

### Animals

Twenty-one Welsh Mountain sheep fetuses of known gestational age were used in this study; all were singleton fetuses (13 male and 8 female). The ewes were housed in individual pens and fed concentrates (200 g day<sup>-1</sup>; 18% protein and 10 MJ kg<sup>-1</sup>; Sheep Nuts #6; H&G Beart, Kings Lynn, UK) with free access to hay, water and a salt-lick block. Overall, the pregnant ewes consumed between 8–11 MJ day<sup>-1</sup> of metabolizable energy. Food, but not water, was withheld for 18–24 h before surgery. All surgical and experimental procedures were conducted in accordance with the UK Animals (Scientific Procedures) Act 1986 and were approved by the University of Cambridge animal ethics committee.

### Surgical and Experimental Procedures

**Fetal cortisol infusion.** Under halothane anaesthesia (1.5% in O<sub>2</sub>-N<sub>2</sub>O), catheters were implanted into the femoral artery and vein of eleven fetuses at 115–117 days of gestation using surgical methods described previously (term 145±2 days) [15]. Blood samples (2 ml) were taken daily throughout the experimental period from the catheterized fetuses to monitor fetal blood gas status. At least six days after catheterization, the fetuses were infused intravenously with either cortisol (2–3 mg kg<sup>-1</sup>day<sup>-1</sup> in 3.0 ml 0.9% saline; EF-Cortelan, GlaxoSmithKline, Brentford, Middlesex, UK; n = 5) or the saline vehicle (3.0 ml day<sup>-1</sup> 0.9% wt/vol; n = 6) for five days beginning at 125–126 days of gestation. The dose of cortisol was chosen to mimic the plasma concentrations normally observed in the immediate parturition period in fetal sheep [1].

**Maternal dexamethasone treatment.** In a separate cohort of animals, ten un-operated pregnant ewes were injected twice i.m. with either dexamethasone (12 mg dexamethasone sodium phosphate; Merck Sharpe and Dohme Ltd, Hoddesdon, Herts, UK; n = 5) in 2 ml saline or the saline vehicle (0.9% w/v NaCl; n = 5) at 24 hour intervals from 125 days of gestation. Tissues were collected 10 hours after the final maternal injection. The dexamethasone treatment regime resembles that recommended for use in human clinical practice [16].

### Blood and Tissue Collection

All fetuses were delivered by Caesarean section under general anaesthesia of the ewe (20 mg kg<sup>-1</sup> sodium pentobarbitone i.v.). At delivery, a 10 ml blood sample was taken by venepuncture of the umbilical artery, and a number of tissues were collected from the fetus after the administration of a lethal dose of barbiturate (200 mg kg<sup>-1</sup> sodium pentobarbitone). Skeletal muscle tissue

samples taken from the hind limb (biceps femoris) were immediately frozen in liquid nitrogen and stored at -80°C until analysis. All blood samples obtained at tissue collection were immediately placed into EDTA-containing tubes and centrifuged for 5 min at 1000 x g and 4°C. The plasma aliquots were stored at -20°C until analysis.

### Biochemical Analyses

Plasma cortisol concentration was measured in the umbilical arterial sample by radioimmunoassay validated for use with ovine plasma [17]. The lower limit of detection was 1.0–1.5 ng.ml<sup>-1</sup>, and the inter-assay coefficient of variation was 12%.

Muscle protein levels were quantified by Western blotting using methods described previously [18,19]. The antibodies used in this study were to IR-β subunit, IGF-1R, PKCζ, Akt1, Akt2, p-Akt Ser473, p-mTOR Ser2448, p-S6K Thr389, calpastatin and GLUT4 (Table 1), and have been used previously in this laboratory to quantify protein levels in the skeletal muscle of fetal sheep [20]. Horseradish peroxidase-conjugated secondary antibodies to the host species were used as appropriate and antibody binding was detected using an enhanced chemiluminescence kit (all products from GE Healthcare Life Sciences, Little Chalfont, Bucks, UK). Protein expression was quantified using Image J software (National Institutes of Health, Bethesda, MD; <http://rsb.info.nih.gov/ij/>).

The glycogen content of skeletal muscle was determined by a biochemical assay as described previously [21]. The inter-assay coefficient of variation was 9%.

### Statistical Analyses

Values are presented as mean values ± SEM. Data were analysed using Student's t-test. Ratios of muscle protein expression were arcsine transformed prior to statistical analysis, and are presented as a fold change from the mean values observed in the respective control group. For all comparisons, statistical significance was accepted when *P*<0.05.

## Results

### Fetal Cortisol Infusion

Plasma cortisol concentration was significantly higher in the cortisol-infused fetuses than in the saline-infused fetuses (*p*<0.05; Table 2). Fetal cortisol infusion for five days led to significant increases in the muscle protein level of p-Akt Ser473 and the expression of Akt2 (*p*<0.05), but not Akt1 (Figure 1A, B and C), and decreases in the protein expression of p-mTOR Ser2448 and p-S6K Thr389 (*p*<0.05; Figure 2A and B). There were no significant differences in muscle protein abundance of IR-β, IGF-1R, PKCζ, calpastatin (Table 2) or GLUT4 (Figure 3B), or muscle glycogen content (Table 2), between the saline and cortisol-treated fetuses.

### Maternal Dexamethasone Treatment

At delivery, plasma cortisol concentration was significantly lower in fetuses whose mothers were injected with dexamethasone compared to those whose mothers were treated with saline (*p*<0.05; Table 2). Exposure of fetuses to dexamethasone caused a significant increase in muscle GLUT4 protein content (*p*<0.05; Figure 3B). However, fetal muscle protein levels of IR-β, IGF-1R, PKCζ, calpastatin (Table 2), Akt 1, Akt2 and p-Akt Ser473 (Figure 1D, E and F), p-mTOR Ser2448 and p-S6K Thr389 (Figure 2C and D), and muscle glycogen content (Table 2) were not significantly different between the saline and dexamethasone-treated animals.

**Table 1.** Details of the antibodies used in Western blotting.


| Protein        | Antibody           | Dilution | Source   | Catalogue Number |
|----------------|--------------------|----------|--|------------------|
| IR-β           | Rabbit polyclonal  | 1:200    | Santa Cruz Biotechnologies, Santa Cruz, CA, USA  | sc711            |
| IGF-1R         | Rabbit polyclonal  | 1:200    | As above.  | sc713            |
| PKCζ           | Rabbit polyclonal  | 1:500    | As above.  | sc216            |
| p-Akt Ser473   | Rabbit monoclonal  | 1:1000   | Cell Signaling Technology, Inc, Beverly, MA, USA | 4058             |
| Akt1           | Mouse monoclonal   | 1:1000   | As above.  | 2967             |
| Akt2           | Rabbit monoclonal  | 1:1000   | As above.  | 3063             |
| p-mTOR Ser2448 | Rabbit polyclonal  | 1:1000   | As above.  | 2971             |
| p-S6K Thr389   | Rabbit polyclonal  | 1:1000   | As above.  | 9205             |
| Calpastatin    | Chicken polyclonal | 1:1000   | Abcam, Cambridge, UK                             | ab16423          |
| GLUT4          | Rabbit polyclonal  | 1:4000   | As above.  | ab654            |

doi:10.1371/journal.pone.0052363.t001


**Discussion**

The results demonstrate that components of the insulin signalling pathways relating to both growth and glucoregulation in skeletal muscle of the ovine fetus are sensitive to glucocorticoid exposure *in utero*. In agreement with the hypothesis of the study, fetal infusion of cortisol led to downregulation of the phosphorylated levels of the anabolic proteins, mTOR and S6K, while maternal dexamethasone treatment increased GLUT4 expression in fetal skeletal muscle.

These findings contribute to the understanding of the regulation of growth by glucocorticoids *in utero*. In fetal sheep, the decline in growth rate seen near term is known to be glucocorticoid-dependent, since it can be prevented by removal of the fetal adrenal gland and induced prematurely by five days of cortisol infusion [2]. Reductions in the phosphorylated levels of mTOR and S6K induced by cortisol in skeletal muscle of the ovine fetus may be one mechanism by which glucocorticoids retard muscle protein synthesis and overall fetal growth. Indeed, glucocorticoids have been shown previously to decrease S6K phosphorylation and



**Figure 1.** Mean (± SEM) protein levels of (A, D) p-Akt, (B, E) Akt1 and (C, F) Akt2 in skeletal muscle of fetuses infused i.v. with either saline or cortisol for 5 days, and fetuses whose mothers were injected i.m. with either saline or dexamethasone. \*, significantly different from saline control group, p<0.05.  
doi:10.1371/journal.pone.0052363.g001



**Figure 2. Mean ( $\pm$  SEM) protein levels of (A, C) p-mTOR and (B, d) p-S6K in skeletal muscle of fetuses infused i.v. with either saline or cortisol for 5 days, and fetuses whose mothers were injected i.m. with either saline or dexamethasone. \*, significantly different from saline control group,  $p < 0.05$ .**

doi:10.1371/journal.pone.0052363.g002

protein synthesis in cultured L6 myoblasts and in rat skeletal muscle *in vivo* [22,23].

In the present study, reductions in muscle p-mTOR and p-S6K in the cortisol-infused fetuses were associated with increased Akt2 and p-Akt protein content. This finding was unexpected given that the PI3K/Akt pathway is a key activator of mTOR phosphory-


lation in adult tissues [7,8]. It is possible that the relative levels of phosphorylation of Akt at Ser473 and Thr308 may change in response to glucocorticoid treatment with consequences for the specificity of downstream targets. Consistent with this possibility, a recent study in the Jeg-3 human choriocarcinoma cell line has shown that an increase in Akt phosphorylation at Ser473, induced

**Table 2. Mean ( $\pm$  SEM) plasma concentrations of cortisol, muscle protein levels of IR- $\beta$ , IGF-1R, PKC $\zeta$  and calpastatin, and muscle glycogen content in fetuses infused i.v. with either saline or cortisol for 5 days, and in fetuses whose mothers were injected i.m. with either saline or dexamethasone.**

|  | Fetal administration |                  | Maternal administration |                  |
|--|----------------------|------------------|-------------------------|------------------|
|  | Saline               | Cortisol         | Saline                  | Dexamethasone    |
| Gestational age (days)                       | 130 $\pm$ 0          | 131 $\pm$ 0.2    | 127 $\pm$ 0             | 127 $\pm$ 0      |
| Plasma cortisol (ng.ml <sup>-1</sup> )       | 14.1 $\pm$ 2.0       | 91.2 $\pm$ 20.9* | 16.1 $\pm$ 2.8          | 8.6 $\pm$ 0.7*   |
| Muscle IR- $\beta$ (au)                      | 1.00 $\pm$ 0.20      | 1.11 $\pm$ 0.10  | 1.00 $\pm$ 0.06         | 0.88 $\pm$ 0.06  |
| Muscle IGF-1R (au)                           | 1.00 $\pm$ 0.21      | 1.06 $\pm$ 0.13  | 1.00 $\pm$ 0.16         | 0.84 $\pm$ 0.13  |
| Muscle PKC $\zeta$ (au)                      | 1.00 $\pm$ 0.09      | 0.86 $\pm$ 0.06  | 1.00 $\pm$ 0.03         | 1.02 $\pm$ 0.04  |
| Muscle calpastatin (au)                      | 1.00 $\pm$ 0.09      | 0.83 $\pm$ 0.10  | 1.00 $\pm$ 0.04         | 1.07 $\pm$ 0.13  |
| Muscle glycogen (mg.g tissue <sup>-1</sup> ) | 46.27 $\pm$ 3.72     | 48.38 $\pm$ 2.21 | 48.70 $\pm$ 1.95        | 50.51 $\pm$ 3.14 |

\*, significantly different from the respective saline control group, Student's t-test,  $p < 0.05$ . IR- $\beta$ , insulin receptor  $\beta$ -subunit; IGF-1R, insulin-like growth factor-1 receptor; PKC $\zeta$ , protein kinase C  $\zeta$ ; au, arbitrary units.

doi:10.1371/journal.pone.0052363.t002



**Figure 3. Mean ( $\pm$  SEM) GLUT4 protein level in skeletal muscle of (a) fetuses infused i.v. with either saline or cortisol for 5 days, and (b) fetuses whose mothers were injected i.m. with either saline or dexamethasone. \***, significantly different from saline control group,  $p < 0.05$ .

doi:10.1371/journal.pone.0052363.g003

by endoplasmic reticulum stress, decreases phosphorylation of mTOR at Ser2448 while increasing phosphorylation of other protein targets [24]. In addition, cortisol may influence p-mTOR and p-S6K expression independently of the PI3K/Akt pathway and/or downstream of these proteins in fetal skeletal muscle. For example, glucocorticoids have been shown to suppress mTOR levels, at least in part, by activation of the translation repressor protein REDD1 in cultured L6 myoblasts and in rat skeletal muscle *in vivo* [25]. There is also evidence to suggest that mTOR has negative feedback effects on Akt phosphorylation. Inhibition of mTOR using rapamycin or its derivatives upregulates Ser473 phosphorylation of Akt in cancer cell lines and human tumours [26,27]. Of the three isoforms of Akt, Akt2 is highly expressed in insulin-sensitive tissues, such as skeletal muscle, and previously, increased Akt2 expression has been associated with increased myoblast differentiation *in vitro* [28,29]. In the present study, however, the tissues were not examined histologically and the effect of cortisol exposure on the structural development of the biceps femoris *in utero* remains unclear.

Muscle GLUT4 protein expression in the ovine fetus was increased by maternal dexamethasone treatment. This observation confirms that previously reported in fetuses of pregnant ewes treated with either single or repeated doses of dexamethasone at an earlier gestational age (106–107 days) [30]. Dexamethasone has also been shown to increase GLUT4 protein levels in cultured L6 myoblasts and in rat skeletal muscle *in vivo* [31,32]. Upregulation

of muscle GLUT4 in the sheep fetus by maternal dexamethasone treatment occurred without any effect on other components of the insulin signalling pathway, such as p-Akt and PKC $\zeta$ . Muscle GLUT4 content in the fetus may be influenced by insulin-independent as well as insulin-dependent mechanisms. For example, in fetal sheep, both experimental conditions of hyperglycaemia-euinsulinaemia and hyperinsulinaemia-euglycaemia increase GLUT4 protein expression in skeletal muscle in a time-dependent manner [33]. Despite the increase in muscle GLUT4 protein level observed in the sheep fetuses exposed to dexamethasone, there were no changes in muscle glycogen although glycogen deposition also depends on intracellular glucose availability, as well as the activities of glycogen synthase and phosphorylase enzymes which were not determined in the present study. Furthermore, the extent to which GLUT4 translocation to the muscle plasma membrane is affected by glucocorticoids *in utero* remains unknown.

Cortisol and dexamethasone exerted different effects on muscle insulin signalling proteins in the fetuses of the two experimental models of intrauterine glucocorticoid exposure. There are several possible explanations for these observations. First, the experimental models differ in the duration and level of glucocorticoid exposure and timing of tissue collection. In the dexamethasone-treated animals, dexamethasone concentrations in the fetal circulation are significantly elevated 1–2 h post-injection and, although still detectable, fall thereafter at tissue and blood collection at 10 hours after the second maternal injection [34], whereas cortisol-treated fetuses were exposed to elevated and stable levels of glucocorticoid for five days prior to tissue collection. In both models, there may have been dynamic changes in insulin signalling protein expression over the time course of each treatment.

Second, natural and synthetic glucocorticoids differ in their properties of action. The synthetic glucocorticoids have longer half-lives and are more potent than the endogenous glucocorticoids, and, unlike the endogenous glucocorticoids that act on both glucocorticoid and mineralocorticoid receptors, synthetic glucocorticoids only bind and activate glucocorticoid receptors. The relative effects of activation of mineralocorticoid receptors by cortisol on insulin signalling pathways in fetal tissues, however, are unknown. The fall in plasma cortisol seen in the dexamethasone-exposed fetuses is in accordance with the negative feedback effects of synthetic glucocorticoids on endogenous glucocorticoid production and has been reported previously in the fetuses of pregnant ewes treated with dexamethasone [21].

Third, some of the effects of glucocorticoid exposure on insulin signalling protein expression in skeletal muscle *in utero* may be due to concomitant changes in other circulating and/or local hormones and growth factors. In both of these fetal sheep models, increases in plasma triiodothyronine and leptin have been demonstrated in response to fetal cortisol infusion and maternal dexamethasone treatment [35,36,37]. Although components of the insulin signalling pathway in fetal skeletal muscle do not appear to be influenced by leptin treatment *in utero* [20], the effects of triiodothyronine are unknown. In addition, previous studies have shown that maternal dexamethasone treatment in pregnant ewes causes increases in circulating glucose and insulin in the fetus [21], while no changes in plasma glucose and insulin are observed after five days of fetal cortisol infusion [2]. The insulin response to dexamethasone may negate some of the direct actions of the glucocorticoid on the insulin signalling proteins as insulin promotes expression of anabolic proteins, such as S6K, in skeletal muscle of the ovine fetus [38]. Furthermore, in fetal sheep, both glucose and insulin are able to upregulate muscle GLUT4 protein

expression [33] and, therefore, the dexamethasone-induced increments in plasma glucose and insulin are likely to contribute to the increase in muscle GLUT4 seen in the present study.

Suppression of mTOR and S6K in skeletal muscle of cortisol-treated sheep fetuses may also be due, at least in part, to local changes in IGF synthesis. Previous studies have shown that cortisol infusion for five days in fetal sheep downregulates both IGFI and IGFI mRNA abundance in skeletal muscle [39,40] which may reduce the anabolic actions of these paracrine agents in the absence of any change in muscle IGF-1R expression. Muscle IGF expression has not been measured in the experimental model of dexamethasone exposure used in the present study, although in other studies, synthetic glucocorticoid treatment in the pregnant ewe has been shown to reduce circulating IGFI and IGF binding proteins in the fetus only after repeated, but not single, doses [41]. Furthermore, in fetal sheep and human infants exposed to synthetic glucocorticoids *in utero*, multiple doses lead to impaired body size at birth while a single dose often has little effect on growth [42,43].

In summary, the current study demonstrated that endogenous and synthetic glucocorticoids alter components of the insulin signalling pathways in skeletal muscle of the ovine fetus in different ways, with potentially different consequences for the regulation of muscle growth in the fetus and insulin sensitivity in postnatal life.

## References

- Fowden AL, Li J, Forhead AJ (1998) Glucocorticoids and the preparation for life after birth: are there long-term consequences of the life insurance? *Proc Nutr Soc* 57: 113–122.
- Fowden AL, Szemere J, Hughes P, Gilmour RS, Forhead AJ (1996) The effects of cortisol on the growth rate of the sheep fetus during late gestation. *J Endocrinol* 151: 97–105.
- Roberts D, Dalziel SR (2006) Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Db Syst Rev* CD004454.
- Challis JR, Sloboda D, Matthews SG, Holloway A, Alfaidy N, et al. (2001) The fetal placental hypothalamic-pituitary-adrenal (HPA) axis, parturition and post natal health. *Mol Cell Endocrinol* 185: 135–144.
- Sloboda DM, Moss TJM, Li S, Doherty DA, Nitsos I, et al. (2005) Hepatic glucose regulation and metabolism in adult sheep: effects of prenatal betamethasone. *Am J Physiol* 289: E721–E728.
- Kelly BA, Lewandowski AJ, Worton SA, Davis EF, Lazdam M, et al. (2012) Antenatal glucocorticoid exposure and long-term alterations in aortic function and glucose metabolism. *Pediatrics* 129: e1282–1290.
- Franke TF (2008) PI3K/Akt: getting it right matters. *Oncogene* 27: 6473–6488.
- Wu M, Falasca M, Blough ER (2010) Akt/protein kinase B in skeletal muscle physiology and pathology. *J Cell Physiol* 226: 29–36.
- Schmelzle T, Hall MN (2000) TOR, a central controller of cell growth. *Cell* 103: 253–262.
- Sengupta S, Peterson TR, Sabatini DM (2010) Regulation of the mTOR complex 1 pathway by nutrients, growth factors, and stress. *Mol Cell* 40: 310–322.
- Molina RD, Carver TD, Hay WW (1993) Ontogeny of insulin effect in fetal sheep. *Pediatr Res* 34: 654–660.
- Shen W, Wisniewski P, Ahmed L, Boyle DW, Denne SC, et al. (2003) Protein anabolic effects of insulin and IGF-I in the ovine fetus. *Am J Physiol* 284: E748–E756.
- Blanco CL, Liang H, Joya-Galeana J, DeFronzo RA, McCurmin D, et al. (2010) The ontogeny of insulin signaling in the preterm baboon model. *Endocrinology* 151: 1990–1997.
- Santalucia T, Camps M, Castelló A, Muñoz P, Nuel A, et al. (1992) Developmental regulation of GLUT-1 (erythroid/Hep G2) and GLUT-4 (muscle/fat) glucose transporter expression in rat heart, skeletal muscle, and brown adipose tissue. *Endocrinology* 130: 837–846.
- Comline RS, Silver M (1972) The composition of foetal and maternal blood during parturition in the ewe. *J Physiol* 222: 233–256.
- Royal College of Obstetricians and Gynaecologists (2010) Antenatal corticosteroids to prevent neonatal morbidity and mortality. Green-top guideline No.7.
- Robinson PM, Comline RS, Fowden AL, Silver M (1983) Adrenal cortex of fetal lamb: changes after hypophysectomy and effects of Synacthen on cytoarchitecture and secretory activity. *Quart J Exp Physiol* 68: 15–27.
- Ozanne SE, Nave BT, Wang CL, Shepherd PR, Prins J, et al. (1997) Poor fetal nutrition causes long-term changes in expression of insulin signaling components in adipocytes. *Am J Physiol* 273: E46–E51.
- Zhu MJ, Ford SP, Nathanielsz PW, Du M (2004) Effect of maternal nutrient restriction in sheep on the development of fetal skeletal muscle. *Biol Reprod* 71: 1968–1973.
- Forhead AJ, Lamb CA, Franko KL, O'Connor DM, Wooding FB, et al. (2008) Role of leptin in the regulation of growth and carbohydrate metabolism in the ovine fetus during late gestation. *J Physiol* 586: 2393–2403.
- Franko KL, Giussani DA, Forhead AJ, Fowden AL (2007) Effects of dexamethasone on the glucogenic capacity of fetal, pregnant, and non-pregnant adult sheep. *J Endocrinol* 192: 67–73.
- Shah OJ, Kimball SR, Jefferson LS (2000a) Acute attenuation of translation initiation and protein synthesis by glucocorticoids in skeletal muscle. *Am J Physiol* 278: E76–E82.
- Shah OJ, Kimball SR, Jefferson LS (2000b) Glucocorticoids abate p70(S6k) and eIF4E function in L6 skeletal myoblasts. *Am J Physiol* 279: E74–E82.
- Yung HW, Charnock-Jones DS, Burton GJ (2011) Regulation of AKT phosphorylation at Ser473 and Thr308 by endoplasmic reticulum stress modulates substrate specificity in a severity dependent manner. *PLoS One* 6(3): e17894.
- Wang H, Kubica N, Ellisen LW, Jefferson LS, Kimball SR (2006) Dexamethasone represses signaling through the mammalian target of rapamycin in muscle cells by enhancing expression of REDD1. *J Biol Chem* 281: 39128–39134.
- Sun SY, Rosenberg LM, Wang X, Zhou Z, Yue P, et al. (2005) Activation of Akt and eIF4E survival pathways by rapamycin-mediated mammalian target of rapamycin inhibition. *Cancer Res* 65: 7052–7058.
- O'Reilly KE, Rojo F, She QB, Solit D, Mills GB, et al. (2006) mTOR inhibition induces upstream receptor tyrosine kinase signaling and activates Akt. *Cancer Res* 66: 1500–1508.
- Vandromme M, Rochat A, Meier R, Carnac G, Besser D, et al. (2001) Protein kinase B beta/Akt2 plays a specific role in muscle differentiation. *J Biol Chem* 276: 8173–8179.
- Heron-Milhavet L, Khouya N, Fernandez A, Lamb NJ (2011) Akt1 and Akt2: differentiating the aktion. *Histol Histopathol* 26: 651–662.
- Gray S, Stonestreet BS, Thamotharan S, Sadowska GB, Daood M, et al. (2006) Skeletal muscle glucose transporter protein responses to antenatal glucocorticoids in the ovine fetus. *J Endocrinol* 189: 219–229.
- Ewart HS, Somwar R, Klip A (1998) Dexamethasone stimulates the expression of GLUT1 and GLUT4 proteins via different signalling pathways in L6 skeletal muscle cells. *FEBS Lett* 425: 179–183.
- Coderre L, Vallega GA, Pilch PF, Chipkin SR (1996) In vivo effects of dexamethasone and sucrose on glucose transport (GLUT-4) protein tissue distribution. *Am J Physiol* 271: E643–E648.
- Anderson MS, He J, Flowers-Ziegler J, Devaskar SU, Hay WW Jr (2001) Effects of selective hyperglycemia and hyperinsulinemia on glucose transporters in fetal ovine skeletal muscle. *Am J Physiol* 281: R1256–R1263.
- Jellyman JK, Gardner DS, McGarrigle HH, Fowden AL, Giussani DA (2009) Antenatal glucocorticoid therapy increases glucose delivery to cerebral circulations during acute hypoxemia in fetal sheep during late gestation. *Am J Obstet Gynecol* 201: 82.e1–8.

## Acknowledgments

The authors thank all the members of technical staff in the Department of Physiology, Development and Neuroscience, University of Cambridge who cared for the animals.

## Author Contributions

Conceived and designed the experiments: JKJ DAG ALF AJF. Performed the experiments: JKJ MSM-G RLC DAG SEO QWS MD ALF AJF. Analyzed the data: JKJ AJF. Contributed reagents/materials/analysis tools: DAG SEO MD ALF AJF. Wrote the paper: JKJ AJF.

35. Forhead AJ, Curtis K, Kaptein E, Visser TJ, Fowden AL (2006) Developmental control of iodothyronine deiodinases by cortisol in the ovine fetus and placenta near term. *Endocrinology* 147: 5988–5994.
36. Forhead AJ, Jellyman JK, Gardner DS, Giussani DA, Kaptein E, et al. (2007) Differential effects of maternal dexamethasone treatment on circulating thyroid hormone concentrations and tissue deiodinase activity in the pregnant ewe and fetus. *Endocrinology* 148: 800–805.
37. O'Connor DM, Blache D, Hoggard N, Brookes E, Wooding, et al. (2007) Developmental control of plasma leptin and adipose leptin messenger ribonucleic acid in the ovine fetus during late gestation: role of glucocorticoids and thyroid hormones. *Endocrinology* 148: 3750–3757.
38. Brown LD, Rozance PJ, Barry JS, Friedman JE, Hay WW (2009) Insulin is required for amino acid stimulation of dual pathways for translational control in skeletal muscle in the late-gestation ovine fetus. *Am J Physiol* 296: E56–E63.
39. Li J, Saunders JC, Gilmour RS, Silver M, Fowden AL (1993) Insulin-like growth factor-II messenger ribonucleic acid expression in fetal tissues of the sheep during late gestation: effects of cortisol. *Endocrinology* 132: 2083–2089.
40. Li J, Forhead AJ, Dauncey MJ, Gilmour RS, Fowden AL (2002) Control of growth hormone receptor and insulin-like growth factor-I expression by cortisol in ovine fetal skeletal muscle. *J Physiol* 541: 581–589.
41. Gatford KL, Owens JA, Moss TJ, Newnham JP, Challis, et al. (2008) Repeated betamethasone treatment of pregnant sheep programs persistent reductions in circulating IGF-I and IGF-binding proteins in progeny. *Am J Physiol* 295: E170–E178.
42. Moss TJM, Sloboda DM, Gurrin LC, Harding R, Challis JRG, et al. (2001) Programming effects in sheep of prenatal growth restriction and glucocorticoid exposure. *Am J Physiol* 281: R960–R970.
43. French NP, Hagan R, Evans SF, Godfrey M, Newnham JP (1999) Repeated antenatal corticosteroids: Size at birth and subsequent development. *Am J Obstet Gynecol* 180: 114–121.