

## **Managing shoulder pain in manual wheelchair users: A scoping review of conservative treatment interventions**

**Mason BS,<sup>1</sup> Warner MB,<sup>2,3</sup> Briley S,<sup>1</sup> Goosey-Tolfrey VL,<sup>1</sup> Vegter RJK.<sup>4</sup>**

<sup>1</sup> Peter Harrison Centre for Disability Sport, School for Sport, Exercise & Health Sciences, Loughborough University, Loughborough, UK.

<sup>2</sup> School of Health Sciences, University of Southampton, UK.

<sup>3</sup> Arthritis Research UK Centre for Sport, Exercise and Osteoarthritis.

<sup>4</sup> Faculty of Medical Sciences, University of Groningen, The Netherlands.

### **Corresponding author:**

**Dr Barry Mason**, Peter Harrison Centre for Disability Sport, School for Sport, Exercise & Health Sciences, Loughborough University, Loughborough, UK.

**Email:** [b.mason@lboro.ac.uk](mailto:b.mason@lboro.ac.uk)

## **ABSTRACT**

**Objective:** To review the literature that has explored conservative treatments for the management of shoulder pain in manual wheelchair users.

**Methods:** Five databases were systematically searched in February 2020 for terms related to shoulder pain and manual wheelchair use. Articles were screened and included if they investigated the conservative treatment of shoulder pain in wheelchair users. Participants' physical characteristics, experimental design and primary and secondary outcome measures were extracted from studies. Studies were grouped according to treatment type to identify gaps in the literature and guide future research.

**Results:** The initial search identified 407 articles, of which 21 studies met the inclusion criteria. Exercise-based treatment interventions were most prevalent (n=12). A variety of exercise modalities were employed such as strengthening and stretching (n=7), ergometer training (n=3), Pilates classes (n=1) and functional electrical stimulation (n=1). Only 3 studies supplemented exercise with an additional treatment type. The Wheelchair Users Shoulder Pain Index was used by 18 studies as the primary measure of shoulder pain. Only 7 of these included an objective measure of shoulder function. Participant characteristics varied amongst studies and physical activity levels were frequently not reported.

**Conclusions.** Despite the high prevalence of shoulder pain in manual wheelchair users, the number of studies to have explored conservative treatment types is low. Exercise is the most commonly used treatment, which is encouraging as physical inactivity can exacerbate other health conditions. Few studies have adopted interdisciplinary treatment strategies or included objective secondary measures to better understand the mechanisms of pain.

## 1 INTRODUCTION

2 Manual wheelchair use places considerable stress on the upper limbs, particularly the  
3 shoulder, due to the repetitive loading induced by wheelchair propulsion in addition to other  
4 activities of daily living, such as transferring and weight relief tasks. Given the limited muscle  
5 mass and low stability, yet high mobility of the shoulder girdle,<sup>1</sup> these activities often lead to  
6 pain, with up to 71% of manual wheelchair users reported to have experienced shoulder pain  
7 at some point in their life.<sup>2,3,4</sup>

8 The most common pathologies associated with shoulder pain are shoulder impingement  
9 syndrome, rotator cuff tears and tendinopathy, bursitis, joint oedema and glenohumeral  
10 instability.<sup>5-7</sup> The consequences of such pathologies can be incredibly severe for wheelchair  
11 users, as it may prevent individuals from being physically active, which can negatively affect  
12 their independence and quality of life.<sup>8,9</sup> This lack of physical activity can also lead to  
13 secondary health conditions such as obesity and cardiovascular disease.<sup>10</sup> Structural changes  
14 as a result of injury within the shoulder may also develop into chronic conditions such as  
15 osteoarthritis, where joint degeneration can take place and may ultimately require shoulder  
16 arthroplasty to repair.<sup>11</sup> Such invasive, surgical techniques are not without risk and should be  
17 considered a last resort given the prolonged post-operative immobilisation imposed.<sup>12</sup>

18 A variety of conservative treatment options are available as an alternative to surgery  
19 for the management of shoulder pain, including exercise, massage, ultrasound, electrical nerve  
20 stimulation, neuromuscular retraining and corticosteroid injections.<sup>13</sup> Conservative treatment  
21 has shown to have beneficial effects on shoulder pain in non-wheelchair users, however,  
22 evidence is rated as low quality.<sup>20</sup> In addition, it cannot be assumed that treatments for non-  
23 wheelchair users will also be appropriate for wheelchair users due to differences in upper and  
24 lower limb function, perceptions of pain and tasks of everyday life that might be affected by

25 shoulder pain. A systematic review on treatment options for wheelchair users found positive  
26 outcomes on shoulder pain following conservative treatment.<sup>14</sup> However, this review only  
27 explored the effectiveness of exercise-based treatments and concluded that exercise was  
28 important for managing shoulder pain without being able to offer suggestions on type,  
29 frequency or duration of exercise. Considering the varied nature and range of conservative  
30 treatments available, it is important to consider all options in addition to exercise to help  
31 determine the most appropriate treatment. Subsequently, the aim of the current scoping review  
32 was to map the existing literature that has explored conservative, non-invasive solutions for the  
33 treatment of shoulder pain in manual wheelchair users to identify gaps in the evidence-base  
34 and to direct future research in this area.

35

## 36 **METHODS**

37 The scoping review was conducted according to previously developed guidelines.<sup>15,16</sup> The  
38 selection process of identification, screening, eligibility and inclusion was performed in  
39 accordance to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses  
40 (PRISMA) guidelines for scoping reviews.<sup>17</sup>

### 41 **Data Sources and Systematic Search**

42 An initial search of relevant databases (MEDLINE, PubMed, PsychINFO, SPORTDiscus and  
43 Web of Science) was performed using ‘shoulder’ AND ‘pain’ AND ‘wheelchair’ as the search  
44 terms. Having reviewed the abstracts of the studies identified by this initial search, it was  
45 decided that the terms ‘pathology’ (patholog\*) and ‘injury’ (injur\*) were also added to the  
46 search. The search was conducted in March 2020 using the aforementioned databases to  
47 identify studies published up until the end of February 2020. The reference lists of suitable

48 studies and review papers identified by the search were also examined to identify any additional  
49 records.

## 50 **Study Selection**

51 The following inclusion / exclusion criteria were applied to determine the eligibility of the  
52 identified articles, developed by BM, RV and MW:

### 53 *Inclusion criteria*

- 54 • Manual wheelchair users with shoulder pain
- 55 • All ages, genders, health conditions and activity levels
- 56 • Research design must include a conservative treatment intervention – either  
57 longitudinal or within-subject measures

### 58 *Exclusion criteria*

- 59 • Case reports or review articles
- 60 • Not available in English
- 61 • Involve invasive/surgical procedures

62

63 Studies identified by the search strategy were imported into Mendeley reference  
64 management software where any duplicate articles were removed. The titles and abstracts of  
65 all studies were reviewed by one author (BM) and evaluated against the eligibility criteria. A  
66 second reviewer (SB) performed the same process on a random sample of 25% of the articles,  
67 with a concordance of 98% between included and excluded articles. Where an agreement was  
68 not reached, the article proceeded to full-text review where all articles were examined by two  
69 authors independently (BM & MW). The level of agreement between the two authors after the

70 first review was 96%. Articles that resulted in a disagreement were then revisited and resolved  
71 by direct communication between authors.

72

### 73 **Data Extraction and Synthesis**

74 A database was developed in Microsoft Excel to document and assimilate extracted data from  
75 all included studies. Database design was agreed by BM, RV and MW and the list of extraction  
76 categories is detailed below:

- 77 i) Author(s);
- 78 ii) Year of publication;
- 79 iii) Purpose;
- 80 iv) Population characteristics (age, disability, years of manual wheelchair use, physical  
81 activity) and sample size;
- 82 v) Methodology and design
- 83 vi) Type of intervention;
- 84 vii) Duration of the intervention;
- 85 viii) Outcome measures;

86 Two authors (BM & MW) then extracted data from 10 different articles each. An  
87 independent reviewer (SB) then checked 20% of both authors extractions for accuracy. Studies  
88 were then grouped and reported according to the type of intervention performed.

89

## 90 **RESULTS**

91 Of the 407 articles identified by the initial search, a total of 21 studies met the inclusion criteria  
92 (Figure 1). Studies were categorised according to the type of conservative treatment

93 intervention. The most common treatment intervention was exercise-based (Table 1), which  
94 formed 12/21 of the studies included.<sup>18-29</sup> Home-based strengthening and stretching  
95 programmes were the most common modality of exercise prescribed (7/12 studies).  
96 Cardiovascular ergometer training was prescribed by 3 studies.<sup>20,21,25</sup> Other studies explored  
97 strengthening and stretching in the form of supervised Pilates classes<sup>26</sup> and functional electrical  
98 stimulation assisted rowing.<sup>28</sup> Remaining studies were categorised as therapeutic-based  
99 (3/21),<sup>30-32</sup> which included acupuncture, Trager Psychophysical Integration and transdermal  
100 nitroglycerine patches, equipment-based (1/21),<sup>33</sup> and educational interventions (2/21),<sup>34,35</sup> or  
101 interventions associated with lifestyle (3/21) assistance<sup>36-38</sup> (Table 2). The majority of  
102 interventions were monodisciplinary. An interdisciplinary treatment approach was adopted by  
103 only 3 studies, where exercise was accompanied by either movement retraining or real-time  
104 electromyographical biofeedback.<sup>22-24</sup>

105

106

\*\*\*FIGURE 1\*\*\*

107

108

\*\*\*TABLE 1 & 2\*\*\*

109

110 Sample sizes ranged from as little as 7 participants<sup>21</sup> to as many as 66 participants.<sup>37</sup>

111 The age range of participants was quite spread, yet similar across studies. Manual wheelchair

112 users with a wide range of health conditions were included in the studies, including individuals

113 with both paraplegia and tetraplegia as well as amputations and neuromuscular impairments.

114 Years' experience of manual wheelchair use was also quite spread, although similar across

115 studies, yet not reported by all. The physical activity levels of participants was only provided

116 by 5 studies and the level of detailed was limited where only hours per week were typically  
117 reported.

118 Of the included studies, 11 adopted an experimental study design, of which 8 were  
119 randomised control trials and 3 were quasi-experimental. The remaining 10 studies were  
120 observational prospective cohort studies. Interventions lasted from as little as 6 weeks up to as  
121 much as 12 months. All but 3 studies<sup>26,29,34</sup> measured shoulder pain according to the Wheelchair  
122 Users Shoulder Pain Index, of which 7 reported a performance corrected version of this  
123 questionnaire.<sup>18,19,24,30,31,33,35</sup> Only 9 studies included an objective measure of shoulder  
124 function, such as strength, range of movement and muscular activity.

125

## 126 **DISCUSSION**

127 The current systematic scoping review revealed that a total of 21 studies have investigated  
128 conservative treatment interventions for managing shoulder pain in wheelchair users. This is  
129 considerably lower than a similar review conducted in non-wheelchair users, where 177 studies  
130 were identified.<sup>13</sup> This illustrates the paucity of research specific to manual wheelchair users  
131 and highlights the need for an increase in well-designed studies investigating the conservative  
132 treatment of shoulder pain, given the high prevalence within this population.<sup>2-4</sup>

### 133 *Treatments*

134 Exercise-based interventions were the most popular type of treatment. The majority  
135 involved a programme of strengthening and stretching exercises using elastic training bands or  
136 weights.<sup>18,19,21-24,27,29</sup> Arm-crank<sup>20,21</sup> and double-poling<sup>25</sup> ergometry interventions were also  
137 trialled, in addition to rowing assisted with functional electrical stimulation as additional means  
138 for strengthening rotator cuff muscles.<sup>28</sup> One study used an alternative approach to reducing  
139 shoulder pain by focusing less on the shoulders and more on core strengthening through a



140 Pilates exercise programme.<sup>26</sup> The structure and supervision provided by exercise classes, such  
141 as Pilates, could prove to be a topic worthy of future investigation due to issues around  
142 adherence in home-based exercise programmes. Activity logs implemented by two studies  
143 noted that good adherence (>75% of all sessions completed) was only reported in 36% to 73%  
144 of participants during home-based exercise programmes.<sup>19,27</sup> Programme duration (6 weeks to  
145 6 months) and frequency of exercise (daily to 3 times/week) also varied amongst studies.  
146 Subsequently further work is required to determine not only the optimal type, but also the  
147 dosage of exercise prescribed when attempting to reduce shoulder pain.

148         Aside from exercise, therapeutic interventions were the second most popular choice of  
149 treatment within the scientific literature, although only three such studies were performed.<sup>30-32</sup>  
150 These studies explored the use of acupuncture,<sup>30,31</sup> Trager Psychophysical Integration,<sup>30</sup> and  
151 transdermal nitroglycerine patches.<sup>32</sup> Acupuncture refers to the insertion of fine needles into  
152 specific locations around the body to correct energy flow imbalances thought to lead to pain  
153 and illness.<sup>30</sup> Trager Psychophysical Integration is a technique that involves hands-on  
154 manipulation and movement re-education, anecdotally thought to minimise joint pain and  
155 improve mobility in individuals with a musculoskeletal disorder.<sup>30</sup> Finally, Transdermal  
156 nitroglycerine patches emit nitroglycerine through the skin, which is transformed into nitric  
157 oxide in the bloodstream and has been reported to be advantageous for the repair and  
158 regeneration of damaged tendons.<sup>39,40</sup> However, detrimental side effects, such as headaches,  
159 were frequently reported with this type of treatment.<sup>32,39,40</sup> Irrespective of the effectiveness of  
160 these individual treatment types, a broad range of therapeutic options exist, such as massage,  
161 ultrasound, manual therapy and corticosteroid injections,<sup>13</sup> that have yet to be explored in  
162 manual wheelchair users and could be worthy of future investigation. It was noted that three  
163 studies had explored the effectiveness of gluco-corticoid or corticosteroid injections. However,

164 these had to be excluded from the review since each study was a single sample case report,  
165 which did not satisfy the inclusion criteria.

166 The remaining six studies explored equipment,<sup>33</sup> educational<sup>34,35</sup> and lifestyle  
167 assistance<sup>36-38</sup> interventions. The only study to investigate equipment-based interventions,  
168 studied the effect of 2-gearred MAGIC Wheels on shoulder pain.<sup>33</sup> The gearing system of  
169 MAGIC Wheels allows participants to select between two different diameter push rims,  
170 depending on the task and can subsequently minimise the force and frequency of pushes  
171 performed by the user.<sup>33</sup> Hoenig et al.<sup>34</sup> and Rice et al.<sup>35</sup> explored the effects of educating users  
172 on aspects including wheelchair fitting, technique and upper limb preservation. However, it  
173 could be argued that this type of specialist education and training is best provided to prevent  
174 shoulder pain rather than as a treatment. Three studies examined the use of mobility service  
175 dogs for managing shoulder pain in wheelchair users.<sup>36-38</sup> Mobility service dogs can be secured  
176 to the front or side of a wheelchair to pull the user and assist with activities of daily living that  
177 can be challenging when experiencing pain, such as pushing uphill, over rough terrain or  
178 negotiating kerbs.<sup>38</sup> Concerns over the lack of cardiorespiratory stimulation reported when  
179 using a mobility service dog and the implications of such must be acknowledged.<sup>41,42</sup>  
180 Therefore, this type of intervention could be of greater use to users suffering from severe  
181 shoulder pain to help maintain their independence, since the lack of physical activity  
182 experienced whilst using a mobility service dog could lead to other contraindications and health  
183 problems.

184 A lack of physical activity and cardiorespiratory stimulation could actually be a  
185 common issue associated with a number of the non-exercise-based interventions. Subsequently  
186 interdisciplinary approaches may be advisable in the management of shoulder pain, which has  
187 previously been advocated for the preservation of upper limb function.<sup>13,43</sup> However, very few  
188 studies identified by the current review adopted interdisciplinary interventions. Kemp et al.<sup>22</sup>

189 and Mulroy et al.<sup>23</sup> both included ‘movement optimisation’ training alongside strengthening  
190 and stretching. The ‘movement optimisation’ training consisted of a series of recommendations  
191 provided by physical therapists to optimise skills that often provoke shoulder pain in  
192 wheelchair users (namely wheelchair propulsion and transfers) and received frequent  
193 reinforcement on these tasks over the duration of the programme.<sup>22,23</sup> Middaugh et al.<sup>24</sup> utilised  
194 electromyographical biofeedback sessions to accompany the home exercise programme they  
195 had prescribed. Individuals who report musculoskeletal pain during repetitive tasks often  
196 struggle with the ‘rest’ part of the cycle where muscle relaxation is required.<sup>44</sup> Subsequently,  
197 electromyographical biofeedback could be used to assist with muscle retraining and effectively  
198 relax overactive muscles during repetitive tasks such as wheelchair propulsion.<sup>24</sup> Although  
199 biofeedback would appear a potentially feasible means for the treatment of shoulder pain, it  
200 remains to be seen whether this is a clinically viable option since access to specialist  
201 electromyographical equipment is unlikely to be widespread. That said, more studies of this  
202 nature attempting to incorporate other treatment modalities alongside an exercise-based  
203 programme are encouraged for the management of shoulder pain in wheelchair users.<sup>13,43</sup>

#### 204 *Participants*

205 Studies included participants with varied physical characteristics. The majority of  
206 studies were male dominant and although a broad range of disabilities were investigated across  
207 studies, most focused on a specific health condition, rather than combining multiple. Although  
208 this approach guarantees homogeneity amongst participants to maximise internal validity, it  
209 can do so at the expense of external validity. This can cause problems for clinicians, as it  
210 prevents them and other practitioners from understanding which populations certain treatments  
211 may be generalised to.

212           The age range of participants was very broad, which implies that wheelchair users of  
213 varying experience levels have been accounted for, however this information was not always  
214 provided. Future research must include details about the number of years participants have  
215 been using a manual wheelchair when examining shoulder pain, as different treatment types  
216 may be more appropriate for someone who has recently acquired an injury compared to  
217 someone who has spent numerous years pushing a wheelchair. This also raises another point  
218 for future consideration. Although it was not an original criterion for data extraction, studies  
219 should also consider how long participants have been experiencing pain, as again different  
220 treatment options may be required for acute and chronic symptoms. Many studies referred to  
221 this, however as a bare minimum, future studies must include more detailed information  
222 regarding participants physical characteristics to assist clinicians with the treatment of shoulder  
223 pain for specific populations.

224           Another characteristic frequently not reported by studies was the physical activity levels  
225 of participants. Recreational activities outside of those performed for daily living could also  
226 predispose to a certain treatment type being more effective than another. For instance,  
227 sedentary individuals may respond better to an exercise-based treatment programme, whereas  
228 for individuals already accustomed to exercise, this might not be the case. Only one study  
229 identified by the current review investigated wheelchair athletes.<sup>29</sup> During the initial search a  
230 further two studies were identified that sampled wheelchair athletes.<sup>45,46</sup> However, one study  
231 was excluded since it included wheelchair athletes asymptomatic of shoulder pain and used  
232 changes in shoulder range of motion to infer changes in pain rather than a direct measure.<sup>46</sup>  
233 Whereas the second study was a one sample case study with a paratriathlete.<sup>45</sup> Although mixed  
234 findings have previously been reported as to whether wheelchair athletes are at a greater or  
235 reduced risk of developing shoulder pain than non-athletic wheelchair users,<sup>47-49</sup>  
236 musculoskeletal differences are likely between these two populations as a result of their

237 differing physical workloads. Subsequently, it should not be assumed that effective treatment  
238 methods for one population would be transferable to another and in particular, athletic  
239 populations require further research.

#### 240 *Measures*

241 The Wheelchair Users Shoulder Pain Index was by far the most common tool used to  
242 quantify shoulder pain and was used by 18 of the 21 studies. Of the three studies not using this  
243 questionnaire, Hoenig et al.<sup>34</sup> simply quantified shoulder pain as nominally present or not,  
244 whereas van der Linden et al.<sup>26</sup> and Garcia-Gomez et al.<sup>29</sup> adopted an alternative visual  
245 analogue scale questionnaire. The use of a nominal scale fails to account for the magnitude of  
246 pain, which should be an important consideration for interventions. Given that the Wheelchair  
247 Users Shoulder Pain Index has been established as a valid and reliable instrument for reporting  
248 shoulder pain in wheelchair users,<sup>50</sup> it is recommended that this questionnaire is reported to  
249 quantify pain wherever possible preferably in its performance corrected format. The  
250 performance corrected version is more applicable to all impairment types of wheelchair users  
251 since not all impairment types may perform all 15 activities themselves and by performing a  
252 correction, comparisons can be made between individuals and studies if necessary.<sup>4</sup> Clinicians  
253 would then be able to compare the relative effectiveness of different treatment options.

254 Although the Wheelchair Users Shoulder Pain Index is a good clinical tool for  
255 monitoring self-reported shoulder pain, pain itself can be considered a relatively subjective  
256 concept. Subsequently, future studies would be encouraged to include more objective measures  
257 of shoulder function alongside the presence of pain. Measures including range of movement,  
258 strength, muscular activity and propulsion kinetics were explored pre and post intervention by  
259 a limited number of studies. These objective measures could enable an insight into the

260 mechanisms responsible for either causing or reducing shoulder pain and may further facilitate  
261 the identification of effective conservative treatment types for clinicians.

## 262 *Design*

263 Of the available literature 9 of the 21 studies included randomised control trials. Although the  
264 aim of the current review was to simply map the available literature and the methodological  
265 designs adopted, future research into the effectiveness of the treatment interventions adopted  
266 will be warranted. In that case, reliable cause and effect relationships between the treatment  
267 and its effect on shoulder pain are paramount, for which randomised control trials remain the  
268 gold standard.<sup>51</sup> Although there are many challenges associated with implementing randomised  
269 control trials, such as cost, time and loss of participants to follow-up,<sup>51</sup> more of these studies  
270 are required to establish the effectiveness of conservative treatment types for reducing shoulder  
271 pain in wheelchair users in future.

272 A limitation associated with the current study was that the effectiveness of each  
273 treatment type was not provided. Although this information could be extremely valuable for  
274 clinicians, to assist with their treatment selection, the current review was a scoping review  
275 designed to identify gaps in the literature to help stimulate further research. Subsequently, it  
276 was not appropriate to conduct a detailed appraisal of included studies design and quality, nor  
277 the effectiveness of the interventions, as would have been expected for a systematic review.  
278 That said, this is still something of interest for future research. A subsequent limitation may lie  
279 within the search terms or inclusion / exclusion criteria adopted. Treatments such as injections  
280 could not be documented since the limited number of studies conducted in wheelchair users  
281 were all case reports. The only study to explore shoulder pain in athletic wheelchair users was  
282 also a case report. Subsequently, future research should consider including single sample case

283 reports so that clinicians can gain a broader understanding of effective treatment types and how  
284 they may differ in different wheelchair user populations.

285 In conclusion, despite the prevalence of shoulder pain amongst manual wheelchair  
286 users, previous research into conservative treatments to help manage this problem have been  
287 scarce. Future research would be recommended to adopt interdisciplinary / multifaceted  
288 interventions, with exercise at the heart of the study. Studies of this nature are important so that  
289 shoulder pain can be treated without neglecting other factors such as physical activity, which  
290 are equally important yet are often overlooked during monodisciplinary studies. Future studies  
291 must also report the physical characteristics of the participants investigated. These steps will  
292 enable clinicians to optimise their treatment strategies and to establish which strategies can be  
293 transferable to specific patients.

294

### 295 **Clinical messages**

- 296 • Exercise was the conservative treatment most frequently used to manage shoulder pain  
297 in wheelchair users.
- 298 • Few studies have explored multidisciplinary treatment strategies for reducing shoulder  
299 pain in wheelchair users.
- 300 • The Wheelchair Users Shoulder Pain Index was the commonly used tool for quantifying  
301 shoulder pain.

302

### 303 **Conflicts of Interest: none declared**

304

### 305 **References**

- 306 1. Veeger HEJ and van der Helm FCT. Shoulder function: the perfect compromise between  
307 mobility and stability. *J Biomech.* 2007;40(10):2119-2129.
- 308 2. Nichols PJ, Norman PA and Ennis JR. Wheelchair user's shoulder? Shoulder pain in  
309 patients with spinal cord lesions. *Scand J Rehabil Med.* 1979;11:29-32.
- 310 3. Pentland WE and Twomey LT. The weight-bearing upper extremity in women with long  
311 term paraplegia. *Paraplegia.* 1991;29:521-530.
- 312 4. Curtis KA, Drysdale GA, Lanza RD, Kolber M, Vitolo RS and West R. Shoulder pain in  
313 wheelchair users with tetraplegia and paraplegia. *Arch Phys Med Rehabil.* 1999;80:453-  
314 457.
- 315 5. Brose SW, Boninger ML, Fullerton B, et al. Shoulder ultrasound abnormalities, physical  
316 examination findings and pain in manual wheelchair users with spinal cord injury. *Arch*  
317 *Phys Med Rehabil.* 2008;89:2086-2093.
- 318 6. Medina GIS, Jesus CLM, Ferreira DM, et al. Is sport practice a risk factor for shoulder  
319 injuries in tetraplegic individuals? *Spinal Cord.* 2015;53:461-466.
- 320 7. Morrow MM, van Straaten MG, Murthy NS, Braman JP, Zanella E and Zhao KD.  
321 Detailed shoulder MRI findings in manual wheelchair users with shoulder pain. *Biomed*  
322 *Res Int.* 2017;76:1-7.
- 323 8. Gutierrez DD, Thompson L, Kemp B and Mulroy SJ. The relationship of shoulder pain  
324 intensity to quality of life, physical activity, and community participation in persons with  
325 paraplegia. *J Spinal Cord Med.* 2011;30(3):251-255.
- 326 9. Wang JC, Chan RC, Tsai YA, et al. The influence of shoulder pain on functional  
327 limitation, perceived health, and depressive mood in patients with traumatic paraplegia.  
328 *J Spinal Cord Med.* 2015;38(5):587-592.
- 329 10. Hoffman MD. Cardiorespiratory fitness and training in quadriplegics and paraplegics.  
330 *Sports Med.* 1986;3(5):312-330.



- 331 11. Kerr J, Borbas P, Meyer DC, Gerber C, Buitrago Tellez C and Wieser K. Arthroscopic  
332 rotator cuff repair in the weight-bearing shoulder. *J Shoulder Elbow Surg.*  
333 2015;24(12):1894-1899.
- 334 12. Jordan RW, Sloan R and Saithna A. Should we avoid shoulder surgery in wheelchair  
335 users? A systematic review of the outcomes and complications. *Orthop Traumatol Surg*  
336 *Res.* 2018;104(6):839-846.
- 337 13. Steuri R, Sattelmayer M, Elsig S et al. Effectiveness of conservative interventions  
338 including exercise, manual therapy and medical management in adults with shoulder  
339 impingement: a systematic review and meta-analysis of RCTs. *Brit J Sports Med.*  
340 2016;51:1340-1347.
- 341 14. Cratsenberg KA, Deitrick CE, Harrington TK et al. Effectiveness of exercise programs  
342 for management of shoulder pain in manual wheelchair users with spinal cord injury. *J*  
343 *Neurol Phys Ther.* 2015;39:197-203.
- 344 15. Arksey H and O'Malley L. Scoping studies: towards a methodological framework. *Int J*  
345 *Soc Res Methodol.* 2005;8(1):19-32.
- 346 16. Peters MDJ, Godrey CM, Khalil H, McInerney P, Parker D and Baldini C. Guidance for  
347 conduction systematic scoping reviews. *Int J Evid Based Healthc.* 2015;13:141-146.
- 348 17. Tricco AC, Lillie E, Zarin W et al. PRISMA extension for scoping reviews (PRISMA-  
349 ScR): checklist and explanation. *Ann Intern Med.* 2018;169(7):467-473.
- 350 18. Curtis KA, Tyner TM, Zachary L et al. Effect of a standard exercise protocol on shoulder  
351 pain in long-term wheelchair users. *Spinal Cord* 1999;37:421-429.
- 352 19. Nawoczenski DA, Ritter-Soronon JM, Wilson CM, Howe BA and Ludewig PM. Clinical  
353 trial of exercise for shoulder pain in chronic spinal injury. *Phys Ther.* 2006;86(12):1604-  
354 1618.


- 355 20. Dyson-Hudson TA, Sisto SA, Bond Q, Emmons R and Kirshblum SC. Arm crank  
356 ergometry and shoulder pain in persons with spinal cord injury. *Arch Phys Med Rehabil.*  
357 2007;88:1727-1729.
- 358 21. Nash MS, van de Ven I, van Elk N and Johnson BM. Effects of circuit resistance training  
359 on fitness attributes and upper-extremity pain in middle-aged men with paraplegia. *Arch*  
360 *Phys Med Rehabil.* 2007;88:70-75.
- 361 22. Kemp BJ, Bateham AL, Mulroy SJ, Thompson L, Adkins RH and Kahan JS. Effects of  
362 reduction in shoulder pain on quality of life and community activities among people  
363 living long-term with SCI paraplegia: a randomized control trial. *J Spinal Cord Med.*  
364 2011;34(3):278-284.
- 365 23. Mulroy SJ, Thompson L, Kemp B et al. Strengthening and optimal movements for  
366 painful shoulders (STOMPS) in chronic spinal cord injury: a randomised controlled trial.  
367 *Phys Ther.* 2011;91(3):305-324.
- 368 24. Middaugh S, Thomas KJ, Smith AR, McFall TL and Klingmueller J. EMG biofeedback  
369 and exercise for treatment of cervical and shoulder pain in individuals with a spinal cord  
370 injury: a pilot study. *Top Spinal Cord Inj Rehabil.* 2013;19(4):311-323.
- 371 25. Norrbrink C, Lindberg T, Wahman K and Bjerkfors A. Effects of an exercise programme  
372 on musculoskeletal and neuropathic pain after spinal cord injury – results from a seated  
373 double-poling ergometer study. *Spinal Cord.* 2012;50:457-461.
- 374 26. van der Linden ML, Bulley C, Geneen LJ, Hooper JE, Cowan P and Mercer TH. Pilates  
375 for people with multiple sclerosis who use a wheelchair: feasibility, efficacy and  
376 participant experiences. *Disabil Rehabil.* 2014;36(11):932-939.
- 377 27. van Straaten M, Cloud BA, Morrow MM, Ludewig PM and Zhao KD. Effectiveness of  
378 home exercise on pain, function and strength of manual wheelchair users with spinal cord

- 379 injury: a high-dose shoulder program with telerehabilitation. *Arch Phys Med Rehabil.*  
380 2014;95:1810-1817.
- 381 28. Wilbanks SR, Rogers R, Pool S and Bickel S. Effects of functional electrical stimulation  
382 assisted rowing on aerobic fitness and shoulder pain in manual wheelchair users with  
383 spinal cord injury. *J Spinal Cord Med.* 2016;39(6):645-654.
- 384 29. Garcia-Gomez S, Perez-Tajero J, Hoozemans M and Barakat R. Effect of a home-based  
385 exercise program on shoulder pain and range of motion in elite wheelchair basketball  
386 players: a non-randomised controlled trial. *Sports.* 2019;7(8):180.
- 387 30. Dyson-Hudson TA, Shiflett SC, Kirshblum SC, Bowen JE and Druin EL. Acupuncture  
388 and trager psychophysical integration in the treatment of wheelchair user's shoulder pain  
389 in individuals with spinal cord injury. *Arch Phys Med Rehabil.* 2001;82:1038-1046.
- 390 31. Dyson-Hudson TA, Kadar P, LaFontaine M, et al. Acupuncture for chronic shoulder  
391 pain in persons with spinal cord injury: a small scale clinical trial. *Arch Phys Med*  
392 *Rehabil.* 2007;88:1276-83.
- 393 32. Giner-Pasqual M, Alcanyis-Alberola M, Querol F, Salinas-Huertas S, Garcia-Masso X  
394 and Gonzalez LM. Transdermal nitroglycerine treatment of shoulder tendinopathies in  
395 patients with spinal cord injuries. *Spinal Cord.* 2011;49:1014-1019.
- 396 33. Finley MA and Rodgers MM. Effect of 2-speed geared manual wheelchair propulsion on  
397 shoulder pain and function. *Arch Phys Med Rehabil.* 2007;88:1622-1627.
- 398 34. Hoenig H, Landerman LR, Shipp KM, et al. A clinical trial of a rehabilitation expert  
399 clinician versus usual care for providing manual wheelchairs. *J Am Geriatr Soc.*  
400 2005;53:1712-1720.
- 401 35. Rice LA, Smith I, Kelleher AR, Greenwald K and Boninger ML. Impact of a wheelchair  
402 education protocol based on practice guidelines for preservation of upper-limb function:  
403 a randomized trial. *Arch Phys Med Rehabil.* 2014;95:10-19.

- 404 36. Hubert G, Tousignant M, Routhier F, Corriveau H and Champagne N. Effect of service  
405 dogs on manual wheelchair users with spinal cord injury: a pilot study. *J Rehabil Res*  
406 *Dev.* 2013;50(3):341-350.
- 407 37. Vincent C, Gagnon DH, Routhier F et al. Service dogs for people with spinal cord injury:  
408 outcomes regarding functional mobility and important occupations. *Assist Technol.*  
409 2015;217:847-851.
- 410 38. Vincent C, Gagnon DH, Dumont F and the ADMI group. Pain, fatigue, function and  
411 participation among long-term manual wheelchair users partnered with a mobility service  
412 dog. *Disabil Rehabil: Assist Technol.* 2019;14(2):99-108.
- 413 39. Berrazueta JR, Losada A, Poveda J, et al. Successful treatment of shoulder pain syndrome  
414 due to supraspinatus tendinitis with transdermal nitroglycerin: a double blind study. *Pain.*  
415 1996;66:63-67.
- 416 40. Kane TP, Ismail M and Calder JD. Topical glycerol trinitrate and noninsertional achilles  
417 tendinopathy: a clinical and cellular investigation. *Am J Sports Med.* 2008;36:1160-1163.
- 418 41. Nash MS. Cardiovascular fitness after spinal cord injuries. In: Lin V, ed. *Spinal Cord*  
419 *Medicine.* New York: Demos; 2002:637-646.
- 420 42. Champagne A, Gagnon D, Vincent C, et al. Comparison of cardiorespiratory demand and  
421 rate of perceived exertion during propulsion in a natural environment with and without  
422 the use of a mobility assistance dog in manual wheelchair users. *Am J Phys Med Rehabil.*  
423 2016;95:685-691.
- 424 43. Paralyzed Veterans of America Consortium for Spinal Cord Medicine. Preservation of  
425 upper limb function following spinal cord injury: A clinical practice guideline for  
426 healthcare professionals. *J Spinal Cord Med.* 2005;28(5):433-470.

- 427 44. Hagg GM and Astrom A. Load pattern and pressure pain threshold in the upper trapezius  
428 muscle and psychosocial factors in medical secretaries with and without shoulder/neck  
429 disorders. *Int Arch Occup Environ Health*. 1997;69(6):423-432.
- 430 45. Diaz R, Stoll AH, Rho ME and Blauwet CA. Preserving the shoulder function of an elite  
431 para triathlete: a case report. *Am J Phys Med Rehabil*. 2018;97(8):69-72.
- 432 46. Wilroy J, Hibberd E. Evaluation of a shoulder injury prevention program in wheelchair  
433 basketball. *J Sport Rehabil*. 2018;27(6):554-559.
- 434 47. Nyland J, Snouse SL, Anderson M, Kelly T and Sterling JC. Soft tissue injuries to USA  
435 Paralympians at the 1996 Summer Games. *Arch Phys Med Rehabil*. 2000;81:368-373.
- 436 48. Finley MA and Rodgers MM. Prevalence and identification of shoulder pathology in  
437 athletic and nonathletic wheelchair users with shoulder pain: a pilot study. *J Rehabil Res*  
438 *Dev*. 2004;41:395-402.
- 439 49. Heyward OW, Vegter RJK, de Groot S and van der Woude LHV. Shoulder complaints  
440 in wheelchair athletes: a systematic review. *PlosOne*. 2017;12(11):e0188410.
- 441 50. Curtis KA, Roach KE, Applegate EB, et al. Development of the wheelchair user's  
442 shoulder pain index (WUSPI). *Paraplegia*. 1995;33:290-293.
- 443 51. Sibbald B and Roland M. Understanding controlled trials – Why are randomised  
444 controlled trials so important? *Brit Med J*. 1998;316:201-201.

445



**Figure 1** – PRISMA flow diagram of the study selection process.

**Table 1** Exercise-based interventions for the treatment of shoulder pain in manual wheelchair users

Authors	Intervention	Duration (weeks)	Sample (n)	Age (yrs)	Participants			Measures		Design
					Disability	Experience (yrs)	Activity (hr/wk)	Pain	Secondary	
Curtis et al. (1999) <sup>18</sup>	HEP strengthening and stretching. 3x15 reps daily	24	42 35 M; 7 F	35 ± 8	SCI, CP, MS & amputees	14 ± 9	Comm X̄ 12	PC- WUSPI	n/a	RCT
Dyson-Hudson et al. (2007) <sup>20</sup>	Arm crank ergometer training. 3x20min / wk	12	23 19 M; 4 F	41 ± 9	SCI (tetra & para)	15 ± 9	Comm 5 ± 4	WUSPI	n/a	RCT
Garcia-Gomez et al. (2019) <sup>29</sup>	HEP strengthening and stretching. 3x30min / wk	10	36 15 M; 21 F	26 ± 8	Not stated	Not stated	Athletes > 6	SPI-WB	Impingement tests & RoM	Quasi
Kemp et al. (2011) <sup>22</sup>	HEP & movement training 3 / wk vs. 1hr educational video	12	58 Not stated	22-72	SCI (all para)	20 ± 11	Comm Not stated	WUSPI	n/a	RCT
Middaugh et al. (2013) <sup>24</sup>	HEP & EMG biofeedback. 4 / wk exercise. 5 EMG sessions	12	15 12 M; 3 F	23-56	SCI (tetra & para)	X̄ 16	Not stated	PC- WUSPI	n/a	RCT
Mulroy et al. (2011) <sup>23</sup>	HEP & movement training 3 / wk vs. 1hr educational video	12	58 Not stated	45 ± 11	SCI (all para)	22 ± 12	Comm Not stated	WUSPI	Shoulder torque & RoM	RCT
Nash et al. (2007) <sup>21</sup>	Resistance & arm crank ergometer. 3x45min / wk	16	7 7 M; 0 F	39-58	SCI (all para)	13 ± 7	Comm Not stated	WUSPI	Strength & power	Coh
Nawoczenski et al. (2006) <sup>19</sup>	HEP strengthening and stretching daily	8	41 28 M; 13 F	47 ± 12	SCI (tetra & para)	17 ± 13	Comm Not stated	PC- WUSPI	n/a	Quasi
Norrbrink et al. (2012) <sup>25</sup>	Double-poling ergometer training	10	8 6 M; 2 F	51 ± 11	SCI (all para)	18 ± 8	Comm Not stated	WUSPI	n/a	Coh
van der Linden et al. (2014) <sup>26</sup>	Supervised Pilates classes. 1-2x60min / wk	12	15 8 M; 7 F	51 ± 8	MS	Not stated	Comm Not stated	VAS	Interscapular distances	Coh
van Straaten et al. (2014) <sup>27</sup>	HEP strengthening and stretching. 3x30 reps, 3 / wk	16	16 13 M; 3 F	25-64	SCI / polio	X̄ 16	Comm Not stated	WUSPI	Isometric strength	Coh
Wilbanks et al. (2016) <sup>28</sup>	FES assisted rowing programme. 3x30min / wk	6	10 8 M; 2 F	47 ± 12	SCI (all para)	18 ± 14	Comm Not stated	WUSPI	Isokinetic strength, EMG	Coh

Nb. HEP – home exercise programme, EMG – Electromyography, FES – Functional Electrical Stimulation, SCI – spinal cord injury, tetra – tetraplegia, para – paraplegia, CP – cerebral palsy, MS – multiple sclerosis, Comm – community users, RCT – randomised controlled trial, Coh – cohort, Quasi – quasi-experimental, WUSPI – wheelchair users shoulder pain index, PC-WUSPI – performance corrected wheelchair users shoulder pain index, VAS – visual analogue scale, RoM – range of movement.

**Table 2** Additional treatment interventions conducted in manual wheelchair users with shoulder pain

Authors	Intervention	Duration (weeks)	Sample (n)	Age (yrs)	Participants			Measures		Design
					Disability	Experience (yrs)	Activity (hr/wk)	Pain	Secondary	
<b>Therapeutic:</b>										
Dyson-Hudson et al. (2001) <sup>30</sup>	Acupuncture vs. TPI. 10 treatments over 5 weeks	15	18 14 M; 4 F	45 ± 11	SCI (tetra & para)	15 ± 8	Comm 6 ± 7	PC- WUSPI	n/a	Quasi
Dyson-Hudson et al. (2007) <sup>31</sup>	Acupuncture vs. placebo. 10 treatments over 5 weeks	15	17 15 M; 2 F	39 ± 11	SCI (tetra & para)	11 ± 9	Comm 8 ± 13	PC- WUSPI	n/a	RCT
Giner-Pasqual et al. (2011) <sup>32</sup>	Transdermal nitroglycerine patch vs. placebo. Daily	24	41 Not stated	42-54	SCI (all para)	Not stated	Athletes Not stated	WUSPI	RoM	RCT
<b>Equipment:</b>										
Finley & Rodgers (2007) <sup>33</sup>	2-gearred, non-powered MAGIC Wheels – 5 months	28	13 7 M; 6 F	46 ± 14	SCI / polio	15 ± 10	Not stated	PC- WUSPI	Impingement tests & RoM	Coh
<b>Educational:</b>										
Hoening et al. (2005) <sup>34</sup>	Education on fitting & propulsion vs. standard care	24	57 Not stated	65 ± 14	Not stated	13 ± 7	Comm Not stated	Yes / No	n/a	
Rice et al. (2014) <sup>35</sup>	Upper limb preservation guidance vs. standard care	52	37 28 M; 9 F	38 ± 16	SCI (tetra & para)	Not stated	Comm Not stated	PC- WUSPI	Propulsion kinetics	RCT
<b>Lifestyle:</b>										
Hubert et al. (2015) <sup>36</sup>	19 days training with mobility service dog	28	11 Not stated	Not stated	SCI (not stated)	Not stated	Comm Not stated	WUSPI	n/a	Coh
Vincent et al. (2015) <sup>37</sup>	Mobility service dog to provide lifestyle assistance	54	66 45 M; 21 F	$\bar{X}$ 41	SCI (not stated)	Not stated	Comm Not stated	WUSPI	n/a	Coh
Vincent et al. (2019) <sup>38</sup>	Mobility service dog to provide lifestyle assistance	54	17 9 M; 8 F	42 ± 15	SCI (not stated)	Not stated	Comm Not stated	WUSPI	n/a	Coh

Nb. TPI – Trager Psychophysical Integration, SCI – spinal cord injury, tetra – tetraplegia, para – paraplegia, Comm – Community users, RCT – randomised controlled trial, Coh – cohort, Quasi – quasi-experimental, WUSPI – wheelchair users shoulder pain index, PC-WUSPI – performance corrected wheelchair users shoulder pain index, RoM – range of movement.