

Quantum spin liquid at finite temperature: proximate dynamics and persistent typicality

I. Rousochatzakis,^{1,2} S. Kourtis,³ J. Knolle,⁴ R. Moessner,⁵ and N. B. Perkins²

¹*Department of Physics and Centre for Science and Materials,
Loughborough University, Loughborough LE11 3TU, United Kingdom*

²*School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA*

³*Department of Physics, Boston University, Boston, MA, 02215, USA*

⁴*Blackett Laboratory, Imperial College London, London SW7 2AZ, United Kingdom*

⁵*Max Planck Institute for the Physics of Complex Systems, D-01187 Dresden, Germany*

(Dated: November 6, 2018)

Quantum spin liquids are long-range entangled states of matter with emergent gauge fields and fractionalized excitations. While candidate materials, such as the Kitaev honeycomb ruthenate α -RuCl₃, show magnetic order at low temperatures T , here we demonstrate numerically a dynamical crossover from magnon-like behavior at low T and frequencies ω to long-lived fractionalized *fermionic* quasiparticles at higher T and ω . This crossover is akin to the presence of spinon continua in quasi-1D spin chains. It is further shown to go hand in hand with persistent *typicality* down to very low T . This, which we argue is a signature of proximate spin liquidity and emergent gauge degrees of freedom more generally, will be useful for the numerical study of many finite-temperature properties of putative quantum spin liquids.

I. Introduction

Quantum spin liquids (QSLs) have been one of the central themes in condensed matter for many years. [1–11] Unlike conventional phases of matter, the characteristic correlations in QSLs are non-local and cannot be detected directly by standard probes. As such, identifying the experimental signatures of QSLs is one of the most challenging tasks in the field. [12] This task has become all the more pressing in the last decade, with the discovery of several candidate materials, including isotropic layered kagome systems [8, 13–15] and the strong spin-orbit coupled iridates and ruthenates. [16–33]

A promising route to detect QSLs is to look for signatures of fractionalization in dynamical probes, such as inelastic neutron scattering (INS), [34–40] Raman scattering, [41–47] resonant inelastic x-ray scattering (RIXS), [48–50] and ultrafast spectroscopy. [51] Such probes couple to multiple fractionalized quasiparticles, leading to characteristic broad scattering profiles. However, potential QSLs tend to be sensitive to perturbations [17, 52–58] and are in fact, in many if not most cases, preempted by magnetic order at low temperatures T . [24–33]

Despite this, here we establish numerically that long-lived fractionalized quasi-particles are still present in the spectrum at finite energies, in the same way that spinon continua survive in quasi-1D spin chains, which also order at low T due to weak interchain interactions. [34–36, 59–63] Fig. 1 illustrates the qualitative picture for the case of honeycomb Kitaev materials, like α -RuCl₃, which are proximate to a gapless spin liquid characterized by emergent magnetic fluxes and Majorana fermions. [7] Here, the characteristic ordering temperature T_N and magnon excitation frequencies ω_m are set by the perturbations that drive the magnetic order, and are therefore much smaller than the dominant energy scale K responsible for stabilizing the Kitaev QSL. As such, dynamical signatures of incipient spin liquidity are generally expected in some range of T and frequency ω , sufficiently above T_N and ω_m , respectively.

As a dynamical probe we analyze the Raman scattering intensity, which measures the scattering of light off the magnetic degrees of freedom, as a function of energy and light polariza-

tion. [64–66] By monitoring the evolution of the intensity as we drive the honeycomb Kitaev QSL toward a number of different instabilities, we show that, irrespective of the nature of the phase the system is driven to, the response indeed follows closely the one originating from fractionalized quasiparticles in a wide range of T and ω , while the magnon-like response – characteristic of the given magnetic order – appears only at low T and ω . This crossover, which can be identified even relatively deep inside the ordered phase, is akin to the confinement of spinons in quasi-1D spin chains at low T . [34–36, 59–63]

Besides the above dynamical crossover, our results reveal yet another manifestation of incipient spin liquidity, that of *persistent typicality* [67–73]: both static and dynamic properties can be captured down to surprisingly low T by propagating, in real and imaginary time, a *single*, randomly chosen many-body quantum state. This remarkable property arises from




FIG. 1. **Dynamical crossover near a Kitaev QSL (schematic).** Magnon excitations, characteristic of the long-range magnetic order at low T and ω , give way to long-lived fractionalized quasi-particles (here Majorana fermions) at higher T and ω . This crossover is accompanied by a remarkable persistence of *typicality* to very low T near the QSL.




FIG. 2. **Persistence of typicality near the Kitaev QSL, I.** Logarithm of contributions to the partition function ζ_r [Eq. (3)] from 320 random states for three points inside the Néel state, stabilized by a Heisenberg coupling J : (a) $\theta=0$, (b) $\theta=0.4\pi$ and (c) $\theta=0.49\pi$ [$J=\cos\theta$, $K=\sin\theta$]. All data refer to the symmetric 24-site cluster of App. VI.

the spectral weight downshift and the large, low- T entropy characteristic of frustrated systems with a large number of competing low energy states and emergent gauge fields. [74] The success of the typicality method is therefore inherently linked to the strong fluctuations present in quantum liquids. This therefore opens a unique route to study, on a quantitative level, an abundance of strongly correlated materials, like, for example, the layered kagome antiferromagnet $\text{ZnCu}_3(\text{OH})_6\text{Cl}_2$ or the honeycomb iridates and ruthenates, $\alpha\text{-Li}_2\text{IrO}_3$, Na_2IrO_3 and $\alpha\text{-RuCl}_3$.

II. Model, Raman vertex and methods

We consider the generic situation of Fig. 1 for the honeycomb magnet with Hamiltonian

$$\mathcal{H} = K \sum_{\alpha} \sum_{\langle ij \rangle \in \alpha} S_i^{\alpha} S_j^{\alpha} + \mathcal{V}. \quad (1)$$

Here, \mathbf{S}_i and \mathbf{S}_j are nearest-neighbor (NN) (pseudo-)spins-1/2, K is the Kitaev coupling, $\alpha = x, y$, or z , depending on the orientation of the bond $\langle ij \rangle$ (see inset of Fig. 1), and \mathcal{V} is a generic perturbation that destabilizes the Kitaev QSL. We will study here two such perturbations that are believed to be relevant in the available Kitaev materials, the NN Heisenberg exchange coupling, J , and the symmetric part of the NN off-diagonal exchange, Γ . [55, 58, 75–79] The corresponding models will be referred to below as the JK - and KT -model, respectively.

In the absence of \mathcal{V} , the Hamiltonian can be reduced to a quadratic problem of fermions moving in the background of static magnetic fluxes, and the ground state is a gapless QSL that lives in the zero-flux sector. [7] As shown by Nasu *et al.*, [80] this state is adiabatically connected to the high- T paramagnetic phase, but there are two characteristic crossover temperature scales, $T_{\ell} \sim K/100$ and $T_h \sim K$, see Fig. 2 of Ref. [80] and Fig. 1. Below T_{ℓ} , the system is locked into the flux-free sector, while for $T_{\ell} < T < T_h$ the Majorana fermions move on top of a thermally disordered flux background.

Raman experiments measure the dynamical response function associated with the Raman vertex \mathcal{R} , which, in the Loudon-

Fleury approximation, [46, 64, 65] is given by

$$\mathcal{R} = \sum_{\langle ij \rangle} (\epsilon_{\text{in}} \cdot \mathbf{d}_{ij})(\epsilon_{\text{out}} \cdot \mathbf{d}_{ij}) \mathcal{H}_{ij}, \quad (2)$$

where \mathbf{d}_{ij} is the displacement from site i to j , ϵ_{in} and ϵ_{out} are the polarizations of the incoming and outgoing light, respectively, and \mathcal{H}_{ij} denotes the interactions between \mathbf{S}_i and \mathbf{S}_j . The Raman intensity is the Fourier transform $I(\omega) = \int d\omega e^{i\omega t} \langle \mathcal{R}(t) \mathcal{R}_0 \rangle$, where $\langle \dots \rangle = \text{Tr}[e^{-\beta \mathcal{H}} \dots] / \text{Tr}[e^{-\beta \mathcal{H}}]$ denotes the statistical average over the Hilbert space, $\beta = 1/T$ is the inverse temperature, and $\mathcal{R}(t) = e^{i\mathcal{H}t} \mathcal{R} e^{-i\mathcal{H}t}$ is the time-evolved Raman vertex. In the 3-fold symmetric models we consider here, the Raman intensity comes entirely from the ‘xy’ and ‘ x^2-y^2 ’ polarization channels. The two channels give identical contributions, as they belong to the two-dimensional irreducible representation E_g , so it suffices to consider the ‘xy’ channel only. [46, 81]

The stochastic method used here amounts to replacing the thermodynamic trace over the Hilbert space of dimension D , with a sampling over r_m randomly chosen states $|r\rangle$, with $r_m \ll D$. [82–87] The Raman correlator is then given by

$$\langle \mathcal{R}(t) \mathcal{R}(0) \rangle \approx \sum_{r=1}^{r_m} d_r f_r(t) / \sum_{r=1}^{r_m} d_r \zeta_r, \quad \text{where} \quad (3)$$

$$f_r(t) = \langle r | e^{-\beta \mathcal{H}/2} \mathcal{R}(t) \mathcal{R}(0) e^{-\beta \mathcal{H}/2} | r \rangle, \quad \zeta_r = \langle r | e^{-\beta \mathcal{H}} | r \rangle,$$

and d_r is the dimensionality of the symmetry sector the given state $|r\rangle$ belongs to. The stochastic sampling is the basis of finite- T Lanczos, [85, 87] Chebyshev polynomial methods, [86] and the standard typicality approach, [67–73] for which $r_m = 1$. Here we take $r_m = 320$ random states (10 for each of the 32 irreducible representations of the symmetry group exploited for the symmetric 24-site cluster, see App. VI). These states represent configurations at $T = \infty$, which then need to be propagated in real and imaginary time in order to evaluate their contribution, $f_r(t)$ and ζ_r , to the dynamical correlation function and the partition function, respectively. This is done here using the standard Lanczos method. [88–91] The results are cross-checked with the complementary low- T Lanczos method, [92] which approaches the problem from low T , see App. VI.




FIG. 3. **Persistence of typicality near the Kitaev QSL, II.** (a-b) Real time evolution of the correlator $f_r(t)$ [Eq. (3)] for 320 random ($T = \infty$) states $|r\rangle$, for the JK model at $\theta = 0.47\pi$, with $T = |K|$ (a) and $0.2|K|$ (b). (c) T -dependence of the maximum standard deviation in $\langle \mathcal{R}(t)\mathcal{R}(0) \rangle = \sum_r d_r f_r(t) / \sum_r d_r \zeta_r$ [see Eq. (3)] for various values of θ . All data refer to the symmetric 24-site cluster of App. VI.

III. Typicality

We begin by showing explicitly that the typicality hypothesis remains valid down to very low T in the vicinity of the Kitaev QSL points. Fig. 2 shows the logarithm of ζ_r , defined in Eq. (3) for 320 states $|r\rangle$, for the JK -model at $\theta = 0$ (a), 0.4π (b) and 0.49π (c), where $J = \cos \theta$ and $K = \sin \theta$. All three panels correspond to points inside the Néel phase (the transition to the Kitaev QSL occurs at $\theta \approx 0.493\pi$, see Ref. [53] and App. VI). At the Heisenberg point (a), the 320 random states give almost identical results for ζ_r for $T \gtrsim |K|$. This is the essence of the typicality hypothesis which tell us that states in the middle part of the energy spectrum are ‘typical’ to each other, and therefore the statistical average can be equivalently obtained by looking at the evolution of a single random state. On general grounds, this hypothesis can be shown [85, 87, 93] to work extremely well at very high T . On cooling down, the system begins to sample the lower end of the energy spectrum, where finite-size effects begin to play a role and the eigenstates are usually in practise no longer typical. As a result, deviations between the different ζ_r become apparent. Quite remarkably, however, the characteristic temperature, T_{typ} , below which the typicality hypothesis breaks down gets lower and lower as we approach the Kitaev QSL point, see panels (b-c) and sketch in Fig. 1.

This persistent typicality can also be seen in the time dependent quantities $f_r(t)$ defined in Eq. (3). Figs. 3 (a-b) show the real and imaginary parts of $f_r(t)$ for 320 states $|r\rangle$, for the representative point $\theta = 0.47\pi$ of the JK -model. For $T = |K|$ (a), the 320 random states give almost identical results for $f_r(t)$, in the entire time region shown. As above, deviations between the curves become visible at lower T , see panel b. Nevertheless, as shown in panel c, the maximum standard deviation between the different results for $\langle \mathcal{R}(t)\mathcal{R}(0) \rangle$ remains small down to very low T , in the vicinity of the Kitaev point.

The success of the typicality method reflects, in essence,

a fundamental property of the proximate Kitaev QSL, the presence of low-energy emergent gauge degrees of freedom. For conventional phases, the energy spectrum of a finite-size cluster is typically very dense in the middle of the spectrum but becomes sparse below some energy scale, proportionate to the bare interaction strength. For the Kitaev QSL, by contrast, the presence of magnetic flux sectors on one hand (with $2^{N/2}$ fermionic states each) and the very low flux gap ($\Delta \approx 0.065K$ [7]) on the other, give rise to finite-size spectra that remain dense down to very low energies. This is demonstrated in Fig. 4 which shows the energy spectrum of the JK -model on the 24-site cluster in the full parameter range of θ . A direct consequence of the spectral downshift around the Kitaev points ($\theta = \pm\pi/2$) is that the system releases as much as half of its entropy only when cooled below $T \sim K/50$. [80]

The exponential number of competing low-energy states in systems with emergent gauge fields quantifies what we mean by ‘very dense’ spectrum. Indeed, as we show in App. VI, the density of states $\rho(E)$ scales exponentially with system size N ,

$$\rho(E) \propto e^{Ns(E)}, \quad (4)$$

where $s(E)$ is the microcanonical entropy per site, down to very low energies, already for clusters with 24 or 32 spins. This is to be contrasted with the magnetically ordered regions of the phase diagram, where the spectral weight is concentrated near a restricted set of configurations (i.e. corresponding to elementary spin flips or magnons).

IV. Dynamical crossover

We now turn to the dynamical crossover announced above. Fig. 5 shows the Raman intensities $I(\omega)$ for six different points in parameter space. Let us first focus on the first row (panels




FIG. 4. **Low-energy spectrum of the JK -model on the 24-site cluster.** Energies are measured from the ground state energy $E_0(\theta)$. The non-linear horizontal axis is used to better highlight what happens in the narrow regions around the Kitaev QSL points $\theta = \pi/2$ and $3\pi/2$. The various symbols correspond to the irreducible representations of the symmetry exploited here. The latter includes: i) the 12 translations, with the symbols Γ , K^* and M^* corresponding, respectively, to zero momentum, the corners of the first Brillouin zone (BZ), and the midpoints of the BZ edges. ii) real space inversion through the middle of the hexagons, with the letters ‘e’ and ‘o’ standing for the even and the odd sectors, respectively. iii) spin inversion (global rotation around the x-axis in spin-space), with the symbols ‘Sze’ and ‘Szo’ standing for the even and odd sectors, respectively. The numbers in parentheses give the degeneracy of each level. The spectrum is obtained by the standard Lanczos method. Specifically, we show the 100 lowest energy states in each sector. Among these, only the lowest five are converged to the requested high precision (here relative precision in the 9th digit). The rest give a good representation with precision that lowers as we go up in energy.

a-c), which show the evolution of $I(\omega)$ as we move deeper inside one particular magnetic phase, here the Néel state stabilized by a Heisenberg coupling J . As above, the JK -model is parametrized as $J = \cos \theta$ and $K = \sin \theta$, with $\theta = \pi/2$ (panel **a**, QSL phase), 0.47π (panel **b**, Néel) and 0.45π (panel **c**, Néel). For the Kitaev point (**a**), the results are consistent with both the exact $T = 0$ result of Knolle *et al* [46] (reference dashed lines) and the T -evolution data reported by Nasu *et al.* [47] The broad profile with the main peak around $\omega \sim |K|$ (related to the Van Hove singularity of the two-fermion density of states) and the bandwidth of $3|K|$ are all reproduced, and the same is true for the characteristic growth of the zero-frequency intensity with T . [47]

The crucial point is that the main features of the intensity remain the same in a wide T and ω regime also inside the Néel phase. Additional structure, characteristic of this phase (and the given finite size cluster), does become visible at low enough T , and becomes sharper as we move deeper inside the phase. In panel **c**, for example, the main peak of the Kitaev point has been depleted, and the spectral weight has been asymmetrically transferred into two new peaks (one around $0.75|K|$ and the other around $1.5|K|$). At the same time, there is an appreciable gap developing at low ω , while a fourth peak appears a little below $3|K|$. While these sharper features may depend in detail on the finite-size cluster at hand, [94] they are washed out with

increasing T , and the response eventually resembles that at the Kitaev point (panel **a**).

Panels **d-f** show that this picture remains essentially the same irrespective of the nature of the phase proximate to the QSL. Here we show the intensity for three more phases, the zigzag (**d**), the ferromagnetic (**e**), and the phase stabilized by the off-diagonal exchange Γ (for the nature of this phase see Refs. [95–97]). As before, at sufficiently high T the response is the same with that of the proximate Kitaev QSL, while the response at low T is distinctive for each of the three phases. For example, the FM phase shows a relatively large zero-frequency weight, and the same is true for the phase stabilized by Γ . Results from the independent low- T Lanczos method lead to the same conclusion, see App. VI. Altogether, these results demonstrate the presence of a dynamical crossover from a conventional, magnon-like picture at low T and ω , characteristic of the low- T ordered phase, to long-lived fractionalized quasi-particles, characteristic of the proximate Kitaev QSL.

V. Fermionic character

We now turn to the nature of these fractionalized quasiparticles and show evidence for their fermionic character. To this end, we follow the analysis of Ref. [47] for the Raman intensity




FIG. 5. **Raman scattering intensity in the ‘xy’ polarization channel for various phases proximate to the Kitaev QSL.** (a) Ideal AF Kitaev point, (b) JK model with $\theta=0.47\pi$ (Néel phase), (c) JK model with $\theta=0.45\pi$ (Néel phase), (d) JK model with $\theta=0.53\pi$ (zigzag phase), (e) JK model with $\theta=1.43\pi$ (FM phase), (f) KT model with $K=-1$ and $\Gamma=0.2$. Dashed lines show the analytical $T=0$ result at the Kitaev QSL point [46]. T and ω are in units of $\sqrt{K^2+J^2}=1$ (a-e) or $|K|$ (f). Results are obtained for the symmetric 24-site cluster of App. VI, by propagating 320 random ($T=\infty$) states in real and imaginary time.

at the ideal QSL point. At this point, the Raman vertex is diagonal in the emergent fluxes and only excites/deexcites pairs of Majorana fermions. In particular, there are two processes contributing to the intensity, one (type A) corresponding to the creation of two fermions, and another (type B) to the creation of one fermion and the annihilation of another. The amplitudes of these processes then scale as $A \propto [1 - f(\epsilon_1)][1 - f(\epsilon_2)]$ and $B \propto f(\epsilon_1)[1 - f(\epsilon_2)]$, respectively, where $\epsilon_{1,2}$ are the energies of the two fermions involved in the process and $f(\epsilon) = 1/[1 + \exp(-\epsilon/T)]$ is the Fermi-Dirac distribution. Furthermore, type (A) processes were shown to dominate the response at high frequencies, while type (B) dominate the response at low frequencies. Following the steps outlined in Ref. [47] then, we consider the T -dependence of the integrated intensities, $n_L = \int_0^{\omega_1} d\omega I(\omega)$ and $n_H = \int_{\omega_2}^{\infty} d\omega I(\omega)$, with $\omega_1=0.25$ and $\omega_2=0.5$, and fit our numerical results for n_L and n_H to the expressions, respectively,

$$y_L = a_L f(\epsilon_L^*) [1 - f(\epsilon_L^*)] + b_L, \quad y_H = a_H [1 - f(\epsilon_H^*)]^2 + b_H, \quad (5)$$

where a_L , b_L , a_H , and b_H are fitting parameters, and $\epsilon_L^* = 0.42$, $\epsilon_H^* = 0.58$ (see detailed justification for the choice of these parameters (and $\omega_{1,2}$ above) in Ref. [47]).

Figs. 6 (a-b) show two representative fits of n_L and n_H , obtained from a least-squared procedure, for two parameter points inside the Néel phase, one very close to the boundary with the Kitaev QSL phase (a) and another further away (b). The agreement between our numerical data for the integrated intensities and the approximate fermionic expressions

of Eq. (5) is very satisfactory close to the boundary, for both n_L and n_H . As we depart further away from the boundary, however, the agreement remains good only for n_H , showing that the low-frequency part of the response is now controlled by magnon-like processes with different quantum statistical properties, while the high-frequency part still tracks the behavior expected for fractionalized fermionic excitations.

This picture is more vividly presented in panel c which shows the evolution of the normalized standard deviations for the two fits, σ_L and σ_H (defined in the caption of Fig. 6), in the JK -model, as a function of θ .

VI. Discussion

We have studied the Kitaev-Heisenberg honeycomb model, also with an additional Γ -term, using high- and low-temperature numerical approaches. We have found signatures of the fermionic Majorana quasi-particles, characteristic of the proximate Kitaev QSL, in the energy spectrum even when long-range magnetic ordering at very low T is present. This is rather direct evidence that proximate quantum spin liquid physics can be observed at intermediate and high energies/frequencies even when the spectrum at low frequencies is strongly rearranged by a phase transition.

From a point of view of basic quantum statistical mechanics, it is worth commenting on the success of the typicality approach for describing the Kitaev quantum spin liquid. Even for system sizes as large as 24 spins, we had little trouble reaching




FIG. 6. **Fermionic character of the Raman response.** (a-b) T -dependence of the integrated Raman intensities $n_{\text{tot}} = \int_0^\infty d\omega I(\omega)$ (black circles), $n_L = \int_0^{\omega_1} d\omega I(\omega)$ (blue triangles) and $n_H = \int_{\omega_2}^\infty d\omega I(\omega)$ (red diamonds), where $\omega_1 = 0.25$, $\omega_2 = 0.5$ (as in Ref. [47]), for the JK model with $\theta = 0.49\pi$ (a) and 0.46π (b). Solid lines are fits to Eqs. (5), with $\varepsilon_L^* = 0.42$ and $\varepsilon_H^* = 0.58$ (as in Ref. [47]) with a_L, b_L, a_H and b_H determined by a least squared fitting procedure. (c) The standard deviations of the fits, $\sigma_L = \sqrt{\sum_{i=1}^p (n_L(T_i) - y_L(T_i))^2 / p}$, and similarly for σ_H [where y_L and y_H are defined in Eq. (5)], as we depart from the AF Kitaev point in the JK model parametrised as $K = \sin \theta$, $J = \cos \theta$.

reduced coupling strengths K/T well in excess of 10. As we explain in more detail in the appendix, to which we have relegated a proper technical discussion, the success of this method is based on two main ingredients. The first is the generation of a vector $(\frac{1}{\sqrt{L}} e^{-\beta H/2} |r\rangle)$ corresponding to the coupling strength (or, in microcanonical terms, energy density) in question. This is achieved by subjecting a state $|r\rangle$, random in a local spin basis (Eq. 6), to an imaginary time evolution (Eq. 9).

The second ingredient is essentially equivalent with the eigenstate thermalisation hypothesis. [98–102] This states that such a randomly chosen eigenvector at a given energy reproduces correlations characteristic of a thermodynamic ensemble at that energy.

Both of these ingredients are present if, for the finite-size cluster under consideration, the density of states remains sufficiently large at the energy in question. This is evidently a question of ‘detail’, as the size of the eigenvectors that fit into the computer memory in practise correspond to sizes of a few dozen spins at best, a long way from the thermodynamic limit. The important general insight here is that highly frustrated systems – by their very nature – are ideal platforms for satisfying this condition. In these systems, competing interactions lead to huge ground state degeneracies in simple idealised model systems. [103]

It is this huge ground state degeneracy which underpins much of the interest in frustrated magnets, as it renders them unstable to a host of different correlated magnetic phases. Crucially, such instabilities only lift the degeneracy of the idealised model on the scale of the perturbation generating them. The entropy of this degenerate manifold gets spread over only this scale. Compared to a conventional magnet, this shows up in a strong downshift of spectral weight to energies below that of the leading term in the Hamiltonian. In fact, this distinguishes frustrated magnetism from ‘low-dimensional’ routes to the sup-

pression of magnetic ordering, and indeed, this has been proposed as a practical diagnostic for this class of materials. [74].

From the present perspective, this spectral weight downshift is precisely what is needed for the typicality method to work. This therefore accounts for both the good convergence of the method in the case of the pure Kitaev model; and for its robustness in the case of proximate spin liquidity, where it successfully accounts for the physics above the energy scale of the perturbation. Amusingly, the breakdown of the typicality method therefore coincides with that of the proximate spin liquid.

This immediately implies that for the study of the ‘correlated paramagnetic’ behaviour of frustrated magnets, the typicality method should be useful much more generally – what’s needed are fluctuations over an large manifold of low-lying states, e.g. those that obey some kind of local constraint. Well-known examples are the low-lying singlets seen in the kagome antiferromagnet [104, 105] and their interpretation in terms of nearest-neighbour valence bond states, [5, 106–109] or the two-in/two-out states in pyrochlore spin ice. [110–113]

Finally, adapting a less methodological viewpoint, it is perhaps also interesting to try to extend the phenomena observed here beyond highly frustrated magnets, by viewing these phenomena more generally from the perspective of confinement of fractional quasiparticles. For instance, this could be a useful analogy for the confinement of spinons in quasi-1D spin chain materials, like $\text{SrCo}_2\text{V}_2\text{O}_8$, [63] at very low T . Given that the elementary magnon excitations associated with magnetic order carry integer quantum numbers, it is tempting to think of magnons as bound states of fractionalized quasi-particles, and the ordering as a condensation of such bound states. In such a scenario, the confining potential between the quasi-particles would be most effective below T_N whereas, above T_N , deconfined quasi-particles could remain evident in the spec-

trum. This latter regime may also be accessible to the typicality method. The capacity of the typicality method to approach the critical coupling of such a transition, here or in the 2D case, potentially relevant to iridates and ruthenates like α -RuCl₃, also remains to be explored.

We hope that our study will motivate further detailed studies to a broader range of candidate quantum spin liquids, frustrated magnets and disordered quantum magnets, opening a window to their finite- but low-temperature properties that are otherwise inaccessible.

Appendix A: Typicality method

In the following, we shed some more light into the mechanism behind the success of the typicality method. We start with a brief description of the method. For a given finite-size cluster (see next section), we exploit symmetries to reduce the size of the Hilbert space. Then, for each given symmetry sector, we choose random initial states

$$|r\rangle = \sum_{\ell=1}^D d_{\ell} |\ell\rangle / \sqrt{\xi}, \quad (6)$$

where D is the dimension of the Hilbert space inside the given sector, $\{|\ell\rangle, \ell = 1-D\}$ is a basis of product-like states, and $\xi=D/3$ is a normalization constant. The coefficients d_{ℓ} are drawn with a uniform distribution in the unit circle, with

$$\overline{d_{\ell}^* d_{\ell'}} = \overline{|d|^2} \delta_{\ell\ell'}, \quad \overline{|d|^2} = 1/3. \quad (7)$$

The analogous relation for the correlations between the coefficients $r_i = \langle i|r\rangle$, where $|i\rangle$ are the eigenstates of \mathcal{H} with $\mathcal{H}|i\rangle = E_i|i\rangle$, is

$$\overline{r_i^* r_{i'}} = \frac{1}{\xi} \sum_{\ell\ell'} \overline{d_{\ell}^* d_{\ell'}} \langle \ell|i\rangle \langle i'|\ell'\rangle = 1/D \delta_{ii'}. \quad (8)$$

We then propagate the random state $|r\rangle$ in imaginary time using the standard Lanczos method, [88–91] to obtain the state

$$|r, \beta\rangle \equiv e^{-\beta\mathcal{H}/2}|r\rangle. \quad (9)$$

Roughly speaking, during the imaginary time evolution, the coefficients r_i are amplified (suppressed) depending on whether the corresponding energy E_i is below (above) $T = 1/\beta$, respectively. Ideally, at large enough β , the normalized state $|r, \beta\rangle / \sqrt{\zeta_r}$ converges to the ground state of \mathcal{H} within the given sector. The speed of convergence to the ground state is related to the sparseness of the spectrum, and the size of the first excitation gap in particular. [91]

Fig. 7 shows the expectation value of \mathcal{H} in $|r, \beta\rangle / \sqrt{\zeta_r}$,

$$\langle \mathcal{H} \rangle_{r, \beta} \equiv \langle r, \beta | \mathcal{H} | r, \beta \rangle / \zeta_r, \quad (10)$$

divided by system size N , for a large number of random states $|r\rangle$, for three system sizes ($N = 16, 24$ and 32 sites), and for two special points in parameter space, the ideal AF Heisenberg point (panel **a**) and the ideal AF Kitaev point (panel **b**). For the Néel state (**a**), the curves converge very quickly with increasing

β , and the converged values correspond to the ground state energy for each given symmetry sector. In particular, the large spreading in the converged values for each given cluster arises: i) from the fact that we have taken random states in many different sectors (24, 32 and 20 sectors for the 16-, 24- and 32-site cluster, respectively) and, more importantly, ii) from sparseness of the energy spectrum at the Néel point, see Fig. 4 of the main text.

The situation is very different for the Kitaev point (panel **b**). Here, the convergence to the ground state is much slower compared to the Néel state, which reflects the very dense spectrum at the Kitaev point and, in particular, the tiny energy gap to the first excited state (see Fig. 4). Furthermore, the curves from all different initial states and all different clusters fall on top of each other, and the width of the distribution becomes narrower as we increase the system size.

Let us now discuss why typicality works so well in the vicinity of the Kitaev points. There are two main ingredients behind the success of the typicality method. The first is that the state $|r, \beta\rangle$ is dominated by eigenstates $|i\rangle$, with E_i lying inside a small window around the canonical expectation value \mathcal{H} , and in particular, that

$$\langle \mathcal{H} \rangle_{r, \beta} \approx \text{Tr}[e^{-\beta\mathcal{H}} \mathcal{H}] / \text{Tr}[e^{-\beta\mathcal{H}}]. \quad (11)$$

The second ingredient for the success of typicality is that the expectation value of some local operators A (static or dynamic) in the state $|r, \beta\rangle$ is the same with the result we get with the canonical ensemble,

$$\langle A \rangle_{r, \beta} \equiv \langle r, \beta | A | r, \beta \rangle / \zeta_r \approx \text{Tr}[A e^{-\beta\mathcal{H}}] / \text{Tr}[e^{-\beta\mathcal{H}}]. \quad (12)$$

Given the first ingredient, Eq. (12) is equivalent with the so-called eigenstate thermalization hypothesis (ETH). [98–102] We will now show how both these ingredients are guaranteed if the density of states is sufficiently large. To this end, we rewrite

$$\zeta_r = \sum_i |r_i|^2 e^{-\beta E_i}, \quad \langle \mathcal{H} \rangle_{r, \beta} = \sum_i |r_i|^2 e^{-\beta E_i} E_i / \zeta_r, \quad (13)$$

and then break up the sum over i into a sum over energy windows $[E - \delta E/2, E + \delta E/2]$ around E and a sum over states j in each window,

$$\zeta_r \rightarrow \sum_E \delta N(E) e^{-\beta E} g_r(E), \quad (14)$$

$$\langle \mathcal{H} \rangle_{r, \beta} \rightarrow \sum_E \delta N(E) E e^{-\beta E} g_r(E) / \zeta_r,$$

where $\delta N(E) = \rho(E)\delta E$ and the quantity

$$g_r(E) = \sum_{j=1}^{\delta N(E)} |r_{E,j}|^2 / \delta N(E), \quad (15)$$

is the average of $|r_i|^2$ inside the window around E . The above way of rewriting the sums in (13) as sums over energy windows has a meaning only if most of the coefficients $|r_{E,j}|^2$ are finite. If, in addition, $\delta N(E)$ is large enough then we can use the central limit theorem to make a statement for the distribution of $g_r(E)$ (for fixed E). According to that theorem, in




FIG. 7. Expectation value of the energy per site, $\langle H \rangle_{r,\beta} / N$, in the normalized state $|r, \beta\rangle / \sqrt{\zeta_r}$ [see definition in Eq. (10)] for a number of initial random states $|r\rangle$ (10 in each irreducible representation; In total, 240 states for the 16-site cluster, 320 for the 24-site cluster, and 200 for the 32-site cluster; for the latter we only show data from the sectors that are even under spin inversion). Panels (a) and (b) correspond to the AF Heisenberg and the AF Kitaev point, respectively.

the limit of large $\delta N(E)$, the distribution of $g_r(E)$ (for fixed E) approaches a normal distribution with standard deviation $\sigma_E^2 / \delta N(E)$, where σ_E^2 is the variance of the individual coefficients $|r_{E,j}|^2$. Therefore, if $\delta N(E)$ is large we can replace

$$|r_{E,j}|^2 \rightarrow \overline{|r|^2} = 1/D \quad (16)$$

in (15), which in turn gives $g_r(E) \approx 1/D$, i.e. $g_r(E)$ becomes independent of the random state $|r\rangle$, see below. The equivalence with the canonical ensemble follows immediately.

We have already emphasised the need for a large enough density of states. In practice, we only need this condition to hold in a narrow region of energy that is fixed by β . Indeed, for large enough system sizes (and finite energy density E/N), the density of states $\rho(E)$ scales exponentially with system size N , namely, $\rho(E) \propto e^{Ns(E)}$, where $s(E)$ is the microcanonical entropy per site. Then, the sum over E in (14) is dominated by a very narrow region around a characteristic energy E^* fixed by $s'(E^*) = \beta$. Using the steepest descent method then gives, for example,

$$\zeta_r(\beta) \propto \rho(E^*) e^{-\beta E^*}. \quad (17)$$

We can test this numerically by checking when the quantity $\ln(\zeta_r)/N$ is independent of system size and r . This is




FIG. 8. Evolution of the quantity $\ln(\zeta_r)/N$, where $\zeta_r = \langle r, \beta | r, \beta \rangle$, with inverse temperature for a number of initial random states $|r\rangle$ (same as in Fig. 7). Panels (a) and (b) correspond to the AF Heisenberg and the AF Kitaev point, respectively.

demonstrated in Fig. 8, which shows this quantity for the same clusters and number of initial states considered in Fig. 7. For the Néel state (panel a), the various curves begin to deviate from each other around $J/T \sim 1$, which roughly coincides with the coupling at which the quantities $\langle H \rangle_{r,\beta}$ begin to converge to the ground state energy of the given sector (compare with Fig. 7).

For the Kitaev point (panel b), on the other hand, the various curves for the quantity $\ln(\zeta_r)/N$ converge to each other, and the spreading of the distribution becomes again narrower and narrower as we increase the system size. This picture is in essence a numerical proof of Eq. (17) and the statement that the density of states is exponentially large with system size, at least for the 24- and 32-site clusters. Equation (17) also says that most of the weight of $|r, \beta\rangle$ comes from eigenstates within the window around E^* , which is the first ingredient behind the success of the typicality method.

We can proceed in a similar way to show that the second ingredient, Eq. (12), is also guaranteed if (4) holds. We have

$$\langle A \rangle_{r,\beta} = \frac{1}{\zeta_r} \langle r, \beta | A | r, \beta \rangle = \frac{1}{\zeta_r} \sum_{i,i'} r_i^* r_{i'} e^{-\beta E_i/2} e^{-\beta E_{i'}/2} \langle i | A | i' \rangle. \quad (18)$$

As above, if the number of finite terms $r_i^* r_{i'}$ involved in the above sum (and in the relevant energy regime) is large enough

we can replace with their mean value given in Eq. (8) to get

$$\langle r, \beta | A | r, \beta \rangle \approx 1/D \sum_i e^{-\beta E_i} \langle i | A | i \rangle. \quad (19)$$

The equivalence with the canonical ensemble follows immediately and, as above, the sum over i is dominated by states with $E_i \sim E^*$. Altogether, the success of the typicality method boils down to the presence of a large density of states $\rho(E)$. As discussed in the main text, this is typical for finite-size spectra in the middle of the spectrum, but ceases to hold below a characteristic energy scale that depends on system size and scales with the strength of the (perturbing) interactions. The distinctive feature of the Kitaev QSL that renders the typicality method successful down to very low T is the remarkable spectral downshift (see Fig. 4), that occurs generically in highly frustrated systems.

Appendix B: Finite-size clusters used in our simulations

Fig. 9 shows the four main finite-size clusters with periodic boundary conditions used in our exact diagonalizations. The one shown in the left bottom panel has 24 sites and the full point group symmetry of the infinite system. The stochastic method results and the spectra shown in the main text are taken on this cluster. The clusters shown in the left top panel and the right panel have 16 and 32 sites, respectively, and lower point groups than that of the infinite system. The results shown in Figs. 7 and 8 are taken on the 16-, 24- and 32-site clusters. The cluster in the middle panel has 24 sites but has lower point group symmetry than the infinite system. The LTLM results of Fig. 11 are taken on this cluster. The symmetries exploited on these clusters include the translations and real space inversion. For the JK -model, we have also exploited the global two-fold rotation around the \mathbf{x} -axis in spin space.

Appendix C: Boundaries of the Kitaev QSL in the JK model

Fig. 10 shows the ground state expectation value of Kitaev's six-body flux operator [7] $W = 2^6 S_1^x S_2^y S_3^z S_4^x S_5^y S_6^z$ as a function of θ , where $K = \sin \theta$ and $J = \cos \theta$. The results are obtained from exact diagonalizations on the three clusters of Fig. 9. The Kitaev QSL phases correspond to the regions around the ideal Kitaev points $\theta = \pm\pi/2$, where $\langle W \rangle$ is very close to 1. [7] The transitions to the magnetically ordered phases correspond to the points where $\langle W \rangle$ drops abruptly to very small values. The shaded QSL regions in Fig. 10 correspond to the symmetric 24-site cluster of Fig. 9 which has the full point group symmetry of the model. The boundaries are consistent with those reported in Ref. [53] and the reorganization of the low-lying excitation spectra in Fig. 4 of the main text.

Appendix D: Results from low- T Lanczos method

The low- T Lanczos method (LTLM) [92] is an adaptation of the earlier, finite- T Lanczos algorithm [85, 87] that correctly captures the zero temperature limit. This method delivers

the response directly in frequency domain and the thermodynamic trace is performed exactly at low enough T , by keeping contributions from all low-lying excitations below an energy cutoff. Specifically, one starts again with r_m random states $|r\rangle$ and obtain, for each state, a set of M approximate eigenstates $\{|\epsilon_i^{(r)}\rangle, i=1-M\}$ by a standard Lanczos iteration scheme. [88–91] The partition function is then approximated by

$$Z \simeq \frac{D}{r_m} \sum_r \sum_i^M |\langle \epsilon_i^{(r)} | r \rangle|^2 e^{-\beta \epsilon_i^{(r)}}, \quad (20)$$

and any given spectral density $C(\omega) \equiv \frac{1}{\pi} \int dt e^{i\omega t} \langle A^\dagger(t) A(0) \rangle$ by

$$C(\omega) \simeq \frac{D}{Z r_m} \sum_r \sum_{i,l,k}^M e^{-\beta(\epsilon_i^{(r)} + \epsilon_k^{(r)})/2} \langle r | \epsilon_i^{(r)} \rangle \langle \epsilon_i^{(r)} | A^\dagger | \epsilon_l^{(r)} \rangle \\ \times \langle \epsilon_l^{(r)} | A | \epsilon_k^{(r)} \rangle \langle \epsilon_k^{(r)} | r \rangle \delta\left(\omega - \tilde{\epsilon}_l^{(r)} + \frac{1}{2}(\epsilon_i^{(r)} + \epsilon_k^{(r)})\right). \quad (21)$$

Here, D is the dimension of the full Hilbert space and $|\tilde{\epsilon}_l^{(r)}\rangle$ are eigenstates of \mathcal{H} with corresponding eigenenergies $\tilde{\epsilon}_l^{(r)}$, obtained by an additional Lanczos run with initial state $A|\epsilon_k^{(r)}\rangle$




FIG. 9. **Finite-size clusters used in our simulations.** All clusters have periodic boundary conditions, and $N = 16$ (left top), 24 (left bottom and middle) or 32 sites (right). The first has the full point group symmetry of the model.




FIG. 10. **Phase boundaries in the JK -model.** Ground state expectation value of Kitaev's hexagonal plaquette operator [7] $W = 2^6 S_1^x S_2^y S_3^z S_4^x S_5^y S_6^z$ (site labeling shown in the inset) in the JK -model, with $K = \sin \theta$, $J = \cos \theta$. Note the nonlinear horizontal axis.

for each k . In practice, $M \sim O(100)$ is sufficient to obtain accurate estimates for observables of a sparse matrix \mathcal{H} , even for $D \sim O(10^8)$. Nevertheless, the computational cost is prohibitive if all three sums are over Lanczos bases with $M \sim 100$. Since higher-energy eigenvalues are exponentially suppressed by their Boltzmann factors at low T , one can further restrict the summations over i and k , so that $\epsilon_{i,k} < \epsilon_c$ and the energy threshold ϵ_c is in turn set by the condition

$$e^{-\beta_c(\epsilon_c - \epsilon_0)} < \epsilon_c, \quad (22)$$

where ϵ_c is the high-energy fraction of the Boltzmann factors to be truncated and β_c is the lowest inverse temperature one is interested in. For example, $\epsilon_c = 0.05$ means that one ignores all eigenstates that contribute 5% of the high-energy tail of the Boltzmann weight. A good accuracy is thus guaranteed for all $\beta \geq \beta_c$. Furthermore, the calculation can be performed independently in each symmetry sector (as in the stochastic method), and the final expectation value is then obtained after an additional averaging over all sectors. Finally, the δ -peaks in Eq. (21) are broadened into Lorentzians by a common broadening parameter η . Here, the value of η has been chosen to facilitate a comparison with results from the typicality method, which misses the finer finite-size structure of the spectra at very low T (due to the breakdown of typicality below T_{typ}).

We have used the LTLM to study the xy -channel Raman response of the Kitaev-Heisenberg model for periodic clusters of up to 24 sites in the window $\beta \geq \beta_c = 3$, within which

numerical calculations can be performed with available resources. Two representative set of results are shown in Fig. 11 for the cluster labeled as ‘24b’ in Fig. 9, along with the values of the LTLM parameters r_m , M , η and ϵ_c used. We have verified that the results do not change appreciably when these parameters are varied. The spectra share the same qualitative features with those from the typicality method. [114] Close to the Kitaev point, but on either side of its boundary with the Néel phase, the Raman response has two main features, namely, a primary peak at $\omega \sim 1$ and a secondary peak/shoulder at $\omega \sim 2$, which disperse with increasing T to lower and higher frequencies respectively (more markedly for $\theta = 0.485\pi$). These observations are also in accordance with published Monte Carlo results at the Kitaev point. [47]

Acknowledgements

We thank W. Brenig, K. Burch, M. Daghofer and A. Honecker for fruitful discussions. IR acknowledges the hospitality of the Max Planck Institute for the Physics of Complex Systems (MPI-PKS) of Dresden, where part of this work was done. This work was in part supported by the Deutsche Forschungsgemeinschaft via grant SFB 1143. SK was partially supported through the Boston University Center for Non-Equilibrium Systems and Computation. IR and NBP were supported by the U.S. Department of Energy, Office of Science, Basic Energy Sciences under Award #DE-SC0018056.

-
- [1] P.W. Anderson, “Resonating valence bonds: A new kind of insulator?” *Mat. Res. Bull.* **8**, 153 – 160 (1973).
 - [2] P. Fazekas and P. W. Anderson, “On the ground state properties of the anisotropic triangular antiferromagnet,” *Phil. Mag.* **30**, 423–440 (1974).
 - [3] V. Kalmeyer and R. B. Laughlin, “Equivalence of the resonating-valence-bond and fractional quantum Hall states,” *Phys. Rev. Lett.* **59**, 2095–2098 (1987).
 - [4] X. G. Wen, Frank Wilczek, and A. Zee, “Chiral spin states and superconductivity,” *Phys. Rev. B* **39**, 11413–11423 (1989).
 - [5] R. Moessner, S. L. Sondhi, and Eduardo Fradkin, “Short-ranged resonating valence bond physics, quantum dimer models, and Ising gauge theories,” *Phys. Rev. B* **65**, 024504 (2001).
 - [6] A.Yu. Kitaev, “Fault-tolerant quantum computation by anyons,” *Annals of Physics* **303**, 2 – 30 (2003).
 - [7] Alexei Kitaev, “Anyons in an exactly solved model and beyond,” *Annals of Physics* **321**, 2 – 111 (2006).
 - [8] Leon Balents, “Spin liquids in frustrated magnets,” *Nature* **464**, 199–208 (2010).
 - [9] Lucile Savary and Leon Balents, “Quantum spin liquids: a review,” *Rep. Prog. Phys.* **80**, 016502 (2017).
 - [10] Yi Zhou, Kazushi Kanoda, and Tai-Kai Ng, “Quantum spin liquid states,” *Rev. Mod. Phys.* **89**, 025003 (2017).
 - [11] Ioannis Rousochatzakis, Yuriy Szytyuk, and Natalia B. Perkins, “Quantum spin liquid in the semiclassical regime,” *Nat. Commun.* **9**, 1575 (2018).
 - [12] Johannes Knolle and Roderich Moessner, “A Field Guide to Spin Liquids,” *arXiv:1804.02037* (2018).
 - [13] Matthew P. Shores, Emily A. Nytko, Bart M. Bartlett, and Daniel G. Nocera, “A Structurally Perfect $S = 1/2$ Kagome Antiferromagnet,” *J. Am. Chem. Soc.* **127**, 13462–13463 (2005).
 - [14] Tian-Heng Han, Joel S. Helton, Shaoyan Chu, Daniel G. Nocera, Jose A. Rodriguez-Rivera, Collin Broholm, and Young S. Lee, “Fractionalized excitations in the spin-liquid state of a kagome-lattice antiferromagnet,” *Nature* **492**, 406–410 (2012).
 - [15] M. R. Norman, “Colloquium,” *Rev. Mod. Phys.* **88**, 041002 (2016).
 - [16] G. Jackeli and G. Khaliullin, “Mott Insulators in the Strong Spin-Orbit Coupling Limit: From Heisenberg to a Quantum Compass and Kitaev Models,” *Phys. Rev. Lett.* **102**, 017205 (2009).
 - [17] J. Chaloupka, George Jackeli, and Giniyat Khaliullin, “Kitaev-Heisenberg Model on a Honeycomb Lattice: Possible Exotic Phases in Iridium Oxides $A_2\text{IrO}_3$,” *Phys. Rev. Lett.* **105**, 027204 (2010).
 - [18] Gang Cao and Lance DeLong, eds., *Frontiers of 4d- and 5d-Transition Metal Oxides* (World Scientific Publishing Co. Pte. Ltd., 2013).
 - [19] William Witczak-Krempa, Gang Chen, Yong Baek Kim, and Leon Balents, “Correlated Quantum Phenomena in the Strong Spin-Orbit Regime,” *Ann. Rev. Cond. Matt. Phys.* **5**, 57–82 (2014).
 - [20] Jeffrey G. Rau, Eric Kin-Ho Lee, and Hae-Young Kee, “Spin-Orbit Physics Giving Rise to Novel Phases in Correlated Systems: Iridates and Related Materials,” *Ann. Rev. Cond. Matt. Phys.* **7**, 195 (2016).
 - [21] Simon Trebst, “Kitaev Materials,” *arXiv:1701.07056* (2017).
 - [22] M. Hermanns, I. Kimchi, and J. Knolle, “Physics of the Kitaev




FIG. 11. **Results for the LTLM method.** Raman response in the xy -channel at (a) the Kitaev QSL point, and (b) $\theta = 0.485\pi$ (within the Néel phase). Standard error due to the sampling involved in Eq. (21) is shown as error bars. Dashed grey lines mark the shift of the main peak with T . Peaks are broadened with a Lorentzian of width $\eta = 0.2$.

Model: Fractionalization, Dynamic Correlations, and Material Connections,” *Annual Review of Condensed Matter Physics* **9**, 17–33 (2018), <https://doi.org/10.1146/annurev-conmatphys-033117-053934>.

- [23] Stephen M. Winter, Alexander A. Tsirlin, Maria Daghofer, Jeroen van den Brink, Yogesh Singh, Philipp Gegenwart, and Roser Valenti, “Models and Materials for Generalized Kitaev Magnetism,” *J. Phys.: Condens. Matter* **29**, 493002 (2017).
- [24] Yogesh Singh and P. Gegenwart, “Antiferromagnetic Mott insulating state in single crystals of the honeycomb lattice material Na_2IrO_3 ,” *Phys. Rev. B* **82**, 064412 (2010).
- [25] Yogesh Singh, S. Manni, J. Reuther, T. Berlijn, R. Thomale, W. Ku, S. Trebst, and P. Gegenwart, “Relevance of the Heisenberg-Kitaev Model for the Honeycomb Lattice Iridates A_2IrO_3 ,” *Phys. Rev. Lett.* **108**, 127203 (2012).
- [26] X. Liu, T. Berlijn, W.-G. Yin, W. Ku, A. Tsvelik, Young-June Kim, H. Gretarsson, Yogesh Singh, P. Gegenwart, and J. P. Hill, “Long-range magnetic ordering in Na_2IrO_3 ,” *Phys. Rev. B* **83**, 220403 (2011).
- [27] J. A. Sears, M. Songvilay, K. W. Plumb, J. P. Clancy, Y. Qiu, Y. Zhao, D. Parshall, and Young-June Kim, “Magnetic order in $\alpha - \text{RuCl}_3$: A honeycomb-lattice quantum magnet with strong spin-orbit coupling,” *Phys. Rev. B* **91**, 144420 (2015).
- [28] R. D. Johnson, S. C. Williams, A. A. Haghighirad, J. Singleton, V. Zapf, P. Manuel, I. I. Mazin, Y. Li, H. O. Jeschke, R. Valentí, and R. Coldea, “Monoclinic crystal structure of $\alpha - \text{RuCl}_3$ and the zigzag antiferromagnetic ground state,” *Phys. Rev. B* **92**, 235119 (2015).
- [29] S. C. Williams, R. D. Johnson, F. Freund, Sungkyun Choi, A. Jesche, I. Kimchi, S. Manni, A. Bombardi, P. Manuel, P. Gegenwart, and R. Coldea, “Incommensurate counterrotating magnetic order stabilized by Kitaev interactions in the layered honeycomb $\alpha - \text{Li}_2\text{IrO}_3$,” *Phys. Rev. B* **93**, 195158 (2016).
- [30] A. Biffin, R. D. Johnson, I. Kimchi, R. Morris, A. Bombardi, J. G. Analytis, A. Vishwanath, and R. Coldea, “Noncoplanar and Counterrotating Incommensurate Magnetic Order Stabilized by Kitaev Interactions in $\gamma - \text{Li}_2\text{IrO}_3$,” *Phys. Rev. Lett.* **113**, 197201 (2014).
- [31] KA Modic, Tess E Smidt, Itamar Kimchi, Nicholas P Breznay, Alun Biffin, Sungkyun Choi, Roger D Johnson, Radu Coldea, Pilanda Watkins-Curry, Gregory T McCandess, *et al.*, “Realization of a three-dimensional spin-anisotropic harmonic honeycomb iridate,” *Nature communications* **5** (2014), 10.1038/ncomms5203.
- [32] A. Biffin, R. D. Johnson, Sungkyun Choi, F. Freund, S. Manni, A. Bombardi, P. Manuel, P. Gegenwart, and R. Coldea, “Unconventional magnetic order on the hyperhoneycomb Kitaev lattice in $\beta - \text{Li}_2\text{IrO}_3$: Full solution via magnetic resonant x-ray diffraction,” *Phys. Rev. B* **90**, 205116 (2014).
- [33] T. Takayama, A. Kato, R. Dinnebier, J. Nuss, H. Kono, L. S. I. Veiga, G. Fabbris, D. Haskel, and H. Takagi, “Hyperhoneycomb Iridate $\beta - \text{Li}_2\text{IrO}_3$ as a Platform for Kitaev Magnetism,” *Phys. Rev. Lett.* **114**, 077202 (2015).
- [34] S. E. Nagler, D. A. Tennant, R. A. Cowley, T. G. Perring, and S. K. Satija, “Spin dynamics in the quantum antiferromagnetic chain compound KCuF_3 ,” *Phys. Rev. B* **44**, 12361–12368 (1991).
- [35] D. A. Tennant, T. G. Perring, R. A. Cowley, and S. E. Nagler, “Unbound spinons in the $S = 1/2$ antiferromagnetic chain KCuF_3 ,” *Phys. Rev. Lett.* **70**, 4003–4006 (1993).
- [36] Martin Mourigal, Mechthild Enderle, Axel Kloppepiper, Jean-Sebastien Caux, Anne Stunault, and Henrik M. Ronnow, “Fractional spinon excitations in the quantum Heisenberg antiferromagnetic chain,” *Nat. Phys.* **9**, 435 – 441 (2013).
- [37] J. Knolle, D. L. Kovrizhin, J. T. Chalker, and R. Moessner, “Dynamics of fractionalization in quantum spin liquids,” *Phys. Rev. B* **92**, 115127 (2015).
- [38] J. Knolle, D. L. Kovrizhin, J. T. Chalker, and R. Moessner, “Dynamics of a Two-Dimensional Quantum Spin Liquid: Signatures of Emergent Majorana Fermions and Fluxes,” *Phys. Rev. Lett.* **112**, 207203 (2014).
- [39] A Banerjee, CA Bridges, J-Q Yan, AA Aczel, L Li, MB Stone, GE Granroth, MD Lumsden, Y Yiu, J Knolle, *et al.*, “Proximate Kitaev quantum spin liquid behaviour in a honeycomb magnet,” *Nature materials* (2016), 10.1038/nmat4604.
- [40] A. Banerjee, J.-Q. Yan, J. Knolle, C. A. Bridges, M. B. Stone, M. D. Lumsden, D. G. Mandrus, D. A. Tennant, R. Moessner, and S. E. Nagler, “Neutron scattering in the proximate quantum spin liquid $\alpha - \text{RuCl}_3$,” *Science* (2017), 10.1126/science.aah6015.
- [41] Dirk Wulferding, Peter Lemmens, Patric Scheib, Jens Röder, Philippe Mendels, Shaoyan Chu, Tianheng Han, and Young S. Lee, “Interplay of thermal and quantum spin fluctuations in the kagome lattice compound herbertsmithite,” *Phys. Rev. B* **82**, 144412 (2010).
- [42] Wing-Ho Ko, Zheng-Xin Liu, Tai-Kai Ng, and Patrick A. Lee,

- “Raman signature of the U(1) Dirac spin-liquid state in the spin- $\frac{1}{2}$ kagome system,” *Phys. Rev. B* **81**, 024414 (2010).
- [43] Luke J. Sandilands, Yao Tian, Kemp W. Plumb, Young-June Kim, and Kenneth S. Burch, “Scattering Continuum and Possible Fractionalized Excitations in α -RuCl₃,” *Phys. Rev. Lett.* **114**, 147201 (2015).
- [44] Luke J. Sandilands, Yao Tian, Anjan A. Reijnders, Heung-Sik Kim, K. W. Plumb, Young-June Kim, Hae-Young Kee, and Kenneth S. Burch, “Spin-orbit excitations and electronic structure of the putative Kitaev magnet α -RuCl₃,” *Phys. Rev. B* **93**, 075144 (2016).
- [45] A. Glamazda, P. Lemmens, S.-H. Do, Y. S. Choi, and K.-Y. Choi, “Raman spectroscopic signature of fractionalized excitations in the harmonic-honeycomb iridates β - and γ -Li₂IrO₃,” *Nat. Commun.* **7**, 12286 (2016).
- [46] J. Knolle, Gia-Wei Chern, D. L. Kovrizhin, R. Moessner, and N. B. Perkins, “Raman Scattering Signatures of Kitaev Spin Liquids in A₂IrO₃ Iridates with A = Na or Li,” *Phys. Rev. Lett.* **113**, 187201 (2014).
- [47] J. Nasu, J. Knolle, D. L. Kovrizhin, Y. Motome, and R. Moessner, “Fermionic response from fractionalization in an insulating two-dimensional magnet,” *Nat. Phys.* **12**, 912–915 (2016).
- [48] Gábor B. Halász, Natalia B. Perkins, and Jeroen van den Brink, “Resonant Inelastic X-Ray Scattering Response of the Kitaev Honeycomb Model,” *Phys. Rev. Lett.* **117**, 127203 (2016).
- [49] Gábor B. Halász, Brent Perreault, and Natalia B. Perkins, “Probing spinon nodal structures in three-dimensional kitaev spin liquids,” *Phys. Rev. Lett.* **119**, 097202 (2017).
- [50] L. Savary and T. Senthil, “Probing Hidden Orders with Resonant Inelastic X-Ray Scattering,” *arXiv:1506.04752* (2015).
- [51] Zhanybek Alpichshev, Fahad Mahmood, Gang Cao, and Nuh Gedik, “Confinement-Deconfinement Transition as an Indication of Spin-Liquid-Type Behavior in Na₂IrO₃,” *Phys. Rev. Lett.* **114**, 017203 (2015).
- [52] Robert Schaffer, Subhro Bhattacharjee, and Yong Baek Kim, “Quantum phase transition in Heisenberg-Kitaev model,” *Phys. Rev. B* **86**, 224417 (2012).
- [53] J. Chaloupka, G. Jackeli, and G. Khaliullin, “Zigzag Magnetic Order in the Iridium Oxide Na₂IrO₃,” *Phys. Rev. Lett.* **110**, 097204 (2013).
- [54] Eric Kin-Ho Lee, Robert Schaffer, Subhro Bhattacharjee, and Yong Baek Kim, “Heisenberg-Kitaev model on the hyperhoneycomb lattice,” *Phys. Rev. B* **89**, 045117 (2014).
- [55] Vamshi M Katukuri, S. Nishimoto, V. Yushankhai, A. Stoyanova, H. Kandpal, Sungkyun Choi, R. Coldea, I. Rousochatzakis, L. Hozoi, and Jeroen van den Brink, “Kitaev interactions between $j=1/2$ moments in honeycomb Na₂IrO₃ are large and ferromagnetic: insights from ab initio quantum chemistry calculations,” *New J. Phys.* **16**, 013056 (2014).
- [56] V. M. Katukuri, N. Satoshi, I. Rousochatzakis, H. Stoll, J. van den Brink, and L. Hozoi, “Strong magnetic frustration and anti-site disorder causing spin-glass behavior in honeycomb Li₂RhO₃,” *Sci Rep* (2015), 10.1038/srep14718.
- [57] Ioannis Rousochatzakis, Johannes Reuther, Ronny Thomale, Stephan Rachel, and N. B. Perkins, “Phase Diagram and Quantum Order by Disorder in the Kitaev $K_1 - K_2$ Honeycomb Magnet,” *Phys. Rev. X* **5**, 041035 (2015).
- [58] Satoshi Nishimoto, Vamshi M. Katukuri, Viktor Yushankhai, Hermann Stoll, Ulrich K. Roessler, Liviu Hozoi, Ioannis Rousochatzakis, and Jeroen van den Brink, “Strongly frustrated triangular spin lattice emerging from triplet dimer formation in honeycomb Li(2)IrO(3),” *Nat Commun* **7** (2016), 10.1038/ncomms10273.
- [59] A. Zheludev, M. Kenzelmann, S. Raymond, E. Ressouche, T. Masuda, K. Kakurai, S. Maslov, I. Tsukada, K. Uchinokura, and A. Wildes, “Energy Separation of Single-Particle and Continuum States in an $S = 1/2$ Weakly Coupled Chains Antiferromagnet,” *Phys. Rev. Lett.* **85**, 4799–4802 (2000).
- [60] I. A. Zaliznyak, H. Woo, T. G. Perring, C. L. Broholm, C. D. Frost, and H. Takagi, “Spinons in the Strongly Correlated Copper Oxide Chains in SrCuO₂,” *Phys. Rev. Lett.* **93**, 087202 (2004).
- [61] Bella Lake, D. Alan Tennant, Chris D. Frost, and Stephen E. Nagler, “Quantum criticality and universal scaling of a quantum antiferromagnet,” *Nat. Mater.* **4**, 329–334 (2005).
- [62] M. Enderle, B. Fåk, H.-J. Mikeska, R. K. Kremer, A. Prokofiev, and W. Assmus, “Two-Spinon and Four-Spinon Continuum in a Frustrated Ferromagnetic Spin-1/2 Chain,” *Phys. Rev. Lett.* **104**, 237207 (2010).
- [63] A. K. Bera, B. Lake, F. H. L. Essler, L. Vanderstraeten, C. Hubig, U. Schollwöck, A. T. M. N. Islam, A. Schneidewind, and D. L. Quintero-Castro, “Spinon confinement in a quasi-one-dimensional anisotropic heisenberg magnet,” *Phys. Rev. B* **96**, 054423 (2017).
- [64] P. A. Fleury and R. Loudon, “Scattering of Light by One- and Two-Magnon Excitations,” *Phys. Rev.* **166**, 514–530 (1968).
- [65] B. Sriram Shastry and Boris I Shraiman, “Raman scattering in Mott-Hubbard systems,” *Int. J. Mod. Phys. B* **5**, 365–388 (1991).
- [66] Thomas P. Devereaux and Rudi Hackl, “Inelastic light scattering from correlated electrons,” *Rev. Mod. Phys.* **79**, 175–233 (2007).
- [67] Sandu Popescu, Anthony J. Short, and Andreas Winter, “Entanglement and the foundations of statistical mechanics,” *Nat. Phys.* **2**, 754–758 (2006).
- [68] Sheldon Goldstein, Joel L. Lebowitz, Roderich Tumulka, and Nino Zanghì, “Canonical Typicality,” *Phys. Rev. Lett.* **96**, 050403 (2006).
- [69] Peter Reimann, “Typicality for Generalized Microcanonical Ensembles,” *Phys. Rev. Lett.* **99**, 160404 (2007).
- [70] Tarek A. Elsayed and Boris V. Fine, “Regression Relation for Pure Quantum States and Its Implications for Efficient Computing,” *Phys. Rev. Lett.* **110**, 070404 (2013).
- [71] Robin Steinigeweg, Jochen Gemmer, and Wolfram Brenig, “Spin-Current Autocorrelations from Single Pure-State Propagation,” *Phys. Rev. Lett.* **112**, 120601 (2014).
- [72] R. Steinigeweg, A. Khodja, H. Niemeyer, C. Gogolin, and J. Gemmer, “Pushing the Limits of the Eigenstate Thermalization Hypothesis towards Mesoscopic Quantum Systems,” *Phys. Rev. Lett.* **112**, 130403 (2014).
- [73] Robin Steinigeweg, Jacek Herbrych, Xenophon Zotos, and Wolfram Brenig, “Heat Conductivity of the Heisenberg Spin-1/2 Ladder: From Weak to Strong Breaking of Integrability,” *Phys. Rev. Lett.* **116**, 017202 (2016).
- [74] Ramirez A. P., “Geometrical Frustration,” in *Handbook of Magnetic Materials*, Vol. 13, edited by Buschow K. H. J. (Elsevier, Amsterdam, 2001) Chap. 4, pp. 423–520.
- [75] Jeffrey G. Rau, Eric Kin-Ho Lee, and Hae-Young Kee, “Generic Spin Model for the Honeycomb Iridates beyond the Kitaev Limit,” *Phys. Rev. Lett.* **112**, 077204 (2014).
- [76] Yuriy Sizyuk, Craig Price, Peter Wölfe, and Natalia B. Perkins, “Importance of anisotropic exchange interactions in honeycomb iridates: Minimal model for zigzag antiferromagnetic order in Na₂IrO₃,” *Phys. Rev. B* **90**, 155126 (2014).
- [77] Stephen M. Winter, Ying Li, Harald O. Jeschke, and Roser Valentí, “Challenges in design of Kitaev materials: Magnetic interactions from competing energy scales,” *Phys. Rev. B* **93**, 214431 (2016).

- [78] Heung-Sik Kim and Hae-Young Kee, “Crystal structure and magnetism in α -RuCl₃: An *ab initio* study,” *Phys. Rev. B* **93**, 155143 (2016).
- [79] Ioannis Rousochatzakis and Natalia B. Perkins, “Classical spin liquid instability driven by off-diagonal exchange in strong spin-orbit magnets,” *Phys. Rev. Lett.* **118**, 147204 (2017).
- [80] Joji Nasu, Masafumi Udagawa, and Yukitoshi Motome, “Thermal fractionalization of quantum spins in a Kitaev model: Temperature-linear specific heat and coherent transport of Majorana fermions,” *Phys. Rev. B* **92**, 115122 (2015).
- [81] Brent Perreault, Johannes Knolle, Natalia B. Perkins, and F. J. Burnell, “Theory of Raman response in three-dimensional Kitaev spin liquids: Application to β - and γ -Li₂IrO₃ compounds,” *Phys. Rev. B* **92**, 094439 (2015).
- [82] John Skilling, “Classic Maximum Entropy,” in *Maximum Entropy and Bayesian Methods: Cambridge, England, 1988*, edited by J. Skilling (Springer Netherlands, Dordrecht, 1989) pp. 45–52.
- [83] David A. Drabold and Otto F. Sankey, “Maximum entropy approach for linear scaling in the electronic structure problem,” *Phys. Rev. Lett.* **70**, 3631–3634 (1993).
- [84] R. N. Silver and H. Röder, “Densities of States of Mega-Dimensional Hamiltonian Matrices,” *Int. J. Mod. Phys. C* **5**, 735–753 (1994).
- [85] J. Jaklič and P. Prelovšek, “Finite-temperature properties of doped antiferromagnets,” *Advances in Physics* **49**, 1–92 (2000).
- [86] Alexander Weiße, Gerhard Wellein, Andreas Alvermann, and Holger Fehske, “The kernel polynomial method,” *Rev. Mod. Phys.* **78**, 275–306 (2006).
- [87] P. Prelovšek and J. Bonča, “Ground State and Finite Temperature Lanczos Methods,” in *Strongly Correlated Systems: Numerical Methods*, edited by Adolfo Avella and Ferdinando Mancini (Springer Berlin Heidelberg, Berlin, Heidelberg, 2013) pp. 1–30.
- [88] C Lanczos, “An iteration method for the solution of the eigenvalue problem of linear differential and integral operators,” *J. Res. Nat’l Bur. Std.* **45**, 255–282 (1950).
- [89] C. C. Paige, *The computation of eigenvalues and eigenvectors of very large sparse matrices*, Ph.D. thesis, U. London (1971).
- [90] J. Cullum and R. Willoughby, *Lanczos Algorithms for Large Symmetric Eigenvalue Computations* (Society for Industrial and Applied Mathematics, 2002).
- [91] Y. Saad, *Numerical Methods for Large Eigenvalue Problems* (Society for Industrial and Applied Mathematics, 2011).
- [92] Markus Aichhorn, Maria Daghofer, Hans Gerd Evertz, and Wolfgang von der Linden, “Low-temperature Lanczos method for strongly correlated systems,” *Phys. Rev. B* **67**, 161103 (2003).
- [93] Anthony Hams and Hans De Raedt, “Fast algorithm for finding the eigenvalue distribution of very large matrices,” *Phys. Rev. E* **62**, 4365–4377 (2000).
- [94] In a finite-size cluster, there are only a certain number of discrete momenta available. The low- T Raman response arises then from pairs of magnon modes with total momentum zero, leading to a series of peaks characteristic of the finite-size cluster.
- [95] Andrei Catuneanu, Youhei Yamaji, Gideon Wachtel, Yong Baek Kim, and Hae-Young Kee, “Path to stable quantum spin liquids in spin-orbit coupled correlated materials,” *Quantum Materials* **3**, 23 (2018).
- [96] Matthias Gohlke, Gideon Wachtel, Youhei Yamaji, Frank Pollmann, and Yong Baek Kim, “Quantum spin liquid signatures in kitaev-like frustrated magnets,” *Phys. Rev. B* **97**, 075126 (2018).
- [97] I. Rousochatzakis and N. B. Perkins, In preparation.
- [98] Mark Srednicki, “Chaos and quantum thermalization,” *Phys. Rev. E* **50**, 888–901 (1994).
- [99] J. M. Deutsch, “Quantum statistical mechanics in a closed system,” *Phys. Rev. A* **43**, 2046–2049 (1991).
- [100] Marcos Rigol, Vanja Dunjko, and Maxim Olshanii, “Thermalization and its mechanism for generic isolated quantum systems,” *Nature* **452**, 854 (2008).
- [101] Marcos Rigol and Mark Srednicki, “Alternatives to eigenstate thermalization,” *Phys. Rev. Lett.* **108**, 110601 (2012).
- [102] Luca D’Alessio, Yariv Kafri, Anatoli Polkovnikov, and Marcos Rigol, “From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics,” *Advances in Physics* **65**, 239–362 (2016), <https://doi.org/10.1080/00018732.2016.1198134>.
- [103] Roderich Moessner and Arthur P. Ramirez, “Geometrical Frustration,” *Physics Today* **59**, 24 (2006).
- [104] P. Lecheminant, B. Bernu, C. Lhuillier, L. Pierre, and P. Sindzingre, “Order versus disorder in the quantum Heisenberg antiferromagnet on the kagomé lattice using exact spectra analysis,” *Phys. Rev. B* **56**, 2521–2529 (1997).
- [105] C. Waldtmann, H.-U. Everts, B. Bernu, C. Lhuillier, P. Sindzingre, P. Lecheminant, and L. Pierre, “First excitations of the spin 1/2 Heisenberg antiferromagnet on the kagomé lattice,” *The European Physical Journal B - Condensed Matter and Complex Systems* **2**, 501–507 (1998).
- [106] Chen Zeng and Veit Elser, “Quantum dimer calculations on the spin-1/2 kagome Heisenberg antiferromagnet,” *Phys. Rev. B* **51**, 8318–8324 (1995).
- [107] M. Mambrini and F. Mila, “RVB description of the low-energy singlets of the spin 1/2 kagomé antiferromagnet,” *The European Physical Journal B - Condensed Matter and Complex Systems* **17**, 651–659 (2000).
- [108] G. Misguich, D. Serban, and V. Pasquier, “Quantum dimer model with extensive ground-state entropy on the kagome lattice,” *Phys. Rev. B* **67**, 214413 (2003).
- [109] Arnaud Ralko, Frédéric Mila, and Ioannis Rousochatzakis, “Microscopic theory of the nearest-neighbor valence bond sector of the spin- $\frac{1}{2}$ kagome antiferromagnet,” *Phys. Rev. B* **97**, 104401 (2018).
- [110] Steven T. Bramwell and Michel J. P. Gingras, “Spin Ice State in Frustrated Magnetic Pyrochlore Materials,” *Science* **294**, 1495–1501 (2001).
- [111] M J P Gingras and P A McClarty, “Quantum spin ice: a search for gapless quantum spin liquids in pyrochlore magnets,” *Reports on Progress in Physics* **77**, 056501 (2014).
- [112] Fennell, T., “Neutron scattering studies of spin ices and spin liquids,” *JDN* **13**, 04001 (2014).
- [113] Claudine Lacroix, Philippe Mendels, and Frédéric Mila, eds., *Introduction to Frustrated Magnetism*, Springer Series in Solid-State Sciences, Vol. 164 (Springer Berlin Heidelberg, 2011).
- [114] Finer differences arise due to the different point group symmetries of the clusters ‘24b’ and ‘24’ of Fig. 9.