

Running head: Caffeine and performance in ball games

Effects of caffeine supplementation on performance in ball games

Jingyi Shannon Chia

Physical Education & Sports Science, Nanyang Technological University, Singapore
Institute for Sports Research, Nanyang Technological University, Singapore
School of Sport, Exercise and Health Sciences, Loughborough University, United Kingdom

Laura Ann Barrett

School of Sport, Exercise and Health Sciences, Loughborough University, United Kingdom

Jia Yi Chow

Physical Education & Sports Science, Nanyang Technological University, Singapore

&

Stephen Francis Burns

Physical Education & Sports Science, Nanyang Technological University, Singapore

Correspondence should be addressed to: Stephen Francis Burns, Physical Education and Sports Science,
National Institute of Education, Nanyang Technological University, 1 Nanyang Walk, Singapore 637616.
Email: stephen.burns@nie.edu.sg

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Abstract

Although a large body of evidence exists documenting caffeine’s ergogenic properties, most studies have focused on endurance performance. Findings from endurance sports, however, cannot be generalized to performance in ball games where, apart from having a high level of endurance, successful athletic performances require a combination of physiological, technical and cognitive capabilities. The purpose of this review was to critically evaluate studies that have examined the effect of a single dose of caffeine in isolation on one or more of these performance measures: total distance, sprint performance, agility, vertical jump performance and accuracy in ball games. Searches on three major databases resulted in 19 studies (invasion games: 13, net-barrier games: 6) that evaluated the acute effects of caffeine on human participants, provided the caffeine dose administered and included a ball games specific task or simulated match. Improvements in sprint performance were observed in 8 out of 10 studies (80 %) and vertical jump in 7 out of 8 studies (88 %). Equivocal results were reported for distance covered, agility and accuracy. Minor side effects were reported in 4 of 19 studies reviewed. Pre-exercise caffeine ingestion between 3.0 and 6.0 mg per kg of body mass appears to be a safe ergogenic aid for athletes in ball games. However, the efficacy of caffeine varies depending on various factors, including but not limited to the nature of the game, physical status and caffeine habituation. More research is warranted to clarify the effects of caffeine on performance measures unique to ball games such as agility and accuracy. It is essential that athletes, coaches and practitioners evaluate the risk-benefit ratio of caffeine ingestion strategies on an individual case-by-case basis.

Key Points

- Caffeine may be ergogenic in ball games by sparing muscle glycogen, increasing force production, decreasing perceptions of pain and stimulating the central nervous system which delays fatigue and enhances performance.
- Acute ingestion of caffeine doses between 3.0 and 6.0 mg per kg of body mass improved sprint performance and vertical jump in both net-barrier and invasion games without any major side effects.
- Caffeine ingestion strategies should be tailored to the individual athlete to determine any ergogenic benefits and minimize potential side effects.

1 **1. Introduction**

2 Ball game players require a combination of aerobic and anaerobic endurance, speed, power, agility and strength
3 [1]. Depending on constraints such as the rules of the game, playing area, number of players, playing positions as
4 well as tactical strategies, the metabolic demands of each game are highly specific [2, 3]. However, successful
5 performance in ball games stresses not only the physiological systems but also imposes a high degree of technical
6 and cognitive demands [4-6]. In competitive settings, the margin between winning and losing can be small. As
7 such, the effects of ergogenic aids such as caffeine on sporting performance are of great interest to athletes,
8 coaches and practitioners.

9
10 Caffeine (1, 3, 7-trimethylxanthine) is a stimulant commonly consumed by athletes due to its reported ergogenic
11 effects and ease of availability either in supplement form or in the diet in tea leaves, coffee or chocolate [7].
12 Several reviews have documented the ergogenic properties of caffeine in endurance sports as well as in repeated
13 or sustained high-intensity events such as sprinting, swimming or rowing [8, 9]. In terms of endurance
14 performance, it is clear that caffeine has the ability to increase both time to exhaustion and performance in time
15 trials [10-13]. Although the benefits of caffeine ingestion in endurance activities are well established, its
16 effectiveness for short-term high-intensity exercise performance is equivocal. In a recent systematic review by
17 Astorino and Roberson [8], it was found that 6 of 11 studies revealed benefits of caffeine for resistance training
18 and 11 of 17 studies revealed improvements in team sports exercise and power-based sports after caffeine
19 supplementation. Here, the studies that found performance improvements comprised elite and trained athletes as
20 opposed to recreationally active and untrained individuals.

21
22 Caffeine is rapidly absorbed and metabolized in the liver to di-methylxanthines (paraxanthine, theobromine,
23 theophylline), each of which has its own effects on the body [14]. Paraxanthine has been shown to stimulate
24 lipolysis, theobromine to cause vasodilation and increase in urine volume and theophylline to cause smooth
25 muscle relaxation [8]. Due to its hydro- and lipophilic properties, caffeine can easily move across cellular
26 membranes including the blood-brain barrier; hence it has the potential to affect all systems in the body including
27 the central nervous system (CNS) [15]. Caffeine is typically administered orally either as a tablet or capsule [16].
28 More recently, alternative delivery methods via caffeine encapsulated chewing gum or mouth rinses have become
29 widely available with direct absorption of caffeine into the blood stream via the buccal mucosa bypassing hepatic
30 metabolism and providing an advantage in terms of absorption rate over traditional delivery methods [17]. The

1 faster delivery of caffeine may result in a more rapid onset of ergogenic effects as the reaction of a drug is limited
2 by the rate at which it reaches target tissues [18].

3

4 It was previously purported that caffeine's ergogenic effects were associated with glycogen sparing and the
5 increase in free-fatty acid oxidation in endurance events [19, 20]. However, the evidence for increased fat
6 metabolism has been questioned not least because this proposed mechanism would not explain the ergogenic
7 effect of caffeine observed in short, high-intensity exercise which involves oxygen-independent metabolic
8 pathways and is unlikely to be limited by muscle glycogen availability [21, 22]. Other proposed mechanisms
9 include an increased secretion of β -endorphins [23] or an increased mobilization of intracellular calcium [14].
10 However, it is now believed that most of caffeine's ergogenic properties stem from its effects on the central
11 nervous system through adenosine receptor antagonism [24-29]. Adenosine and caffeine have opposite effects to
12 each other in regulating cellular activities. Hence, the ingestion of caffeine inhibits the effects of adenosine on
13 neurotransmission, perceived exertion and arousal, resulting in enhanced performance.

14

15 While the exact mechanism remains unclear, it is evident that at least a portion of individuals involved in top level
16 competition, benefit from the improved endurance and short, high-intensity performance seen with caffeine
17 ingestion. However, it cannot be assumed that similar performance improvements with caffeine would be
18 observed in an applied setting of ball games where physiological stress interacts with a high degree of technical
19 and cognitive demands [4-6]. Therefore, the purpose of the present review was to analyze the effect of caffeine
20 on specific performance measures of ball games: total distance covered, sprint performance, agility, vertical jump
21 performance and accuracy.

22

23

2. Methodological Aspects

24 This review is reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-
25 analyses (PRISMA) guidelines. A standardized literature search strategy was performed using the following
26 keyword combinations: Caffeine AND (sports games OR field games OR sports performance OR simulated sports
27 OR badminton OR baseball OR basketball OR bowling OR floorball OR football OR golf OR handball OR hockey
28 OR netball OR rugby OR soccer OR squash, OR table tennis OR tennis OR volleyball) via the following 3
29 databases; PubMed, SPORTDiscus and Web of Science from January 1980 to November 2016. The search terms
30 were agreed upon by JSC and LAB. For the PubMed search, an example for the first search term is as follows:

1 ("caffeine"[MeSH Terms] OR "caffeine"[All Fields]) AND ("sports"[MeSH Terms] OR "sports"[All Fields])
2 AND "games"[All Fields]. All titles and abstracts from the search were downloaded to Endnote X7.7 (Clarivate
3 Analytics). Manual cross-referencing from the reference lists of the articles was performed to identify any
4 potential missing studies. Thereafter, duplicate articles were removed and titles and abstracts were screened for
5 relevance. The identification and subsequent assessment of articles was done by two independent reviewers (JSC
6 and LAB). The two authors met and discussed any discrepancies until a consensus was achieved.

7
8 The articles were included for review and assessment based on the following criteria: (1) studies which tested the
9 acute effect(s) of caffeine on sport-specific tasks or specifically designed sport scenarios or simulated matches
10 relating to ball games; (2) included a caffeine only condition; (3) examined caffeine ingestion in isolation (without
11 other compounds such as carbohydrate); (4) used a single administration of caffeine; (5) included human
12 participants only; (6) included information on caffeine delivery method and administration; (7) administered
13 caffeine when participants were not in a fatigued state; (8) the article was available in English and (9) were
14 available as full articles (not abstracts). Ball games were defined as any form of game which includes a ball or
15 similar striking object such as a shuttle in badminton.

16
17 The Grading of Recommendations Assessment, Development and Evaluation (GRADE) method was used to
18 assess the quality of evidence for the main physiological and performance outcomes [30]. The GRADE method
19 rates the quality of outcomes into 1 of 4 levels (very low, low, moderate, and high) by evaluating five domains:
20 risk of bias, (in-)consistency of results (heterogeneity), (in-)directness of evidence, (im-)precision of results, and
21 publication bias. Using the GRADE approach, the evidence from the articles were rated as high quality given that
22 all studies were randomized controlled trials. If appropriate, the quality of outcomes was then downgraded based
23 on deficiencies in the above domains, lowering the quality from high to moderate to low or very low, depending
24 on the severity of the deficiency. The GRADEpro GDT online software (Evidence Prime) was used [31].

25
26 In addition, quality assessment was completed independently for each individual study included using the
27 Physiotherapy Evidence Database (PEDro) scale on the final selection of articles by 2 authors. This scale was
28 selected because of its ability to objectively assess the internal validity of each study which is essential when
29 evaluating the treatment effects of caffeine on performance. The PEDro scale yields a total score of 10, with more
30 points corresponding to higher quality [32]. Studies with a PEDro score of less than 6 were deemed as low quality

1 and not included in the review. This approach has been applied previously in similar reviews on caffeine [8, 9].
2 For each study, the following information was extracted: study design, number of participants, sex, level of
3 playing experience, caffeine habits, caffeine administration (dosage, amount and form), and comparisons of
4 physiological and specific performance outcome measures between the caffeine and placebo conditions. Cohen's
5 *d* was used as an estimate of effect size for the pairwise comparisons described above; it was calculated by dividing
6 the difference between the mean values with the standard deviation of the placebo condition. An effect size of
7 0.20 was considered a small effect, 0.50 a moderate effect and 0.80 a large effect [33].

8

9

3. Findings

10 We found 2154 articles through database searches. Of the 1028 that remained after the removal of 1126 duplicates,
11 we excluded 949 articles which were not relevant (Figure 1). Based on the inclusion criteria, 19 met the full set
12 of criteria and were included for review. Results were presented for a total of 279 participants, 74 % of whom
13 were male. The number of participants in each study ranged from 5 [34] to 26 [35]. The ball games within the
14 included studies were separated into 2 categories; invasion ($n = 13$) and net-barrier ($n = 6$) games. All participants
15 were competing actively and the majority trained for ~ 2 hours per day for 4 to 5 times a week. Apart from 2
16 studies which did not provide information relating to the participants' caffeine consumption habits [36, 37],
17 participants were mostly classified as light (< 100 mg per day) caffeine users. Participants stated that they were
18 regular consumers of caffeine (no information regarding the amount was provided) in the study by Stuart and
19 colleagues [38]. Higher ceiling levels of caffeine consumption were reported for participants in the study by
20 Foskett and colleagues [39] at < 350 mg of caffeine per day and for participants in the study by Tucker and
21 colleagues [34] at < 500 mg of caffeine per day.

22

23 All studies were randomized and included a placebo condition for comparative purposes. Participants were
24 instructed to standardize their diet, fluid intake and activity levels prior to the experimental trials. The duration
25 and degree of the standardization protocols differed among studies. The duration whereby participants had to
26 abstain from all dietary sources of alcohol, caffeine and other stimulants ranged from the day itself [36, 40] to 24
27 hours [35-37, 41-45], 48 hours [34, 38, 39, 46-50] and the entire study duration (2 sessions separated by 1 week)
28 [51]. Participants were also asked to adopt a similar diet and fluid intake level the day before each experimental
29 trial. Several studies also included a pre-competition meal either 2 [46] or 3 hours before each trial to mimic their
30 habitual routines [35, 36, 40, 44, 48, 49, 51]. In terms of activity levels, participants were advised to avoid all

1 strenuous exercise the day before each trial to minimize any muscle soreness and fatigue. Only in 3 studies were
2 participants involved in a light and standardized training session the day before each trial [44, 49, 50]. No
3 information on pre-trial standardization and procedures was reported in 3 studies [34, 37, 45]. The overall rating
4 for quality of evidence determined using the GRADE approach for both physiological and performance outcomes
5 was low. The quality of evidence was downgraded primarily for the following reasons: firstly, the heterogenous
6 nature of the evidence in terms of differences in the population (e.g. expertise level, caffeine consumption habits,
7 nature of sport involved in) and measurement methods of the outcomes (e.g. sport-specific modifications of test
8 protocols), making it difficult to compare across the studies reviewed. For example, comparing the results for
9 sprint performance between rugby players and other ball games such as tennis, would not be the most reliable as
10 rugby players would be more accustomed to longer sprints as compared to shorter distances. Thus, ratings for
11 inconsistency and imprecision were higher which resulted in a downgrading in the quality of evidence. Secondly,
12 although blood glucose and blood lactate concentrations were mentioned in 2 and 3 studies respectively, mean
13 comparisons were only performed in 1 study [43] and hence the evidence for these 2 outcomes were considered
14 as “sparse” and subsequently labelled as “low quality”. However, the quality of individual studies reviewed from
15 the perspective of internal validity was high with PEDro quality scores being ≥ 8 . Only 1 study had a score of 8
16 as a single-blinded procedure was used [41]. The kappa score for inter-rater reliability of PEDro scores was 1,
17 indicating perfect agreement between the 2 reviewers.

18

19 **3.1 Caffeine administration**

20 Most studies utilized a protocol whereby caffeine was administered to the participants 60 minutes before exercise.
21 Exceptions were 4 studies where participants were given the caffeine dose 30 [41, 52], 50 minutes [43] and 70
22 minutes [38] prior to the trials. The timing of caffeine ingestion relates directly to the pharmacokinetics of caffeine.
23 Caffeine, when ingested, is rapidly and completely absorbed by the gastrointestinal tract, reaching peak plasma
24 concentrations within approximately 1 hour after ingestion (99 % of the ingested caffeine is absorbed within 45
25 minutes) [14, 53]. Moreover, this rate of absorption does not seem to be affected by the dose of caffeine consumed
26 at least at amounts of caffeine consumption ≤ 10 mg of caffeine per kg of body mass (BM) [54]. The absorption
27 rate may, however, be affected by the form of administration (e.g. capsule, gum, drink solution) although a quicker
28 absorption rate does not necessarily mean a greater response to or effect of caffeine [54]. Its half-life in the body
29 is approximately 4 to 6 hours, although the rate of caffeine clearance differs between individuals [14].

30

1 In all the studies reviewed, caffeine was administered based on the individual's body mass as opposed to using an
2 absolute dose. Using an absolute dose creates large variability in responses which may affect the results of the
3 study. Caffeine doses ranging from $3.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$ [34-36, 40, 41, 44, 46-51] up to $6.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$ were reviewed
4 [37-39, 42, 43, 45]. Within the reviewed studies, it seems that the ingestion of $3.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$ of caffeine was
5 sufficient to elicit improvements in at least 1 performance variable in all but 1 study by Tucker and colleagues
6 [34]. It is possible that the sample size in the study ($n = 5$) could have been too small to detect any significant
7 performance improvements. However, as no dose less than $3.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$ of caffeine was given in any study, it
8 is uncertain if a minimal threshold for performance improvement exists. Of the 12 studies which administered a
9 dose of $3.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$ of caffeine, most ($n = 10$) used a beverage form (9 = 250 ml; 1 = 650 ml). The remaining
10 2 used a gelatin capsule filled with caffeine.

11

12 One study administered a moderate dose of $4.4 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$ of caffeine [52] and performance improvements were
13 reported. However, at a higher dosage of $6.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$, results were equivocal. Six studies administered 6.0
14 $\text{mg}\cdot\text{kg}^{-1} \text{ BM}$, either as a 500 ml beverage ($n = 2$) or in capsule format ($n = 4$). Surprisingly, performance
15 improvements were not as evident with half of these studies reporting no change in performance when compared
16 with placebo. This suggests that a threshold exists for the amount of caffeine ingested, warranting the need to
17 investigate the dose-response effect of caffeine in ball games. At present, current data suggest that a dose of 3.0
18 $\text{mg}\cdot\text{kg}^{-1} \text{ BM}$ caffeine is sufficient to elicit a positive effect on performance markers in ball games.

19

20 **3.2 Physiological effects**

21 *3.2.1 Heart rate responses*

22 Considering the stimulatory effect of caffeine on the central nervous system, it is commonly assumed that caffeine
23 increases heart rate (HR) during exercise [55]. Heart rate responses were reported for 12 studies in this review.
24 There was little evidence suggesting that caffeine increased exercise HR. Mean HR remained unchanged in 10
25 studies (~ 83 %); 2 studies reported a significant increase in mean HR with caffeine during the simulated match
26 as compared to the placebo [36, 49]. For instance, Del Coso and colleagues [49] reported that mean HR during 3
27 simulated 15-minute rugby matches increased significantly with caffeine as compared to the placebo (168 ± 7
28 $\text{b}\cdot\text{min}^{-1}$ vs. $164 \pm 6 \text{ b}\cdot\text{min}^{-1}$). Changes in mean HR could be a result of many interacting factors, such as the physical
29 status of the individuals or the presence of any stressors. It is interesting to note here that the participants in these

1 2 studies were females [36, 49]. However, there appears to be no available evidence documenting sex-related
2 differences in cardiovascular responses to caffeine ingestion during exercise.

3

4 Mixed findings were observed for changes in peak or maximal HR following caffeine ingestion. Of the 8 studies
5 which reported peak or maximal HR, there were no significant changes in these values in 6 of the studies [35, 40,
6 46, 48, 49, 51]. Conversely, increases in peak HR were observed in 2 studies [36, 44]. Del Coso and colleagues
7 [44] reported an increase in peak HR from $181 \pm 9 \text{ b}\cdot\text{min}^{-1}$ to $185 \pm 11 \text{ b}\cdot\text{min}^{-1}$ during a field hockey simulated
8 match. Interestingly, this increase, was not maintained as the game progressed, with no differences in peak HR in
9 the second half of the match. The increase in HR observed was related closely to the stimulant effect of caffeine
10 in increasing high speed movements. Specifically, the increase in peak HR was accompanied by an increase in
11 distance covered at high speed ($15.0 - 18.9 \text{ km}\cdot\text{h}^{-1}$). The players, however, were unable to maintain an increased
12 speed in the second half of the match; thus, caffeine was not effective in reducing the magnitude of fatigue
13 experienced. There was no difference in peak HR between the caffeine and placebo conditions during the second
14 half of the match.

15

16 *3.2.2 Blood lactate and glucose concentrations*

17 Likewise, higher blood lactate and glucose concentrations were reported after the first half of a soccer match in
18 the caffeine trial as compared to the placebo (blood lactate: $7.9 \pm 4.8 \text{ mmol}\cdot\text{L}^{-1}$ vs. $6.1 \pm 3.6 \text{ mmol}\cdot\text{L}^{-1}$; blood
19 glucose: $7.5 \pm 2.1 \text{ mmol}\cdot\text{L}^{-1}$ vs. $6.3 \pm 1.6 \text{ mmol}\cdot\text{L}^{-1}$) [43]. Higher blood lactate concentrations can be a reflection
20 of a higher contribution from anaerobic glycolysis to energy provision in the first half of the match. Considering
21 the length of the match, the subsequent disappearance of this difference in blood lactate concentration between
22 the 2 trials in the second half of the match appears to be consistent with caffeine's alleged glycogen sparing
23 mechanism and a shift toward using FFAs as the main fuel source during the match. Consequently, this may
24 explain why blood lactate concentrations were similar to placebo in the second half of the match. However,
25 caffeine has also been shown to consistently increase blood lactate levels without increasing muscle lactate [56,
26 57]. This suggests that caffeine's effect is likely a result of decreased clearance of blood lactate rather than an
27 increase in glycolytic flux. Alternatively, as peak concentration of caffeine is achieved 1 hour post-ingestion, there
28 may have been a decreasing influence of caffeine over the course of the match which was most obvious in the
29 second half of the game and which led to a subsequent decrease in high-intensity work. In terms of blood glucose,
30 the higher concentrations could indicate a larger catecholamine response in the first half of the caffeine trial [43].

1 In summary, in a match setting, the physiological effects of caffeine appear to be more pronounced during the
2 first than second half particularly for blood lactate and glucose concentrations.

3

4 *3.2.3 Sweat rate and dehydration levels*

5 Sweat rate and dehydration levels were unaffected by caffeine [35, 44, 46, 48, 49, 51] except for 1 study by Gallo-
6 Salazar and colleagues [40] on elite junior tennis players. Sweat rate was 33.5 ± 10.6 % higher in the caffeine trial
7 (0.7 ± 0.3 L·h⁻¹ vs. 0.5 ± 0.3 L·h⁻¹), resulting in greater dehydration levels (0.2 ± 0.4 % vs. 0.1 ± 0.5 %). This level
8 of dehydration, however, is acceptable given that dehydration levels > 2 % of body mass are reported to result in
9 significant impairments in exercise and skill performance [58, 59]. The ingestion of caffeine does not seem to
10 affect sweat rate or hydration levels.

11

12 **3.3 Performance effects**

13 *3.3.1 Total distance covered*

14 Exercise capacity is expected to increase with caffeine possibly through its purported effect on glycogen sparing
15 but this could also be through other mechanisms. One way to quantify the effect of caffeine in resisting fatigue
16 and enhancing exercise capacity is by measuring the total distance covered by the individual during match play.
17 In recent years, the advancement in global positioning satellite (GPS) technology has made it easy and convenient
18 to quantify movement patterns without hindering game actions. No information regarding total distance covered
19 in net-barrier games has been reported. Of the 5 studies on invasion games which reported total distance covered
20 during a match, increases were observed in 3 studies, 2 in soccer [46, 48] and 1 in rugby [35]. In contrast, Del
21 Coso and colleagues [44] reported no change in total distance covered during a simulated 2 x 25-minute hockey
22 match. Despite this, there was an increase in overall distance covered during high-intensity running (358 ± 117 m
23 vs. 303 ± 67 m) and sprinting (117 ± 55 m vs. 85 ± 41 m); running actions at moderate intensity were found to be
24 reduced and transformed into higher speed movements. Similar changes in movement patterns were reported in
25 the studies where total distance covered with ingestion of caffeine increased. This increased distance at high speed
26 represents a meaningful advantage for athletes and coaches.

27

28 One exception is the study by Pettersen and colleagues [43] where match activity and fatigue resistance during a
29 90-minute soccer match were examined in 22 youth players. Neither total distance covered nor movement patterns
30 were altered with the ingestion of caffeine. The activity pattern in soccer can be influenced by several factors apart

1 from the ability to resist fatigue. For instance, it is possible that teams with a high technical ability might not have
2 to work as hard during a game [60]. As such, technical differences between the 2 teams involved in the study may
3 have meant that the physical capacities of the players were not stressed to a great extent and therefore the effect
4 of caffeine not as evident. The age of the participants might have also affected the results of this study. With the
5 exception of the study by Gallo-Salazar and colleagues [40], most other studies in this review tested adult athletes,
6 whereas participants in this study on soccer were youth athletes (17.6 ± 1.1 years). Little research has been done
7 on the effects of caffeine in younger athletes and the paucity of evidence makes it difficult to draw conclusions or
8 recommendations. Performance benefits observed in the adult population, however, should not be generalized to
9 youth athletes. As such, more studies are needed in youth athletes whilst carefully weighing up any relevant ethical
10 considerations surrounding the use of caffeine in this population.

11 12 *3.3.2 Sprint performance*

13 The effect of caffeine on sprint performance was clearly evidenced as there was an increase in the number of
14 sprint bouts performed by individuals in a match setting in hockey [44], rugby [35], soccer [46, 48] as well as in
15 tennis [40]. The definition of a sprint differed across studies. A velocity greater than $23 \text{ km}\cdot\text{h}^{-1}$ was considered a
16 sprint in hockey by Del Coso and colleagues [44] whilst, any movement with a velocity above $18 \text{ km}\cdot\text{h}^{-1}$ was
17 considered a sprint in soccer [46, 48]. Sprint distance was also found to increase during a hockey ($117 \pm 55 \text{ m}$ vs.
18 $85 \pm 41 \text{ m}$) [44] and rugby match ($208 \pm 38 \text{ m}$ vs. $184 \pm 38 \text{ m}$) [35], indicating enhanced sprint ability with
19 caffeine.

20
21 Sprint velocity and sprint time were commonly measured with individuals performing tests of repeated ($n = 6-8$)
22 short sprints of either 15 m or 40 m. In 3 out of 4 studies which measured average peak and maximal speed across
23 repeated sprints, increases were observed in the caffeine trials [40, 46, 48]. Only 1 study by Del Coso and
24 colleagues [49] reported no change in maximal speed during a 6 x 30 m sprint test in rugby players. Despite this,
25 sprint performance in terms of pace at sprint velocity ($> 20 \text{ km}\cdot\text{h}^{-1}$) during the match increased with caffeine (6.1
26 $\pm 3.4 \text{ m}\cdot\text{min}^{-1}$ vs. $4.6 \pm 3.3 \text{ m}\cdot\text{min}^{-1}$). Conversely, during actual match play, Lara and colleagues [48] reported that
27 the increase in average peak ($24.5 \pm 1.7 \text{ km}\cdot\text{h}^{-1}$ vs. $24.2 \pm 1.6 \text{ km}\cdot\text{h}^{-1}$) and maximal speed ($25.6 \pm 1.4 \text{ km}\cdot\text{h}^{-1}$ vs.
28 $25.0 \pm 1.4 \text{ km}\cdot\text{h}^{-1}$) which had been observed during the sprint test was not present. Maximal speed attained during
29 the 2 x 40-minute match was unaffected with caffeine. The reason for this contradictory finding is not apparent;
30 however, it highlights the importance of measuring sprint velocity in a match setting as opposed to simply

1 conducting sprint tests to evaluate the ergogenic properties of caffeine. This is crucial as players and coaches are
2 most interested in performance enhancements during a match.

3

4 The effect of caffeine on sprint time, however, was not clear. Only 3 studies reported sprint times; 1 on floorball
5 [52], rugby [38] and soccer [37]. Moreover, the findings were mixed. In the study by Stuart and colleagues [38],
6 20 m, 30 m, offensive, defensive and tackle sprint times were measured using the Rugby test which was designed
7 based on time-motion analysis of first-class level rugby matches to simulate match activity. The positive effects
8 of caffeine on sprint time were clear, with mean percentage improvements of 0.5 to 2.9 %. However, Krasnanova
9 and colleagues [52] only found faster timings with the longer 6 x 40 m sprint tests (52.1 ± 4.4 s vs. 54.0 ± 5.3 s)
10 in floorball players, whereas time to completion for the 6 x 9 m test was not improved with caffeine. In addition,
11 sprint time measured for both the dominant and non-dominant side of soccer players was not improved with
12 caffeine [37]. Here, participants performed 3 trials, beginning from the same starting position where they reacted
13 to the tester's movement (step forward with right foot and change direction to the left and vice versa) and sprinted
14 to either the left or right side. It is important to note that the sprint distance for this study was only 4.5 m. Moreover,
15 reactive agility rather than sprint performance was the primary component of interest. As such, care must be taken
16 when interpreting these results.

17

18 3.3.3 Agility

19 Movement patterns in ball games typically involve many directional changes, commonly termed *agility*. Two
20 studies on net-barrier games have investigated the effects of caffeine on pre-planned directional sprints assessed
21 by the agility T-test, with equivocal results [50, 51]. Given the relatively small playing area in net-barrier games
22 such as badminton and volleyball, quick changes in direction and subsequent acceleration are critical for
23 successful performances. The agility T-test requires individuals to move through a T-shaped circuit as fast as
24 possible to simulate the fast directional changes and short sprints in a game. A reduction in total time taken (10.3
25 ± 0.4 s vs. 10.8 ± 0.4 s) to complete the test was reported by Del Coso and colleagues when the participants
26 ingested a $3.0 \text{ mg}\cdot\text{kg}^{-1}$ BM caffeine beverage pre-exercise as compared to the placebo [51]. However, with the
27 same dosage of caffeine administered, Abian and colleagues [50] found no statistically significant improvement
28 in time taken to complete the test, despite a 0.4 ± 1.4 % improvement in agility with caffeine. A limitation of these
29 studies was that neither tested the effect of caffeine on agility in a game setting - that is in combination with a
30 perceptual component which requires the participant to initiate the movement response as opposed to such pre-

1 planned tests. Thus, at present the limited available evidence prevents any firm conclusion being drawn on the
2 ability of caffeine to improve the agility of individuals in a game setting.

3

4 *3.3.4 Vertical jump performance*

5 Vertical jump is important for successful athletic performance in ball games [61, 62]. Countermovement jumps
6 (CMJ) and the fifteen seconds maximal jump test (15 RJ) are common tests used as measures of lower body power
7 in athletes [62]. The 15 RJ requires the individual to jump vertically repeatedly over 15 seconds, landing and
8 jumping with both feet. Countermovement jump height was higher after caffeine ingestion as compared to the
9 placebo in all 6 studies which measured CMJ height. This included both invasion and net-barrier games, namely
10 badminton [50], basketball [47], soccer [39, 48] and volleyball [36, 51]. The greatest reported increase in jump
11 height with caffeine was approximately 5.0 % observed in badminton (37.7 ± 4.5 cm to 39.5 ± 5.1 cm) [50] and
12 volleyball (35.9 ± 4.4 cm to 37.7 ± 4.6 cm) [51] players. Likewise, the ingestion of caffeine significantly increased
13 jump height measured during the 15 RJ test performed by basketball (30.2 ± 3.6 cm vs. 28.8 ± 3.4 cm) [47], soccer
14 (35.8 ± 5.5 cm vs. 34.7 ± 4.7 cm) [46] and volleyball (30.7 ± 4.5 cm vs. 29.3 ± 4.8 cm) [51] players.

15

16 Three studies also measured game-specific variations of vertical jumps such as smash, spike, block or rebound
17 jump heights [34, 36, 50]. Specific to volleyball, Perez-Lopez and colleagues [36] reported significant increases
18 in spike (43.3 ± 4.7 cm vs. 44.4 ± 5.0 cm) and block jump heights (35.2 ± 5.1 cm vs 36.1 ± 5.1 cm) with pre-
19 exercise ingestion of caffeine. However, the study by Tucker and colleagues [34] failed to find any difference in
20 rebound jump height in basketball players in the caffeine trial as compared to the placebo trial. The authors
21 calculated a reactive strength index (RSI) based on the jump height and time spent on the mat before rebounding
22 as quickly as possible upwards. The failure to observe any performance benefit may be attributed to the small
23 sample size in this study ($n = 5$). Closer examination of the individual performances revealed that RSI was higher
24 in the caffeine trial for 3 out of the 5 participants. Moreover, this improvement was approximately 9 % for 1
25 individual participant (119 ± 9 cm·s⁻¹ vs. 109 ± 6 cm·s⁻¹). As such, it appears that pre-exercise ingestion of caffeine
26 has some ergogenic effect on jump height.

27

28 *3.3.5 Accuracy*

29 Accuracy can be a key factor in successful sporting performances. Though not studied extensively, caffeine as a
30 stimulant purported to increase mental alertness and cognitive functioning may possibly enhance task accuracy

1 [24]. On the other hand, caffeine consumption might be associated with several side effects such as tremors and
2 nervousness which could negatively affect accuracy [63]. Depending on the nature of the game, the effect of
3 caffeine on accuracy can be measured in terms of (a) passing, (b) shooting and (c) serving accuracy. Only 3 studies
4 assessed passing accuracy using sport-specific passing tests [38, 39, 42]. The Loughborough Soccer Passing Test
5 (LSPT), for instance, requires individuals to complete a circuit of 16 passes from a dynamic starting position,
6 against coloured targets as quickly as possible while minimizing error. The next desired target is denoted by an
7 audible signal immediately after each pass. Apart from a high level of accuracy, enhanced proprioceptive and
8 cognitive functioning is essential in this test. A trend for improved passing accuracy was observed in the caffeine
9 trial ($p = 0.06$). Likewise, in rugby players, Stuart and colleagues [38] reported a 10 % improvement in passing
10 accuracy in players who were pressured to pass rapidly in a rugby test simulating match conditions. Participants
11 in these 2 studies were given a caffeine dose of $6.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$, suggesting that caffeine does not lead to over
12 arousal and impaired accuracy at this dosage, at least in the participants in these 2 studies [38, 39]. Surprisingly,
13 despite the same dosage of caffeine ($6.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$) being administered, Assi and Bottoms [42] found no effect
14 of caffeine on passing accuracy in a rugby passing test. Each participant had three attempts with each hand; a
15 successful hit earned them 2 points, making 6 the maximum score attainable. Although accuracy scores improved,
16 no statistically significant differences were found. Given the nature of the test and scoring system, it is possible
17 that the scoring system was not sensitive enough to reflect these improved scores or that the test failed to impart
18 sufficient pressure upon the participants for the effects of caffeine to become evident.

19

20 In terms of shooting accuracy, Abian and colleagues [47] reported that caffeine ingestion had no effect on accuracy
21 in both free throw and 3-point shots in basketball. Likewise, accuracy in floorball shooting was also unaffected
22 by caffeine [52]. Participants performed repeated 8 m shots with the aim of scoring as many goals from 10 attempts
23 as possible. The failure to find any improvement in accuracy could be related to the task design. In this case,
24 performing repeated shots is not realistic and representative of an actual shot taken in a match situation. Given the
25 limited evidence on shooting accuracy in ball games, it is difficult to draw a conclusion on the efficacy of caffeine,
26 warranting the need for more studies.

27

28 Lastly, in net-barrier games such as tennis, the serve represents one of the most important shots and has a
29 significant impact on the match result [64]. Findings from the 2 studies reviewed on tennis were mixed; first,
30 Hornery and colleagues [41] reported that a pre-exercise caffeine ingestion of $3.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$ had no effect on

1 stroke accuracy. Yet, when a higher dose of caffeine of $6.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$ was administered by Klein and colleagues
2 [45], stroke accuracy as assessed by a tennis skill test (TST) improved. The number of successful shots out of 324
3 during the TST increased in the caffeine trial (295 ± 11 shots) as compared to the placebo (289 ± 10 shots) trial.
4 This was the only study reviewed which examined sex-related differences in the effects of caffeine and the
5 investigators reported a similar effect in both male and female participants. Both studies, however, only
6 administered a single dose of caffeine; hence it is not possible to determine if the increased accuracy is attributable
7 to the higher dose of caffeine.

8

9 *3.3.6 Summary*

10 The weight of evidence supports the ingestion of an acute dose of caffeine ($3.0 - 6.0 \text{ mg}\cdot\text{kg}^{-1} \text{ BM}$) approximately
11 60 minutes before ball games (Table 1). In all 19 studies reviewed, no negative effects on performance were
12 reported. Three out of 5 studies showed improvement in total distance covered (rugby: 1 [35]; soccer: 2 [48, 46]).
13 Four aspects of sprint performance were evaluated in this review; sprint count increased in all 5 studies (hockey:
14 1 [44]; rugby: 1 [35]; soccer: 2 [46, 48]; tennis: 1 [40]), sprint time was faster in 2 out of 3 studies (floorball: 1
15 [52]; rugby: 1 [38]), sprint velocity was higher in 3 out of 4 studies (soccer: 2 [48, 46]; tennis: 1 [40]) and sprint
16 distance increased in 2 out of 3 studies (hockey: 1 [44]; rugby: 1 [35]). Two studies examined the effects of
17 caffeine on agility with only 1 study on volleyball reporting a faster time to completion in the agility test [51].
18 With respect to vertical jump performance, all 6 studies (badminton: 1 [50]; basketball: 1 [47]; soccer: 2 [48, 39];
19 volleyball: 2 [36, 51]) that measured CMJ heights reported increases in the caffeine condition and 3 out of 3
20 studies (basketball: 1 [47]; soccer: 1 [46]; volleyball: 1 [51]) reported higher jump heights in the 15 RJ test. Lastly,
21 in terms of accuracy; 2 out of 3 studies (rugby: 2 [42, 38]) observed higher passing accuracy. However, none of
22 the 2 studies which measured shooting accuracy reported any positive effects and 1 out of 2 studies reported
23 increased serve accuracy in tennis [45].

24

25 **3.4 Possible mechanisms of caffeine action**

26 While the exact mechanism for caffeine's ergogenic effect remains unclear, several possible reasons are examined
27 in this section [14] and are related to the potential performance outcomes in ball games highlighted in Section 3.3.
28 Caffeine has been shown to increase FFA mobilization which could result in muscle glycogen sparing [20, 65].
29 Sparing of glycogen may enhance exercise capacity as glycogen depletion has been shown to result in fatigue.
30 This increase in reliance on FFA mobilization could greatly impact both invasion and net-barrier games which

1 involve intense and long running distances such as basketball, hockey, rugby, soccer and tennis. However,
2 glycogen sparing has not been found in all studies and this mechanism has been questioned [23, 56] as it does not
3 explain the ergogenic effect of caffeine observed in performances such as sprinting or vertical jumps which
4 involve oxygen-independent metabolic pathways [21, 22]. Caffeine has also been found to increase the secretion
5 of β -endorphins [23]. The elevated endorphin concentrations in the blood have analgesic properties which
6 decrease the perception of pain, allowing for greater work to be performed at the same level of muscle pain [66].
7 This delays the fatigue response of individuals and increases their exercise capacity, allowing them to exercise
8 longer, especially in games with few and/or short rest periods and minimal substitutions such as soccer. In
9 addition, the reduction in the perception of pain may be of particular importance in contact games such as rugby
10 given the high number of impacts associated with the game. There is also evidence that caffeine increases motor
11 unit recruitment [67] and muscle activation [68] during a forceful contraction; this direct intracellular action may
12 also be ergogenic. The increased formation of muscle cross bridges results in greater force production and hence,
13 increased strength and power can be generated. The rate of calcium release from the sarcoplasmic reticulum
14 following caffeine ingestion also increases the force of muscular contraction [69]. This is not only advantageous
15 in ball games with a high level of contact (e.g. rugby), but also in stop-and-go games (e.g. basketball and
16 badminton) which require multiple accelerations and short sprints and could result in improved jump performance
17 and agility. Moreover, increases in the force of muscular contraction could lead to a more powerful stroke and
18 shot in net-barrier games (e.g. tennis and volleyball) which is critical for successful performance. Finally, caffeine
19 acts as an adenosine antagonist and inhibits its original function which facilitates sleep [15]. This increased
20 wakefulness [70] and heightened attention levels stimulates the central nervous system, delays fatigue and has the
21 potential to enhance performance. This could lead to improvements in total distance run, agility and to
22 improvements in cognitive processing which is important for motor skill execution such as passing rugby balls,
23 hitting badminton shuttles or shooting accuracy in soccer.

24

25 **3.5 Potential side effects**

26 Seven studies reported in the present review utilized a 7 item (yes/no) questionnaire (items: headache, abdominal
27 or gut discomfort, muscle soreness, increased vigour or activeness, tachycardia and heart palpitations, insomnia
28 and increased anxiety) to assess the side effect(s) of caffeine. Four studies also included an added item - increased
29 urine production - in their survey. The most common side effect with the ingestion of caffeine compared with the
30 placebo was insomnia, with significantly higher prevalence rates reported in 2 studies; [31.2 % (5 out of 16) vs.

1 6.3 %] [50] and [27.0 % (4 out of 15) vs. 7.0 %] [51]. Prevalence of insomnia with caffeine also approached
2 statistical significance in 3 other studies [44, 48, 49]. However, even within these studies, there was substantial
3 variation in individual responses in that some participants did not experience insomnia. Theoretically, increased
4 insomnia may affect performance during training or competition the next day and hence, depending on the
5 individual's competition schedule, the use of caffeine should be carefully evaluated.

6
7 Apart from increased insomnia, the other side effect commonly reported the morning after the test session was
8 increased vigour and activeness. In 2 of these studies, there was a significant increase in the prevalence of
9 increased vigour and activeness with caffeine as reported by [37.5 % (6 out of 16) vs. 0 %] [47] and [15.0 % (2
10 out of 13) vs. 0 %] [36] of the participants as compared to the placebo. There was also a tendency for increased
11 vigour and activeness in the study by Del Coso and colleagues [49], with a prevalence rate of 50 % (8 out of 16).
12 Furthermore, increased vigour and activeness was also accompanied by increased nervousness in 15 % (2 of 13)
13 of participants in the study by Perez-Lopez and colleagues [36]. As the participants were females, it is important
14 to note that caffeine metabolism may be related to hormone concentrations. The systemic clearance of caffeine
15 has been found to be slower in the luteal phase of the menstrual cycle [71] and women taking oral contraceptives
16 have also been found to have decreased clearance of caffeine [72] which could potentially increase the likelihood
17 of experiencing side effects. However, another possible reason for the increased nervousness and activeness could
18 be associated with the amount of caffeine these individuals were accustomed to consuming. Unlike the other two
19 studies where participants were light users of caffeine (< 60 mg/day), no information regarding the caffeine
20 consumption habits of the participants was reported in the study. Hence, it is possible that the individuals were
21 naïve caffeine users and this subsequently resulted in greater stimulatory effects of caffeine.

22
23 It appears that caffeine is generally well tolerated but some individuals may be more susceptible to experiencing
24 adverse side effects from caffeine consumption and these adverse effects are important to understand as they could
25 have a negative effect on performance [73]. Genetic differences have been shown to determine the rate of caffeine
26 metabolism. The CYP1A2 gene is responsible for the building of the cytochrome P450 enzyme which controls
27 the metabolism of caffeine. There are two forms of the CYP1A2 gene and individuals who express the
28 CYP1A2*1A form are rapid caffeine metabolisers and individuals who carry the CYP1A2*1F form are slow
29 caffeine metabolisers which may leave them more susceptible to adverse side effects [74]. Interestingly a recent
30 paper found CYP1A2 genotype variations did not affect the ergogenic effects or the prevalence of side effects

1 after the ingestion of a moderate dose of caffeine although the sample size was small which may have influenced
2 the results [75].

3

4 Overall, these data indicate that the acute ingestion of caffeine results in only minor side effects and hence is safe
5 for athletes to consume. More importantly, although side effects were experienced, the ingestion of caffeine
6 resulted in performance improvements in these studies. Individuals are thus strongly advised to practice their
7 intake of caffeine before their competition periods to identify any side effects.

8

9 **3.6 Recommendations for future work**

10 The present review has highlighted potential issues surrounding the use of caffeine in ball games which should be
11 addressed by future research. Firstly, given the ergogenic properties of caffeine, it is likely that caffeine may play
12 an important role in performance during periods of increased physical and cognitive demand. In a sport setting,
13 this is particularly relevant when one is fatigued (i.e., second half of matches) [76]. However, there is limited
14 research investigating the ability of caffeine in attenuating performance impairment(s) with fatigue in ball games
15 [77] even though athletes can consume caffeine during rest periods. A novel study by Dunvnjak-Zaknich and
16 colleagues [6] examined the effect of caffeine on reactive agility and decision making accuracy when fresh and
17 fatigued. The various performance tests were interspersed between each exercise quarter of an 80-minute (4 x 20-
18 minute) simulated game. Unlike typical agility studies which involved pre-planned sprints, reactive agility is a
19 unique measure requiring participants to respond to an external stimulus and then initiate their movement,
20 simulating actual game settings. Findings from this study showed that a 6.0 mg·kg⁻¹ BM dose of caffeine resulted
21 in improvements in both reactive agility and decision making accuracy. These effects, however, were similar in
22 both fresh and fatigued conditions. It is important to note that this study did not simulate any particular sport,
23 highlighting the need for further sport-specific research in this area in order for athletes and coaches to make
24 informed decisions. Another area of interest to examine is the difference between divided or repeated doses of
25 caffeine versus a single dose given that any benefit of caffeine appears to be more pronounced during the first half
26 of matches. Currently, multiple doses ingested over time have only been investigated with concurrent
27 carbohydrate ingestion [78]. Lastly, the recent emergence of caffeine mouth-rinses and gums has opened the
28 possibility of faster performance effects through relatively small amounts of caffeine.

29

30

4. Conclusion

1 This review is the first to focus on the effects of caffeine on ball games specific performance measures (e.g. sprint
2 performance, vertical jump performance, shooting/passing accuracy) unlike previous reviews which examined
3 solely endurance or sprint performance. The physiological effects of caffeine were more evident in the first half
4 of matches, with elevated heart rate responses, blood lactate and glucose concentrations compared to the second
5 half. Contrary to popular belief that caffeine is diuretic, sweat rate and hydration levels were unaffected by caffeine
6 ingestion. The current body of research suggests that pre-exercise caffeine ingestion of doses ranging from 3.0 to
7 6.0 mg·kg⁻¹ BM, which translates to approximately 1.5 to 3.0 and 2.0 to 4.0 8 -oz (~ 227 g) cups (50 kg female
8 and 70 kg male, respectively) of regular drip brewed coffee (~ 100 mg of caffeine) [15], can result in increases in
9 vertical jump and sprint performance during a simulated match. However, equivocal results were found for total
10 distance covered, agility as well as accuracy, warranting the need for more research in this area. Future studies
11 should test the effect of caffeine using representative task designs for the most accurate evaluation. The influence
12 of training status and level of caffeine habituation should also be considered. Across the studies reviewed, only
13 minor side effects with caffeine ingestion were reported. Considering individual differences in caffeine sensitivity,
14 athletes are strongly advised to try caffeine while training before using it during their competition period to ensure
15 tolerance and to identify any potential side effects.

16

17 **Compliance with Ethical Standards**

18 **Funding**

19 Jingyi Shannon Chia is supported by the NIE Research Scholarship and by the Institute for Sports Research at
20 Nanyang Technological University. No other sources of funding were used to assist in the preparation of this
21 article.

22 **Conflicts of Interest**

23 Jingyi Shannon Chia, Laura Barrett, Jia Yi Chow and Stephen Burns declare that they have no conflicts of interest
24 relevant to the content of this review.

References

- 1 1. Bishop D. Dietary supplements and team-sport performance. *Sports Med.* 2010;40(12):995-1017.
- 2 2. Williams C, Rollo I. Carbohydrate nutrition and team sport performance. *Sports Med.* 2015;45 Suppl 1:S13-
- 3 22. doi:10.1007/s40279-015-0399-3.
- 4 3. Holway FE, Spriet LL. Sport-specific nutrition: practical strategies for team sports. *J Sports Sci.* 2011;29(Suppl
- 5 1):S115-S25. doi:10.1080/02640414.2011.605459.
- 6 4. Williams AM, Ford PR. Expertise and expert performance in sport. *Int Rev Sport Exerc Psychol.* 2008;1(1):4-
- 7 18. doi:10.1080/17509840701836867.
- 8 5. Damian F, David P, Tim G. Skill and physiological demands of open and closed training drills in Australian
- 9 football. *Int J Sports Sci Coach.* 2008;3(4):489-99. doi:doi:10.1260/174795408787186512.
- 10 6. Duvnjak-Zaknich DM, Dawson BT, Wallman KE, et al. Effect of caffeine on reactive agility time when fresh
- 11 and fatigued. *Med Sci Sports Exerc.* 2011;43(8):1523-30. doi:10.1249/MSS.0b013e31821048ab.
- 12 7. Del Coso J, Munoz G, Munoz-Guerra J. Prevalence of caffeine use in elite athletes following its removal from
- 13 the World Anti-Doping Agency list of banned substances. *Appl Physiol Nutr Metab.* 2011;36(4):555-61.
- 14 doi:10.1139/h11-052.
- 15 8. Astorino TA, Roberson DW. Efficacy of acute caffeine ingestion for short-term high-intensity exercise
- 16 performance: a systematic review. *J Strength Cond Res.* 2010;24(1):257-65. doi:10.1519/JSC.0b013e3181c1f88a.
- 17 9. Ganio MS, Klau JF, Casa DJ, et al. Effect of caffeine on sport-specific endurance performance: a systematic
- 18 review. *J Strength Cond Res.* 2009;23(1):315-24. doi:10.1519/JSC.0b013e31818b979a.
- 19 10. Bell DG, McLellan TM. Effect of repeated caffeine ingestion on repeated exhaustive exercise endurance. *Med*
- 20 *Sci Sports Exerc.* 2003;35(8):1348-54. doi:10.1249/01.mss.0000079071.92647.f2.
- 21 11. Pasman WJ, van Baak MA, Jeukendrup AE, et al. The effect of different dosages of caffeine on endurance
- 22 performance time. *Int J Sports Med.* 1995;16(4):225-30. doi:10.1055/s-2007-972996.
- 23 12. Graham TE, Spriet LL. Metabolic, catecholamine, and exercise performance responses to various doses of
- 24 caffeine. *J Appl Physiol* 1995;78(3):867-74.
- 25 13. Bridge CA, Jones MA. The effect of caffeine ingestion on 8 km run performance in a field setting. *J Sports*
- 26 *Sci.* 2006;24(4):433-9. doi:10.1080/02640410500231496.
- 27 14. Graham TE. Caffeine and exercise: metabolism, endurance and performance. *Sports Med.* 2001;31(11):785-
- 28 807.

- 1 15. Goldstein ER, Ziegenfuss T, Kalman D, et al. International Society of Sports Nutrition position stand: caffeine
2 and performance. *J Int Soc Sports Nutr.* 2010;7(1):5. doi:10.1186/1550-2783-7-5.
- 3 16. Doherty M, Smith PM. Effects of caffeine ingestion on exercise testing: a meta-analysis. *Int J Sport Nutr*
4 *Exerc Metab.* 2004;14(6):626-46.
- 5 17. Leon Shargel, Yu ABC. *Applied biopharmaceutics and pharmacokinetics.* Appleton & Lange; 1999.
- 6 18. Kamimori GH, Karyekar CS, Otterstetter R, et al. The rate of absorption and relative bioavailability of caffeine
7 administered in chewing gum versus capsules to normal healthy volunteers. *Int J Pharm.* 2002;234(1-2):159-67.
- 8 19. Costill DL, Dalsky GP, Fink WJ. Effects of caffeine ingestion on metabolism and exercise performance. *Med*
9 *Sci Sports.* 1978;10(3):155-8.
- 10 20. Spriet LL, MacLean DA, Dyck DJ, et al. Caffeine ingestion and muscle metabolism during prolonged exercise
11 in humans. *Am J Physiol.* 1992;262(6 Pt 1):E891-8.
- 12 21. Bruce CR, Anderson ME, Fraser SF, et al. Enhancement of 2000-m rowing performance after caffeine
13 ingestion. *Med Sci Sports Exerc.* 2000;32(11):1958-63.
- 14 22. Collomp K, Ahmaidi S, Chatard JC, et al. Benefits of caffeine ingestion on sprint performance in trained and
15 untrained swimmers. *Eur J Appl Physiol Occup Physiol.* 1992;64(4):377-80.
- 16 23. Laurent D, Schneider KE, Prusaczyk WK, et al. Effects of caffeine on muscle glycogen utilization and the
17 neuroendocrine axis during exercise. *J Clin Endocrinol Metab.* 2000;85(6):2170-5. doi:10.1210/jcem.85.6.6655.
- 18 24. Sokmen B, Armstrong LE, Kraemer WJ, et al. Caffeine use in sports: considerations for the athlete. *J Strength*
19 *Cond Res.* 2008;22(3):978-86. doi:10.1519/JSC.0b013e3181660cec.
- 20 25. Spriet LL, Gibala MJ. Nutritional strategies to influence adaptations to training. *J Sports Sci.* 2004;22(1):127-
21 41. doi:10.1080/0264041031000140608.
- 22 26. Davis JM, Zhao Z, Stock HS, et al. Central nervous system effects of caffeine and adenosine on fatigue. *Am*
23 *J Physiol Regul Integr Comp Physiol.* 2003;284(2):R399-404. doi:10.1152/ajpregu.00386.2002.
- 24 27. Boulenger JP, Marangos PJ. Caffeine withdrawal affects central adenosine receptors but not benzodiazepine
25 receptors. *J Neural Transm Gen Sect.* 1989;78(1):9-15.
- 26 28. Fredholm BB, Battig K, Holmen J, et al. Actions of caffeine in the brain with special reference to factors that
27 contribute to its widespread use. *Pharmacol Rev.* 1999;51(1):83-133.
- 28 29. Davis JK, Green JM. Caffeine and anaerobic performance: ergogenic value and mechanisms of action. *Sports*
29 *Med.* 2009;39(10):813-32. doi:10.2165/11317770-000000000-00000.

- 1 30. Balshem H, Helfand M, Schunemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin*
2 *Epidemiol.* 2011;64(4):401-6. doi:10.1016/j.jclinepi.2010.07.015.
- 3 31. Prime E. GRADEpro GDT. McMaster University. 2015. grade.pro.org. Accessed 9th June 2017.
- 4 32. PEDro scale. <https://www.pedro.org.au/>. Accessed April 3, 2017.
- 5 33. Cohen J. *Statistical power analysis for the behavioural sciences*. 2nd ed. Hillsdale (NJ): Lawrence Erlbaum
6 Associates; 1988.
- 7 34. Tucker MA, Hargreaves JM, Clarke JC, et al. The effect of caffeine on maximal oxygen uptake and vertical
8 jump performance in male basketball players. *J Strength Cond Res.* 2013;27(2):382-7.
9 doi:10.1519/JSC.0b013e31825922aa.
- 10 35. Del Coso J, Ramirez JA, Munoz G, et al. Caffeine-containing energy drink improves physical performance of
11 elite rugby players during a simulated match. *Appl Physiol Nutr Metab.* 2013;38(4):368-74. doi:10.1139/apnm-
12 2012-0339.
- 13 36. Perez-Lopez A, Salinero JJ, Abian-Vicen J, et al. Caffeinated energy drinks improve volleyball performance
14 in elite female players. *Med Sci Sports Exerc.* 2015;47(4):850-6. doi:10.1249/mss.0000000000000455.
- 15 37. Jordan JB, Korgaokar A, Farley RS, et al. Caffeine supplementation and reactive agility in elite youth soccer
16 players. *Pediatr Exerc Sci.* 2014;26(2):168-76. doi:10.1123/pes.2013-0134.
- 17 38. Stuart GR, Hopkins WG, Cook C, et al. Multiple effects of caffeine on simulated high-intensity team-sport
18 performance. *Med Sci Sports Exerc.* 2005;37(11):1998-2005.
- 19 39. Foskett A, Ali A, Gant N. Caffeine enhances cognitive function and skill performance during simulated soccer
20 activity. *Int J Sport Nutr Exerc Metab.* 2009;19(4):410-23.
- 21 40. Gallo-Salazar C, Areces F, Abian-Vicen J, et al. Enhancing physical performance in elite junior tennis players
22 with a caffeinated energy drink. *Int J Sports Physiol Perform.* 2015;10(3):305-10. doi:10.1123/ijsp.2014-0103.
- 23 41. Hornery DJ, Farrow D, Mujika I, et al. Caffeine, carbohydrate, and cooling use during prolonged simulated
24 tennis. *Int J Sports Physiol Perform.* 2007;2(4):423-38.
- 25 42. Assi HN, Bottoms L. The effects of caffeine on rugby passing accuracy while performing the Reactive Agility
26 Test. *Science & Sports.* 2014;29(5):275-81. doi:<http://dx.doi.org/10.1016/j.scispo.2014.07.012>.
- 27 43. Pettersen SA, Krstrup P, Bendiksen M, et al. Caffeine supplementation does not affect match activities and
28 fatigue resistance during match play in young football players. *J Sports Sci.* 2014;32(20):1958-65.
29 doi:10.1080/02640414.2014.965189.

- 1 44. Del Coso J, Portillo J, Salinero JJ, et al. Caffeinated energy drinks improve high-speed running in elite field
2 hockey players. *Int J Sport Nutr Exerc Metab.* 2016;26(1):26-32. doi:10.1123/ijsnem.2015-0128.
- 3 45. Klein C, Clawson A, Martin M, et al. The effect of caffeine on performance in collegiate tennis players. *J*
4 *Caffeine Res.* 2012;2(3):111-6.
- 5 46. Del Coso J, Munoz-Fernandez VE, Munoz G, et al. Effects of a caffeine-containing energy drink on simulated
6 soccer performance. *PLoS One.* 2012;7(2):e31380. doi:10.1371/journal.pone.0031380.
- 7 47. Abian-Vicen J, Puente C, Salinero JJ, et al. A caffeinated energy drink improves jump performance in
8 adolescent basketball players. *Amino Acids.* 2014;46(5):1333-41. doi:10.1007/s00726-014-1702-6.
- 9 48. Lara B, Gonzalez-Millan C, Salinero JJ, et al. Caffeine-containing energy drink improves physical
10 performance in female soccer players. *Amino Acids.* 2014;46(5):1385-92. doi:10.1007/s00726-014-1709-z.
- 11 49. Del Coso J, Portillo J, Munoz G, et al. Caffeine-containing energy drink improves sprint performance during
12 an international rugby sevens competition. *Amino Acids.* 2013;44(6):1511-9. doi:10.1007/s00726-013-1473-5.
- 13 50. Abian P, Del Coso J, Salinero JJ, et al. The ingestion of a caffeinated energy drink improves jump performance
14 and activity patterns in elite badminton players. *J Sports Sci.* 2015;33(10):1042-50.
15 doi:10.1080/02640414.2014.981849.
- 16 51. Del Coso J, Perez-Lopez A, Abian-Vicen J, et al. Enhancing physical performance in male volleyball players
17 with a caffeine-containing energy drink. *Int J Sports Physiol Perform.* 2014;9(6):1013-8. doi:10.1123/ijsp.2013-
18 0448.
- 19 52. Krasňanová I, Sedliak M, Lednický A, editors. Effect of caffeine on motoric performance of female floorball
20 players. *Acta Facultatis Educationis Physicae Universitatis Comenianae;* 2014.
- 21 53. Magkos F, Kavouras SA. Caffeine use in sports, pharmacokinetics in man, and cellular mechanisms of action.
22 *Crit Rev Food Sci Nutr.* 2005;45(7-8):535-62. doi:10.1080/1040-830491379245.
- 23 54. Bonati M, Latini R, Galletti F, et al. Caffeine disposition after oral doses. *Clin Pharmacol Ther.* 1982;32(1):98-
24 106.
- 25 55. Sinclair CJ, Geiger JD. Caffeine use in sports. A pharmacological review. *J Sports Med Phys Fitness.*
26 2000;40(1):71-9.
- 27 56. Graham TE, Helge JW, MacLean DA, et al. Caffeine ingestion does not alter carbohydrate or fat metabolism
28 in human skeletal muscle during exercise. *J Physiol.* 2000;529 Pt 3:837-47.
- 29 57. Chesley A, Howlett RA, Heigenhauser GJ, et al. Regulation of muscle glycogenolytic flux during intense
30 aerobic exercise after caffeine ingestion. *Am J Physiol.* 1998;275(2 Pt 2):R596-603.

- 1 58. Wall BA, Watson G, Peiffer JJ, et al. Current hydration guidelines are erroneous: dehydration does not impair
2 exercise performance in the heat. *Br J Sports Med.* 2015;49(16):1077-83. doi:10.1136/bjsports-2013-092417.
- 3 59. Arnaoutis G, Kavouras SA, Angelopoulou A, et al. Fluid balance during training in elite young athletes of
4 different sports. *J Strength Cond Res.* 2015;29(12):3447-52. doi:10.1519/jsc.0000000000000400.
- 5 60. Bradley PS, Carling C, Gomez Diaz A, et al. Match performance and physical capacity of players in the top
6 three competitive standards of English professional soccer. *Hum Mov Sci.* 2013;32(4):808-21.
7 doi:10.1016/j.humov.2013.06.002.
- 8 61. Markovic G. Does plyometric training improve vertical jump height? A meta-analytical review. *Br J Sports*
9 *Med.* 2007;41(6):349-55; discussion 55. doi:10.1136/bjism.2007.035113.
- 10 62. Kraska JM, Ramsey MW, Haff GG, et al. Relationship between strength characteristics and unweighted and
11 weighted vertical jump height. *Int J Sports Physiol Perform.* 2009;4(4):461-73.
- 12 63. Salinero JJ, Lara B, Abian-Vicen J, et al. The use of energy drinks in sport: perceived ergogenicity and side
13 effects in male and female athletes. *Br J Nutr.* 2014;112(9):1494-502. doi:10.1017/s0007114514002189.
- 14 64. Gillet E, Leroy D, Thouwarecq R, et al. A notational analysis of elite tennis serve and serve-return strategies
15 on slow surface. *J Strength Cond Res.* 2009;23(2):532-9. doi:10.1519/JSC.0b013e31818efe29.
- 16 65. Essig D, Costill DL, Van Handel PJ. Effects of caffeine ingestion on utilization of muscle glycogen and lipid
17 during leg ergometer cycling. *Int J Sports Med.* 1980;01(02):86-90. doi:10.1055/s-2008-1034637.
- 18 66. Gonglach AR, Ade CJ, Bemben MG, et al. Muscle pain as a regulator of cycling intensity: effect of caffeine
19 ingestion. *Med Sci Sports Exerc.* 2016;48(2):287-96. doi:10.1249/mss.0000000000000767.
- 20 67. Bazzucchi I, Felici F, Montini M, et al. Caffeine improves neuromuscular function during maximal dynamic
21 exercise. *Muscle Nerve.* 2011;43(6):839-44. doi:10.1002/mus.21995.
- 22 68. Behrens M, Mau-Moeller A, Weippert M, et al. Caffeine-induced increase in voluntary activation and strength
23 of the quadriceps muscle during isometric, concentric and eccentric contractions. *Sci Rep.* 2015;5:10209.
24 doi:10.1038/srep10209.
- 25 69. McPherson PS, Kim YK, Valdivia H, et al. The brain ryanodine receptor: a caffeine-sensitive calcium release
26 channel. *Neuron.* 1991;7(1):17-25.
- 27 70. Landolt HP, Retey JV, Tonz K, et al. Caffeine attenuates waking and sleep electroencephalographic markers
28 of sleep homeostasis in humans. *Neuropsychopharmacology.* 2004;29(10):1933-9. doi:10.1038/sj.npp.1300526.
- 29 71. Lane JD, Steege JF, Rupp SL, et al. Menstrual cycle effects on caffeine elimination in the human female. *Eur*
30 *J Clin Pharmacol.* 1992;43(5):543-6.

- 1 72. Ali A, O'Donnell JM, Starck C, et al. The effect of caffeine ingestion during evening exercise on subsequent
2 sleep quality in females. *Int J Sports Med.* 2015;36(6):433-9. doi:10.1055/s-0034-1398580.
- 3 73. Tarnopolsky MA. Caffeine and endurance performance. *Sports Med.* 1994;18(2):109-25.
- 4 74. Mckim WA. Caffeine and the methylxanthines. In: Mckim WA, editor. *Drugs and Behavior: an introduction*
5 *to behavioral pharmacology.* New Jersey: Prentice Hall; 1996. p. 191-210.
- 6 75. Salinero JJ, Lara B, Ruiz-Vicente D, et al. CYP1A2 genotype variations do not modify the benefits and
7 drawbacks of caffeine during exercise: a pilot study. *Nutrients.* 2017;9(3). doi:10.3390/nu9030269.
- 8 76. Russell M, West DJ, Harper LD, et al. Half-time strategies to enhance second-half performance in team-sports
9 players: a review and recommendations. *Sports Med.* 2015;45(3):353-64. doi:10.1007/s40279-014-0297-0.
- 10 77. Duncan MJ, Taylor S, Lyons M. The effect of caffeine ingestion on field hockey skill performance following
11 physical fatigue. *Res Sports Med.* 2012;20(1):25-36. doi:10.1080/15438627.2012.634686.
- 12 78. Stevenson EJ, Hayes PR, Allison SJ. The effect of a carbohydrate-caffeine sports drink on simulated golf
13 performance. *Appl Physiol Nutr Metab.* 2009;34(4):681-8. doi:10.1139/h09-057.

14

1 **Figure Captions**

2 **Fig. 1** Flowchart summarizing search strategy and study selection process

1 Table Captions

Table 1. Acute effects of caffeine supplementation in ball games

Table 1. Acute effects of caffeine supplementation in ball games

| Study | Subject profile | Standard | Intervention | Mode | Measure (units) and change | CAF (mean ± SD) | PLA (mean ± SD) | <i>d</i> | P |
|----------------------|--|---|---|-------------------|---|---|--|--|--|
| Invasion | | | | | | | | | |
| Del Coso et al. [44] | 13 males/ hockey Low caffeine users: < 100 mg/day | Spanish first division hockey league At least 6 years of experience | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (250 ml) | 2 x 25-min simulated match: Total distance (m): NC Distance at moderate-intensity (m): ↓* Distance at high-intensity (m): ↑* Sprinting distance (m): ↑* Number of body accelerations and decelerations: NC Sprint count: ↑* Physiological measures: Mean HR (b·min ⁻¹): NC Peak HR (b·min ⁻¹): ↑* Sweat rate (L·h ⁻¹): NC Dehydration rate (%): NC | 6055 ± 499 712 ± 116 358 ± 117 117 ± 55 618 ± 221 6.3 ± 2.9 158 ± 16 185 ± 11 0.9 ± 0.4 -0.2 ± 0.8 | 6035 ± 451 793 ± 135 303 ± 67 85 ± 41 697 ± 285 4.5 ± 1.9 154 ± 14 181 ± 9 1.0 ± 0.3 -0.1 ± 0.7 | 0.04 -0.60 0.80 0.78 -0.28 0.95 0.29 0.44 -0.33 -0.14 | 0.87 0.03 0.05 0.02 0.15 0.05 0.34 0.03 0.30 0.50 |
| Abian et al. [47] | 16 males/ basketball Light caffeine users: < 60 mg/day | Spanish first division of National Spanish League At least 6 years of experience | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (250 ml) | Accuracy tests: Free throw (%): NC Three-point (%): NC Jump tests: CMJ height (cm): ↑* CMJ peak power (W·kg ⁻¹): NC 15s maximal jump test: Mean jump height (cm): ↑* Mean power (kW): ↑* Yo-yo test: Total distance (m): NC | 70.7 ± 11.8 39.9 ± 11.8 38.3 ± 4.4 53.9 ± 5.0 30.2 ± 3.6 51.4 ± 5.7 2000 ± 706 | 70.3 ± 11.0 38.1 ± 12.8 37.5 ± 4.4 53.8 ± 5.5 28.8 ± 3.4 49.4 ± 4.6 1925 ± 702 | 0.04 0.14 0.18 0.02 0.41 0.43 0.11 | 0.45 0.33 < 0.05 0.45 < 0.05 < 0.05 0.19 |
| Assi et al. [42] | 9 males/ rugby Consumed < 300 mg caffeine/day | University level Trained ≥ 4 times/week | 6.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (500 ml) | PAT: Total score: NC Left hand: ↑* Right hand: NC Physiological measures: Mean HR (b·min ⁻¹): NC | 9 ± 1 5 ± 1 4 ± 1 135 ± 9 | 8 ± 1 4 ± 1 4 ± 1 133 ± 7 | 1.00 1.00 0.00 0.29 | 0.24 0.01 0.75 0.15 |

Table 1 continued.

| Study | Subject profile | Standard | Intervention | Mode | Measure (units) and change | CAF (mean ± SD) | PLA (mean ± SD) | <i>d</i> | P |
|-------------------------------------|--|--|--|----------------------|--|--------------------|--------------------|--------------------|--------|
| Krasnanov a et al. [52] | 14 females/ floorball Light caffeine users: not more than 1 serving of caffeinated beverage per day | Slovakia league team | 4.4 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross- over design | Capsule | Standing long jump test: | 186.8 ± 23.5 | 189.9 ± 20.8 | -0.15 | > 0.05 |
| | | | | | Distance (cm): NC | | | | |
| | | | | | Shuttle run test: | 15.5 ± 0.9 | 15.7 ± 1.0 | -0.20 ^a | > 0.05 |
| | | | | | 9 m (s): NC | | | | |
| | | | | | 40 m (s): ↓* | 52.1 ± 4.4 | 54.0 ± 5.3 | -0.36 ^a | < 0.01 |
| | | | | | Reaction speed test: | 671.4 ± 68.9 | 694.1 ± 104.3 | -0.22 ^a | > 0.05 |
| Time to complete (ms): ↓ | | | | | | | | | |
| Shooting accuracy test: | 2.7 ± 1.5 | 2.8 ± 1.6 | -0.06 | > 0.05 | | | | | |
| Successful attempts: NC | | | | | | | | | |
| Lara et al. [48] | 18 females/ soccer Light caffeine users: not more than 1 coffee or 1 serving of energy drink per day | At least 3 years of soccer experience Trained ~ 2 hrs/day | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross- over design | Beverage (250 ml) | Jump tests: | 27.4 ± 3.8 | 26.6 ± 4.0 | 0.20 | < 0.05 |
| | | | | | CMJ height (cm): ↑* | | | | |
| | | | | | CMJ peak power (W·kg ⁻¹): ↑ | 43.0 ± 5.2 | 42.2 ± 4.5 | 0.18 | 0.08 |
| | | | | | Sprint test: | 25.6 ± 1.4 | 25.0 ± 1.4 | 0.43 | < 0.05 |
| | | | | | Maximal running speed test (km·h ⁻¹): ↑* | | | | |
| | | | | | 2 x 40-min simulated match: | 7087 ± 1501 | 6631 ± 1618 | 0.28 | < 0.05 |
| | | | | | Total distance (m): ↑* | | | | |
| | | | | | Sprint count: ↑* | | | | |
| | | | | | Maximal speed (km·h ⁻¹): NC | 24.2 ± 2.4 | 23.8 ± 2.5 | 0.16 | 0.25 |
| | | | | | Physiological measures: | 158 ± 12 | 152 ± 13 | 0.46 | 0.10 |
| | | | | | Mean HR (b·min ⁻¹): ↑ | | | | |
| | | | | | Peak HR (b·min ⁻¹): ↑ | | | | |
| Sweat rate (L·h ⁻¹): NC | 0.5 ± 0.1 | 0.6 ± 0.1 | -1.00 | 0.47 | | | | | |
| Dehydration rate (%): NC | 0.2 ± 0.8 | 0.2 ± 0.8 | 0.00 | 0.86 | | | | | |

Table 1 continued.

| Study | Subject profile | Standard | Intervention | Mode | Measure (units) and change | CAF (mean ± SD) | PLA (mean ± SD) | <i>d</i> | P |
|----------------------|--|--|--|----------------------|---|--|--|--|--|
| Petersen et al. [43] | 19 males/ soccer (outfield players) None were regular coffee- drinkers | Norwegian professional football club | 6.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross- over design | Capsule | 2 x 90-min simulated match: Total distance (m): NC Number of accelerations: NC Sprint distance (m): NC Yo-yo test: Total distance (m): NC Physiological measures: Blood glucose (mmol·L ⁻¹): ↑* Blood lactate (mmol·L ⁻¹): ↑* Mean HR (b·min ⁻¹): NC | 10062 ± 916 123 ± 31 109 ± 58 829 ± 328 7.5 ± 2.1 7.9 ± 4.8 166 ± 11 | 9854 ± 901 126 ± 24 112 ± 69 819 ± 289 6.3 ± 1.6 6.1 ± 3.6 168 ± 8 | 0.23 0.11 0.05 0.03 0.75 0.50 -0.25 | 0.13 0.67 0.84 0.91 < 0.05 < 0.05 > 0.05 |
| Del Coso et al. [35] | 26 males/ rugby Light caffeine users: < 60 mg/day | Spanish first division rugby league At least 5 years of experience Trained ~ 2 hrs, 4-5 times/week | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross- over design | Beverage (250 ml) | 2 x 30-min simulated match: Total distance (m): ↑* Distance covered at > 20 km·h ⁻¹ (m): ↑* Sprint count: ↑* Number of impacts: ↑* Physiological measures: Mean HR (b·min ⁻¹): NC Peak HR (b·min ⁻¹): NC Sweat rate (L·h ⁻¹): NC Dehydration rate (%): NC | 5139 ± 475 208 ± 38 12 ± 7 641 ± 366 151 ± 11 189 ± 12 1.8 ± 0.8 1.3 ± 0.9 | 4749 ± 589 184 ± 38 10 ± 7 481 ± 352 145 ± 8 185 ± 12 1.8 ± 0.6 1.3 ± 0.6 | 0.66 0.63 0.29 0.45 0.75 0.33 0.00 0.00 | 0.01 0.01 < 0.05 0.01 > 0.05 > 0.05 > 0.05 > 0.05 |
| Jordan et al. [37] | 17 males/ soccer No history of using caffeine as ergogenic aid | Premier league youth Trained 3 times/week | 6.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross- over design | Capsule | Reaction time: Dominant side (s): NC Non-dominant side (s): ↓* Sprint time: Dominant side (s): NC Non-dominant side (s): NC Total time to completion: Dominant side (s): NC Non-dominant side (s): ↓ | 0.801 ± 0.106 0.820 ± 0.095 1.001 ± 0.062 0.998 ± 0.046 1.806 ± 0.139 1.819 ± 0.116 | 0.847 ± 0.106 0.911 ± 0.153 1.001 ± 0.048 1.007 ± 0.060 1.854 ± 0.105 1.918 ± 0.186 | -0.43 ^a -0.59 ^a 0.00 -0.15 ^a -0.46 -0.53 | 0.35 0.04 0.94 0.81 0.33 0.05 |

Table 1 continued.

| Study | Subject profile | Standard | Intervention | Mode | Measure (units) and change | CAF (mean ± SD) | PLA (mean ± SD) | <i>d</i> | P |
|----------------------|---|---|---|---------------------|--|--|--|--|--|
| Del Coso et al. [49] | 16 females/ rugby Light caffeine users: < 60 mg/day | Spanish rugby sevens National team At least 4 years of experience Trained ~ 2 hrs, 4-5 times/week | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (250 ml) | 15s maximal jump test: Mean power (kW): ↑* Sprint test: Maximal running speed (km·h ⁻¹): NC 3 x 15-min simulated match: Mean running pace (km·h ⁻¹): ↑* Pace at sprint velocity (m·min ⁻¹): ↑* Physiological measures: Mean HR (b·min ⁻¹): ↑* Peak HR (b·min ⁻¹): NC Sweat rate (L·h ⁻¹): NC Dehydration rate (%): NC | 25.6 ± 11.8 25.0 ± 1.7 95.4 ± 12.7 6.1 ± 3.4 168 ± 7 189 ± 10 1.4 ± 0.3 0.5 ± 0.5 | 23.5 ± 10.1 25.0 ± 1.5 87.5 ± 8.3 4.6 ± 3.3 164 ± 6 188 ± 9 1.4 ± 0.5 0.6 ± 0.6 | 0.21 0.00 0.95 0.45 0.67 0.11 0.00 0.00 | 0.05 0.91 < 0.05 < 0.05 < 0.05 0.62 0.99 0.67 |
| Tucker et al. [34] | 5 males/ basketball Consumed < 500 mg caffeine/day | Elite level | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Capsule | Endurance performance: MVO ₂ (ml·min ⁻¹ ·kg ⁻¹): NC Anaerobic power: RSI (cm·s ⁻¹): NC Physiological measures: Blood lactate (mmol·L ⁻¹): ↑ | NS 111.6 ± 14.5 ^b NS | NS 111.6 ± 17.2 ^b NS | - 0.00 NS | > 0.05 > 0.05 NS |
| Del Coso et al. [46] | 19 males/ soccer Light caffeine users: < 60 mg/day | Semi-professional At least 5 years of experience Trained ~ 2 hrs, 4-5 times/week | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (~ 650 ml) | 15s maximal jump test: Mean jump height (cm): ↑* Mean power (kW): ↑* Repeated sprint ability test: Mean peak running speed (km·h ⁻¹): ↑* 2 x 40-min simulated match: Total distance (m): ↑* Sprint counts: ↑* | 35.8 ± 5.5 61.8 ± 7.3 26.3 ± 1.8 7782 ± 878 30 ± 10 | 34.7 ± 4.7 59.5 ± 6.9 25.6 ± 2.1 7352 ± 881 24 ± 8 | 0.23 0.33 0.33 0.49 0.75 | < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 |

Table 1 continued.

| Study | Subject profile | Standard | Intervention | Mode | Measure (units) and change | CAF (mean ± SD) | PLA (mean ± SD) | <i>d</i> | P |
|---------------------------------|---|--|---|-------------------|---|---|---|--|--|
| Del Coso et al. [46] continued. | | | | | Physiological measures: Mean HR (b·min ⁻¹): NC Peak HR (b·min ⁻¹): NC Sweat rate (L·h ⁻¹): NC | 160 ± 10 196 ± 7 1.1 ± 0.3 | 161 ± 12 197 ± 12 1.0 ± 0.3 | -0.08 -0.08 0.33 | > 0.05 > 0.05 > 0.05 |
| Foskett et al. [39] | 12 males/ soccer Consumed < 350 mg caffeine/day | Regional premier division | 6.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (500 ml) | LSPT: Total time (s): ↓* Number of perfect passes: ↑ Jump tests: CMJ height (cm): ↑* Physiological measures: Mean HR (b·min ⁻¹): NC | 51.6 ± 7.7 6.4 ± 0.4 57.1 ± 5.1 164 ± 11 | 53.9 ± 8.5 5.7 ± 0.6 55.6 ± 5.1 166 ± 11 | -0.27 ^a 1.17 0.29 -0.18 | 0.02 0.06 0.01 > 0.05 |
| Stuart et al. [38] | 9 males/ rugby Regular caffeine users | High level amateur players Trained ~ 5 hrs/week | 6.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Capsule | Rugby specific test: Passing accuracy: ↑ Offensive sprint time (s): ↓ Defensive sprint time (s): ↓ Tackle sprint time (s): ↓ 20 m sprint time (s): ↓ 30 m sprint time (s): ↓ First-drive power (W): ↑ Second-drive power (W): ↓ | 9.6 %; ± 6.1 ^c 1.3 %; ± 4.1 ^c 2.4 %; ± 2.8 ^c 2.9 %; ± 1.3 ^c 0.5 %; ± 1.7 ^c 2.3 %; ± 2.5 ^c 5.0 %; ± 2.5 ^c -1.2 %; ± 6.8 ^c | 4.2 ^d 6.0 ^d 13.6 ^d 9.3 ^d 3.3 ^d 5.0 ^d 1690 ^d 1470 ^d | - - - - - - - - | NS NS NS NS NS NS NS NS |
| Net | | | | | | | | | |
| Abian et al. [50] | 16 males/ badminton Light caffeine users: < 60 mg/day | Elite (8 international level players) At least 5 years of experience Trained ~ 2 hrs, 5 times/week | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (250 ml) | Handgrip maximal force: Left (N): ↑ Right (N): ↑ Jump variables: Squat jump height (cm): ↑* Smash jump height w shuttle (cm): ↑ Smash jump height w/o shuttle (cm): ↑ Squat jump peak power (W·kg ⁻¹): ↑* CMJ height (cm): ↑* CMJ peak power (W·kg ⁻¹): ↑* | 429 ± 87 477 ± 69 36.4 ± 4.3 39.4 ± 8.5 45.1 ± 5.5 55.4 ± 7.2 39.5 ± 5.1 56.1 ± 7.4 | 414 ± 75 465 ± 67 34.5 ± 4.7 38.8 ± 9.7 44.3 ± 6.4 52.4 ± 6.6 37.7 ± 4.5 54.2 ± 6.9 | 0.20 0.18 0.40 0.06 0.13 0.45 0.40 0.28 | 0.06 0.12 0.01 0.40 0.19 0.01 0.01 0.03 |

Table 1 continued

| Study | Subject profile | Standard | Intervention | Mode | Measure (units) and change | CAF (mean ± SD) | PLA (mean ± SD) | <i>d</i> | P |
|------------------------------|--|---|---|-------------------|---|--|--|--|--|
| Abian et al. [50] continued. | | | | | Agility T-test (s): ↓ 45-min simulated match: Total impacts: ↑* | 10.0 ± 0.4 7707 ± 2033 | 10.1 ± 0.4 7395 ± 1594 | -0.25 ^a 0.20 | 0.31 0.01 |
| Gallo-Salazar et al. [40] | 10 males, 4 females/ tennis Light caffeine users: < 1 can of soda or energy drink | Elite junior | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (250 ml) | Handgrip maximal force: Left (N): ↑* Right (N): ↑* Maximal velocity serve test (m/s): NC Sprint test: Maximal running speed (km·h ⁻¹): ↑ 3 set simulated match: Total distance (m): NC Sprint counts: ↑* Peak running velocity (m·s ⁻¹): ↑ Points won on service (%): ↑ Physiological measures: Mean HR (b·min ⁻¹): ↑ Peak HR (b·min ⁻¹): ↓ Sweat rate (L·h ⁻¹): ↑* Dehydration rate (%): ↑* | 361 ± 74 402 ± 83 42.7 ± 5.0 22.9 ± 2.1 2904 ± 430 13.2 ± 1.7 20.5 ± 2.8 51.8 ± 6.8 144 ± 4 178 ± 4 0.7 ± 0.3 0.2 ± 0.4 | 348 ± 76 387 ± 83 42.6 ± 4.8 22.3 ± 2.0 3058 ± 620 12.1 ± 1.7 19.5 ± 2.3 48.3 ± 7.2 143 ± 3 181 ± 3 0.5 ± 0.3 0.1 ± 0.5 | 0.17 0.18 0.02 0.30 -0.25 0.65 0.43 0.49 0.33 -1.00 0.67 0.20 | 0.03 0.03 0.49 0.07 0.24 0.05 0.44 0.16 0.35 0.44 0.04 0.04 |
| Perez-Lopez et al. [36] | 13 females/ volleyball No information on caffeine consumption habits | Spanish national league division 2 At least 6 years of experience Trained ~ 2 hrs, 5 times/week | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (250 ml) | Handgrip maximal force: Left (N): ↑* Right (N): ↑* Volleyball specific tests: Spike jump height (cm): ↑* Block jump height (cm): ↑* Block jump peak power (W·kg ⁻¹): ↑* Squat jump height (cm): ↑* Squat jump peak power (W·kg ⁻¹): ↑* CMJ height (cm): ↑* CMJ peak power (W·kg ⁻¹): ↑* | 335 ± 32 340 ± 45 44.4 ± 5.0 36.1 ± 5.1 50.7 ± 6.1 29.4 ± 3.6 47.6 ± 4.9 33.1 ± 4.5 46.9 ± 5.0 | 295 ± 47 303 ± 38 43.3 ± 4.7 35.2 ± 5.1 46.6 ± 6.4 28.1 ± 3.2 46.4 ± 4.0 32.0 ± 4.6 46.2 ± 5.1 | 0.85 0.97 0.15 0.18 0.64 0.36 0.30 0.24 0.14 | 0.02 0.00 0.02 0.04 0.01 0.03 0.01 0.02 0.05 |

Table 1 continued.

| Study | Subject profile | Standard | Intervention | Mode | Measure (units) and change | CAF (mean ± SD) | PLA (mean ± SD) | <i>d</i> | P |
|------------------------------------|--|---|---|-------------------|---|--|---|--|--|
| Perez-Lopez et al. [36] continued. | | | | | Physiological measures: Mean HR (b·min ⁻¹): ↑* Peak HR (b·min ⁻¹): ↑* | 142 ± 14 185 ± 10 | 132 ± 14 180 ± 5 | 0.71 1.00 | 0.04 0.05 |
| Del Coso et al. [51] | 15 males/ volleyball Light caffeine users: < 30 mg/day | Collegiate level At least 4 years of experience Trained ~ 2 hrs, 4 times/week | 3.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Beverage (250 ml) | Standing spike test: Maximal ball velocity (m·s ⁻¹): ↑* Jump tests: Squat jump height (cm): ↑* Squat jump peak power (W·kg ⁻¹): ↑* CMJ height (cm): ↑* CMJ peak power (W·kg ⁻¹): ↑* RJ-15: Mean jump height (cm): ↑* Mean peak power (W·kg ⁻¹): ↑ Agility T-test (s): ↓* 3 set simulated match: Frequency of successful actions: ↑* Physiological measures: Mean HR (b·min ⁻¹): NC Peak HR (b·min ⁻¹): NC Sweat rate (L·h ⁻¹): NC Dehydration rate (%): NC | 21.2 ± 2.3 32.7 ± 4.2 53.4 ± 5.5 37.7 ± 4.6 52.7 ± 5.0 30.7 ± 4.5 303 ± 65 10.3 ± 0.4 34.3 ± 16.5 134 ± 8 180 ± 12 1.0 ± 0.4 0.3 ± 0.2 | 20.6 ± 2.3 31.0 ± 4.3 51.6 ± 7.5 35.9 ± 4.4 51.4 ± 5.0 29.3 ± 4.8 295 ± 67 10.8 ± 0.4 24.6 ± 14.3 137 ± 13 184 ± 18 1.0 ± 0.3 0.1 ± 0.2 | 0.26 0.40 0.24 0.41 0.26 0.29 0.12 -1.25 ^a 0.68 -0.23 -0.22 0.00 1.00 | < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 > 0.05 < 0.05 < 0.05 > 0.05 > 0.05 > 0.05 > 0.05 |
| Klein et al. [45] | 8 males, 8 females/ tennis Consumed average 97 mg caffeine/day | Collegiate | 6.0 mg·kg ⁻¹ BM CAF or PLA in a double-blinded, randomized cross-over design | Capsule | TST: Number of successful shots: ↑* Physiological measures: Mean HR during TST (b·min ⁻¹): NC Mean HR during TM: (b·min ⁻¹): NC | 294.9 ± 10.9 153 ± 1 152 ± 3 | 288.8 ± 10.4 152 ± 1 151 ± 3 | 0.59 1.00 0.33 | 0.03 > 0.05 > 0.05 |

Table 1 continued.

| Study | Subject profile | Standard | Intervention | Mode | Measure (units) and change | CAF (mean ± SD) | PLA (mean ± SD) | <i>d</i> | P |
|---|--|---|--|---------|--|-----------------|-----------------|----------|--------|
| Hornery et al. [41] | 12 males/ tennis Non-habitual caffeine consumers | At least 5 years of competitive experience Trained 15- 20 hrs/week | 3.0 mg·kg ⁻¹ BM CAF or PLA in a single-blinded, randomized cross- over design | Capsule | Serve: | 165 ± 15 | 159 ± 15 | 0.40 | 0.01 |
| | | | | | Velocity in final set of match (km·h ⁻¹): ↑* | | | | |
| | | | | | Groundstroke: | NS | NS | - | > 0.05 |
| | | | | | Stroke accuracy (%): NC | | | | |
| Physiological measures: | NS | NS | - | NS | | | | | |
| Blood glucose (mmol·L ⁻¹): NS | | | | | | | | | |
| Blood lactate (mmol·L ⁻¹): NS | NS | NS | - | NS | | | | | |

Note. BM = body mass, CAF = caffeine, CMJ = countermovement jump, *d* = effect size, HR = heart rate, b·min⁻¹ = beats per minute, MVO₂ = maximal oxygen consumption, LSPT = Loughborough Soccer Passing Test, PAT = passing accuracy test, PLA = placebo, RJ-15 = 15s rebound jump, RPE = rating of perceived exertion, RSI = reactive strength index, TM = treadmill exercise, TST = tennis skill test, w = with, w/o = without, NC = no change, NS = not specified, ↑ = increased, ↓ = decreased, * = significant change, - = not applicable, ^a = a negative effect size (*d*) indicates performance improvement, ^b = mean values calculated from data provided in study, ^c = mean percentage change ± 90 % confidence limits, ^d = standard deviation values not provided in study.