



Omega 3 fatty acids, inflammation and DNA methylation: an overview

Bethan Hussey, Martin R. Lindley & Sarabjit S. Mastana

To cite this article: Bethan Hussey, Martin R. Lindley & Sarabjit S. Mastana (2017) Omega 3 fatty acids, inflammation and DNA methylation: an overview, *Clinical Lipidology*, 12:1, 24-32

To link to this article: <http://dx.doi.org/10.1080/17584299.2017.1319454>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 02 May 2017.



Submit your article to this journal [↗](#)



Article views: 266



View related articles [↗](#)



View Crossmark data [↗](#)

Omega 3 fatty acids, inflammation and DNA methylation: an overview

Bethan Hussey , Martin R. Lindley  and Sarabjit S. Mastana 

Translational Chemical Biology Research Group, School of Sport Exercise and Health Sciences, Loughborough University, Loughborough, UK

ABSTRACT

Omega-3 polyunsaturated fatty acids (n-3 PUFAs) are known to be anti-inflammatory and to alter gene expression within the cells. Emerging evidence indicates that one of the mechanisms for this process involves the alteration of epigenetic markers, such as DNA methylation. The focus of this overview is to document the current evidence for n-3 PUFA effects on DNA methylation and how these may impact on the inflammatory processes.

ARTICLE HISTORY

Received 24 December 2016
Revised 22 February 2017
Accepted 11 April 2017

KEYWORDS

Fish oils; n-3 PUFAs;
inflammation; epigenetics;
DNA methylation

Introduction

n-3 polyunsaturated fatty acids and inflammation

Populations with a diet rich in omega-3 (n-3) polyunsaturated fatty acids (PUFAs) have a lower incidence of chronic non-communicable diseases [1]. Evidence for n-3 PUFAs, such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), on their ability to modulate inflammation has come from *in vitro* work [2–4], animal models [5–8] and human studies [4,9,10]; although the latter evidence is weaker as a consequence of more complex environmental factors. The mechanisms through which n-3 PUFAs promote an anti-inflammatory environment within the body are multiple and complex [11] (Figure 1). Early evidence suggests that n-3 PUFAs exert these effects through changes in gene expression as a result of varied transcription factor activation [12,13], but which may also arise from changes in epigenetic markers, such as DNA methylation.

Epigenetics

Epigenetics investigates heritable chemical modifications to the genome which are independent of the DNA sequence [14–16]. These chemical changes, including DNA methylation, histone modifications, chromatin remodelling and non-coding RNAs [17–23], are vital for normal cellular development, cellular processes and cell-specific gene expression profiles [16,24]. Interactions between the environment, particularly diet, and the genome through epigenetic mechanisms lead to changes in phenotype and

are implicated in many diseases [25–28]. They also provide a target for therapeutic interventions [29].

DNA methylation, a methyl group covalently attached to the fifth position of the pyrimidine ring to give 5-methylcytosine (5mC), is most commonly found on a cytosine positioned next to a guanine (CpG) [30]. DNA methylation levels within the genome are cell and tissue specific, with changes observed during the human lifespan [31–33]. Dysregulation of 5mC has been associated with diseases, including cancer and cardiovascular disease [24,34,35]. DNA methylation is the most accessible and widely studied epigenetic mark [36] and has been shown to vary with nutrition, disease and age [37–39]. Here, we will discuss the emerging evidence for the interaction of n-3 PUFAs and DNA methylation, specifically discussing their impact on inflammation.

Cross-sectional epigenome-wide association studies

Epigenome-wide associations studies (EWAS), which despite their name, focus mainly on DNA methylation, have linked n-3 PUFA consumption with differential DNA methylation. However, there is a lack of EWAS that use biochemical analysis to assess dietary n-3 PUFA, with a preference towards the readily available, but less accurate, food frequency questionnaire (FFQ).

One prominent EWAS within a distinct population, the *Yup'ik*, from Alaska ($n = 185$) utilised biochemical analysis to investigate associations with DNA methylation [40]. The

Yup'ik have a traditional diet rich in fish which is changing with westernisation [41], allowing for stratification of this population into high and low PUFA consumption. Using red blood cell (RBC) nitrogen stable isotope ($\delta^{15}\text{N}$), previously associated with EPA and DHA RBC levels [42], to separate into these high and low PUFA groups, associations with CpG methylation of biologically relevant targets, including genes involved in T-cell homeostasis were found [40]. Increased DNA methylation with high PUFA consumption was observed in 78% of the significant associations [40]; with genomic instability occurring with hypomethylation [43], the increased methylation observed may, therefore, be beneficial to the stability of the genome. A separate cohort, Greek pre-adolescents ($n = 69$), also found associations between dietary fats and DNA methylation, including sites within pathways linked to inflammation; nuclear factor kappa B (*NFkB*), peroxisome proliferator-activated receptor (*PPAR α*), leptin (*LEP*) and interleukin (*IL*)-6 [44]. However, the dietary fat intake for the cohort was assessed using FFQs, which did not directly measure the levels of n-3 PUFAs that have made it to the target tissue from the food.

Consistent with previous supplementation studies [45,46] and dietary salmon consumption [47], the *Yup'ik* cohort found individuals with higher n-3 PUFA intake had significantly lower levels of plasma triglyceride (TG), increased high-density lipoproteins (HDL), and higher but not significant levels of low-density lipoproteins (LDL) and

total cholesterol [40]. However, Dekkers *et al.* [48] found that there was an effect of blood lipids (TG, HDL and LDL) on differential DNA methylation in the genes coding for six key regulators of lipid metabolism. It therefore, needs to be questioned if the differential methylation observed is due to the n-3 PUFAs in the diet, differences in blood lipid profiles, or both, and one method of addressing this is through closely controlled intervention studies.

Gestation and infant n-3 PUFA supplementation intervention studies

Epigenome-wide association studies

The impact of nutrition during early life is of increasing interest due to the plasticity of epigenetic regulation during development and the links to diseases in later life [49,50]. Supplementation during gestation or early infancy with n-3 PUFAs has therefore been studied to a greater extent than it has been in adults.

EWAS in combination pre-and post-intervention can be used to investigate the impact of n-3 PUFA supplementation on differentially methylated regions (DMRs) throughout the genome. Two intervention studies took slightly different approaches, the first supplemented the mothers with 3.7 g/day EPA + DHA ($n = 36$) or placebo ($n = 34$, placebo details not provided) from 20 weeks of gestation to delivery [51]. The second study supplemented




Figure 1. Mechanisms Omega-3 polyunsaturated fatty acids (n-3 PUFAs) action to promote an anti-inflammatory environment.

Notes: GPR120, G-coupled protein receptor 120; NFkB, nuclear factor kappa B; n-6 PUFAs, omega 6 polyunsaturated fatty acids; SPM, Specialised pro-resolving mediators; TLR4, toll like receptor 4.

9-month-old infants with either 1.6 g/day EPA + DHA ($n = 6$) or placebo 3.1 g/day linoleic acid ($n = 6$, LA in the form of sunflower oil) for nine months [52]. Neither investigation found a significant difference in DNA methylation between those taking the n-3 PUFA and those taking the placebo after adjusting for multiple testing [51,52].

However, a third more recent large randomised control trial (RCT) supplementing mothers during gestation with 0.8 g/day DHA + 0.1 g/day EPA ($n = 190$) or vegetable oil ($n = 179$), found that there were significant differences between DMRs, including those relating to immune function, between the two experimental groups [53]. The dose of n-3 PUFAs in this third study was much lower, in addition, due to the larger cohort it was possible to split the analysis between the sexes, with males being found to have a greater number of DMRs than females. The lack of findings in the first two studies referred to above may be due to the small, mixed sex samples used and future studies could be designed to eliminate this possibility. There are significant differences in the way males and females metabolise and store n-3 PUFAs [54–56] as well as differences in global methylation levels between the sexes [57]; as seen by the clear effect observed between sexes in the third study [53]. It is therefore, important to note that when using small sample sizes with n-3 PUFA and DNA methylation studies, a lack of separation of the sexes may result in critical results being overlooked.

Notably, all studies presented thus far have utilised the Infinium Human Methylation 450 k arrays. This technology measures DNA methylation at over 450,000 individual CpG sites throughout the genome. And although a powerful tool in explorative work in determining DMRs, there are limitations to the use of 450 k array data, including multiple testing, the need for complicated normalisation and the generation of artefactual data [58]. Therefore, targeted analysis of specific targets including candidate genes must be completed to confirm findings, and investigate differential methylation and biological relevance.

Specific targets and candidate genes

Long interspersed nucleotide element 1 (LINE1)

Measuring the levels of DNA methylation in repetitive elements, such as long interspersed nucleotide elements (LINEs) can provide a proxy for global DNA methylation [59]. It has been estimated that repetitive transposable elements account for 45% of the genome [60] and they contain one-third of the genomes DNA methylation [61,62] to repress their transcription [63]. Associations between *LINE1* methylation and lifestyle factors, including physical activity and nutrition [57,64–67] have been found and have also been shown to be associated with both disease [68–70] and ageing [71,72].

Lee *et al.* investigated the DNA methylation of *LINE1* in cord blood mononuclear cells (CBMCs) from mothers who received 0.4 g/day DHA ($n = 131$) or placebo ($n = 130$, olive oil) during gestation [73]. Neither this study or the third RCT above found any differences in *LINE1* methylation between supplementation and control groups [53,73]. However, Lee *et al.* did find an interaction between smoking status and DHA supplementation; CBMCs of mothers who smoked and took the DHA during gestation had significantly higher *LINE1* methylation levels than the CBMCs of smoking mothers in the control group [73]. This suggests a complex interaction between multiple environmental stimuli and the epigenome.

Cytokine genes

In addition to *LINE1* methylation, cytokine gene DNA methylation was also investigated by Lee *et al.*, with no significant differences in methylation of *TNFA*, *IL13*, *GATA3*, *STAT3*, *IL10* and *FOXP3* being observed. However, the promoter methylation was lower (not significantly) for *IFN γ* in those supplemented with DHA [73]. The study was novel in that the supplementation only contained DHA and was also at a low dose compared to many other studies. As observed in cell membranes where there is a dose-response increase in the incorporation of n-3 PUFAs, it is likely that an increase in the dose may have led to more significant changes in DNA methylation. *In vitro* studies provide evidence of differing effects of EPA and DHA on inflammation [74,75], it would be interesting to see if there are differing effects on the epigenome had a separate EPA supplementation group had been included. The comparison between EPA and DHA is often lacking in human studies due to the increased cost and a greater availability of mixed n-3 PUFA supplements commercially.

Adult n-3 PUFA supplementation intervention studies

Candidate targets

PUFAs are important structural components of cells membranes. Consumption of foods rich in n-3 PUFAs, such as salmon, result in increased incorporation and modification of the cell phospholipid membrane [47]. Composition of the membrane is altered by clustering of lipids rafts containing the n-3 PUFAs, resulting in the formation of large raft domains [76]. As shown in murine obesity models [77] and HeLa cells [78] large lipid rafts can suppress the cell activation by impaired signalling, ultimately affecting cell function through suppression of downstream pathways, including those involved in inflammation. Leptin is localised to these lipid rafts and its expression is reduced by n-3 PUFAs [79]. In humans, cross-sectional evidence indicates

differential methylation in the leptin pathway genes with varying n-3 PUFA status [44], and epigenetic control of *LEP* has been demonstrated in a murine model [80]. However, using a candidate gene approach, n-3 PUFA supplementation has been shown to have no effect on *LEP*, leptin receptor (*LEPR*) or pro-opiomelanocortin (*POMC*) promoter methylation in a murine model [81].

The membrane glycoprotein which promotes inflammation in monocytes and macrophages, cluster of differentiation 36 (*CD36*), has been found to be increased as the result of n-3 PUFA *in vitro* [82] and in animals [83]. *CD36* promoter methylation was significantly reduced, when adjusted for baseline body weight, in a weight loss and n-3 PUFA supplementation study in Spanish young adult overweight females [84]. However, changes in DNA methylation in cluster of differentiation (*CD14*), pyruvate dehydrogenase kinase 4 (*PDK4*) and fatty acid desaturase 1 (*FADS1*) was only significant as a result of the low-calorie diet and not the supplementation. Had the gene expression been studied we may have expected to see an increase in the mRNA for these genes. This effect is the counter to that expected from n-3 PUFAs where in human studies, decreased expression of *CD36*, *CD14* and *PDK4* mRNA is observed in peripheral blood mononuclear cells (PBMCs) isolated after supplementation [13]. It is possible that the low-calorie diet impacted on the DNA methylation to a greater extent than the n-3 PUFAs, which only attenuated the decrease in methylation of *CD36* once adjusted for baseline body weight [84].

There are relatively few studies that have investigated the effect of n-3 PUFA supplementation on DNA methylation in adults, and of these many have confounding factors. Addressing the confounding factors of this study [84], the participants were asked not to consume seafood and to follow a detailed diet plan during the intervention period to minimise variability. However, the Spanish diet is high in shellfish/fish, on average 88.6 g/person/day are consumed [85], and therefore, the researchers may have simply replaced the n-3 PUFAs that the participants obtained from their diets with the supplementation. Although no biochemical measurements for EPA and DHA were provided in this paper [84], earlier work from the same cohort found only a small difference in EPA and DHA after the intervention, with no data comparing baseline to endpoint [86]. Future studies should include measurement of n-3 PUFA incorporation in to the investigated cell membranes, for example, into PBMCs.

Endogenous metabolism of PUFAs occurs within a cell, the initial step in the metabolic pathway takes places in the endoplasmic reticulum where LA and alpha-linolenic acid (ALA) undergo elongation of the fatty acid carbon chain by fatty acid elongase (ELOVL) and desaturation by insertion of double bonds by $\Delta 6$ -desaturase and $\Delta 5$ -desaturase [87]. The $\Delta 6$ -desaturase and $\Delta 5$ -desaturase enzymes, encoded

by the *FADS2* and *FADS1* genes respectively, are rate-limiting steps in this metabolic pathway [88,89]. Genetic polymorphisms of these genes and others related to the PUFA metabolism can significantly alter the levels of PUFAs within the body [90] and differential methylation is likely to have a similar effect. The methylation of *FADS1* is altered by a low-calorie diet with a small interaction with n-3 PUFA supplementation [84]. When investigated alone, with a higher dose of n-3 PUFA supplementation, no significant changes in methylation were observed in *FADS1* or the elongation gene *ELOVL-2* [91]. However, the methylation of two other fatty acid metabolism genes, *FADS2* and *ELOVL-5*, increased with supplementation; with a larger number of CpG sites changing methylation in females than males and a negative relationship between the DNA methylation and the gene mRNA levels observed [91].

Future targets: inflammatory gene methylation

Modulation of inflammatory gene expression occurs through transcription factors, such as peroxisome proliferator-activated receptor gamma (PPAR γ). Interactions between PPAR γ and fatty acids result in a decrease in cytokine expression [92]. In a cancer cell line, treatment with EPA and DHA has been shown to increase expression of PPAR γ gene, *PPARG* [93]. In murine models of diabetes, expression of *PPARG* is known to be modulated by DNA methylation within its promoter [94] and differential methylation is observed in type two diabetes mellitus (T2DM) [95], however the impact of n-3 PUFAs on *PPARG* DNA methylation has not been measured. Epigenetic regulation of the PPAR coactivator 1 alpha (PCG-1 α) gene, *PPARGC1A*, has been implicated in T2DM in humans [96]. The regulation of the *PPARGC1A* appears to be influenced by changes in DNA methylation and has been shown to interact with PPAR γ . In men, DNA methylation of the *PPARGC1A* gene promoter is significantly increased after a high-fat over-feeding diet [97]. Whereas, in obese patients fed a low-calorie diet, the methylation of *PPARGC1A* has been shown to decrease [98].

Inflammatory diseases are often characterised with chronic activation of NF κ B transcription factor and release of inflammatory cytokines [99] and therefore, an important target for changes in DNA methylation by n-3 PUFAs. Associations between dietary fats and DNA methylation in the NF κ B pathway, measured using the Infinium 450 k array, were observed in the Greek pre-adolescent cross-sectional cohort [44], and therefore, may provide valuable targets within intervention studies.

As previously described, there was no change in DNA methylation being observed in cytokine DNA methylation levels with gestational supplementation [73]. However, the study used low doses of DHA and it has not been investigated in adult intervention studies or with combined EPA

and DHA supplementation. Taking TNF α as an example, EPA treatment does not affect the stability of TNF α mRNA in cytoplasm and therefore, reductions in TNF α mRNA as a result of EPA treatment is likely to be the result of decreased formation rather than degradation of the cytokine [100]. The influence of n-3 PUFAs on TNF α expression may be the result of alterations in epigenetic mechanism, such as DNA methylation. Epigenetic regulation plays an important role in the control of TNF α expression [101–103]. Specifically, TNF α has been shown to be hypomethylated in Crohn's disease patients [104], is related to adipogenesis [105], and that CpG methylation interacts with histone methylation to silence TNF α expression during endotoxin tolerance [106]. Cross-sectional studies employing FFQ have found associations between TNF α methylation and dietary intake of certain nutrients including fats, however, no associations with n-3 PUFAs [107,108]. As seen with TNF α , studies examining DNA methylation must choose the tissue of interest carefully as more variability is observed between tissues from the same individual, than within tissue between individuals in the TNF α promoter [109].

Implications and considerations for future research

Most of evidence for the control of gene expression by CpG methylation comes from genes with CpG islands [30]. However, not all genes have CpG islands and it is not only the DNA methylation that occurs in the promoter region near the transcription start site (TSS) that plays an important role in regulation of gene expression, with evidence suggesting that 5mC in gene bodies and non-promoter regions plays an important role [110]. In genes which do not have a CpG island, including inflammatory genes *NOS2*, *IL-2*, *IFN- γ* , *MMP13*, *IL-1 β* and *TNF- α* ; CpG methylation proximal to the TSS has been shown to influence transcription [106,111–115]. Careful considerations must therefore be taken when designing assays to measure the DNA methylation of candidate genes.

As we have seen, investigations of the effects of n-3 PUFAs on DNA methylation are in their infancy and as such there are some points to consider when conducting this research. Within the n-3 PUFAs literature there is not yet a consensus on the required dose in humans required to produce an anti-inflammatory effect. Many of the existing supplementation studies so far investigating the epigenetic effect of n-3 PUFAs have used low doses with only small changes observed. In a review by Calder [11], it is suggested that a dose between 1.35 and 2.7 g/day of EPA is required for the anti-inflammatory effects. It is likely that there will be a dose-response interaction between n-3 PUFA supplementation and epigenetic mechanisms, as is seen with cytokine production [116]. There is also no consensus on the time frame required for epigenetic changes

to occur. The largest incorporation of n-3 PUFAs into erythrocyte membranes is within the first 4 weeks of supplementation and it is not known if the epigenetic changes occur within the same time frame. There are fewer studies that measure the incorporation into PBMCs and none that consider CpG methylation alongside incorporation.

The majority of intervention studies investigating epigenetic changes do not account for the impact of the intervention on the cell population from which the DNA was extracted. This results in multiple complications, firstly if the influence of n-3 PUFAs is specific to one cell type it may not be observed if the cells are not separated prior to analysis. Secondly, changes may be falsely observed or masked with variations in cell populations. Differentiation into the cell subtypes found within blood requires changes within the epigenetic signatures and therefore, the observed changes in percentage methylation for an individual CpG may be the result of a change in the cell population rather than a result of the intervention. Data can be normalised to account for the proportion of different cells [52,91] and this is a consideration future studies should make.

Conclusion

Associations between n-3 PUFAs and differential DNA methylation have been demonstrated, with further evidence of n-3 PUFA impact on DNA methylation seen in supplementation interventions. A more targeted approach to the effects of n-3 PUFAs, and use of well-controlled supplementation studies, both *in vitro* and *in vivo*, could elucidate a mechanism through which n-3 PUFAs alter the DNA methylation within the inflammatory landscape.

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

Bethan Hussey  <http://orcid.org/0000-0002-1827-5489>
 Martin R. Lindley  <http://orcid.org/0000-0001-7686-9421>
 Sarabjit S. Mastana  <http://orcid.org/0000-0002-9553-4886>

References

- [1] Simopoulos AP. Essential fatty acids in health and chronic disease. *Am J Clin Nutr.* 1999;70:560S–569S.
- [2] Verlengia R, Gorjao R, Kanunfre C, et al. Comparative effects of eicosapentaenoic acid and docosahexaenoic acid on proliferation, cytokine production, and pleiotropic gene expression in Jurkat cells. *J Nutr Biochem.* 2004;15:657–665.
- [3] Trebble T, Arden NK, Stroud MA, et al. Inhibition of tumour necrosis factor- α and interleukin 6 production by mononuclear cells following dietary fish-oil

- supplementation in healthy men and response to antioxidant co-supplementation. *Br J Nutr.* **2003**;90:405–412.
- [4] Endres S, Ghorbani R, Kelley VE, et al. The effect of dietary supplementation with n-3 polyunsaturated fatty acids on the synthesis of interleukin-1 and tumor necrosis factor by mononuclear cells. *N Engl J Med.* **1989**;320:265–271.
- [5] Yaqoob P, Calder P. Effects of dietary lipid manipulation upon inflammatory mediator production by murine macrophages. *Cell Immunol.* **1995**;163:120–128.
- [6] Richard C, Lewis ED, Goruk S, et al. A dietary supply of docosahexaenoic acid early in life is essential for immune development and the establishment of oral tolerance in female rat offspring. *J Nutr.* **2016**;146:2398–2406.
- [7] Olson MV, Liu Y-C, Dangi B, et al. Docosahexaenoic acid reduces inflammation and joint destruction in mice with collagen-induced arthritis. *Inflamm Res.* **2013**;62:1003–1013.
- [8] Al-Khalifa H, Givens DI, Rymer C, et al. Effect of n-3 fatty acids on immune function in broiler chickens. *Poult Sci.* **2012**;91:74–88.
- [9] Itariu BK, Zeyda M, Hochbrugger EE, et al. Long-chain n-3 PUFAs reduce adipose tissue and systemic inflammation in severely obese nondiabetic patients: a randomized controlled trial. *Am J Clin Nutr.* **2012**;96:1137–1149.
- [10] Tartibian B, Maleki BH, Abbasi A. Omega-3 fatty acids supplementation attenuates inflammatory markers after eccentric exercise in untrained men. *Clin J Sport Med.* **2011**;21:131–137.
- [11] Calder PC. Marine omega-3 fatty acids and inflammatory processes: effects, mechanisms and clinical relevance. *Biochem Biophys Acta – Mol Cell Biol Lipids.* **2015**;1851:469–484.
- [12] Rudkowska I, Paradis A-M, Thifault E, et al. Transcriptomic and metabolomic signatures of an n-3 polyunsaturated fatty acids supplementation in a normolipidemic/normocholesterolemic Caucasian population. *J Nutr Biochem.* **2013**;24:54–61.
- [13] Bouwens M, van de Rest O, Dellschaft N, et al. Fish-oil supplementation induces antiinflammatory gene expression profiles in human blood mononuclear cells. *Am J Clin Nutr.* **2009**;90:415–424.
- [14] Holliday R. Epigenetics: a historical overview. *Epigenetics.* **2006**;1:76–80.
- [15] Huang B, Jiang C, Zhang R. Epigenetics: the language of the cell? *Epigenomics.* **2014**;6:73–88.
- [16] Bernstein BE, Meissner A, Lander ES. The mammalian epigenome. *Cell.* **2007**;128:669–681.
- [17] Barski A, Cuddapah S, Cui K, et al. High-resolution profiling of histone methylations in the human genome. *Cell.* **2007**;129:823–837.
- [18] Kouzarides T. Chromatin modifications and their function. *Cell.* **2007**;128:693–705.
- [19] Berger SL. The complex language of chromatin regulation during transcription. *Nature.* **2007**;447:407–412.
- [20] Jenuwein T. Translating the histone code. *Science (80-).* **2001**;293:1074–1080.
- [21] Bird A. DNA methylation patterns and epigenetic memory. *Genes Dev.* **2002**;16:6–21.
- [22] Goll MG, Bestor TH. Eukaryotic cytosine methyltransferases. *Ann Rev Biochem.* **2005**;74:481–514.
- [23] Mattick JS, Makunin IV. Non-coding RNA. *Hum Mol Genet.* **2006**;15:R17–R29.
- [24] Jones PA, Baylin SB. The epigenomics of cancer. *Cell.* **2007**;128:683–692.
- [25] Park LK, Friso S, Choi S-W. Nutritional influences on epigenetics and age-related disease. *Proc Nutr Soc.* **2012**;71:75–83.
- [26] Brunet A, Berger SL. Epigenetics of aging and aging-related disease. *J Gerontol Ser A Biol SciMed Sci.* **2014**;69:S17–S20.
- [27] Benayoun BA, Pollina EA, Brunet A. Epigenetic regulation of ageing: linking environmental inputs to genomic stability. *Nat Rev Mol Cell Biol.* **2015**;16:593–610.
- [28] Portela A, Esteller M. Epigenetic modifications and human disease. *Nat Biotechnol.* **2010**;28:1057–1068.
- [29] Kelly TK, De Carvalho DD, Jones PA. Epigenetic modifications as therapeutic targets. *Nat Biotechnol.* **2010**;28:1069–1078.
- [30] Jones PA, Takai D. The role of DNA methylation in mammalian epigenetics. *Science.* **2001**;293:1068–1070.
- [31] Gilbert SF. Ageing and cancer as diseases of epigenesis. *J Biosci.* **2009**;34:601–604.
- [32] Wilson VL, Smith RA, Ma S, et al. Genomic 5-methyldeoxycytidine decreases with age. *J Biol Chem.* **1987**;262:9948–9951.
- [33] Jones MJ, Goodman SJ, Kobor MS. DNA methylation and healthy human aging. *Aging Cell.* **2015**;14:924–932.
- [34] Ehrlich M. DNA methylation in cancer: too much, but also too little. *Oncogene.* **2002**;21:5400–5413.
- [35] Muka T, Koromani F, Portilla E, et al. The role of epigenetic modifications in cardiovascular disease: a systematic review. *Int J Cardiol.* **2016**;212:174–183.
- [36] Lam LL, Emberly E, Fraser HB, et al. Factors underlying variable DNA methylation in a human community cohort. *Proc Nat Acad Sci.* **2012**;109:17253–17260.
- [37] Gomes MVM, Toffoli LV, Arruda DW, et al. Age-related changes in the global DNA methylation profile of leukocytes are linked to nutrition but are not associated with the MTHFR C677T genotype or to Functional capacities. Suzuki H, editor. *PLoS ONE.* **2012**;7:e52570.
- [38] Choi S-W, Friso S. Epigenetics: a new bridge between nutrition and health. *Adv Nutr.* **2010**;1:8–16.
- [39] Bacalini MG, Friso S, Olivieri F, et al. Present and future of anti-ageing epigenetic diets. *Mech Ageing Dev.* **2014**;136–137:101–115.
- [40] Aslibekyan S, Wiener HW, Havel PJ, et al. DNA methylation patterns are associated with n-3 fatty acid intake in Yup'ik people. *J Nutr.* **2014**;144:425–430.
- [41] Bersamin A, Luick BR, King IB, et al. Westernizing diets influence fat intake, red blood cell fatty acid composition, and health in remote alaskan native communities in the center for alaska native health study. *J Am Diet Assoc.* **2008**;108:266–273.
- [42] O'Brien DM, Kristal AR, Jeannet MA, et al. Red blood cell 15 N: a novel biomarker of dietary eicosapentaenoic acid and docosahexaenoic acid intake. *Am J Clin Nutr.* **2009**;89:913–919.
- [43] Chen RZ, Pettersson U, Beard C, et al. DNA hypomethylation leads to elevated mutation rates. *Nature.* **1998**;395:89–93.
- [44] Voisin S, Almén MS, Moschonis G, et al. Dietary fat quality impacts genome-wide DNA methylation patterns in a cross-sectional study of Greek preadolescents. *Eur J Hum Genet.* **2015**;23:654–662.

- [45] Harris WS. n-3 fatty acids and serum lipoproteins: human studies. *Am J Clin Nutr.* 1997;65:1645S–1654S.
- [46] Das UN. Beneficial effect(s) of n-3 fatty acids in cardiovascular diseases: but, why and how? *Prostaglandins. Leukot Essent Fat Acids.* 2000;63:351–362.
- [47] Raatz SK, Johnson LK, Rosenberger TA, et al. Twice weekly intake of farmed Atlantic salmon (*Salmo salar*) positively influences lipoprotein concentration and particle size in overweight men and women. *Nutr Res.* 2016;36:899–906.
- [48] Dekkers KF, van Iterson M, Slieker RC, et al. Blood lipids influence DNA methylation in circulating cells. *Genome Biol.* 2016;17:138–150.
- [49] Gluckman PD, Hanson MA, Cooper C, et al. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med.* 2008;359:61–73.
- [50] Gluckman PD, Hanson MA. Living with the past: evolution, development, and patterns of disease. *Science.* 2004;305:1733–1736.
- [51] Amarasekera M, Noakes P, Strickland D, et al. Epigenome-wide analysis of neonatal CD4 + T-cell DNA methylation sites potentially affected by maternal fish oil supplementation. *Epigenetics.* 2014;9:1570–1576.
- [52] Lind MV, Martino D, Harsløf LBS, et al. Genome-wide identification of mononuclear cell DNA methylation sites potentially affected by fish oil supplementation in young infants: a pilot study. *Prostaglandins Leukot Essent Fat Acids.* 2015;101:1–7.
- [53] van Dijk SJ, Zhou J, Peters TJ, et al. Effect of prenatal DHA supplementation on the infant epigenome: results from a randomized controlled trial. *Clin Epigenetics.* 2016;8:114–127.
- [54] Burdge GC, Wootton SA. Conversion of α -linolenic acid to eicosapentaenoic, docosapentaenoic and docosahexaenoic acids in young women. *Br J Nutr.* 2002;88:411–422.
- [55] Burdge GC, Jones AE, Wootton SA. Eicosapentaenoic and docosapentaenoic acids are the principal products of α -linolenic acid metabolism in young men. *Br J Nutr.* 2002;88:355–363.
- [56] Lohner S, Fekete K, Marosvölgyi T, et al. Gender differences in the long-chain polyunsaturated fatty acid status: systematic review of 51 publications. *Ann Nutr Metab.* 2013;62:98–112.
- [57] Zhang FF, Cardarelli R, Carroll J, et al. Significant differences in global genomic DNA methylation by gender and race/ethnicity in peripheral blood. *Epigenetics.* 2011;6:623–629.
- [58] Dedeurwaerder S, Defrance M, Bizet M, et al. A comprehensive overview of Infinium human methylation 450 data processing. *Brief Bioinform.* 2014;15:929–941.
- [59] Zhu Z-Z, Hou L, Bollati V, et al. Predictors of global methylation levels in blood DNA of healthy subjects: a combined analysis. *Int J Epidemiol.* 2012;41:126–139.
- [60] Lander ES, Linton LM, Birren B, et al. Initial sequencing and analysis of the human genome. *Nature.* 2001;409:860–921.
- [61] Schmid CW. Does SINE evolution preclude Alu function? *Nucleic Acids Res.* 1998;26:4541–4550.
- [62] Kochanek S, Renz D, Doerfler W. DNA methylation in the Alu sequences of diploid and haploid primary human cells. *EMBO J.* 1993;12:1141–1151.
- [63] Kemp JR, Longworth MS. Crossing the LINE toward genomic instability: LINE-1 retrotransposition in cancer. *Front Chem.* 2015;3:68.
- [64] Agodi A, Barchitta M, Quattrocchi A, et al. Low fruit consumption and folate deficiency are associated with LINE-1 hypomethylation in women of a cancer-free population. *Genes Nutr.* 2015;10:30.
- [65] Garcia-Lacarte M, Milagro FI, Zulet MA, et al. LINE-1 methylation levels, a biomarker of weight loss in obese subjects, are influenced by dietary antioxidant capacity. *Redox Rep.* 2016;21:67–74.
- [66] Marques-Rocha JL, Milagro FI, Mansego ML, et al. LINE-1 methylation is positively associated with healthier lifestyle but inversely related to body fat mass in healthy young individuals. *Epigenetics.* 2016;11:49–60.
- [67] Zhang FF, Morabia A, Carroll J, et al. Dietary patterns are associated with levels of global genomic DNA methylation in a cancer-free population. *J Nutr.* 2011;141:1165–1171.
- [68] Schulz WA, Steinhoff C, Florl AR. Methylation of endogenous human retroelements in health and disease. In: Doerfler W, Böhm P, editors. *DNA Methylation Development Genetic Disease, and Cancer.* Berlin, Heidelberg: Springer; 2006. p. 211–250.
- [69] Joyce BT, Gao T, Zheng Y, et al. Prospective changes in global DNA methylation and cancer incidence and mortality. *Br J Cancer.* 2016;115:465–472.
- [70] Baccarelli A, Wright R, Bollati V, et al. Ischemic heart disease and stroke in relation to blood DNA methylation. *Epidemiology.* 2010;21:819–828.
- [71] Bollati V, Schwartz J, Wright R, et al. Decline in genomic DNA methylation through aging in a cohort of elderly subjects. *Mech Ageing Dev.* 2009;130:234–239.
- [72] Jintaridh P, Mutirangura A. Distinctive patterns of age-dependent hypomethylation in interspersed repetitive sequences. *Physiol Genomics.* 2010;41:194–200.
- [73] Lee H-S, Barraza-Villarreal A, Hernandez-Vargas H, et al. Modulation of DNA methylation states and infant immune system by dietary supplementation with -3 PUFA during pregnancy in an intervention study. *Am J Clin Nutr.* 2013;98:480–487.
- [74] Mickleborough TD, Tecklenburg SL, Montgomery GS, et al. Eicosapentaenoic acid is more effective than docosahexaenoic acid in inhibiting proinflammatory mediator production and transcription from LPS-induced human asthmatic alveolar macrophage cells. *Clin Nutr.* 2009;28:71–77.
- [75] Gorjão R, Azevedo-Martins AK, Rodrigues HG, et al. Comparative effects of DHA and EPA on cell function. *Pharmacol Ther.* 2009;122:56–64.
- [76] Turk HF, Chapkin RS. Membrane lipid raft organization is uniquely modified by n-3 polyunsaturated fatty acids. *Prostaglandins Leukot Essent Fat Acids.* 2013;88:43–47.
- [77] Kim W, Fan Y-Y, Barhoumi R, et al. n-3 polyunsaturated fatty acids suppress the localization and activation of signaling proteins at the immunological synapse in murine CD4+ T cells by affecting lipid raft formation. *J Immunol.* 2008;181:6236–6243.
- [78] Chapkin RS, Wang N, Fan Y-Y, et al. Docosahexaenoic acid alters the size and distribution of cell surface microdomains. *Biochim Biophys Acta Biomembr.* 2008;1778:466–471.

- [79] Reseland JE, Haugen F, Hollung K, et al. Reduction of leptin gene expression by dietary polyunsaturated fatty acids. *J Lipid Res.* 2001;42:743–750.
- [80] Shen W, Wang C, Xia L, et al. Epigenetic modification of the leptin promoter in diet-induced obese mice and the effects of N-3 polyunsaturated fatty acids. *Sci Rep.* 2014;4:5282.
- [81] Fan C, Liu X, Shen W, et al. The regulation of leptin, leptin receptor and pro-opiomelanocortin expression by N-3 Pufas in diet-induced obese mice is not related to the methylation of their promoters. *Nutr Metab (Lond).* 2011;8:31.
- [82] Vallvé J-C, Ullaque K, Girona J, et al. Unsaturated fatty acids and their oxidation products stimulate CD36 gene expression in human macrophages. *Atherosclerosis.* 2002;164:45–56.
- [83] Alexander Aguilera A, Hernández Díaz G, Lara Barcelata M, et al. Induction of Cd36 expression elicited by fish oil PUFA in spontaneously hypertensive rats. *J Nutr Biochem.* 2006;17:760–765.
- [84] Do Amaral CL, Milagro FI, Curi R, et al. DNA methylation pattern in overweight women under an energy-restricted diet supplemented with fish oil. *Biomed Res Int.* 2014;2014:1–10.
- [85] Varela-Moreiras G, Ruiz E, Valero T, et al. The Spanish diet: an update. *Nutr Hosp.* 2013;28:13–20.
- [86] Parra D, Ramel A, Bandarra N, et al. A diet rich in long chain omega-3 fatty acids modulates satiety in overweight and obese volunteers during weight loss. *Appetite.* 2008;51:676–680.
- [87] Benatti P, Peluso G, Nicolai R, et al. Polyunsaturated fatty acids: biochemical, nutritional and epigenetic properties. *J Am Coll Nutr.* 2004;23:281–302.
- [88] Nakamura MT, Nara TY. Structure, function, and dietary regulation of $\Delta 6$, $\Delta 5$, and $\Delta 9$ desaturases. *Ann Rev Nutr.* 2004;24:345–376.
- [89] Marquardt A, Stöhr H, White K, et al. cDNA cloning, genomic structure, and chromosomal localization of three members of the human fatty acid desaturase family. *Genomics.* 2000;66:175–183.
- [90] Glaser C, Heinrich J, Koletzko B. Role of FADS1 and FADS2 polymorphisms in polyunsaturated fatty acid metabolism. *Metabolism.* 2010;59:993–999.
- [91] Hoile SP, Clarke-Harris R, Huang R-C, et al. Supplementation with N-3 long-chain polyunsaturated fatty acids or olive oil in men and women with renal disease induces differential changes in the DNA methylation of FADS2 and ELOVL5 in peripheral blood mononuclear cells. Schunck W-H, editor. *PLoS ONE.* 2014;9:e109896.
- [92] Jaudszus A, Gruen M, Watzl B, et al. Evaluation of suppressive and pro-resolving effects of EPA and DHA in human primary monocytes and T-helper cells. *J Lipid Res.* 2013;54:923–935.
- [93] Rovito D, Giordano C, Vizza D, et al. Omega-3 PUFA ethanolamides DHEA and EPEA induce autophagy through PPAR γ activation in MCF-7 breast cancer cells. *J Cell Physiol.* 2013;228:1314–1322.
- [94] Fujiki K, Kano F, Shiota K, et al. Expression of the peroxisome proliferator activated receptor gamma gene is repressed by DNA methylation in visceral adipose tissue of mouse models of diabetes. *BMC Biol.* 2009;7:38.
- [95] Nilsson E, Jansson PA, Perflyev A, et al. Altered DNA methylation and differential expression of genes influencing metabolism and inflammation in adipose tissue from subjects with type 2 diabetes. *Diabetes.* 2014;63:2962–2976.
- [96] Ling C, Del Guerra S, Lupi R, et al. Epigenetic regulation of PPARGC1A in human type 2 diabetic islets and effect on insulin secretion. *Diabetologia.* 2008;51:615–622.
- [97] Brøns C, Jacobsen S, Nilsson E, et al. Deoxyribonucleic acid methylation and gene expression of PPARGC1A in human muscle is influenced by high-fat overfeeding in a birth-weight-dependent manner. *J Clin Endocrinol Metab.* 2010;95:3048–3056.
- [98] Kirchner H, Nylen C, Laber S, et al. Altered promoter methylation of PDK4, IL1 B, IL6, and TNF after Roux-en Y gastric bypass. *Surg Obes Relat Dis.* 2014;10:671–678.
- [99] Epstein FH, Barnes PJ, Karin M. Nuclear factor- κ B – a pivotal transcription factor in chronic inflammatory diseases. *N Engl J Med.* 1997;336:1066–1071.
- [100] Zhao Y, Joshi-Barve S, Barve S, et al. Eicosapentaenoic acid prevents LPS-induced TNF- α expression by preventing NF- κ B activation. *J Am Coll Nutr.* 2004;23:71–78.
- [101] Sullivan KE, Reddy ABM, Dietzmann K, et al. Epigenetic regulation of tumor necrosis factor alpha. *Mol Cell Biol.* 2007;27:5147–5160.
- [102] Kochanek S, Radbruch A, Tesch H, et al. DNA methylation profiles in the human genes for tumor necrosis factors alpha and beta in subpopulations of leukocytes and in leukemias. *Proc Nat Acad Sci USA.* 1991;88:5759–5763.
- [103] Takei S, Fernandez D, Redford A, et al. Methylation status of 5'-regulatory region of tumor necrosis factor α gene correlates with differentiation stages of monocytes. *Biochem Biophys Res Commun.* 1996;220:606–612.
- [104] Nimmo ER, Prendergast JG, Aldhous MC, et al. Genome-wide methylation profiling in Crohn's disease identifies altered epigenetic regulation of key host defense mechanisms including the Th17 pathway. *Inflamm Bowel Dis.* 2012;18:889–899.
- [105] Zhu J, Xia L, Ji C, et al. Differential DNA methylation status between human preadipocytes and mature adipocytes. *Cell Biochem Biophys.* 2012;63:1–15.
- [106] Gazzar ME, Yoza BK, Chen X, et al. G9a and HP1 couple histone and DNA methylation to TNF transcription silencing during endotoxin tolerance. *J Biol Chem.* 2008;283:32198–32208.
- [107] Bollati V, Favero C, Albetti B, et al. Nutrients intake is associated with DNA methylation of candidate inflammatory genes in a population of obese subjects. *Nutrients.* 2014;6:4625–4639.
- [108] Hermsdorff HH, Mansego ML, Campión J, et al. TNF-alpha promoter methylation in peripheral white blood cells: relationship with circulating TNF α , truncal fat and n-6 PUFA intake in young women. *Cytokine.* 2013;64:265–271.
- [109] Kochanek S, Toth M, Dehmel A, et al. Interindividual concordance of methylation profiles in human genes for tumor necrosis factors alpha and beta. *Proc Nat Acad Sci USA.* 1990;87:8830–8834.
- [110] Lou S, Lee H-M, Qin H, et al. Whole-genome bisulfite sequencing of multiple individuals reveals complementary roles of promoter and gene body methylation in transcriptional regulation. *Genome Biol.* 2014;15:408.

- [111] Gross TJ, Kremens K, Powers LS, et al. Epigenetic silencing of the human NOS2 gene: rethinking the role of nitric oxide in human macrophage inflammatory responses. *J Immunol.* [2014](#);192:2326–2338.
- [112] Hashimoto K, Otero M, Imagawa K, et al. Regulated transcription of human matrix metalloproteinase 13 (MMP13) and interleukin-1 (IL1B) genes in chondrocytes depends on methylation of specific proximal promoter CpG sites. *J Biol Chem.* [2013](#);288:10061–10072.
- [113] Berkley AM, Hendricks DW, Simmons KB, et al. Recent thymic emigrants and mature naive T Cells exhibit differential DNA methylation at key cytokine loci. *J Immunol.* [2013](#);190:6180–6186.
- [114] Northrop JK, Thomas RM, Wells AD, et al. Epigenetic remodeling of the IL-2 and IFN-gamma loci in memory CD8 T cells is influenced by CD4 T cells. *J Immunol.* [2006](#);177:1062–1069.
- [115] Williams CL, Schilling MM, Cho SH, et al. STAT4 and T-bet are required for the plasticity of IFN- expression across Th2 ontogeny and influence changes in Ifng promoter DNA methylation. *J Immunol.* [2013](#);191:678–687.
- [116] Caughey GE, Mantzioris E, Gibson RA, et al. The effect on human tumor necrosis factor alpha and interleukin 1 beta production of diets enriched in n-3 fatty acids from vegetable oil or fish oil. *Am J Clin Nutr.* [1996](#);63:116–122.