



**Fear of falling and activities of daily living function:
Mediation effect of dual-task ability**

Journal:	<i>Aging and Mental Health</i>
Manuscript ID	CAMH-2016-0022.R2
Manuscript Type:	Original Article
Keywords:	Dual Task, Fear of Falling, ADL, ageing

SCHOLARONE™
Manuscripts

Running header: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 1

Fear of falling and activities of daily living function: Mediation effect of dual-task ability

Abstract

Objective: The aim of the study was to explore the association between fear of falling (FOF), dual-task ability during a mobility task, and the activities of daily living (ADL) in a sample of older adults.

Methods: Seventy-six older adults (mean age $M = 70.87 \pm 5.16$ years) participated in the study. Data on FOF (using the Falls Self-Efficacy Scale-International), walking ability during both single- and dual-task performances and ADL were collected.

Results: Mediation analysis demonstrated the mediation effect of dual-task ability ($\beta = 0.238, p = 0.011$) between FOF and ADL level ($\beta = 0.559, p < 0.001$). Moreover, significantly lower performances were observed during dual-task condition [$F(2, 73) = 7.386, p < 0.001$], and lower ADL levels were also found in older adults with FOF [$F(2, 73) = 13.734, p < 0.001$].

Conclusion: The study underlines the relationship between FOF, dual-task ability and ADL level. These results could be used to develop specific intervention programmes for successful ageing.

Keywords: Dual Task, Fear of Falling, ADL, ageing

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 2

Introduction

Changes in physical domains, such as reduction in aerobic endurance, lower body muscular strength and decreased mobility (Brustio, Magistro, & Liubicich, 2015; Chodzko-Zajko et al., 2009; Magistro, Candela, Brustio, Liubicich, & Rabaglietti, 2014), and cognitive domains, such as executive function, visual-spatial, verbal and working memory (Park, O'Connell, & Thomson, 2003), observed in advancing ageing are among the major factors that increase the risk of falls (Montero-Odasso, Muir, & Speechley, 2012; Rubenstein, 2006). Approximately one-third of individuals aged 65 years or more experience falls at least once a year (Montero-Odasso et al., 2012) with adverse consequences in both physical and psychological domains. This could lead to a decrease in independence of Activities of Daily Living (ADL), which often require multitasking, and in quality of life (Gillespie et al., 2009; Lord, Sherrington, Menz, & Close, 2007).

Existing literature has referred to Fear Of Falling (FOF) as a person's loss of confidence in his/her balance capabilities, or as a general concept of low confidence of avoiding falls and being afraid of falling (Legters, 2002). According to the different measures and constructs used to assess it, FOF is an important psychological factor that is highly prevalent in independent older adults, too (Scheffer, Schuurmans, van Dijk, van der Hooft, & de Rooij, 2008).

The reviews of Denkinger (2015) and Scheffer (2008) have shown that FOF is associated with an experience of falling; FOF is also commonly found among older adults without a history of falls. Moreover, these reviews reported some risk factors from cross-sectional and longitudinal studies that have been associated with FOF, such as female gender, older age (Kempen, van Haastregt, McKee, Delbaere, & Zijlstra, 2009; Scheffer et al., 2008; Zijlstra, van Haastregt, van Eijk, & Kempen, 2005), physiologic outcomes, including mobility decline (Lach, Ball, & Birge, 2012; Patil, Uusi-Rasi, Kannus, Karinkanta, &

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 3

1
2
3 Sievanen, 2014) and restriction of ADL (Murphy, Williams, & Gill, 2002), increased
4
5 requirement for health care (Zijlstra et al., 2007), increased risk of falling (Friedman, Munoz,
6
7 West, Rubin, & Fried, 2002) and institutionalisation (Cumming, Salkeld, Thomas, & Szonyi,
8
9 2000; Hsu, Alfermann, Lu, & Lin, 2013). Therefore, in some older adults with impaired
10
11 mobility a FOF may cause specific decrease in mobility and ADL (Viljanen et al., 2012).
12
13 Indeed, one of the major consequences of fear of falling is the restriction and avoidance of
14
15 activities (Cumming et al., 2000; Delbaere, Crombez, Vanderstraeten, Willems, & Cambier,
16
17 2004). Many older adults with FOF restrict the range of their activities in an attempt to avoid
18
19 fall-related morbidity and possible social embarrassment. While in the short term the
20
21 reduction or avoidance of activities may protect against falls, in the long term activity
22
23 restriction (leading to decrease in mobility and ADL) may contribute to the reduction of
24
25 physical and psychological health of older adults (Choi & Ko, 2015; Delbaere et al., 2004)
26
27 and eventually result in loss of independence, falling accidents and increased FOF (Delbaere
28
29 et al., 2004; Friedman et al., 2002).
30
31
32
33

34 Recently, different studies have used dual-task paradigms to investigate the
35
36 motor/cognitive interference during simple tasks of ADL and its associations with FOF.
37
38 Previous studies on the dual-task paradigm have showed that mobility tasks are complex
39
40 tasks that require a constant recruitment of motor, sensory and cognitive resources in relation
41
42 to different everyday environments (Al-Yahya et al., 2011; Yogev-Seligmann, Hausdorff, &
43
44 Giladi, 2008). Specifically, these studies have showed that walking tasks with a secondary
45
46 attention-demanding task (e.g. cognitive tasks) may be difficult for older adults (Hall, Echt,
47
48 Wolf, & Rogers, 2011; Hausdorff, Schweiger, Herman, Yogev-Seligmann, & Giladi, 2008;
49
50 Hollman, Kovash, Kubik, & Linbo, 2007) due to the additional attention and executive
51
52 function required to perform the different tasks (Al-Yahya et al., 2011; Yogev-Seligmann et
53
54 al., 2008).
55
56
57
58
59
60

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 4

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Some studies have underlined that dual-task activities during walking may be particularly challenging for older adults with FOF (Asai, Misu, Doi, Yamada, & Ando, 2014; Donoghue, Cronin, Savva, O'Regan, & Kenny, 2013; Uemura, Yamada, et al., 2012) or those concerned or anxious about balance (Gage, Sleik, Polych, McKenzie, & Brown, 2003). FOF may increase the cognitive resources involved during locomotion. Consequently, older adults with FOF usually have lesser cognitive resources available for the control of gait and balance compared with older adults without FOF (Gage et al., 2003). Indeed, older adults with FOF under dual-task conditions have a slower walking speed and shorter stride length with lesser variability in these parameters and increased double-support phase than those seen in older adults without FOF (Donoghue et al., 2013).

FOF may be considered a potential health problem comparable to a fall due to the adverse consequences on ADL function, leading to an increased requirement for health care and its related costs (Zijlstra et al., 2007). Moreover, FOF may have important implications for the successful performance of dual-task activities, which have been associated with an increased risk of falling (Muir-Hunter & Wittwer, 2015). Taken together, the above findings suggest that FOF is a complex problem in ageing people, with negative outcomes for both their independence in ADL and their management of dual-task activities. ADLs too may be considered as complex tasks, as they require people to perform more activities at the same time. During these multi task activities, executive functions (referring to the cognitive ability involved in the planning and execution of goal-directed behaviours, attention, abstract reasoning, and judgment (Lezak, 2004)) are highly involved. Indeed, in older adults the decline in executive functions has been associated with ADL difficulties (Johnson, Lui, & Yaffe, 2007) and linked with higher FOF (Uemura, Shimada, et al., 2012). Consequently, the ability to divide attention between two or more simultaneous tasks is an important aspect of functional movement during ADL (Shin & An, 2014). For example, in everyday life, people

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 5

1
2
3 may walk and talk at the same time while carrying an object or moving from one location to
4
5 another and simultaneously monitoring the environment. Thus, dual-task ability may be
6
7 representative of actual daily situations, due to the ecological need in the performance of
8
9 ADL (Guedes et al., 2014). According to this interpretation, we believe that in these
10
11 situations, the ability to manage different tasks and, in particular, dual-task ability is essential
12
13 not to decrease the performance in the ADL. In fact, dual-task activities depend on executive
14
15 function (Chu, Tang, Peng, & Chen, 2013), which includes the individual's ability to carry
16
17 out and successfully perform and manage ADL (Yogev-Seligmann et al., 2008). We
18
19 hypothesised that dual-task activity may play a role in the relationship between FOF and
20
21 ADL. However, to our knowledge, no studies have evaluated the relationships between FOF,
22
23 dual-task performance and ADL limitations. To address this question, this study explored the
24
25 associations between FOF, dual-task performance of a mobility task and ADL. Our
26
27 hypothesis is that older adults with a high level of concern of falling have poorer mobility
28
29 performance and are more likely to report ADL difficulties than those with lower levels of
30
31 concern. Moreover, a poor performance in dual-task condition mediates this effect.

Method

Participants

Originally, 89 older adults agreed to participate in the present study. The participants were recruited in Northern Italy. The inclusion criteria were as follows: age between 60 and 80 years; ability to walk independently; no self-reported neurological (e.g., Alzheimer's disease, Parkinson's disease) or musculoskeletal (e.g., orthopaedic) conditions affecting mobility or balance; having a mini-mental state examination score (Folstein, Folstein, & McHugh, 1975) higher than 24 and being able to understand the instructions and perform simple arithmetic exercises. Thirteen subjects were excluded because they did not meet the inclusion criteria. Specifically, 5 participants did not meet the age cut-off, 2 used a walking

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 6

aid, 3 reported neurological/musculoskeletal diseases, and 3 had a mini-mental state examination score lower than 24.

Seventy-six older adults (mean age $M = 70.87$ years, $SD \pm 5.16$ years; 51 females) were enrolled in this study. The Ethical Committee of the University of Torino approved the study procedures, and all participants were informed that participation in the study was voluntary and confidential, and did not receive any incentive to participate. All the selected individuals gave their written informed consent before data collection in accordance with Italian law.

Measures

The participants attended a single data collection session. Sociodemographic data, including age, gender, number of years of education, family status, and history of falls, were recorded using a self-report questionnaire. Family status was categorized dividing the sample as living with another person (e.g. married/living as married) or alone (single/separated/divorced). Participants who fell in the previous 12 months were categorized as having a history of falling. Table 1 shows the details of all sociodemographic data of all participants.

FOF was assessed using the Falls Efficacy Scale (FES-International; Delbaere et al., 2010; Yardley et al., 2005), which is a self-report questionnaire containing 16 items about the level of the individual's concern regarding falling in different ADL (Delbaere et al., 2010; Yardley et al., 2005). Each item is scored on a four-point Likert scale (1 = 'not at all concerned'–4 = 'very concerned'), with a total score ranging from 16 to 64. Higher scores indicate a higher concern of falling. The test–retest reliability and intra-class coefficient of this scale have been estimated to be 0.96 and 0.96, respectively (Yardley et al., 2005). Using the cut-off points of the FES-International, the sample was divided into three groups: "no-concern" (16-19), "moderate-concern" (20-27) and "concern" (28-64) (Delbaere et al., 2010).

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 7

1
2
3 A 10m walking test (single task) was used to assess the mobility of the subjects. The
4 subjects were instructed to walk at their self-selected comfortable pace on a 10m long
5 walkway without assistance. The performance time was measured as the time taken to walk
6 the middle 6m (Montero-Odasso et al., 2009), from when the subjects crossed the 2m mark to
7 when the same crossed the 8m mark (Steffen, Hacker, & Mollinger, 2002). We decided to use
8 the middle 6m to give the subjects time to accelerate and decelerate (Asai et al., 2014;
9 Montero-Odasso et al., 2009). For the dual-task test, the participants performed the 10m
10 walking test as above and were instructed to simultaneously count backwards in increments
11 of three from a randomised number from 80 to 99 (dual task). No instructions were given
12 regarding which task to prioritise during the dual-task performance. The same evaluator
13 tested all the participants. A stopwatch was used for the data collection with the time
14 recorded in seconds, which was taken as the task score.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30 The Groningen Activity Restriction Scale (GARS) (Kempen, Miedema, Ormel, &
31 Molenaar, 1996) was used to measure the level of disability (physical impairment) in ADL
32 and the Instrumental Activities of Daily Living (IADL). The GARS consists of 18 items on
33 daily activities. Specifically, the scale contains 11 items referring to ADL and personal care
34 and 7 items to IADL. Each item was scored on a four-point Likert scale (1 = "I cannot do it
35 without someone else's help"-4 = "yes, I can do it fully and independently without any
36 difficulty"), with a total score ranging from 18 to 72. Lower scores indicate a higher level of
37 difficulty experienced by a person in taking care of themselves and the performance of
38 household activities. The internal estimated reliability was 0.98 and 0.92 for older men and
39 women, respectively (Kempen et al., 1996).

50 51 52 **Statistical analyses**

53
54 The sociodemographic characteristics and mobility performance were tabulated for
55 the entire sample and presented as the mean (standard deviation) or frequency (percentage).
56
57
58
59
60

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 8

1
2
3 Using age, gender, education years and history of falls as covariates, a 3×2
4
5 ANCOVA with the group (no-concern; moderate concern; concern) as the between-subject
6
7 factor and walking conditions (single-task test; dual-task test) as the within-subject factor was
8
9 performed to evaluate the difference of FOF on the walking conditions. The different levels
10
11 of FOF were determined by significant group \times condition interactions.

14 Using age, gender, education years and history of fall as covariates, an ANCOVA was
15
16 conducted to determine whether the ADL ability (GARS score) was different for groups with
17
18 different levels of FOF (no-fear; moderate-fear; fear). A post hoc analysis with a Bonferroni
19
20 adjustment was computed to identify statistically significant age comparisons.

23 To quantify the dual-task ability of the subjects, the Dual-Task Cost (DTC) was
24
25 calculated as the difference between the scores for the single-task and dual-task performances
26
27 (Hausdorff et al., 2008). A positive value of DTC indicates better performance during the
28
29 dual-task activity. Conversely, a negative value indicates a slower performance during the
30
31 dual-task activity. To test the relationship between FOF and ADL through DTC, a mediation
32
33 analysis was performed according to the approach described by Baron and Kenny (1986).
34
35 First, the direct effects of the FOF scores on the GARS scores were determined using
36
37 regression analyses. Second, if the relationship was significant, the mediator was included in
38
39 the model and the main effect between the independent variable (FOF scores) and the
40
41 mediator (DTC) was verified. Third, the mediating effect of mobility on the relationship
42
43 between the independent variable (FOF scores) and the outcome (GARS scores) was
44
45 checked. Finally, the Sobel test was used to verify the mediation model.

49 The Statistical Package for Social Sciences (SPSS 20.0 for Windows; IBM Corp.,
50
51 Armonk, NY, USA) was used for all statistical analyses. The statistical significance level was
52
53 set at $p < 0.05$.
54
55
56
57
58
59
60

Results

There were 24 participants (31.6%) in the no-concern group, 37 (48.7%) in the moderate-concern group, and 15 (19.7%) in the concern group (see Table 1 for more details on the sociodemographic data of all participants and of the different groups).

Figure 1 shows the mean scores and standard deviations for the no-concern, moderate-concern, and concern groups, from the single- and dual-task tests. The ANCOVA analysis showed significant differences for the group [$F(2, 67) = 7.358, p < 0.01, \eta^2 = 0.188$], condition [$F(1, 63) = 4.061, p < 0.05, \eta^2 = 0.057$] and for the interaction group \times condition [$F(2, 73) = 7.358, p < 0.01, \eta^2 = 0.180$]. Focusing on the main effect of the group, the post hoc analysis with Bonferroni adjustment revealed significant differences between the no-concern and concern groups ($p < 0.001$) and between the moderate-concern and concern groups ($p < 0.001$), but not between the no-concern and moderate-concern groups ($p > 0.05$). Considering the significance observed for condition, the walking performance was better in the single-task test than in the dual-task test in the concern group [from 5.47, 95% CI (4.61–6.32) to 7.98, 95% CI (6.38–9.59) seconds; $F(1, 10) = 7.399, p < 0.05, \text{partial } \eta^2 = 0.425$]. In contrast, walking performance was not statistically significantly different between the single-task test and the dual-task test both in no-concern [from 4.47, 95% CI (4.19–4.76) to 5.52, 95% CI (5.00–6.05) seconds; $F(1, 18) = 1.730, p > 0.05, \text{partial } \eta^2 = 0.088$] and moderate-concern group [from 4.71, 95% CI (4.43–4.99) to 5.78, 95% CI (5.31–6.25) seconds; $F(1, 31) = 1.499, p > 0.05, \text{partial } \eta^2 = 0.046$]. According to the significant interaction of group \times condition, the effect of the dual-task performance was not uniform among the groups. Indeed, we noticed a larger increase in dual-task performance in the concern group than that in the moderate-concern and no-concern groups.

ANCOVA analysis of GARS score yielded a statistical significance for the different levels of FOF [$F(2, 67) = 10.570, p < 0.001$]. The GARS scores increased from the no-

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 10

concern group (18.95 ± 1.51), to the moderate-concern group (20.38 ± 2.44), to the concern group (24.0 ± 5.12). Bonferroni post hoc analysis revealed that the increase from the no-concern to concern groups [4.6 , 95% CI ($2.02-7.17$)] was statistically significant ($p < 0.001$), as well as the increase from the moderate-concern to concern groups [3.51 , 95% CI ($1.27-5.75$)], $p < 0.001$] but not the increase from the no-concern to moderate-concern groups [1.08 , 95% CI (-0.99 to 3.31)], $p > 0.05$].

Figure 2 shows the results of the mediation model. The main effect of the predictor (FOF scores) on the outcome (GARS scores) was statistically significant ($\beta = 0.640$, $p < 0.001$), and the mediator had a positive effect on the GARS score ($\beta = 0.427$, $p < 0.001$). After introducing the mediator (DTC), we noticed a decrease in the relationship between the predictor and the outcome ($\beta = 0.559$, $p < 0.001$), as well as a significant decrease of the coefficient between the mediator and the outcome ($\beta = 0.238$, $p = 0.011$). The Sobel test indicated that the mediation model was partially mediated ($z = 1.98$, $p < 0.05$).

Discussion

The aim of the study was to explore the cross-sectional associations among FOF, DT performance and ADL in older adults.

We observed a different level of performance between the single- and dual-task tests among the groups. Specifically, we observed that, on average, the walking performance was lower when simultaneously counting backwards (dual-task test) compared with that when walking only (single-task test). The above findings underline that an additional cognitive task may decrease walking ability in older adults (Brustio, Magistro, Rabaglietti, & Liubicich, 2015; Hall et al., 2011; Hollman et al., 2007) due to the involvement of cognitive function during walking (Al-Yahya et al., 2011). Interestingly, the decrease in walking performance under dual-task conditions was more pronounced in the concern group compared with that in the no-concern and moderate-concern groups. These results supported the idea of a negative

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 11

1
2
3 association between FOF and dual-task performance (Asai et al., 2014; Donoghue et al.,
4
5 2013). The decrease in dual-task performance, observed in older adults with concern about
6
7 falling, may indicate a difficulty managing two different required tasks, one being the
8
9 performance of the walking test and the other being the cognitive task. It is possible that older
10
11 people with concern of falling adopted a more cautious walking pattern to avoid fall
12
13 (Donoghue et al., 2013) due to the reduction of the attentional resources available to
14
15 successfully perform two or more different tasks (Gage et al., 2003). Additionally, to the best
16
17 of our knowledge our findings expand current literature providing clear evidence of the role
18
19 of FOF and dual-task performance with regards to the ADL. A strength of our study is that
20
21 FOF was not assessed by a simple question about FOF, as in previous studies (Asai et al.,
22
23 2014; Donoghue et al., 2013; Uemura, Yamada, et al., 2012) but by the FES-I that is able to
24
25 discriminate between different levels of "concerns" about falling.
26
27

28
29
30 Our data were in agreement with the existing literature (Choi & Ko, 2015; Cumming
31
32 et al., 2000; Lawson & Katherine, 2014; Murphy et al., 2002) and showed that the concern
33
34 group has a greater impairment in ADL compared with the other groups. This decline in
35
36 ADLs ability may be related to a possible strategy by older adults with FOF to avoid certain
37
38 activities adopting a more inactive lifestyle. At the same time, as people age, FOF may lead
39
40 to functional decline and, consequently, to a restriction in ADL (Friedman et al., 2002;
41
42 Mendes da Costa et al., 2012; Yogev-Seligmann et al., 2008), which has important
43
44 implications for successful ageing and negative consequences for quality of life.
45
46

47
48 Finally, the mediation model on the associations among FOF, ADL and DTC showed
49
50 that DTC plays a mediation role in the relationship between FOF and ADL. In fact, we found
51
52 that FOF had both direct and indirect effects on ADL through the mediation of DTC. On the
53
54 other hand, our data showed that older adults with a low concern about falling also had a
55
56 lower DTC, with less difficulty with ADL's, a significant index of disability and frailty in
57
58
59
60

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 12

1
2
3 ageing people (Candela, Zucchetti, & Magistro, 2013; Candela, Zucchetti, Ortega,
4
5 Rabaglietti, & Magistro, 2015). Our model could help preventing the drop in ADL typically
6
7 observed in older age. Indeed, previous studies (Friedman et al., 2002; Mendes da Costa et
8
9 al., 2012; Murphy et al., 2002) have reported that restriction in ADL affects about 46%–56%
10
11 older adults with FOF, underlining the necessity of intervention programmes focused on
12
13 decreasing FOF and increasing ADL ability. Physical intervention programmes may decrease
14
15 the FOF (Brustio, Magistro, Ivaldi, et al., 2015; Huang, Chung, Chen, Chin, & Wang, 2016;
16
17 Sjosten, Vaapio, & Kivela, 2008; Zijlstra et al., 2005) and decrease the DTC (Agmon, Belza,
18
19 Nguyen, Logsdon, & Kelly, 2014); thus, we hypothesised, based on our results, that
20
21 interventions focused on decreasing FOF (e.g. physical or cognitive interventions) and DTC
22
23 may have positive effects by maintaining or improving ADL functionality, which is linked
24
25 with successful ageing.
26
27

28
29
30 Despite of these findings, our study presents some limitations. The sample size of the
31
32 study did not allow us to extend our conclusions to the general older population. Indeed, the
33
34 older adults involved in the present study were relatively young (age range: 60–80 years) and
35
36 did not have cognitive or physical impairments. Additionally, our secondary demanding task
37
38 included only the performance of a specific cognitive task (i.e. counting backwards in
39
40 decrements of three). Future studies are needed to confirm the validity of our model in a
41
42 longitudinal way and using a different secondary task (e.g. manual tasks or decision-making
43
44 tasks), common in everyday life, to better understand the role of dual-task conditions in the
45
46 relationship between FOF and ADL ability. Finally, we only focused on the mobility
47
48 performance of the subjects under dual-task conditions and not on the cognitive performance.
49
50 Future studies need to investigate both mobility and cognitive performance to better describe
51
52 the physical and cognitive impairments during dual-task performance among older adults
53
54 with FOF.
55
56
57
58
59
60

1
2
3 In summary, our study highlighted the role of dual-task ability in the relationship
4
5 between FOF and ADL ability during the ageing process. In agreement with the results from
6
7 previous studies, we found a lower performance under dual-task conditions and more ADL
8
9 limitations in older adults with FOF compared with those with no concern or moderate
10
11 concern. During the ageing process, FOF has a complex role in everyday activities and has
12
13 important implications in the successful performance of dual-task activities and ADL.
14
15 Identification of the relationships among FOF, dual-task activity, and ADL may provide
16
17 information that could be used to improve the possibility of a successful, better ageing. A
18
19 clarification of the relationships among these variables may be useful in the implementation
20
21 of specific prevention strategies for preventing FOF and the decline in ADL ability in older
22
23 adults.
24
25
26

27 **Conflicts of interest**

28
29 The authors declare that there are no conflicts of interest.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 14

References

- Agmon, M., Belza, B., Nguyen, H. Q., Logsdon, R. G., & Kelly, V. E. (2014). A systematic review of interventions conducted in clinical or community settings to improve dual-task postural control in older adults. *Cinical Interventions in Aging*, *9*, 477-492.
- Al-Yahya, E., Dawes, H., Smith, L., Dennis, A., Howells, K., & Cockburn, J. (2011). Cognitive motor interference while walking: a systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews*, *35*(3), 715-728.
- Asai, T., Misu, S., Doi, T., Yamada, M., & Ando, H. (2014). Effects of dual-tasking on control of trunk movement during gait: respective effect of manual- and cognitive-task. *Gait & Posture*, *39*(1), 54-59.
- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, *51*(6), 1173-1182.
- Brustio, P. R., Magistro, D., Ivaldi, S., Caglio, M. M., Rabaglietti, E., & Liubicich, M. E. (2015). Neuromotor training in older women living in long-term care setting: A pilot study. *Geriatric Nursing*, *36*(5), 361-366.
- Brustio, P. R., Magistro, D., & Liubicich, M. E. (2015). Changes in temporal parameters during performance of the Step Test in older adults. *Gait Posture*, *41*(1), 217-221.
- Brustio, P. R., Magistro, D., Rabaglietti, E., & Liubicich, M. E. (2015). Age-related differences in dual task performance: a cross sectional study on women *Geriatrics & Gerontology International*, doi: 10.1111/ggi.12700.
- Candela, F., Zucchetti, G., & Magistro, D. (2013). Individual Correlates of Autonomy in Activities of Daily Living of Institutionalized Elderly Individuals: An Exploratory Study in a Holistic Perspective. *Holistic Nursing Practice*, *27*(5), 284-291.

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 15

- 1
2
3 Candela, F., Zucchetti, G., Ortega, E., Rabaglietti, E., & Magistro, D. (2015). Preventing
4
5 Loss of Basic Activities of Daily Living and Instrumental Activities of Daily Living
6
7 in Elderly: Identification of Individual Risk Factors in a Holistic Perspective. *Journal*
8
9 *of Holistic Nursing, 29*(5), 313-322.
10
11 Chodzko-Zajko, W. J., Proctor, D. N., Fiatarone Singh, M. A., Minson, C. T., Nigg, C. R.,
12
13 Salem, G. J., & Skinner, J. S. (2009). American College of Sports Medicine position
14
15 stand. Exercise and physical activity for older adults. *Medicine & Science in Sports &*
16
17 *Exercise, 41*(7), 1510-1530.
18
19 Choi, K., & Ko, Y. (2015). Characteristics associated with fear of falling and activity
20
21 restriction in South Korean older adults. *Journal of Aging and Health,*
22
23 0898264315573519.
24
25
26
27 Chu, Y. H., Tang, P. F., Peng, Y. C., & Chen, H. Y. (2013). Meta-analysis of type and
28
29 complexity of a secondary task during walking on the prediction of elderly falls.
30
31 *Geriatrics & Gerontology International, 13*(2), 289-297.
32
33
34 Cumming, R. G., Salkeld, G., Thomas, M., & Szonyi, G. (2000). Prospective study of the
35
36 impact of fear of falling on activities of daily living, SF-36 scores, and nursing home
37
38 admission. *Journals of Gerontology Series A-Biological Sciences & Medical*
39
40 *Sciences, 55*(5), M299-305.
41
42
43 Delbaere, K., Close, J. C., Mikolaizak, A. S., Sachdev, P. S., Brodaty, H., & Lord, S. R.
44
45 (2010). The Falls Efficacy Scale International (FES-I). A comprehensive longitudinal
46
47 validation study. *Age and Ageing, 39*(2), 210-216.
48
49
50 Delbaere, K., Crombez, G., Vanderstraeten, G., Willems, T., & Cambier, D. (2004). Fear-
51
52 related avoidance of activities, falls and physical frailty. A prospective community-
53
54 based cohort study. *Age and Ageing, 33*(4), 368-373.
55
56
57
58
59
60

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 16

Denkinger, M. D., Lukas, A., Nikolaus, T., & Hauer, K. (2015). Factors associated with fear of falling and associated activity restriction in community-dwelling older adults: a systematic review. *Am J Geriatr Psychiatry, 23*(1), 72-86.

Donoghue, O. A., Cronin, H., Savva, G. M., O'Regan, C., & Kenny, R. A. (2013). Effects of fear of falling and activity restriction on normal and dual task walking in community dwelling older adults. *Gait & Posture, 38*(1), 120-124.

Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research, 12*(3), 189-198.

Friedman, S. M., Munoz, B., West, S. K., Rubin, G. S., & Fried, L. P. (2002). Falls and fear of falling: which comes first? A longitudinal prediction model suggests strategies for primary and secondary prevention. *Journal of the American Geriatrics Society, 50*(8), 1329-1335.

Gage, W. H., Sleik, R. J., Polych, M. A., McKenzie, N. C., & Brown, L. A. (2003). The allocation of attention during locomotion is altered by anxiety. *Experimental Brain Research, 150*(3), 385-394.

Gillespie, L. D., Robertson, M. C., Gillespie, W. J., Lamb, S. E., Gates, S., Cumming, R. G., & Rowe, B. H. (2009). Interventions for preventing falls in older people living in the community. *Cochrane Database of Systematic Reviews, 2*(CD007146).

Guedes, R. C., Dias, R. C., Pereira, L. S., Silva, S. L., Lustosa, L. P., & Dias, J. M. (2014). Influence of dual task and frailty on gait parameters of older community-dwelling individuals. *Brazilian Journal of Physical Therapy, 18*(5), 445-452.

Hall, C. D., Echt, K. V., Wolf, S. L., & Rogers, W. A. (2011). Cognitive and motor mechanisms underlying older adults' ability to divide attention while walking. *Physical Therapy, 91*(7), 1039-1050.

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 17

Hausdorff, J. M., Schweiger, A., Herman, T., Yogev-Seligmann, G., & Giladi, N. (2008).

Dual Task Decrements in Gait among Healthy Older Adults: Contributing Factors.

Journals of Gerontology Series A-Biological Sciences & Medical Sciences, 63(12),

1335-1343.

Hollman, J. H., Kovash, F. M., Kubik, J. J., & Linbo, R. A. (2007). Age-related differences in

spatiotemporal markers of gait stability during dual task walking. *Gait & Posture*,

26(1), 113-119.

Hsu, Y., Alfermann, D., Lu, F. J., & Lin, L. L. (2013). Pathways from fear of falling to

quality of life: the mediating effect of the self-concept of health and physical

independence. *Aging & Mental Health*, 17(7), 816-822.

Huang, T. T., Chung, M. L., Chen, F. R., Chin, Y. F., & Wang, B. H. (2016). Evaluation of a

combined cognitive-behavioural and exercise intervention to manage fear of falling

among elderly residents in nursing homes. *Aging & Mental Health*, 20(1), 2-12.

Johnson, J. K., Lui, L. Y., & Yaffe, K. (2007). Executive function, more than global

cognition, predicts functional decline and mortality in elderly women. *J Gerontol A*

Biol Sci Med Sci, 62(10), 1134-1141.

Kempen, G. I., Miedema, I., Ormel, J., & Molenaar, W. (1996). The assessment of disability

with the Groningen Activity Restriction Scale. Conceptual framework and

psychometric properties. *Social Science & Medicine*, 43(11), 1601-1610.

Kempen, G. I., van Haastregt, J. C., McKee, K. J., Delbaere, K., & Zijlstra, G. A. (2009).

Socio-demographic, health-related and psychosocial correlates of fear of falling and

avoidance of activity in community-living older persons who avoid activity due to

fear of falling. *BMC Public Health*, 9, 170.

Lach, H. W., Ball, L. J., & Birge, S. J. (2012). The Nursing Home Falls Self-Efficacy Scale:

development and testing. *Clinical Nursing Research*, 21(1), 79-91.

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 18

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Lawson, O., & Katherine, A. (2014). The Impact of Fear of Falling on Functional Independence Among Older Adults Receiving Home Health Services. *The Open Journal of Occupational Therapy*, 2(3), 4.
- Legters, K. (2002). Fear of falling. *Phys Ther*, 82(3), 264-272.
- Lezak, M. D. (2004). *Neuropsychological Assessment*: Oxford University Press.
- Lord, S. R., Sherrington, C., Menz, H. B., & Close, J. C. (2007). *Falls in older people: risk factors and strategies for prevention*: Cambridge University Press.
- Magistro, D., Candela, F., Brustio, P. R., Liubicich, M. E., & Rabaglietti, E. (2014). A Longitudinal Study on the Relationship Between Aerobic Endurance and Lower Body Strength in Italian Sedentary Older Adults. *Journal of Aging and Physical Activity*, 23(3), 444-451.
- Mendes da Costa, E., Pepersack, T., Godin, I., Bantuelle, M., Petit, B., & Leveque, A. (2012). Fear of falling and associated activity restriction in older people. results of a cross-sectional study conducted in a Belgian town. *Archives of Public Health*, 70(1), 1.
- Montero-Odasso, M., Bergman, H., Phillips, N. A., Wong, C. H., Sourial, N., & Chertkow, H. (2009). Dual-tasking and gait in people with mild cognitive impairment. The effect of working memory. *BMC Geriatrics*, 9, 41.
- Montero-Odasso, M., Muir, S. W., & Speechley, M. (2012). Dual-task complexity affects gait in people with mild cognitive impairment: the interplay between gait variability, dual tasking, and risk of falls. *Archives of Physical Medicine and Rehabilitation*, 93(2), 293-299.
- Muir-Hunter, S. W., & Wittwer, J. E. (2015). Dual-task testing to predict falls in community-dwelling older adults: a systematic review. *Physiotherapy*.

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 19

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Murphy, S. L., Williams, C. S., & Gill, T. M. (2002). Characteristics associated with fear of falling and activity restriction in community-living older persons. *Journal of the American Geriatrics Society, 50*(3), 516-520.
- Park, H. L., O'Connell, J. E., & Thomson, R. G. (2003). A systematic review of cognitive decline in the general elderly population. *International Journal of Geriatric Psychiatry, 18*(12), 1121-1134.
- Patil, R., Uusi-Rasi, K., Kannus, P., Karinkanta, S., & Sievanen, H. (2014). Concern about Falling in Older Women with a History of Falls: Associations with Health, Functional Ability, Physical Activity and Quality of Life. *Gerontology, 60*(1), 22-30.
- Rubenstein, L. Z. (2006). Falls in older people: epidemiology, risk factors and strategies for prevention. *Age and Ageing, 35 Suppl 2*, ii37-ii41.
- Scheffer, A. C., Schuurmans, M. J., van Dijk, N., van der Hoof, T., & de Rooij, S. E. (2008). Fear of falling: measurement strategy, prevalence, risk factors and consequences among older persons. *Age and Ageing, 37*(1), 19-24.
- Shin, S. S., & An, D. H. (2014). The Effect of Motor Dual-task Balance Training on Balance and Gait of Elderly Women. *Journal of Physical Therapy Science, 26*(3), 359-361.
- Sjosten, N., Vaapio, S., & Kivela, S. L. (2008). The effects of fall prevention trials on depressive symptoms and fear of falling among the aged: a systematic review. *Aging & Mental Health, 12*(1), 30-46.
- Steffen, T. M., Hacker, T. A., & Mollinger, L. (2002). Age- and gender-related test performance in community-dwelling elderly people: Six-Minute Walk Test, Berg Balance Scale, Timed Up & Go Test, and gait speeds. *Physical Therapy, 82*(2), 128-137.

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 20

1
2
3 Uemura, K., Shimada, H., Makizako, H., Yoshida, D., Doi, T., Tsutsumimoto, K., & Suzuki,
4
5 T. (2012). A lower prevalence of self-reported fear of falling is associated with
6
7 memory decline among older adults. *Gerontology*, *58*(5), 413-418.

9
10 Uemura, K., Yamada, M., Nagai, K., Tanaka, B., Mori, S., & Ichihashi, N. (2012). Fear of
11
12 falling is associated with prolonged anticipatory postural adjustment during gait
13
14 initiation under dual-task conditions in older adults. *Gait & Posture*, *35*(2), 282-286.

15
16 Viljanen, A., Kulmala, J., Rantakokko, M., Koskenvuo, M., Kaprio, J., & Rantanen, T.
17
18 (2012). Fear of falling and coexisting sensory difficulties as predictors of mobility
19
20 decline in older women. *Journals of Gerontology Series A-Biological Sciences &*
21
22 *Medical Sciences*, *67*(11), 1230-1237.

23
24 Yardley, L., Beyer, N., Hauer, K., Kempen, G., Piot-Ziegler, C., & Todd, C. (2005).
25
26 Development and initial validation of the Falls Efficacy Scale-International (FES-I).
27
28 *Age and Ageing*, *34*(6), 614-619.

29
30
31
32 Yogev-Seligmann, G., Hausdorff, J. M., & Giladi, N. (2008). The role of executive function
33
34 and attention in gait. *Movement Disorders*, *23*(3), 329-342; quiz 472.

35
36 Zijlstra, G. A., van Haastregt, J. C., van Eijk, J. T., & Kempen, G. I. (2005). Evaluating an
37
38 intervention to reduce fear of falling and associated activity restriction in elderly
39
40 persons: design of a randomised controlled trial [ISRCTN43792817]. *BMC Public*
41
42 *Health*, *5*, 26.

43
44
45 Zijlstra, G. A., van Haastregt, J. C., van Eijk, J. T., van Rossum, E., Stalenhoef, P. A., &
46
47 Kempen, G. I. (2007). Prevalence and correlates of fear of falling, and associated
48
49 avoidance of activity in the general population of community-living older people. *Age*
50
51 *and Ageing*, *36*(3), 304-309.

Table 1. Sociodemographic characteristics of the study participants.

	All (<i>n</i> = 76)	Fear of falling group		
		No-Concern (<i>n</i> = 24)	Moderate- Concern (<i>n</i> = 37)	Concern (<i>n</i> = 15)
Age, years	70.87 (5.16)	68.7 (0.65)	71.88 (5.11)	71.82 (5.29)
Gender				
Male	25 (32.9)	13 (54.2)	9 (24.3)	3 (20.0)
Female	51 (67.1)	11 (45.8)	28 (75.7)	12 (80.0)
Years of education	8.51 (3.57)	10.87 (3.47)	8.16 (3.67)	6.93 (2.63)
History of falling				
Yes	44 (62.0)	15 (68.2)	21 (61.8)	8 (53.3)
No	27 (38.0)	7 (31.8)	13 (38.2)	7 (46.7)
Family status				
Married/living as married	55 (72.4)	22 (91.7)	26 (70.3)	7 (46.7)
Single/separated/divorced	1 (1.3)	0 (0.0)	1 (2.7)	0 (0.0)
Widow	20 (26.3)	2 (8.3)	10 (27.0)	7 (53.3)
Single task, s	4.79 (1.03)	4.50 (0.66)	4.83 (0.86)	5.47 (1.57)
Dual task, s	6.13 (2.04)	5.59 (1.16)	5.74 (1.49)	7.98 (3.12)
GARS score	20.64 (3.41)	18.95 (1.51)	20.38 (2.44)	24.0 (5.12)

Note: Data are presented as the mean (SD) for continuous variables (e.g. age and years of education) and frequency (%) for categorical variables. Single-task indicates the 10-m walking test; dual-task indicates the 10-m walking test while counting backwards in increments of three; GARS score indicates the score on the Groningen Activity Restriction Scale.

Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 22


Figure captions

Figure 1: Results of the 10-m walking test for both the single- and dual-task performances in the no-concern (black solid), moderate-concern (dashed solid) and concern groups (white solid).

Figure 2: Mediation model among GARS score (outcome), FOF score (predictor) and DTC (mediator).


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

Figure 1



Running head: FEAR OF FALLING ADL AND MEDIATION ROLE OF DUAL-TASK 1

Figure 2



Notes: Data are collapsed across the participants. FOF score indicates the score on the Falls Efficacy Scale; DTC indicates the dual-task cost; GARS score indicates the score on the Groningen Activity Restriction Scale. Sobel test: $z = 1.98$; $*p < 0.05$ and $**p < 0.01$.