



# Engineering dissipative channels for realizing Schrödinger cats in SQUIDs

Mark J. Everitt<sup>1\*</sup>, Timothy P. Spiller<sup>2</sup>, Gerard J. Milburn<sup>3</sup>, Richard D. Wilson<sup>1†</sup> and Alexandre M. Zagoskin<sup>1</sup>

<sup>1</sup> Quantum Systems Engineering Research Group and Department of Physics, Loughborough University, Loughborough, UK

<sup>2</sup> Quantum Information Science, School of Physics and Astronomy, University of Leeds, Leeds, UK

<sup>3</sup> Centre for Engineered Quantum Systems, School of Mathematics and Physics, The University of Queensland, St Lucia, QLD, Australia

## Edited by:

Andrew D. Greentree, RMIT  
University, Australia

## Reviewed by:

Jonas Maziero, Universidade Federal  
de Santa Maria, Brazil  
Zoltán Zimborás, University College  
London, UK

## \*Correspondence:

Mark J. Everitt, Quantum Systems  
Engineering Research Group and  
Department of Physics,  
Loughborough University,  
Leicestershire, Loughborough  
LE11 3TU, UK

e-mail: m.j.everitt@physics.org

## † Present address:

Richard D. Wilson, Sonobex Limited,  
Warwick, UK

We show that by engineering the interaction with the environment, there exists a large class of systems that can evolve irreversibly to a cat state. To be precise, we show that it is possible to engineer an irreversible process so that the steady state is close to a pure Schrödinger's cat state by using double well systems and an environment comprising two-photon (or phonon) absorbers. We also show that it should be possible to prolong the lifetime of a Schrödinger's cat state exposed to the destructive effects of a conventional single-photon decohering environment. In addition to our general analysis, we present a concrete circuit realization of both system and environment that should be fabricatable with current technologies. Our protocol should make it easier to prepare and maintain Schrödinger cat states, which would be useful in applications of quantum metrology and information processing as well as being of interest to those probing the quantum to classical transition.

**Keywords:** cat state, quantum metrology, single-photon decoherence, decoherence, two-photon absorbers, quantum to classical transition, double well systems

## 1. INTRODUCTION

The development of many quantum technologies depends on an ability to engineer strongly non-classical states. Such states take the form of either highly entangled states of distinct degrees of freedom or a quantum coherent superposition of macroscopically distinct states in a single degree of freedom (Sanders, 2012), known as Schrödinger's cat states [after a well known thought experiment (Schrödinger, 1935)]. It is these cat states that we consider in this paper. There has been great progress in the production of such states as well as experimentally reconstructing such states through a series of measurements in a process known as quantum state tomography (Monroe et al., 1996; Noel and Stroud, 1996; Leibfried et al., 2005; Ourjoumtsev et al., 2007; Deléglise et al., 2008; Gao et al., 2010). These developments are of great importance as, in addition to their curious nature, Schrödinger cat states can be used as a resource for developing technologies such as quantum computing (Ralph et al., 2003; Gilchrist et al., 2004), quantum communication (Jeong et al., 2001; van Enk and Hirota, 2001), and quantum metrology (Munro et al., 2002; Blatt and Wineland, 2008; Giovannetti et al., 2011). The main obstacle to deploying cat states in such applications is their fragility as they are destroyed by noise in a process termed environmental decoherence. A careful consideration of optical cat states shows that this decoherence may be interpreted as due to Poisson distributed jumps between even and odd cat states whenever a single photon is lost (Carmichael, 1993, 2003; Vitali et al., 1997). Their production and maintenance require very precise quantum control as well as low dissipation.

In this work, we present a possible realization of a protocol for double well system [in this case a Superconducting

Quantum-Interference Device (SQUID)] to create Schrödinger cat states using the interaction of the system with a special kind of environment. To be specific, we engineer an environment comprising a bath of two-photon absorbers, for certain initial states, such that the system relaxes to a steady state, which is close to a pure Schrödinger cat state. The use of open systems as well as the measurement process was proposed by Yurke, Schleich, and Walls (Yurke and Stoler, 1986; Yurke et al., 1990). Gilles, Garraway, and Knight also proposed that it would be possible to engineer an environment of this kind that when paired with a parametric photon pump would exhibit many interesting effects in quantum optical systems, from the generation of Schrödinger cat states to manifestly quantum statistics (McNeil and Walls, 1974; Simaan and Loudon, 1978; Loudon and Knight, 1987; Gilles and Knight, 1993a,b; Gilles et al., 1994a,b; Guerra et al., 1997) (see also Tor-nau and Bach, 1974; Simaan and Loudon, 1975; Hildred and Hall, 1978; Agarwal and Hildred, 1986; Gerry, 1993; Gerry and Hach, 1993; Hach and Gerry, 1994). Two-photon absorption has also been suggested as a powerful resource for application in quantum computing (Franson et al., 2004). We note that there already exist schemes for realizing non-classical states via engineered dissipative channels (Amico et al., 2008; Diehl et al., 2008; Kraus et al., 2008; Schirmer and Wang, 2010; Ticozzi et al., 2010; Zhang et al., 2010; Busch et al., 2011; Pechen, 2011; Scully et al., 2011; Chen et al., 2012; Ticozzi and Viola, 2012; Yamamoto, 2012; Ikeda and Yamamoto, 2013) as well as a number of experimental realizations (Barreiro et al., 2011; Krauter et al., 2011; Leghtas et al., 2013). We also note that SQUIDs are an ideal candidate system for realizing this protocol as they have already been shown able

to support appropriate quantum states (Nakamura et al., 1999; Friedman et al., 2000; Grajcar et al., 2004; Il'ichev et al., 2004) and, as we shall later show, existing circuit designs can be used to engineer an environment with the suitable characteristics (Deng et al., 2010; Kumar and DiVincenzo, 2010).


We note that our scheme is simpler than and different from other driven dissipative bistable systems [for example, the coherently driven optical cavity containing a Kerr medium (Walls and Milburn, 2008), the driven Duffing mechanical resonator (Babourina-Brooks et al., 2008), tapered optical fibers (Hendrickson et al., 2010), and photon pumps (Gilles and Knight, 1993a; Gilles et al., 1994a)], as we do not include driving on either the cavity resonance or the coordinate degree of freedom. The possibility of engineering dissipative channels, such as the one that we present in this work, opens up new opportunities for exploring quantum phenomena from the micro to macroscopic level and in fields as diverse as quantum optics (Haycock et al., 2000), Bose–Einstein condensates (Andrews, 1997), quantum electronics (Friedman et al., 2000), and nano-mechanics (Badzey and Mohanty, 2005) [for which multi-phonon relaxation has already been proposed (Voje et al., 2013a,b)] or any other system in which it is possible to generate a double well potential.

## 2. MODEL SYSTEM: THE SQUID RING

For the results presented in this paper, we have used as an example system a superconducting quantum-interference device (SQUID) ring. Our reason for choosing SQUIDs is that these devices are routinely fabricated and their theory is very well understood. We note that we have investigated a number of other systems (but do not include results here) and our analysis indicates that the key feature of the ring is that it can be made to form a double well potential. Moreover, non-linear systems derived from the Josephson junction in circuit QED exhibit multi photon resonance when driven by an external field (Deppe et al., 2008) and thus we expect two-photon decay to be present in such systems. The real difficulty is making it dominate over single photon effects. We will return to this in Section 4 where we propose a concrete circuit realization of a suitable and realistic environment for a SQUID. Beyond these considerations, we believe that there is nothing particularly special about the exact form of the potential needed to realize our protocol. The potential energy of the SQUID comprising a thick superconducting ring enclosing a Josephson junction weak link takes the form of a harmonic oscillator perturbed by a cosine

$$U(\Phi_x) = \frac{(\Phi - \Phi_x)^2}{2\Lambda} - \frac{\hbar I_c}{2e} \cos\left(2\pi \frac{\Phi}{\Phi_0}\right)$$

where the coordinate  $\Phi$  is the total magnetic flux in the ring and  $\Phi_0 = h/2e$  is the superconducting flux quantum. We have chosen example circuit parameters that are in-line with modern fabrication techniques and suited to experimental realizations:  $\Lambda = 3 \times 10^{-10}$  H for the ring's inductance and  $I_c = 2 \mu\text{A}$  as the critical current of the weak link (although not in the above formula, we also chose a capacitance  $C = 5 \times 10^{-15}$  F). We set the externally applied magnetic flux  $\Phi_x = 0.5\Phi_0$ , so that the ring's potential forms a degenerate double well. It is also convenient to introduce the bosonic annihilation  $a$ , and creation  $a^\dagger$  operators



**FIGURE 1 | Stationary state energy levels: the potential energy of the ring (black) as well as the energy of the ring's stationary states (blue).**

Parameters used here and throughout the paper are inductance  $\Lambda = 3 \times 10^{-10}$  H, capacitance  $C = 5 \times 10^{-15}$  F, critical current of the weak link  $I_c = 2 \mu\text{A}$ , and externally applied magnetic flux  $\Phi_x = 0.5\Phi_0$ . Note that, we have exaggerated the energy difference between the ground and first excited states as well as stationary states two and three in order to make the different energies visible on this plot.

where  $\Phi = \sqrt{\frac{\hbar}{2C\omega_{LC}}}(a + a^\dagger)$  and  $\omega_{LC} = 1/\sqrt{\Lambda C}$ . In **Figure 1**, we show the potential energy of the ring as well as the energy of the ring's stationary states. It is worth noting that the ground state and first excited state approximate, respectively, symmetric and anti-symmetric superpositions of two coherent states centered at the bottom of each well. These two states have very nearly the same energy and the difference in their energy has been exaggerated in this plot (as have those for the second and third excited states).

## 3. THE EFFECT OF IDEAL ENVIRONMENTS

### 3.1. BACKGROUND

We model the effect of the environment on the system using the master equation in the Lindblad form (Viola et al., 1997)

$$\frac{d\rho}{dt} = -\frac{i}{\hbar} [H, \rho] + \frac{1}{2} \sum_j \left\{ [L_j, \rho L_j^\dagger] + [L_j \rho, L_j^\dagger] \right\}$$

where  $\rho$  is the density matrix describing the state of the system (initially  $\rho = |\psi(t=0)\rangle\langle\psi(t=0)|$ ) and  $H$  is the system's Hamiltonian. The non-unitary effect of the environment on the system is contained in the Lindblad operators  $L_j$  with each describing a possible environment. For example, the usual Ohmic (i.e., analogous to friction proportional to velocity) or lossy bath, at zero temperature, would be described by a Lindblad operator proportional to the annihilation operator. For an undriven system, the master equation has steady state solution that, in the presence of an environment, is usually a density operator in a mixed state. In certain circumstances, at zero temperature, these solutions may be pure states such as the vacuum state of the harmonic oscillator. In general, the steady state solutions will not exhibit features such as superpositions of macroscopically distinct states and are

relatively uninteresting. It is precisely this process where the environment essentially removes the system's quantum coherence from de-localized, or more generally non-Gaussian states, that is known as environmental decoherence. The density matrix for a decohered system without these quantum correlations represents a statistical mixture of possible states of the system and, for a single quantum object, can be directly compared with classical probability density distributions (Habib et al., 1998). It should be noted, however, that there are driven dissipative systems, for example, dispersive bistability, for which the steady state is a mixed state with a considerable amount of quantum coherence in the limit of large Kerr non-linearity (Wolinsky and Carmichael, 1988; Carmichael, 1993, 2003).


We find very different behavior if one chooses a different environment comprising two-photon absorbers, described by a Lindblad operator proportional to the square of the annihilation operator. In **Figure 2**, we show the energy expectation values and von-Neumann entropy,  $S = -\text{Tr}[\rho \ln \rho]$  as functions of time for solutions of the master equation for the ring in the presence of such an environment. We used as initial conditions the first twenty energy eigenstates of the ring Hamiltonian. In these plots, the energy behaves just as one would expect the energy of an undriven open quantum system to do – it settles to a single value. When one inspects the dynamics of the entropy, however, the story is quite different. One usually expects the entropy to grow from zero to some asymptotic value as the system evolves into a mixed state. While we see that this is the initial behavior, the entropy does not monotonically increase; instead it decreases until the entropy is nearly negligible. It appears that the system has to a significant extent recohered and the final density matrix is very nearly that of a pure state. While this is not the usual behavior of an open quantum system, based on previous work such as Yurke and Stoler (1986), Yurke et al. (1990), Gilles and Knight (1993a), and Gilles et al. (1994a), it is in-line with our expectations of an environment that “decoheres” a system to an almost pure state that is a very good approximation to a Schrödinger cat state.

### 3.2. PHASE SPACE METHOD: THE WIGNER FUNCTION

In order to demonstrate that the system does indeed decay to a Schrödinger cat state, we will make use of the Wigner function. These pseudo probability density functions in phase space have been of great utility in demonstrating that some quantum states are Schrödinger cats (Deléglise et al., 2008). The Wigner function is


$$W(\Phi, Q) = \frac{1}{2\pi\hbar} \int \langle \Phi + \zeta | \rho | \Phi - \zeta \rangle \exp\left(-\frac{2iQ\zeta}{\hbar}\right) d\zeta$$

where  $Q$  is the charge variable that is conjugate to the magnetic flux  $\Phi$ . In **Figure 3**, we show three Wigner functions. **Figure 3A** shows the initial state and is a coherent state centered at the origin. This is clearly recognizable as the expected Gaussian bell shape associated with coherent states. We have solved the master equation for the ring in a lossy bath, with a Lindblad of  $L = \sqrt{0.2}a$  and allowed the system to reach its steady state to obtain **Figure 3B**. This is the Wigner function of a statistical mixture of two macroscopically distinct states and is in-line with expectations of the effect of a decohering environment on such a device (Everitt et al., 2004). In



**FIGURE 2 | Effect of decoherence on energy and entropy.** We show the dynamical evolution of the ring's energy and entropy using each of its first twenty stationary states as initial conditions. The dynamics have been found by solving the master equation for the ring in the presence of a bath of two-photon absorbers (with  $L = \sqrt{0.2}a^2$ ). We have provided insets for increased resolution of the system's initial dynamics. The top plot shows the dynamics of the ring's total energy. As expected for an open quantum system of this kind, the ring can be seen to decohere to one energy, a little above that of the ground state. The bottom plot shows the dynamics of the von-Neuman entropy for the ring. In each case, the initial entropy is zero as the system starts in a pure state. The entropy grows before dropping off to a low value indicating that the system's steady state solution is very nearly a pure state.

**Figure 3C**, we show the Wigner function that we obtain by solving the master equation, as for (**Figure 3B**), but replacing the damping term with a bath of two-photon absorbers, with  $L = \sqrt{0.2}a^2$ . We notice two things: firstly, the state has rotated, which we believe to be a consequence of a squeezing action associated with the bath and secondly that there are interference terms between the distinct states of the system. These interference terms, indicating quantum



**FIGURE 3 | Cooling for a cat.** In this figure we show, by making use of Wigner functions, the effect of two different environments on a ring prepared in a coherent state biased at zero flux. Each graph contains a top down view with a three dimensional plot of the function as a not to fixed scale inset. The graphs show **(A)**, the initial state, which takes the form of a Gaussian bell. **(B)** The steady state solution to the master equation under the influence of a conventional decohering environment comprising a lossy bath (with a Lindblad proportional to the annihilation operator  $L = \sqrt{0.2}a$ ). The ring has decohered to two distinct macroscopic states

and we do not see the interference terms between them that are characteristic of a Schrödinger cat state. We have instead a statistical mixture, the usual and expected result (Everitt et al., 2004). **(C)** The steady state solution to the master equation for the ring coupled to a bath of two-photon absorbers (with a Lindblad proportional to the square of annihilation operator,  $L = \sqrt{0.2}a^2$ ). In this case, the ring has decohered to a superposition of two macroscopic states but now there are interference terms between these states indicating quantum coherence – the signature of a Schrödinger cat state.

coherence, confirm that this state is indeed a very good approximation to a Schrödinger cat. We note that two-photon decay preserves parity, which can be easily seen if we consider the representation of the systems state vector in the harmonic oscillator basis. Here, we see that  $a^2$  will only couple even states to even states and odd states to odd states. Thus, the action of  $a^2$  on any initial state must preserve its parity. Hence, an environment comprising only two-photon absorbers would ensure that the system will relax to a steady state with same parity as the initial state. It is this symmetry property of the environment together with the symmetry in the Hamiltonian and initial condition that leads to steady state solutions that are Schrödinger cat states.

### 3.3. QUANTIFYING NON-LOCAL CORRELATIONS

In order to examine quantitatively the emergence of this cat from the initial coherent state we introduce, following (Nogues et al., 2000; Białynicki-Birula et al., 2002), a measure of how de-localized the system is in phase space that is the integral of negative parts of the Wigner function

$$N(\rho) = \frac{1}{2} \int \{|W(\Phi, Q)| - W(\Phi, Q)\} d\Phi dQ.$$

In absolute terms, this is a useful measure, but when we know (by inspecting the Wigner function) that the states we are examining are cat-like, a more useful measure may well be a relative cattiness to some reference Schrödinger cat state.


Hence, we define:

$$\text{Cat}(\rho, \rho_{\text{ref}}) = \frac{N(\rho)}{N(\rho_{\text{ref}})} \quad (1)$$


which quantifies the ratio of the de-localization of one cat state against a reference cat and enables us to quantify if one is more [ $\text{Cat}(\rho, \rho_{\text{ref}}) > 1$ ], less [ $\text{Cat}(\rho, \rho_{\text{ref}}) < 1$ ], or just as [ $\text{Cat}(\rho, \rho_{\text{ref}}) = 1$ ] catty than the other. We have chosen to introduce this measure over using existing metrics such as the fidelity as it does not contain any contributions of the type that occur from, for example, correlations between a cat and its related mixed state (which might be thought of as the overlap of the “classical” like parts of the state). Computing relative measures such as the fidelity is further complicated by the fact that the final states, in terms of the size and orientation, for the different environments are very different from each other (having very little overlap). Hence, performing meaningful estimates of fidelity would be quite difficult, perhaps even impossible as we would have to provide different reference states for each environment against which to measure the fidelity. We note that while there are some limitations with the Cat measure and it should be applied with care, for the problem we study here, it suits our purposes very well. In **Figure 4**, we show the dynamics of this quantity for comparison with the results presented in **Figure 3** using as a reference state  $\rho_{\text{ref}}$  the final cat state shown in **Figure 3C**. Here, we can clearly see that the cattiness of the system subject to an environment of two-photon absorbers monotonically increases and asymptotically converges to a steady state.

### 3.4. RESULTS

It is interesting to consider what would happen to a ring that was initially in a Schrödinger cat state under the influence of a bath of two-photon absorbers. For systems with deep enough double well potentials, such as the one considered here the ground




and first excited energy eigenstates are both Schrödinger cats. The ground state is, to good approximation, an even superposition of two macroscopically distinct coherent states while the first excited state is an odd superposition as can be seen from their Wigner functions in **Figures 5A,C**, respectively. The even and odd nature of these superpositions is reflected in the Wigner function by the phase of the interference terms between the two Gaussians of the cat. It is known that such states would decohere under the environment of a lossy bath to a statistical mixture (Everitt et al., 2004). The dynamics of the system coupled to an environment comprising a bath of two-photon absorbers are, once more, found by solving the master equation with an  $L = \sqrt{0.2}a^2$ , until an approximate steady state is reached. The Wigner function of these states is then shown with **Figure 5A** evolving to **Figures 5B–D** and **Figures 5C,D**. We observe that the phase in the final cat reflects that of the initial cat and the system has not simply decohered to the same steady state. The environment thus seems to preserve some of the symmetry of the initial state. We have checked the first twenty stationary states all of which decay to one of these cats or the other. Moreover, the pattern that was observed from the ground and first excited state persists and all even and odd states seem to evolve to cats of the



same form as those shown in **Figures 5B,D** that are out of phase with each other.

This approach seems all very well and good, but an environment of two-photon absorbers is very special. It would be hard to construct such an environment without having any other source of decoherence present. We therefore need to verify that the effects of a two-photon absorbing environment cannot be completely destroyed by the presence of a more traditional environment such as a lossy bath. In **Figure 6**, we show the results of just such a check. For each plot, the system's initial state was the ring's ground state as shown in **Figure 5A**. In **Figure 6A**, we show the effect of a lossy bath. We solve the master equation with a Lindblad  $L = \sqrt{0.02}a$  and allow the system to evolve until it has just decohered to a statistical mixture and we have plotted the Wigner function at this point in time. We use this run as a benchmark for computing the next two cases, which show the Wigner function solutions of the master equation computed over the same interval. In **Figure 6B**, we show the effect of a two-photon absorbing bath once more "decohering" to a Schrödinger cat state ( $L = \sqrt{0.2}a^2$ ). And in **Figure 6C**,

we apply both the lossy bath of in **Figure 6A** and a two-photon absorbing environment of in **Figure 6B** to the ring ( $L_1 = \sqrt{0.02}a$  and  $L_2 = \sqrt{0.2}a^2$ ). We see that in this figure, there remain residual Schrödinger cat state features in the Wigner function. Hence, it seems that not only does a bath of two-photon absorbers create Schrödinger cat states but also enables Schrödinger cat states to be more resilient to other forms of decoherence. In other words, the presence of an environment of two-photon absorbers seems to be prolonging the life of a damped cat. In **Figure 6D**, we quantify the cattiness using  $\text{Cat}(\rho, \rho_{\text{ref}})$  using the initial stationary state as shown in **Figure 5A** as the reference cat. For the three environments considered here, we find that for the system's later evolution the environment of two-photon absorbers does indeed prolong the lifetime of the initial cat even in the presence of a lossy bath. We note that we obtain an almost identical set of results if we start the system off in a coherent state centered at the origin (as in **Figure 3A**). We chose to use the ring's ground state as, in our view, we obtained a more instructive plot of the states cattiness from the systems dynamics. For a direct comparison of the dynamics



of  $\text{Cat}(\rho, \rho_{\text{ref}})$  for these two initial conditions, we now note that the dashed line shown in **Figure 4** was found for a lossy bath and a two-photon absorbing environment with  $L_1 = \sqrt{0.02}a$  and  $L_2 = \sqrt{0.2}a^2$ . The green and blue lines of **Figures 4** and **6D** are directly comparable. The idea that the presence of a two-photon absorbing environment can be used to extend the lifetime (and also generate) Schrödinger cat states that holds equally well for two very different initial conditions.

#### 4. A REALISTIC MODEL OF AN ENVIRONMENT

##### 4.1. OVERVIEW

In order to make our above discussion a reality, we need to engineer a dissipative quantum channel that acts as a two-photon absorber. Here, we suggest a concrete realization that, whilst not perfect, still retains the key feature of environmentally induced “decoherence” to a Schrödinger cat state. Our proposal makes use of non-linearly coupled electromagnetic fields and SQUIDs. Such quantum electrodynamic circuits have already been investigated in the context of weak non-demolition measurement (Deng et al., 2010; Kumar and DiVincenzo, 2010). One example comprises two microwave superconducting resonators coupled via a SQUID, which in addition to a cross Kerr effect also manifests two-photon conversion terms if the cavities are resonant (Kumar and DiVincenzo, 2010). Such systems can be quantized (Everitt et al., 2001a,b; Stiffell et al., 2005; Wallquist et al., 2006) and with a suitable arrangement and choice of circuit parameters can be reduced (Wielinga and Milburn, 1993; Santamore et al., 2004) to the form of a double well system subject to a two-photon absorbing environment (see Section 4 for details). Unavoidably, this process also brings with it an additional dephasing term, that adds to the master equation another Lindblad operator proportional to  $a^\dagger a$ . Nevertheless, we can report that while the dephasing term smears out the Gaussian peaks in the cat the interference terms in the Wigner function representing quantum coherence between the cat states remain strong. The fact that this dephasing term preserves parity is once more the key factor in ensuring the steady state of our engineered dissipative channel is still a Schrödinger cat state. Our proposal could lead to an initial realization of a two-photon absorbing environment and concomitant interesting effects. The engineering of improved dissipative channels, without additional and unwanted decoherence effects, remains an open and interesting problem.

##### 4.2. THE MODEL: DERIVATION OF THE MASTER EQUATION

There are a number of models (Deng et al., 2010; Kumar and DiVincenzo, 2010) whereby two microwave superconducting cavities can be non-linearly coupled using SQUIDs. We will base our discussion on Kumar and DiVincenzo (2010). In that model, the Hamiltonian describing two microwave cavities, a probe (p) cavity and a signal (s) cavity, coupled with a SQUID is

$$H = E_{Cp}n_p^2 + E_{Lp}\phi_p^2 + E_{Cs}n_s^2 + E_{Ls}\phi_s^2 + A \left[ E_{Lp}^4\phi_p^4\cos^4\beta + E_{Ls}^4\phi_s^4\sin^4\beta + 6E_{Lp}^2E_{Ls}^2\phi_p^2\phi_s^2\cos^2\beta\sin^2\beta \right] \quad (2)$$

where  $n_\alpha, \phi_\alpha$  are the standard charge and phase conjugate variables describing the collective electrical degree of freedom in each cavity and  $A = 16\pi^2 L_1/\Phi_0^4$  with  $L_1$  defined as the coefficient of

the leading non-linear current term of the SQUID inductance. We will set  $\cos^2\beta = \sin^2\beta = 1/2$ .

The system can be quantized in the usual way in terms of the bosonic annihilation and creation operators  $b, b^\dagger$  for the probe and  $a, a^\dagger$  and for the signal cavity defined by (Wallquist et al., 2006)

$$\phi_p \rightarrow \left( \frac{E_{Cp}}{4E_{Lp}} \right)^{1/4} (b + b^\dagger) \quad (3)$$

$$n_p \rightarrow -i \left( \frac{E_{Lp}}{4E_{Cp}} \right)^{1/4} (b - b^\dagger) \quad (4)$$

$$\phi_s \rightarrow \left( \frac{E_{Cs}}{4E_{Ls}} \right)^{1/4} (a + a^\dagger) \quad (5)$$

$$n_s \rightarrow -i \left( \frac{E_{Ls}}{4E_{Cs}} \right)^{1/4} (a - a^\dagger) \quad (6)$$

The Hamiltonian may then be written as

$$H = \hbar\omega_p b^\dagger b + \hbar\omega_s a^\dagger a + \hbar\chi_b b^{\dagger 2} b^2 + \hbar\chi_a a^{\dagger 2} a^2 + \hbar\sqrt{\chi_a\chi_b} (b^2 a^{\dagger 2} + b^{\dagger 2} a^2 + 4a^\dagger a b^\dagger b) \quad (7)$$

Unlike Kumar and DiVincenzo (2010), we have *not* neglected the terms like  $b^2 a^{\dagger 2}$  as we will choose  $\omega_p = \omega_s$  so that these terms are resonant<sup>1</sup>.

We now include the dissipative channels for this model in the usual way. The density operator for the total system, in the interaction picture, satisfies

$$\frac{d\rho}{dt} = -i[H_I, \rho] + \kappa_a \mathcal{D}[a]\rho + \kappa_b \mathcal{D}[b]\rho \quad (8)$$

where  $\mathcal{D}[L]\rho = L\rho L^\dagger - \frac{1}{2}(L^\dagger L\rho + \rho L^\dagger L)$  and

$$H_I = \hbar\chi_b b^{\dagger 2} b^2 + \hbar\chi_a a^{\dagger 2} a^2 + \hbar(\epsilon^* b + \epsilon b^\dagger) + \hbar\sqrt{\chi_a\chi_b} (b^2 a^{\dagger 2} + b^{\dagger 2} a^2 + 4a^\dagger a b^\dagger b) \quad (9)$$

and  $\kappa_a, \kappa_b$  are the decay rates of the photon number in the signal and probe cavity, respectively, and we have included a resonant coherent driving of the probe cavity with  $\epsilon = \sqrt{\kappa_b}\epsilon_b$  where  $|\epsilon_b|^2$  is the photon flux of the driving field. We have also assumed that each cavity sees a zero temperature environment.

In the absence of the SQUID mediated interactions, the probe cavity will relax to a coherent state with the steady state amplitude

$$\beta_0 = \frac{-2i\epsilon}{\kappa_b}. \quad (10)$$

<sup>1</sup>Note that the coefficients in this equation are  $\hbar\chi_a = \frac{3A}{8}E_{Cs}E_{Ls}^3, \hbar\chi_b = \frac{3A}{8}E_{Cp}E_{Lp}^3, \hbar\omega_s = 2\sqrt{E_{Cs}E_{Ls}} \left[ 1 + \frac{3A}{8}E_{Ls} \left( \sqrt{E_{Cs}E_{Ls}^3} + \sqrt{E_{Cp}E_{Lp}^3} \right) \right]$  and  $\hbar\omega_p = 2\sqrt{E_{Cp}E_{Lp}} \left[ 1 + \frac{3A}{8}E_{Lp} \left( \sqrt{E_{Cs}E_{Ls}^3} + \sqrt{E_{Cp}E_{Lp}^3} \right) \right]$ .

We will choose the phase of the probe driving as a reference phase and set  $\beta_0$  to be real. If we make a canonical transformation to the displaced picture by

$$b = \bar{b} + \beta_0 \tag{11}$$

we can linearize the Hamiltonian, Equation (9), in  $\bar{b}, \bar{b}^\dagger$  to obtain

$$H_I = H_a + 4\hbar\sqrt{\chi_a\chi_b}\beta_0 (\bar{b} + \bar{b}^\dagger) a^\dagger a + 2\hbar\sqrt{\chi_a\chi_b}\beta_0 (\bar{b}^\dagger a^2 + \bar{b} a^{\dagger 2}) \tag{12}$$

where the effective Hamiltonian for the signal mode alone is

$$H_a = \hbar\chi_a a^{\dagger 2} a^2 + 4\hbar\sqrt{\chi_a\chi_b}\beta_0^2 a^\dagger a + \hbar\sqrt{\chi_a\chi_b}\beta_0^2 (a^2 + a^{\dagger 2}) \tag{13}$$

which is equivalent to a parametrically driven Kerr non-linear cavity. This model was considered by Wielinga and Milburn (1993). It is equivalent to a double well system with a hyperbolic fixed point at the origin in phase space and two elliptic fixed points symmetrically displaced from the origin. The second and third terms in equation (12) can be given a familiar interpretation. The second term is of the same form as the radiation pressure interaction between a mechanical resonator ( $\bar{b}, \bar{b}^\dagger$ ) and a cavity field ( $a, a^\dagger$ ). The last term is equivalent to the quantum description of sub/second harmonic generation considered by Drummond et al. (1980).

We now assume that  $\kappa_b$ , the line width of the probe cavity is large,  $\kappa_b \gg \kappa_a, \sqrt{\chi_a\chi_b}$  and we adiabatically eliminate it from the dynamics. In that case from the point of view of the signal mode, the first term in equation (12) looks like a fluctuating cavity detuning while the last term looks like a two-photon loss term. This can be verified by explicit adiabatic elimination of the probe cavity field. We assume that the probe cavity, in the displaced picture, remains very close to its steady state of zero photons. The method is described in Santamore et al. (2004). The effective master equation for the signal cavity is

$$\frac{d\rho_s}{dt} = -\frac{i}{\hbar} [H_a, \rho_s] + \Gamma_2 D[a^2] \rho_s + \Gamma_\perp D[a^\dagger a] \rho_s + \kappa_a D[a] \rho_s \tag{14}$$


where the two-photon decay rate  $\Gamma_2$  and dephasing rate  $\Gamma_\perp$  are given by

$$\Gamma_2 = \frac{16\chi_a\chi_b\beta_0^2}{\kappa_b} \text{ and } \Gamma_\perp = \Gamma_2/4 = \frac{4\chi_a\chi_b\beta_0^2}{\kappa_b}. \tag{15}$$

### 4.3. RESULTS


A peculiar feature of using SQUID coupled cavities is that the price paid for two-photon decay is an additional dephasing term on the signal cavity field. Using the strong dependence on the steady state amplitude  $\beta_0$  in the two-photon rate, we can make the two-photon decay term that dominate over the single photon decay of the signal cavity over the time scales of interest. In **Figure 7A**, we show that the dephasing term that is introduced in the above (un-damped,  $\kappa_a = 0$ ) master equation has little effect

on the Schrödinger cat nature of the steady state solution associated with the two-photon absorbing bath. In order to consider a worst case scenario, in **Figure 7B**, we go on to consider what would happen if we simultaneously weaken the effect of the two-photon environment and make the dephasing term even stronger. We have not taken these values from our model as the ratio of  $\Gamma_\perp$  to  $\Gamma_2$  has not been preserved. Our reason for presenting this data is to indicate that alternative circuit realizations, where the beneficial effects of the two-photon absorbing environment are reduced and the damaging effects of the dephasing term increased, might still be used to engineer a steady state cat. We therefore believe that



**FIGURE 7 | A persistent cat, here we look at the effect of including the dephasing term in addition to the bath of two-photon absorbers to the ring initially in its ground state. (A)** The steady state solution for the environment as derived in Section 4.2 and described by master equation, equations (14) and (15) with  $\Gamma_2 = 0.2$  (or  $L_2 = \sqrt{0.2}a^2$ ) and  $\Gamma_\perp = 0.05$  (or  $L_\perp = \sqrt{0.05}a^2$ ) and we have set the damping coefficient  $\kappa_a = 0$ . **(B)** We show that even for an environment, other than the one considered in **(A)**, where dephasing dominates over the two-photon absorption process ( $L_2 = \sqrt{0.02}a^2$  and  $L_\perp = \sqrt{0.08}a^2$ ), it is still possible for the steady state of the ring to be a cat.





**FIGURE 8 | A stubborn cat II: the effect of a realistic environment for comparison with Figure 6, in this plot we show the cattiness for a: lossy bath ( $L_1 = \sqrt{0.02}a$ ); a lossy bath and a two-photon absorbing bath ( $L_1 = \sqrt{0.02}a$  and  $L_2 = \sqrt{0.2}a^2$ ); and also the full model including the dephasing term ( $L_1 = \sqrt{0.02}a$ ,  $L_2 = \sqrt{0.2}a^2$ , and  $L_\perp = \sqrt{0.05}a^\dagger a$ ). Even with the dephasing term included the environment of two-photon absorbers still prolongs the lifetime of the initial cat in the presence of a lossy bath.**

the discussion in Section 3 is in-line with the behavior of realistic environments.

Finally, in **Figure 8** we show the cattiness [see equation (1)] of the realistic environment with two of the cases considered in **Figure 6**. Specifically, we consider the cases of a lossy bath ( $L_1 = \sqrt{0.02}a$ ); a lossy bath and a two-photon absorbing ( $L_1 = \sqrt{0.02}a$  and  $L_2 = \sqrt{0.2}a^2$ ) and also the full model, of equation (14), including the dephasing term ( $L_1 = \sqrt{0.02}a$ ,  $L_2 = \sqrt{0.2}a^2$ , and  $L_\perp = \sqrt{0.05}a^\dagger a$ ). Even with the dephasing term included the environment of two-photon absorbers still prolongs the lifetime of the initial cat in the presence of a lossy bath.

## 5. CONCLUSION

There are two phenomena that embody quantum mechanics, namely entanglement and the Schrödinger's cat thought experiment for making macroscopic superposition states (Schrödinger, 1935). The latter was proposed to highlight the difficulties that we have connecting quantum mechanics with everyday experience, as it neatly demonstrates the problems of understanding the emergence of the classical world from quantum theory and the measurement of quantum systems. Schrödinger's cat has become the icon of the subject and evolved to have a well defined meaning. It is an accepted explanation within the popular literature that the reason the original thought experiment does not translate into reality (if conducted with a real cat in a box, etc.) is that the coupling of the environment to the radiation source (which included the cat itself) makes it impossible to observe the coherence between the two superposed macroscopically distinct states – a process known as decoherence. As such, environmental decoherence is something that many deem to be a crucial element in the quantum to classical transition (Bell, 1990; Habib et al., 1998; Schlosshauer, 2005; Everitt, 2009; Everitt et al., 2009). We

have presented an example of an engineered environment that may be used to produce Schrödinger cat states as a steady state. It may well be that system and environment such as the one we have used here could play an interesting role in quantum mechanically enhanced metrology probing foundational aspects of quantum mechanics associated with realizing macroscopic quantum phenomena and the quantum to classical transition. In addition, the two-photon decay channel, if monitored appropriately, enables a measurement of the intensity squared of the number and may also enable novel non-linear feedback protocols. Although it is beyond the scope of the current paper, we conjecture that it may soon be possible, following (Yurke et al., 1990), to make use of an environment to create a conditional Schrödinger cat state by measurement.

## ACKNOWLEDGMENTS

Mark J. Everitt, Richard D. Wilson, and Richard D. Wilson thank the Templeton Foundation for their generous support. Gerard J. Milburn acknowledges the support of the National Science Foundation under Grant No. NSF PHY11-25915 and the Australian Research Council Centre of Excellence for Engineered Quantum Systems grant CE110001013. Mark J. Everitt would like to thank Andrew Archer, Gerry Swallowe, and Richard Giles for their help with the preparation of our manuscript.

## REFERENCES

- Agarwal, G., and Hildred, G. (1986). Time development of squeezing in two photon absorption. *Opt. Commun.* 58, 287–289. doi:10.1016/0030-4018(86)90452-9
- Amico, L., Fazio, R., Osterloh, A., and Vedral, V. (2008). Entanglement in many-body systems. *Rev. Mod. Phys.* 80, 517–576. doi:10.1103/RevModPhys.80.517
- Andrews, M. R. (1997). Observation of interference between two Bose condensates. *Science* 275, 637–641. doi:10.1126/science.275.5300.637
- Babourina-Brooks, E., Doherty, A., and Milburn, G. J. (2008). Quantum noise in a nanomechanical duffing resonator. *New J. Phys.* 10, 105020. doi:10.1088/1367-2630/10/10/105020
- Badzey, R. L., and Mohanty, P. (2005). Coherent signal amplification in bistable nanomechanical oscillators by stochastic resonance. *Nature* 437, 995–998. doi:10.1038/nature04124
- Barreiro, J. T., Müller, M., Schindler, P., Nigg, D., Monz, T., Chwalla, M., et al. (2011). An open-system quantum simulator with trapped ions. *Nature* 470, 486–491. doi:10.1038/nature09801
- Bell, J. (1990). Against “measurement”. *Phys. World* 3, 33.
- Białynicki-Birula, I., Cirone, M., Dahl, J., Fedorov, M., and Schleich, W. (2002). In- and outbound spreading of a free-particle s-wave. *Phys. Rev. Lett.* 89, 060404. doi:10.1103/PhysRevLett.89.060404
- Blatt, R., and Wineland, D. (2008). Entangled states of trapped atomic ions. *Nature* 453, 1008–1015. doi:10.1038/nature07125
- Busch, J., De, S., Ivanov, S. S., Torosov, B. T., Spiller, T. P., and Beige, A. (2011). Cooling atom-cavity systems into entangled states. *Phys. Rev. A* 84, 022316. doi:10.1103/PhysRevA.84.022316
- Carmichael, H. (1993). *An Open Systems Approach to Quantum Optics: Lectures Presented at the Université Libre de Bruxelles, October 28 – November 4, 1991 (Lecture Notes in Physics Series)*. Berlin: Springer-Verlag.
- Carmichael, H. (2003). *Statistical Methods in Quantum Optics 1: Master Equations and Fokker-Planck Equations (Statistical Methods in Quantum Optics)*. Berlin: Springer.
- Chen, X.-Y., Shen, L.-T., Yang, Z.-B., Wu, H.-Z., and Chen, M.-F. (2012). Engineering W-type steady states for three atoms via dissipation in an optical cavity. *J. Opt. Soc. Am. B* 29, 1535. doi:10.1364/JOSAB.29.001535
- Deléglise, S., Dotsenko, I., Sayrin, C., Bernu, J., Brune, M., Raimond, J.-M., et al. (2008). Reconstruction of non-classical cavity field states with snapshots of their decoherence. *Nature* 455, 510–514. doi:10.1038/nature07288

- Deng, C., Gambetta, J. M., and Lupascu, A. (2010). Quantum nondemolition measurement of microwave photons using engineered quadratic interactions. *Phys. Rev. B* 82, 220505. doi:10.1103/PhysRevB.82.220505
- Deppe, F., Mariani, M., Menzel, E. P., Marx, A., Saito, S., Kakuyanagi, K., et al. (2008). Two-photon probe of the Jaynes Cummings model and controlled symmetry breaking in circuit QED. *Nat. Phys.* 4, 686. doi:10.1038/nphys1016
- Diehl, S., Micheli, A., Kantian, A., Kraus, B., Büchler, H. P., and Zoller, P. (2008). Quantum states and phases in driven open quantum systems with cold atoms. *Nat. Phys.* 4, 878–883. doi:10.1038/nphys1073
- Drummond, P., McNeil, K., and Walls, D. (1980). Non-equilibrium transitions in sub/second harmonic generation. *Opt. Acta (Lond.)* 28, 211.
- Everitt, M. J. (2009). On the correspondence principle: implications from a study of the nonlinear dynamics of a macroscopic quantum device. *New J. Phys.* 11, 013014. doi:10.1088/1367-2630/11/1/013014
- Everitt, M. J., Clark, T. D., Stiffell, P., Prance, H., Prance, R. J., Vourdas, A., et al. (2001a). Quantum statistics and entanglement of two electromagnetic field modes coupled via a mesoscopic squid ring. *Phys. Rev. B* 64, 184517. doi:10.1103/PhysRevB.64.184517
- Everitt, M. J., Stiffell, P., Clark, T. D., Vourdas, A., Ralph, J. F., Prance, H., et al. (2001b). Fully quantum-mechanical model of a squid ring coupled to an electromagnetic field. *Phys. Rev. B* 63, 144530. doi:10.1103/PhysRevB.63.144530
- Everitt, M. J., Clark, T. D., Stiffell, P. B., Vourdas, A., Ralph, J. F., Prance, R. J., et al. (2004). Superconducting analogs of quantum optical phenomena: macroscopic quantum superpositions and squeezing in a superconducting quantum-interference device ring. *Phys. Rev. A* 69, 043804. doi:10.1103/PhysRevA.69.043804
- Everitt, M. J., Munro, W. J., and Spiller, T. P. (2009). Quantum-classical crossover of a field mode. *Phys. Rev. A* 79, 032328. doi:10.1103/PhysRevA.79.032328
- Franson, J. D., Jacobs, B. C., and Pittman, T. B. (2004). Quantum computing using single photons and the Zeno effect. *Phys. Rev. A* 70, 062302. doi:10.1103/PhysRevA.70.062302
- Friedman, J. R., Patel, V., Chen, W., Tolpygo, S. K., and Lukens, J. E. (2000). Quantum superposition of distinct macroscopic states. *Nature* 406, 43. doi:10.1038/35017505
- Gao, W.-B., Lu, C.-Y., Yao, X.-C., Xu, P., Günhe, O., Goebel, A., et al. (2010). Experimental demonstration of a hyper-entangled ten-qubit Schrödinger cat state. *Nat. Phys.* 6, 331–335. doi:10.1038/nphys1603
- Gerry, C. C. (1993). Non-classical properties of even and odd coherent states. *J. Mod. Opt.* 40, 1053–1071. doi:10.1080/09500349314551131
- Gerry, C. C., and Hach, E. E. (1993). Generation of even and odd coherent states in a competitive two-photon process. *Phys. Lett. A* 174, 185–189. doi:10.1016/0375-9601(93)90756-P
- Gilchrist, A., Nemoto, K., Munro, W. J., Ralph, T. C., Glancy, S., Braunstein, S. L., et al. (2004). Schrödinger cats and their power for quantum information processing. *J. Opt. B Quantum Semiclassical Opt.* 6, S828. doi:10.1088/1464-4266/6/8/032
- Gilles, L., Garraway, B. M., and Knight, P. L. (1994a). Generation of nonclassical light by dissipative two-photon processes. *Phys. Rev. A* 49, 2785–2799. doi:10.1103/PhysRevA.49.2785
- Gilles, L., Garraway, B., and Knight, P. (1994b). Generation of nonclassical light by dissipative two-photon processes. *Phys. Rev. A* 49, 2785–2799. doi:10.1103/PhysRevA.49.2785
- Gilles, L., and Knight, P. (1993b). Two-photon absorption and nonclassical states of light. *Phys. Rev. A* 48, 1582–1593. doi:10.1103/PhysRevA.48.1582
- Gilles, L., and Knight, P. L. (1993a). Two-photon absorption and nonclassical states of light. *Phys. Rev. A* 48, 1582–1593. doi:10.1103/PhysRevA.48.1582
- Giovannetti, V., Lloyd, S., and Maccone, L. (2011). Advances in quantum metrology. *Nat. Photon.* 23, 222–229. doi:10.1038/nphoton.2011.35
- Grajcar, M., Izmalkov, A., Il'ichev, E., Wagner, T., Oukhanski, N., Hübner, U., et al. (2004). Low-frequency measurement of the tunneling amplitude in a flux qubit. *Phys. Rev. B* 69, 060501. doi:10.1103/PhysRevB.69.060501
- Guerra, E. S., Garraway, B. M., and Knight, P. L. (1997). Two-photon parametric pumping versus two-photon absorption: a quantum jump approach. *Phys. Rev. A* 55, 3842–3857. doi:10.1103/PhysRevA.55.3842
- Habib, S., Shizume, K., and Zurek, W. H. (1998). Decoherence, chaos, and the correspondence principle. *Phys. Rev. Lett.* 80, 4361–4365. doi:10.1103/PhysRevLett.80.4361
- Hach, E. III, and Gerry, C. (1994). Generation of mixtures of Schrödinger-cat states from a competitive two-photon process. *Phys. Rev. A* 49, 490–498. doi:10.1103/PhysRevA.49.490
- Haycock, D. L., Alsing, P. M., Deutsch, I. H., Grondalski, J., and Jessen, P. S. (2000). Mesoscopic quantum coherence in an optical lattice. *Phys. Rev. Lett.* 85, 3365–3368. doi:10.1103/PhysRevLett.85.3365
- Hendrickson, S. M., Lai, M. M., Pittman, T. B., and Franson, J. D. (2010). Observation of two-photon absorption at low power levels using tapered optical fibers in rubidium vapor. *Phys. Rev. Lett.* 105, 173602. doi:10.1103/PhysRevLett.105.173602
- Hildred, G. P., and Hall, A. G. (1978). Antibunching: more than one kind in two-photon absorption. *J. Phys. A Math. Gen.* 11, L209. doi:10.1088/0305-4470/11/8/007
- Ikeda, Y., and Yamamoto, N. (2013). Deterministic generation of Gaussian pure states in a quasiloocal dissipative system. *Phys. Rev. A* 87, 033802. doi:10.1103/PhysRevA.87.033802
- Il'ichev, E., Smirnov, A. Y., Grajcar, M., Izmalkov, A., Born, D., Oukhanski, N., et al. (2004). Radio-frequency method for investigation of quantum properties of superconducting structures. *Fiz. Nizk. Temp.* 30, 823.
- Jeong, H., Kim, M. S., and Lee, J. (2001). Quantum-information processing for a coherent superposition state via a mixed entangled coherent channel. *Phys. Rev. A* 64, 052308. doi:10.1103/PhysRevA.64.052308
- Kraus, B., Büchler, H., Diehl, S., Kantian, A., Micheli, A., and Zoller, P. (2008). Preparation of entangled states by quantum Markov processes. *Phys. Rev. A* 78, 042307. doi:10.1103/PhysRevA.78.042307
- Krauter, H., Muschik, C. A., Jensen, K., Wasilewski, W., Petersen, J. M., Cirac, J. I., et al. (2011). Entanglement generated by dissipation and steady state entanglement of two macroscopic objects. *Phys. Rev. Lett.* 107, 080503. doi:10.1103/PhysRevLett.107.080503
- Kumar, S., and DiVincenzo, D. P. (2010). Exploiting Kerr cross nonlinearity in circuit quantum electrodynamics for nondemolition measurements. *Phys. Rev. B* 82, 014512. doi:10.1103/PhysRevB.82.014512
- Leghtas, Z., Vool, U., Shankar, S., Hatridge, M., Girvin, S. M., Devoret, M. H., et al. (2013). Stabilizing a Bell state of two superconducting qubits by dissipation engineering. *Phys. Rev. A* 88, 5. doi:10.1103/PhysRevA.88.023849
- Leibfried, D., Knill, E., Seidelin, S., Britton, J., Blakestad, R. B., Chiaverini, J., et al. (2005). Creation of a six-atom “Schrödinger cat” state. *Nature* 438, 639–642. doi:10.1038/nature04251
- Loudon, R., and Knight, P. (1987). Squeezed light. *J. Mod. Opt.* 34, 709–759. doi:10.1080/09500348714550721
- McNeil, K. J., and Walls, D. F. (1974). A master equation approach to nonlinear optics. *J. Phys. A Math. Gen.* 7, 617. doi:10.1063/1.4891470
- Monroe, C., Meekhof, D. M., King, B. E., and Wineland, D. J. (1996). A “Schrödinger cat” superposition state of an atom. *Science* 272, 1131–1136. doi:10.1126/science.272.5265.1131
- Munro, W. J., Nemoto, K., Milburn, G. J., and Braunstein, S. L. (2002). Weak-force detection with superposed coherent states. *Phys. Rev. A* 66, 023819. doi:10.1103/PhysRevA.66.023819
- Nakamura, Y., Pashkin, Y., and Tsai, J. (1999). Coherent control of macroscopic quantum states in a single-cooper-pair box. *Nature* 398, 786. doi:10.1038/19718
- Noel, M., and Stroud, J. (1996). Excitation of an atomic electron to a coherent superposition of macroscopically distinct states. *Phys. Rev. Lett.* 77, 1913–1916. doi:10.1103/PhysRevLett.77.1913
- Nogues, G., Rauschenbeutel, A., Osnaghi, S., Bertet, P., Brune, M., Raimond, J. M., et al. (2000). Measurement of a negative value for the Wigner function of radiation. *Phys. Rev. A* 62, 054101. doi:10.1103/PhysRevA.62.054101
- Ourjoumteva, A., Jeong, H., Tualle-Brouiri, R., and Grangier, P. (2007). Generation of optical “Schrödinger cats” from photon number states. *Nature* 448, 784–786. doi:10.1038/nature06054
- Pechen, A. (2011). Engineering arbitrary pure and mixed quantum states. *Phys. Rev. A* 84, 042106. doi:10.1103/PhysRevA.84.042106
- Ralph, T. C., Gilchrist, A., Milburn, G. J., Munro, W. J., and Glancy, S. (2003). Quantum computation with optical coherent states. *Phys. Rev. A* 68, 042319. doi:10.1103/PhysRevA.68.042319
- Sanders, B. C. (2012). Review of entangled coherent states. *J. Phys. A Math. Theor.* 45, 244002. doi:10.1088/1751-8113/45/24/244002
- Santamore, D. H., Goan, H.-S., Milburn, G. J., and Roukes, M. L. (2004). An harmonic effects on a phonon-number measurement of a quantum-mesoscopic-mechanical oscillator. *Phys. Rev. A* 70, 052105. doi:10.1103/PhysRevA.70.052105
- Schirmer, S. G., and Wang, X. (2010). Stabilizing open quantum systems by Markovian reservoir engineering. *Phys. Rev. A* 81, 062306. doi:10.1103/PhysRevA.81.062306

- Schlosshauer, M. (2005). Decoherence, the measurement problem, and interpretations of quantum mechanics. *Rev. Mod. Phys.* 76, 1267–1305. doi:10.1103/RevModPhys.76.1267
- Schrödinger, E. (1935). Die gegenwärtige situation in der quantenmechanik. *Naturwissenschaften* 23, 807–812. doi:10.1007/BF01491891
- Scully, M. O., Chapin, K. R., Dorfman, K. E., Kim, M. B., and Svidzinsky, A. (2011). Quantum heat engine power can be increased by noise-induced coherence. *Proc. Natl. Acad. Sci. U.S.A.* 108, 15097–15100. doi:10.1073/pnas.1110234108
- Simaan, H. D., and Loudon, R. (1975). Quantum statistics of single-beam two-photon absorption. *J. Phys. A Math. Gen.* 8, 539. doi:10.1088/0305-4470/8/4/016
- Simaan, H. D., and Loudon, R. (1978). Off-diagonal density matrix for single-beam two-photon absorbed light. *J. Phys. A Math. Gen.* 11, 435. doi:10.1088/0305-4470/11/2/018
- Stiffell, P. B., Everitt, M. J., Clark, T. D., Harland, C. J., and Ralph, J. F. (2005). Quantum downconversion and multipartite entanglement via a mesoscopic superconducting quantum interference device ring. *Phys. Rev. B* 72, 014508. doi:10.1103/PhysRevB.72.014508
- Ticozzi, F., Schirmer, S. G., and Wang, X. (2010). Stabilizing quantum states by constructive design of open quantum dynamics. *IEEE Trans. Automat. Contr.* 55, 2901–2905. doi:10.1109/TAC.2010.2079532
- Ticozzi, F., and Viola, L. (2012). Stabilizing entangled states with quasi-local quantum dynamical semigroups. *Philos. Trans. A Math. Phys. Eng. Sci.* 370, 5259–5269. doi:10.1098/rsta.2011.0485
- Tornau, N., and Bach, A. (1974). Quantum statistics of two-photon absorption. *Opt. Commun.* 11, 46–49. doi:10.1016/0030-4018(74)90330-7
- van Enk, S. J., and Hirota, O. (2001). Entangled coherent states: teleportation and decoherence. *Phys. Rev. A* 64, 022313. doi:10.1103/PhysRevA.64.022313
- Viola, L., Onofrio, R., and Calarco, T. (1997). Macroscopic quantum damping in squid rings. *Phys. Lett. A* 229, 23–31. doi:10.1016/S0375-9601(97)00154-0
- Vitali, D., Tombesi, P., and Milburn, G. J. (1997). Controlling the decoherence of a “meter” via stroboscopic feedback. *Phys. Rev. Lett.* 79, 2442–2445. doi:10.1103/PhysRevLett.79.2442
- Voje, A., Croy, A., and Isacsson, A. (2013a). Multi-phonon relaxation and generation of quantum states in a nonlinear mechanical oscillator. *New J. Phys.* 15, 053041. doi:10.1088/1367-2630/15/5/053041
- Voje, A., Isacsson, A., and Croy, A. (2013b). Nonlinear-dissipation induced entanglement of coupled nonlinear oscillators. *Phys. Rev. A* 88, 022309. doi:10.1103/PhysRevA.88.022309
- Wallquist, M., Shumeiko, V. S., and Wendin, G. (2006). Selective coupling of superconducting charge qubits mediated by a tunable stripline cavity. *Phys. Rev. B* 74, 224506. doi:10.1103/PhysRevB.74.224506
- Walls, D., and Milburn, G. (2008). *Quantum Optics (Springer Link)*. Berlin: Springer.
- Wielinga, B., and Milburn, G. J. (1993). Quantum tunneling in a Kerr medium with parametric pumping. *Phys. Rev. A* 48, 2494–2496. doi:10.1103/PhysRevA.48.2494
- Wolinsky, M., and Carmichael, H. J. (1988). Quantum noise in the parametric oscillator: from squeezed states to coherent-state superpositions. *Phys. Rev. Lett.* 60, 1836–1839. doi:10.1103/PhysRevLett.60.1836
- Yamamoto, N. (2012). Pure Gaussian state generation via dissipation: a quantum stochastic differential equation approach. *Philos. Trans. A Math. Phys. Eng. Sci.* 370, 5324–5337. doi:10.1098/rsta.2011.0529
- Yurke, B., Schleich, W., and Walls, D. (1990). Quantum superpositions generated by quantum nondemolition measurements. *Phys. Rev. A* 42, 1703–1711. doi:10.1103/PhysRevA.42.1703
- Yurke, B., and Stoler, D. (1986). Generating quantum mechanical superpositions of macroscopically distinguishable states via amplitude dispersion. *Phys. Rev. Lett.* 57, 13–16. doi:10.1103/PhysRevLett.57.13
- Zhang, W., Sun, C. P., and Nori, F. (2010). Equivalence condition for the canonical and microcanonical ensembles in coupled spin systems. *Phys. Rev. E* 82, 041127. doi:10.1103/PhysRevE.82.041127

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 11 August 2014; accepted: 04 October 2014; published online: 30 October 2014.

Citation: Everitt MJ, Spiller TP, Milburn GJ, Wilson RD and Zagoskin AM (2014) Engineering dissipative channels for realizing Schrödinger cats in SQUIDS. *Front. ICT* 1:1. doi: 10.3389/fict.2014.00001

This article was submitted to *Quantum Computing*, a section of the journal *Frontiers in ICT*.

Copyright © 2014 Everitt, Spiller, Milburn, Wilson and Zagoskin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.