

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.



CC creative commons  
COMMONS DEED

**Attribution-NonCommercial-NoDerivs 2.5**

**You are free:**

- to copy, distribute, display, and perform the work

**Under the following conditions:**

 **Attribution.** You must attribute the work in the manner specified by the author or licensor.

 **Noncommercial.** You may not use this work for commercial purposes.

 **No Derivative Works.** You may not alter, transform, or build upon this work.

- For any reuse or distribution, you must make clear to others the license terms of this work.
- Any of these conditions can be waived if you get permission from the copyright holder.

**Your fair use and other rights are in no way affected by the above.**

This is a human-readable summary of the [Legal Code \(the full license\)](#).

[Disclaimer](#) 

For the full text of this licence, please go to:  
<http://creativecommons.org/licenses/by-nc-nd/2.5/>



## **Origin and significance of dispersed facies basal ice: Svínafellsjökull, Iceland**

Simon Cook (1), Darrel Swift (2), David Graham (3), and Nicholas Midgley (4)

(1) Centre for Glaciology, Aberystwyth University, UK (smc@aber.ac.uk), (2) Department of Geography, University of Sheffield, UK, (3) Department of Geography, Loughborough University, UK, (4) School of Animal, Rural & Environmental Sciences, Nottingham Trent University, UK

Dispersed facies basal ice (massive ice with dispersed debris aggregates) outcrops at the margins of many ice masses and is important to glaciologists because of the information it provides about the nature of subglacial conditions and processes in the deep interior of glaciers and ice sheets. There has been little agreement, however, about how it forms with possible mechanisms including regelation and water flow through the intercrystalline vein network, strain-induced metamorphism of firnified glacier ice, shearing of basal debris-rich ice, freeze-on of subglacial water, and incorporation of surface debris into glacier ice. We test these established hypotheses at the temperate glacier Svínafellsjökull, southeast Iceland, and show that none fully account for dispersed facies characteristics here. From analysis of physical, sedimentological and stable isotope ( $\delta^{18}\text{O}$  and  $\delta\text{D}$ ) characteristics we suggest that dispersed facies forms from a combination of regelation and strain-induced metamorphism of debris-laden ice originally entrained by tectonic processes at the base of an icefall. We suggest that a terminal overdeepening may serve to further thicken dispersed facies as the glacier flows against a prominent reverse bed-slope. There may also be a lack of subglacial drainage across the overdeepening which further allows dispersed facies to survive in thicknesses up to  $\sim 20\text{m}$  despite the temperate location. Our results demonstrate that, despite its low sediment content ( $\sim 1.6\%$ ), the thick layer of dispersed facies contributes a higher annual sediment flux than other more debris-rich basal ice types. Hence dispersed facies and the processes that create it should not be overlooked in assessments of glacial sediment budgets.