

1 **Title:** Effect of team rank and player classification on activity profiles of elite wheelchair
2 rugby players

3 **Running Head:** Influence of team rank in wheelchair rugby

4 **Keywords:** Paralympic sports, performance, player tracking, mobility

5

1 **Abstract**

2 The aim of the current study was to establish which indicators of mobility are associated with
3 successful wheelchair rugby performance and determine whether these indicators differed
4 across classification. Data were collected from 11 international teams during 30 matches (353
5 match observations) using a radio-frequency based, indoor tracking system across two
6 tournaments. Players ($n = 111$) were first grouped by team rank as determined by their
7 International Wheelchair Rugby Federation (IWRF) world ranking (LOW, MID, HIGH) and
8 then into one of four groups based on their IWRF classification: group I (0.5), II (1.0-1.5), III
9 (2.0-2.5), IV (3.0-3.5). The volume of activity (relative distance and mean speed), peak speed,
10 and time spent within classification-specific arbitrary speed zones were calculated for each
11 individual. Although no differences were identified in the volume of activity, playing time
12 was significantly reduced in LOW ($34:51 \pm 8:35$) compared to MID ($48:54 \pm 0:51$) and
13 HIGH ($45:38 \pm 9:53$), which was further supported by the greater number of substitutions
14 performed by LOW. HIGH achieved greater peak speeds ($3.55 \pm 0.40 \text{ m}\cdot\text{s}^{-1}$) than LOW (3.27
15 $\pm 0.42 \text{ m}\cdot\text{s}^{-1}$) and MID ($3.45 \pm 0.41 \text{ m}\cdot\text{s}^{-1}$). Peak speed was further shown to be
16 classification-dependent ($P \leq 0.005$), whereby HIGH groups III and IV players achieved
17 greater peak speeds than LOW and MID. The time spent performing high-intensity activities
18 was also greater in HIGH compared to LOW and MID, whilst further influenced by
19 classification ($P \leq 0.0005$). To conclude, peak speed and the ability to perform a greater
20 number of high-intensity activities were associated with successful performance in
21 wheelchair rugby.

1 **Introduction**

2 Wheelchair rugby is an intermittent, court-based team sport characterised by frequent short-
3 term high-intensity demands superimposed on a background of aerobic activity. It consists of
4 4 x 8-minute quarters, played on an indoor wooden sprung surface (15 m x 28 m). The game-
5 clock is started once the ball is in-play and regulations restrict a team to a total of 40 s to
6 score otherwise they concede possession. Participating athletes generally have one of the
7 following conditions: spinal cord injury at the level of their cervical vertebrae, multiple
8 amputations, polio, neurological disorders such as cerebral palsy and some forms of muscular
9 dystrophy (Goosey-Tolfrey & Leicht, 2013). At present, wheelchair rugby players are
10 classified into one of seven classification groups based on their impairment, ranging from 0.5
11 (most impaired) to 3.5 (least impaired). Each team is permitted to field four players at any
12 one time, whereby the total number of points cannot exceed 8.0 points (Molik et al., 2008)
13 with continuous roll-on substitutions permitted.

14 Only recently have the match-play characteristics of elite wheelchair rugby received
15 scientific attention (Molik et al., 2008; Morgulec-Adamowicz et al., 2010; Rhodes, Mason,
16 Perrat, Smith, & Goosey-Tolfrey, 2014a; Rhodes et al., 2014b; Sarro, Misuta, Burkett,
17 Malone, & Barros, 2010). Early notational analysis data suggested that ‘high-point’ players
18 (2.0-3.5 classification group) generally perform better than ‘low-point’ players (0.5-1.5
19 classification group) in most of the ball-handling match activities such as points scored,
20 interceptions, passes made and passes caught (Molik et al., 2008; Morgulec-Adamowicz et al.,
21 2010). The close relationship between classification and on-court role may partially explain
22 such findings. Low-point players typically possess limited shoulder and wrist stability that
23 impede ball-handling capabilities as well as reduce wheelchair manoeuvrability skills,
24 restricting players to defensive on-court roles (Molik et al., 2008). Alternatively, high-point
25 players generally display good shoulder and wrist stability, enabling players to perform ball-
26 handling tasks and wheelchair handling skills effectively, which sees them occupy offensive
27 on-court roles (IWRP, 2014). Through recent developments in technology (Rhodes et al.,
28 2014a), information regarding the activity profiles during wheelchair rugby match-play have
29 been described. During competition, elite wheelchair rugby players typically cover distances
30 between 3500-4600 m (Rhodes et al., 2014b; Sarro, et al., 2010), with an average peak speed
31 of $3.48 \pm 0.36 \text{ m}\cdot\text{s}^{-1}$ (Rhodes et al., 2014b). Match-play has been further characterised by
32 prolonged low-intensity activities ($\leq 50\%$ peak speed) interspersed with frequent periods of
33 short (1.7-1.9 s) high-intensity activities (Rhodes et al., 2014b). Classification-dependant

1 trends in match performance were further highlighted; whereby greater total distance, mean
2 speed (Sarro et al., 2010) and peak speed values (Rhodes et al., 2014b) were reported as
3 classification group increased. Furthermore, when grouped by on-court role, notable trends in
4 the intensity of match-play activity were also evident (Rhodes et al., 2014b). Specifically,
5 defensive players spent a significantly greater amount of time performing very low speed
6 activities compared to offensive players, whilst, a greater number of high-intensity bouts
7 were exhibited by defensive players (~13) compared to offensive players (~9) (Rhodes et al.,
8 2014b). Such results may be attributed to the key requirements for the varying on-court roles.
9 These roles require defensive players to block and trap opponents resulting in longer
10 durations of very low speed activity, yet must perform high-intensity activities more
11 frequently to compete with more functionally able opponents. However, an understanding of
12 which aspects of mobility are associated with successful performance is required to further
13 future monitoring in the training environment.

14 Whilst key indicators of successful performance have been explored using team
15 rank in able-bodied sports, no such information exists for wheelchair rugby. In the only study
16 to investigate the influence of successful performance within an elite wheelchair sporting
17 application, greater peak speeds were reported in high-ranked wheelchair tennis players
18 (Sindall et al., 2013). While high peak speeds may be advantageous for wheelchair tennis
19 performance, it is important to acknowledge the classification and tactical roles associated
20 with individuals in wheelchair rugby that could influence this relationship. Therefore the aim
21 of the current study was to establish which aspects of mobility were associated with
22 successful performance as determined by team rank during elite wheelchair rugby. A
23 secondary aim was to determine whether the impact of mobility on performance was further
24 influenced by classification.

25 **Methods**

26 **Participants**

27 All National teams competing in the 2013 European and Americas Zonal Championships
28 were invited to participate in the current study. Out of the 15 competing teams, 11 agreed to
29 participate giving a sample of 111 elite International wheelchair rugby players (male: $n = 110$;
30 female: $n = 1$; age: 32 ± 7 years). Similar to previous investigations (Di Salvo et al., 2009;
31 Rampinini et al., 2007), players were subdivided into the following three groups according to

1 their teams International Wheelchair Rugby Federation (IWRF) ranking prior to the start of
2 both competitions: the top 3 ranked teams (HIGH); middle 5 ranked teams (MID); and the
3 lowest 3 ranked teams (LOW). Team ranking was taken prior to the start of both competitions
4 based on recommendations made by previous research (Castellano et al., 2014) and did not
5 change between competitions. Each player was assigned into one of four groups according to
6 their IWRF classification (Morgulec-Adamowicz et al., 2010; Rhodes et al., 2014b),
7 defensive players were categorised as groups I (0.5) and II (1.0-1.5), whilst offensive players
8 as groups III (2.0-2.5) and IV (3.0-3.5). Approval for the study was obtained from the IWRF
9 and the organising committee of each tournament in addition to the University's local ethical
10 advisory committee. All participants provided their written informed consent to participate in
11 the current investigation.

12 **Equipment**

13 Data were collected during all matches using a radio-frequency based indoor tracking system
14 (8 Hz; Ubisense, Cambridge, UK) as previously described and validated by Rhodes et al.
15 (2014a). When assessing total distance during a simulated match quarter (999 ± 65 m), the
16 indoor tracking system reported a relative error of $< 0.2\%$ compared against a total laser
17 station as the criterion measure. Furthermore, a mean systematic error of $0.05 \text{ m}\cdot\text{s}^{-1}$ for
18 measuring peak speed was reported during linear sprints in excess of $4.00 \text{ m}\cdot\text{s}^{-1}$. Each player
19 was equipped with a small, lightweight tag (size = $40 \times 40 \times 10$ mm; mass = 25 g), positioned
20 on or near the foot-strap of the players rugby wheelchair. All players were familiarised with
21 the tag locations during training sessions and practice matches prior to the start of the
22 competitions.

23 **Experimental design**

24 The indoor tracking system was installed on the main court of each tournament venue and
25 data were collected from a total of 30 matches. Each participating team was monitored
26 whenever they played on the main court (minimum of 3 matches, range 3-6), with data
27 collected during pool ($n = 20$), crossover ($n = 4$) and placement ($n = 6$) matches. Each match
28 involving a participating team was included for data collection with each team member
29 equipped with a radio-frequency tag. Up to 24 players (12 players from each team) wore a tag
30 during any given match, with a match observation characterised for each individual by the
31 accumulation of activity collected during the respective four quarters of that match (353

1 0.6), moderate (0.6-1.2), large (1.2-2.0), and very large (≥ 2.0) based on previous guidelines
2 (Batterham & Hopkins, 2006).

3 **Results**

4 No significant effect of team rank was observed for relative distance (Figure 1a; $P = 0.532$)
5 and mean speed (Figure 1b; $P = 0.538$). However, there was a significant difference between
6 mean playing time (mm:ss) and team rank ($P \leq 0.0005$), which was significantly reduced in
7 LOW ($34:51 \pm 8:35$) compared to MID ($48:54 \pm 0:51$; $P \leq 0.0005$; 95% CI = -245.7 to -157.8;
8 ES = 1.7) and HIGH ($45:38 \pm 9:53$; $P \leq 0.0005$; 95% CI = -136.1 to -44.0; ES = 1.2). The
9 number of substitutions performed was also shown to be influenced by team rank ($P \leq$
10 0.0005). LOW performed a greater number of substitutions per match (12 ± 4) than both MID
11 (4 ± 3 ; $P \leq 0.0005$; 95% CI = 4.7 to 10.6; ES > 2.0) and HIGH (5 ± 3 ; $P \leq 0.0005$; 95% CI =
12 2.9 to 9.8; ES = 1.7).

13 Peak speed was significantly affected by team rank ($P = 0.002$). As illustrated in
14 Figure 1c, HIGH achieved greater peak speeds ($3.56 \pm 0.40 \text{ m}\cdot\text{s}^{-1}$) compared to LOW ($3.27 \pm$
15 $0.42 \text{ m}\cdot\text{s}^{-1}$; $P \leq 0.0005$; 95% CI = -0.4 to -0.1; ES = 0.7) and MID ($3.45 \pm 0.41 \text{ m}\cdot\text{s}^{-1}$; $P =$
16 0.003 ; 95% CI = 0.1 to 0.2; ES = 0.3). The relative time spent within low, high and very high
17 speed zones were also significantly influenced by team rank ($P \leq 0.0005$). Figure 2 reveals
18 LOW ($52.3 \pm 7.0\%$) spent more time in the low speed zone compared to MID ($46.7 \pm 7.9\%$;
19 $P \leq 0.0005$; 95% CI = 3.0 to 8.1; ES = 0.7) and HIGH ($46.8 \pm 7.6\%$; $P \leq 0.0005$; 95% CI =
20 2.9 to 8.1; ES = 0.8). However, HIGH spent greater time within high ($2.9 \pm 1.6\%$) and very
21 high ($0.7 \pm 0.8\%$) speed zones compared to LOW ($1.5 \pm 1.1\%$ and $0 \pm 0.4\%$; $P \leq 0.0005$; ES
22 = 0.9-1.0) and MID ($2.0 \pm 1.3\%$ and $0.3 \pm 0.5\%$; $P \leq 0.025$; ES = 0.6). High-intensity
23 activities were also significantly influenced by team rank ($P \leq 0.0005$). As shown in Table 2,
24 HIGH performed a greater number of relative high-intensity activities compared to LOW (P
25 ≤ 0.0005 ; 95% CI = -0.5 to -0.2; ES = 1.4) and MID ($P = 0.006$; 95% CI = -0.3 to -0.04; ES =
26 0.8). Whilst HIGH also covered greater mean ($P \leq 0.001$; ES = 0.5-0.8) and max distances (P
27 ≤ 0.006 ; ES = 0.6-1.1), for a longer mean ($P \leq 0.0005$; ES = 0.8-1.0) and max duration ($P \leq$
28 0.008 ; ES = 0.5-1.1) at high-intensities compared to both LOW and MID.

29 ***INSERT FIGURE 1 & 2 HERE***

30 ***INSERT TABLE 2***

1 No significant interaction was observed between team rank and classification group
2 for relative distance ($P = 0.141$) or mean speed ($P = 0.102$). However, classification group
3 was shown to influence peak speed values across team rank ($P = 0.008$). Table 3 reveals
4 HIGH achieved significantly greater peak speeds compared to LOW across all classification
5 groups ($P \leq 0.001$; ES = 0.6-1.5), whilst HIGH groups III and IV players achieved greater
6 peak speeds compared to respective MID players ($P \leq 0.005$; ES = 0.7-0.8). A significant
7 interaction was observed across low ($P = 0.009$), high ($P \leq 0.0005$) and very high ($P \leq$
8 0.0005) speed zones, whilst a significant interaction also existed for the high-intensity
9 activities performed during match-play ($P \leq 0.0005$). Post hoc analyses revealed:

- 10 • Group I: LOW players spent significantly greater time within the low speed zone
11 compared to MID ($P \leq 0.0005$; 95% CI = 3.0 to 12.3; ES = 1.4) and HIGH ($P \leq$
12 0.0005 ; 95% CI = 2.1 to 11.2; ES = 1.3). Whilst LOW and MID spent significantly
13 less time in the high ($P \leq 0.0005$; ES = 1.0-1.3) and very high speed zones ($P \leq 0.029$;
14 ES = 1.1-1.4) compared to HIGH. LOW displayed a significant difference with HIGH
15 for all high-intensity activities ($P \leq 0.005$; ES = 1.0-1.5), whilst the relative number
16 ($P = 0.002$; 95% CI = -0.5 to -0.1; ES = 0.7), max distance ($P = 0.027$; 95% CI = -7.8
17 to -0.9; ES = 0.9) and max duration ($P = 0.038$; 95% CI = 0.5 to 1.8; ES = 0.9) of
18 high-intensity activities significantly differed between LOW and MID. MID
19 performed significantly less high-intensity activities compared to HIGH ($P \leq 0.008$;
20 ES = 0.6-1.0).
- 21 • Group II: LOW players spent significantly less time in high and very high speed
22 zones as opposed to MID ($P = 0.006$; ES = 0.6-0.8) and HIGH ($P = 0.07$; ES = 0.8-
23 1.0), whilst MID spent significantly less time in the high speed zone than HIGH ($P =$
24 0.003 ; 95% CI = 0.3 to 1.3; ES = 0.5). LOW were shown to perform significantly less
25 relative number of high-intensity activities compared to MID ($P = 0.004$; 95% CI = -
26 0.7 to -0.1; ES = 1.1) and HIGH ($P \leq 0.0005$; 95% CI = -0.9 to -0.3; ES = 1.7).
- 27 • Group III: LOW and MID players were found to spend significantly less time in high
28 ($P \leq 0.023$; ES = 0.7-0.8) and very high speed zones ($P \leq 0.026$; ES = 1.1-1.2) as
29 opposed to HIGH. All parameters of high-intensity activities were also shown to be
30 significantly lower in LOW ($P \leq 0.001$; ES = 0.5-0.8) and MID ($P \leq 0.006$; ES = 0.4-
31 0.8) compared to HIGH.
- 32 • Group IV: MID were shown to spend significantly less time within high speed zones
33 ($P = 0.022$; 95% CI = 0.2 to 2.0; ES = 0.9) compared to HIGH. Although the relative

1 number of high-intensity activities did not differ between team ranks ($P \geq 0.174$; $ES \leq$
2 0.2), LOW and MID were found to cover less mean ($P \leq 0.0005$; $ES = 1.1-2.0$) and
3 max distances ($P \leq 0.001$; $ES = 1.0-2.0$), for lower mean ($P \leq 0.0005$; $ES = 1.3-2.0$)
4 and max durations at high-intensities ($P \leq 0.001$; $ES = 1.0-2.0$) compared to HIGH.

5 ***INSERT TABLE 3***

6 **Discussion**

7 The current study was the first to examine the influence of team rank on the activity profiles
8 of elite wheelchair rugby players during competition to establish which aspects of mobility
9 are critical to successful performance. Although the volume of activity was largely
10 uninfluenced by team rank, peak speeds and the capacity to perform at high-intensities were
11 both found to be associated with successful performance in wheelchair rugby. Activity
12 profiles were further influenced by classification and on-court role, as demonstrated by the
13 significantly higher peak speed values observed for HIGH offensive players (2.0-3.5). High-
14 intensity activities were also shown to be important and classification-dependant, with greater
15 time spent within very high speed zones observed in HIGH group I and offensive players.
16 Such results demonstrate which aspects of mobility performance were most associated with
17 successful performance in wheelchair rugby, which may have implications on future training
18 prescription and performance monitoring.

19 The current study revealed large differences in playing time between team ranks. The
20 shorter playing time of LOW suggests that players lack the physical capacity to maintain
21 performance over prolonged durations, which was further supported by the greater number of
22 substitutions performed by LOW. Consequently, coaching strategies designed to maximise
23 physical capacity may improve the match performance of lower-ranked teams. Nevertheless,
24 the relative distance covered, along with mean speed were not significantly different between
25 MID and HIGH. Therefore, it appeared that successful performance in wheelchair rugby was
26 not influenced by the volume of activity performed. Even when categorised by classification,
27 the volume of activity performed was largely unaffected by team rank. The comparable
28 activity volume of all players across team ranks reported here suggests association to
29 successful performance is negligible. Despite this, the performance of Paralympic court-
30 based sport players has previously been shown to be highly dependent upon aerobic fitness
31 (Bernardi et al., 2010). Therefore, elite wheelchair rugby players should be sufficiently

1 prepared so that they can meet the activity demands (3500-4600 m) required for competition
2 (Rhodes et al., 2014b; Sarro et al., 2010).

3 Since opponents can dictate a player's movement on-court, it was anticipated that the
4 ability to frequently reach high speeds and sustain high-intensity activities would be restricted.
5 Previous research has suggested that sprinting performance and the ability to reach high peak
6 speeds to be less of a priority in wheelchair rugby compared to acceleration and
7 manoeuvrability performance (Mason, Porcellato, van der Woude, & Goosey-Tolfrey, 2010).
8 That said, our study found HIGH achieved greater peak speeds ($3.59 \pm 0.44 \text{ m}\cdot\text{s}^{-1}$) than both
9 LOW ($3.31 \pm 0.49 \text{ m}\cdot\text{s}^{-1}$) and MID ($3.46 \pm 0.43 \text{ m}\cdot\text{s}^{-1}$). This supports and extends previous
10 knowledge gleaned from wheelchair tennis (Sindall et al., 2013). Furthermore, although the
11 majority of activity during wheelchair rugby is spent at low-intensities (~75%) (Rhodes et al.,
12 2014b), the current study established that players from HIGH spent a greater proportion of
13 time performing high-intensity activities compared to players from lower ranked teams. One
14 likely explanation that is difficult to quantify from the current data is that HIGH prevented
15 the opposition from achieving high peak speeds and sustaining high-intensity activities by
16 adopting full-court press tactics. Such tactics work by pressurising the ball-handler and
17 reducing the on-court space using 'trapping' techniques (Malone & Orr, 2010). On the other
18 hand, it is possible that HIGH players may be more capable of creating court space in-order
19 to perform higher peak speeds and a greater number of high-intensity activities. Although
20 team efficiency and playing style may account for some differences between team ranks,
21 future notational analysis techniques are required to establish this information with regard to
22 positional transitions, ball possession, and court zones etc. Nevertheless the current findings
23 reveal that success in wheelchair rugby can be characterised by a player's ability to
24 consistently reach high peak speeds, whilst performing at high-intensities and therefore
25 training and game-patterns should be structured to promote this.

26 Our findings were able to distinguish differences between offensive and defensive
27 roles and highlight the increased importance of peak speed for offensive players. As such,
28 HIGH offensive players achieved greater peak speeds than MID offensive players, whilst no
29 differences existed between MID and HIGH defensive players. Previous research has
30 demonstrated that the majority of points (~88%) are scored by offensive players in
31 wheelchair rugby (Molik et al., 2008; Morgulec-Adamowicz et al., 2010). Subsequently the
32 capacity to achieve superior peak speeds could prove pivotal to perform this role effectively
33 and influence team success. Alternatively, peak speed may be less important for defensive

1 players, whereby tactical aspects of performance may be more of a necessity. This could be
2 associated with differences in equipment between roles, whereby defensive players typically
3 use a wheelchair with a substantial rear-wheel camber (Keogh, 2011). While a greater camber
4 increases their stability and blocking ability, this comes at the expense of peak speed (Faupin
5 Campillo, Weissland, Gorce, & Thevenon, 2004; Mason, Woude, Tolfrey, & Goosey-Tolfrey,
6 2011). Such findings therefore further reiterate the need for role-specific training (Rhodes et
7 al., 2014b) and also identify this parameter as one of the key performance indicators for talent
8 identification purposes.

9 The magnitude of differences in high-intensity activities was found to be
10 classification-dependant, whereby HIGH group I and offensive players spent significantly
11 greater time within very high speed zones and were able to sustain these activities for longer
12 compared to respective MID players. It could be suggested that players at the highest level of
13 wheelchair rugby have the physical capacity to maintain repeated high-intensity activities
14 during match-play. Additionally, it is plausible that like peak speeds, HIGH players are more
15 capable of finding court space in-order to maintain repeated high-intensity activities.
16 Subsequently, training strategies aimed at sustaining high-intensity activities under the
17 pressure of opponents may be beneficial for offensive players. Alternatively, the time spent
18 within the very high speed zone and the ability to sustain high-intensity activities were not
19 shown to differ in group II players between MID and HIGH. Subsequently, given that
20 previous research identified no differences in ball-handling patterns between groups I and II
21 (Morgulec-Adamowicz et al., 2010), such results may imply that the mobility characteristics
22 of group I players could be more critical to successful team performance than group II
23 players, whilst subsequently reducing the total on-court classification points (8.0 points
24 permitted at any one time). Nevertheless, a technical analysis of wheelchair rugby is further
25 required to supplement the activity profiles currently presented to gain a holistic appraisal of
26 the sport.

27 Team line-up is an additional factor that may influence the activity profiles during
28 match-play. The current data would suggest LOW and HIGH teams generally utilised group
29 III players, as opposed to MID teams that typically employed more group II and IV players
30 during match-play. It is recommended however that future research investigates the effect of
31 different line-up strategies (i.e. mid-point vs. high- and low-point line-ups) on activity
32 profiles and performance in wheelchair rugby. In elite athletes with tetraplegia, peak heart

1 rate response is generally reduced (Goosey-Tolfrey & Leicht, 2013), consequently the
2 collection of heart rate in wheelchair rugby players is therefore questionable and methods
3 such as rating of perceived exertion (RPE) may be better advocated. Despite this, a limitation
4 of the current study was the inability to examine the individual physiological responses in
5 relation to the determination of speed zones. Future work utilising individualised
6 physiological measures (e.g. blood lactate) alongside the traditional arbitrary approach is
7 recommended (Hunter et al., 2014). Nevertheless, normalising speed zones based on match-
8 play sprinting capacity may reflect an ecologically valid approach to between-player and rank
9 comparisons. Moreover, the categorising of movement into speed zones could further be used
10 to identify individual work:rest ratios. Such information would provide coaches with
11 important information that could implemented into future training strategies and is
12 subsequently worthy of further investigation. Finally, although high-intensity activities were
13 deemed an important indicator of performance in wheelchair rugby, a limitation with the
14 current study is that the indoor tracking system used cannot quantify acceleration
15 performance due to the restricted sampling frequency. Therefore future research would
16 benefit from the incorporation of accelerometry to provide a more in-depth insight into high-
17 intensity activities, which may occur at low speeds, in wheelchair rugby.

18 **Conclusion**

19 The data provide new insights into the possible influence of successful performance on
20 activity profiles and highlights the impact of classification. The capacity to reach higher peak
21 speeds and to perform increased activities at high-intensities was associated with successful
22 performance in wheelchair rugby. These variables were further influenced by classification,
23 specifically in group I and offensive players (groups III & IV). Although the volume of
24 activity appeared uninfluenced by team rank, player conditioning appeared important since
25 LOW performed more substitutions and consequently averaged shorter playing durations.

1 **References**

- 2 Batterham, A., & Hopkins, W. (2006). Making meaningful inferences about magnitudes.
3 *International Journal of Sports Physiology and Performance*, *1*, 50-57.
- 4 Bernardi, M., Guerra, E., Di Giacinto, B. Di Cesare, A. Castellano, V., & Bhambhani Y.
5 (2010). Field evaluation of Paralympic athletes in selected sports: implications for training.
6 *Medicine and Science in Sports and Exercise*, *42*, 1200-1208.
- 7 Cahill, N., Lamb, K., Worsfold, P., Headey, R., & Murray, S. (2013). The movement
8 characteristics of English Premiership rugby union players. *Journal of Sports Sciences*, *31*,
9 229-237.
- 10 Castellano, J., Alvarez, D., & Bradley, P. (2014). Evaluation of research using computerized
11 tracking systems (Amisco and Prozone) to analyse physical performance in elite soccer: a
12 systematic review. *Sports Medicine*, *44*, 701-712.
- 13 Cnaan, A., Laird, N.M., & Slasor, P. (1997). Using the general linear mixed model to analyse
14 unbalanced repeated measures and longitudinal data. *Statistics in Medicine*, *16*, 2349-2380.
- 15 Faupin, A., Campillo, P., Weissland, T., Gorce, P., & Thevenon, A. (2004) The effects of
16 rear-wheel camber on the mechanical parameters produced during wheelchair sprinting of
17 handibasketball athletes. *Journal of rehabilitation research and development*, *41*, 421-428.
- 18 Goosey-Tolfrey, V.L., & Leicht, C.A. (2013). Field-based physiological testing of wheelchair
19 athletes. *Sports Medicine*, *43*, 77-91.
- 20 Hunter, F., Bray, J., Towlson, C., Smith, M., Barrett, S., Madden, J., Abt, G., & Lovell, R.
21 (2014). Individualisation of time-motion analysis: a method comparison and case report
22 series. *International Journal of Sports Medicine*, [Epub ahead of print].
- 23 International Wheelchair Rugby Federation. IWRF (online). <http://www.iwrf.com>. Accessed
24 1 Nov 2014.
- 25 Keogh, J.L. (2011) Paralympic sport: an emerging area for research and consultancy in sports
26 biomechanics. *Sports Biomechanics*, *10*, 234-253.
- 27 Mason, B.S., Porcellato, L., van der Woude, L.H., & Goosey-Tolfrey, V.L. (2010). A
28 qualitative examination of wheelchair configuration for optimal mobility performance in
29 wheelchair sports: A pilot study. *Journal of Rehabilitation Medicine*, *42*, 141-149.
- 30 Mason, B.S., van der Woude, L.H., Tolfrey, K., & Goosey-Tolfrey, V.L. (2011). The effects
31 of rear-wheel camber on maximal effort mobility performance in wheelchair athletes.
32 *International Journal of Sports Medicine*, *33*, 199-204.
- 33 Molik, B., Lubelska, E., Kosmol, A., Bogdan, M., Yilla, A., & Hyla, E. (2008). An
34 examination of the international wheelchair rugby federation classification system utilizing
35 parameters of offensive game efficiency. *Adapted Physical Activity Quarterly*, *25*, 335-351.

- 1 Morgulec-Adamowicz, N., Kosmol, A., Bogdan, M., Molik, B., Rutkowska, I., &
2 Bednarczuk, G. (2010). Game efficiency of wheelchair rugby athletes at the 2008 Paralympic
3 Games with regard to player classification. *Human Movement, 11*, 29-36.
- 4 Rhodes, J.M., Mason, B.S., Perrat, B., Smith, M., & Goosey-Tolfrey, V.L. (2014a). The
5 validity and reliability of a novel indoor player tracking system for use within wheelchair
6 court sports. *Journal of Sports Sciences, 32*, 1639-1647.
- 7 Rhodes, J.M., Mason, B.S., Perrat, B., Smith, M., Malone, L.A., & Goosey-Tolfrey, V.L.
8 (2014b). Activity profiles of elite wheelchair rugby players during competition. *International*
9 *Journal of Sports Physiology and Performance, [Epub ahead of print.]*
- 10 Sarro, K.J., Misuta, M.S., Burkett, B., Malone, L.A., & Barros, R.M.L. (2010). Tracking of
11 wheelchair rugby players in the 2008 demolition derby final. *Journal of Sports Sciences, 28*,
12 193-200.
- 13 Sindall, P., Lenton, J.P., Tolfrey, K., Cooper, R.A., Oyster, M., & Goosey-Tolfrey, V.L.
14 (2013). Wheelchair tennis match-play demands: Effect of player rank and result.
15 *International Journal of Sports Physiology and Performance, 8*, 28-37.
- 16

1 **Figure Legends**

2 **Figure 1.** Relative distance (a), mean speed (b), and peak speed (c) in relation to team rank
3 during match-play. Data presented as means \pm SD. *Significantly different to MID.
4 #Significantly different to HIGH.

5 **Figure 2.** Match intensity in relation to team rank during a typical match. Data presented as
6 means \pm SD. *Significantly different to MID. #Significantly different to HIGH.

Table 1. Classification-specific arbitrary speed zones ($\text{m}\cdot\text{s}^{-1}$) previously proposed for wheelchair rugby

	Classification group			
	I	II	III	IV
Very low	≤ 0.60	≤ 0.69	≤ 0.73	≤ 0.76
Low	0.61-1.50	0.70-1.72	0.74-1.84	0.77-1.91
Moderate	1.51-2.39	1.73-2.75	1.85-2.94	1.92-3.06
High	2.40-2.84	2.76-3.27	2.95-3.49	3.07-3.63
Very High	> 2.84	> 3.27	> 3.49	> 3.63

Classification groups: I = 0.5; II = 1.0-1.5; III = 2.0-2.5; IV = 3.0-3.5.

Table 2. High-intensity activities (mean \pm SD) during a typical wheelchair rugby match in relation to team rank

		Team Rank		
		LOW (<i>n</i> = 79)	MID (<i>n</i> = 145)	HIGH (<i>n</i> = 129)
High-intensity Activities	Relative Number (<i>n</i>)	$0.4 \pm 0.3^{*\#}$	$0.6 \pm 0.4^{\#}$	0.9 ± 0.4
	Mean Distance (m)	$4.5 \pm 2.8^{\#}$	$5.3 \pm 2.6^{\#}$	6.7 ± 2.7
	Max Distance (m)	$7.7 \pm 5.2^{*\#}$	$11.3 \pm 5.7^{\#}$	14.1 ± 6.1
	Mean Duration (s)	$1.4 \pm 0.8^{\#}$	$1.6 \pm 0.8^{\#}$	2.2 ± 0.8
	Max Duration (s)	$2.4 \pm 1.6^{*\#}$	$3.4 \pm 1.7^{\#}$	4.2 ± 1.8

n = number of match observations. *Significantly different to MID. [#]Significantly different to HIGH.

Table 3. Activity profiles during a typical wheelchair rugby match categorised by team rank and classification

	Team Rank											
	LOW				MID				HIGH			
	Defensive		Offensive		Defensive		Offensive		Defensive		Offensive	
	I (n = 13)	II (n = 13)	III (n = 42)	IV (n = 11)	I (n = 22)	II (n = 53)	III (n = 24)	IV (n = 46)	I (n = 23)	II (n = 24)	III (n = 68)	IV (n = 14)
Relative Distance (m·min ⁻¹)	60.1 ± 5.2	66.0 ± 3.9	74.9 ± 15.0	80.6 ± 11.4	59.0 ± 8.1	69.1 ± 8.1	77.4 ± 7.5	76.4 ± 10.4	63.2 ± 6.9	71.7 ± 14.2	76.8 ± 8.2	81.3 ± 6.4
Mean Speed (m·s ⁻¹)	1.00 ± 0.09	1.10 ± 0.07	1.25 ± 0.25	1.34 ± 0.19	0.98 ± 0.13	1.15 ± 0.14	1.29 ± 0.12	1.28 ± 0.17	1.05 ± 0.12	1.17 ± 0.21	1.26 ± 0.14	1.35 ± 0.11
Peak Speed (m·s ⁻¹)	2.60 ± 0.15 ^{*#}	3.13 ± 0.27 [#]	3.45 ± 0.30 [#]	3.57 ± 0.25 ^{*#}	2.91 ± 0.26	3.36 ± 0.35	3.51 ± 0.22 [#]	3.73 ± 0.35 [#]	3.08 ± 0.25	3.45 ± 0.30	3.70 ± 0.32	3.92 ± 0.26
Very Low (%)	33.3 ± 8.0	36.0 ± 5.7	30.0 ± 8.6	27.2 ± 13.2	35.7 ± 9.6	34.6 ± 9.3	26.9 ± 7.5	31.1 ± 11.8	33.7 ± 6.5	34.7 ± 12.1	29.7 ± 8.5	28.5 ± 4.7
Low (%)	48.2 ± 5.5 ^{*#}	46.2 ± 5.5	55.0 ± 6.7	53.1 ± 7.7	40.6 ± 5.7	43.9 ± 9.3	52.4 ± 6.2	50.5 ± 7.1	41.6 ± 4.9	40.4 ± 7.1	49.6 ± 6.5	52.7 ± 4.8
Moderate (%)	20.0 ± 4.1	18.5 ± 4.1	17.6 ± 4.1	19.9 ± 7.6	20.4 ± 4.4	21.0 ± 3.6	20.6 ± 4.6	18.2 ± 4.5	21.2 ± 5.1	21.5 ± 7.3	18.9 ± 4.4	18.4 ± 3.5
High (%)	1.5 ± 1.3 [#]	1.1 ± 0.9 ^{*#}	1.6 ± 1.0 [#]	2.2 ± 2.1	2.6 ± 1.8 [#]	2.1 ± 1.2 [#]	1.6 ± 0.8 [#]	1.9 ± 1.2 [#]	4.1 ± 1.9	3.0 ± 1.8	2.5 ± 1.3	3.0 ± 1.2
Very High (%)	0.1 ± 0.3 [#]	0.1 ± 0.1 ^{*#}	0.3 ± 0.2 [#]	0.5 ± 0.1	0.3 ± 0.3 [#]	0.4 ± 0.4	0.2 ± 0.1 [#]	0.4 ± 0.5	1.1 ± 1.0	0.6 ± 0.7	0.6 ± 0.8	0.7 ± 0.5
Relative Number (min ⁻¹)	0.3 ± 0.3 ^{*#}	0.3 ± 0.2 ^{*#}	0.4 ± 0.3 [#]	0.6 ± 0.5	0.8 ± 0.5 [#]	0.7 ± 0.4	0.5 ± 0.2 [#]	0.6 ± 0.4	1.1 ± 0.5	1.0 ± 0.5	0.8 ± 0.4	0.7 ± 0.2
Mean Distance (m)	3.3 ± 2.5 [#]	4.0 ± 3.2	4.9 ± 3.1 [#]	4.8 ± 1.9 [#]	4.4 ± 3.1 [#]	4.8 ± 2.3	5.2 ± 2.0 [#]	6.1 ± 2.9 [#]	6.5 ± 2.1	5.8 ± 2.2	6.5 ± 3.1	8.6 ± 1.6
Max Distance (m)	5.1 ± 3.8 ^{*#}	6.3 ± 5.4	8.7 ± 5.5 [#]	8.9 ± 4.1 [#]	8.9 ± 4.3 [#]	11.6 ± 5.9	10.1 ± 5.1 [#]	12.0 ± 6.3 [#]	13.5 ± 5.1	13.5 ± 5.8	12.9 ± 6.6	17.2 ± 4.3
Mean Duration (s)	1.3 ± 0.9 [#]	1.3 ± 1.0	1.5 ± 0.9 [#]	1.4 ± 0.5 [#]	1.5 ± 1.1 [#]	1.6 ± 0.7	1.5 ± 0.6 [#]	1.7 ± 0.8 [#]	2.4 ± 0.8	0.6 ± 1.8	1.9 ± 1.0	2.5 ± 0.4
Max Duration (s)	1.9 ± 1.3 ^{*#}	2.0 ± 1.7 ^{*#}	2.6 ± 1.7 [#]	2.5 ± 1.2 [#]	3.3 ± 1.6 [#]	3.7 ± 1.8	2.9 ± 1.5 [#]	3.4 ± 1.7 [#]	4.6 ± 1.7	4.2 ± 1.7	3.8 ± 2.0	4.8 ± 1.0

n = number of match observations.

*Significantly different to MID. #Significantly different to HIGH.