

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.



**CC creative commons**  
COMMONS DEED

**Attribution-NonCommercial-NoDerivs 2.5**

**You are free:**

- to copy, distribute, display, and perform the work

**Under the following conditions:**

**BY:** **Attribution.** You must attribute the work in the manner specified by the author or licensor.

**Noncommercial.** You may not use this work for commercial purposes.

**No Derivative Works.** You may not alter, transform, or build upon this work.

- For any reuse or distribution, you must make clear to others the license terms of this work.
- Any of these conditions can be waived if you get permission from the copyright holder.

**Your fair use and other rights are in no way affected by the above.**

This is a human-readable summary of the [Legal Code \(the full license\)](#).

[Disclaimer](#) 

For the full text of this licence, please go to:  
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

# Zhang&Das\_MNConference\_Abstarct\_Experiment\_FinalMicro

## needle assisted micro-particle delivery: experiments using a skin mimicking agarose gel

Dongwei Zhang, Diganta B Das

Chemical Engineering Department, Loughborough University, Loughborough, UK  
(Emails: [148271236@163.com](mailto:148271236@163.com); [d.b.das@lboro.ac.uk](mailto:d.b.das@lboro.ac.uk))

Micro-particle delivery system is an advanced technology for transfer DNA loaded micro-particles to the target tissue. Cell damage was found as a problem for micro-particle delivery systems due to the impaction of high-velocity micro-particles<sup>1,2</sup>. A concept of combining microneedles with micro-particle delivery system, namely, microneedle assisted micro-particle delivery<sup>3-6</sup> is presented in this case. The preliminary conception is that a number of micro-particles are able to penetrate through the pierced holes (created by microneedles) to reach the target area. Thus, the particle velocity can be reduced appropriately by using this system. The system is consisted of three different stage, which are acceleration, separation and deceleration stage. In the experiment, an agarose gel was chosen as a model target for this system. Two different sizes of solid microneedle patches (Adminpatch) were bought from nanoBioSciences Limited Liability Company (LLC) (Sunnyvale, CA, USA) and used in the experiment. In addition, the penetration depths of micro-particles were analyzed in relation to a number of variables, namely, the operating pressure, the particle size, the MN dimensions. The result of micro-particle penetration in the gel was caught by a digital optical microscope. As shown in Fig.1, a number of stainless steel micro-particles penetrated through the pierced holes and reached a further penetration depth inside gel. In this case, stainless steel micro-particles are accelerated to approximately 130 m/s, which is significantly lower than other systems (e.g. PowderJet<sup>7</sup>, Contoured shock tube<sup>8</sup>). Finally, the results show that the penetration depth increases from an increase in particle size, operating pressure and MN length. The above results directly show the feasibility of microneedle assisted micro-particle delivery.




Fig. 1: Stainless steel micro-particle penetration in the agarose gel<sup>4</sup>

### References

1. O'Brien J.A., Lummis S.C.R., (2011). Nano-biolistics: a method of biolistic transfection of cells and tissues using a gene gun with novel nanometer-sized projectiles, *BMC Biotechnology*, 11; 66

2. Uchida M., Li X.W., Mertens P., Alpar H.O. (2009). Transfection by particle bombardment: Delivery of plasmid DNA into mammalian cells using gene gun, *Biochim. Biophys. Acta.*, **1790**(8): 754-764
3. Zhang D.W., Das D.B., Rielly C.D. (2013). An experimental study of microneedle assisted micro-particle delivery, *Journal of Pharmaceutical Sciences*, 102(10): 3632-3644, DOI: 10.1002/jps.23665.
4. Zhang D.W., Das D.B., Rielly C.D. (2014). Microneedle assisted micro-particle delivery from gene guns: Experiments using skin-mimicking agarose gel. *Journal of Pharmaceutical Sciences*, 103(2): 613-627, DOI: 10.1002/jps.23835.
5. Zhang D.W., Das D.B., Rielly C.D. (2013). Potential of microneedle assisted micro-particle delivery by gene guns: A review. *Drug Delivery*, DOI: 10.3109/10717544.2013.864345.
6. Zhang D.W., Das D.B., Rielly C.D. (2014). Microneedle assisted micro-particle delivery by gene gun: Experiments and modelling on the effects of particle characteristics. *Drug delivery* (accepted), DOI: 10.3109/10717544.2014.887158 (in press).
7. Bellhouse B.J., Greenford J.C., Sarphe D.F. (2006). Particle delivery, Patent: EP 1637173 A2
8. Liu Y., Kendall M.A.F., (2006). Numerical analysis of gas and micro-particle interactions in a hand-held shock tube device, *Biomed Microdevices*, **8**: 341-351