



This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.

 **creative commons**  
C O M M O N S D E E D

**Attribution-NonCommercial-NoDerivs 2.5**

**You are free:**

- to copy, distribute, display, and perform the work

**Under the following conditions:**

 **Attribution.** You must attribute the work in the manner specified by the author or licensor.

 **Noncommercial.** You may not use this work for commercial purposes.

 **No Derivative Works.** You may not alter, transform, or build upon this work.

- For any reuse or distribution, you must make clear to others the license terms of this work.
- Any of these conditions can be waived if you get permission from the copyright holder.

**Your fair use and other rights are in no way affected by the above.**

This is a human-readable summary of the [Legal Code \(the full license\)](#).

[Disclaimer](#) 

For the full text of this licence, please go to:  
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368


Editors: W.T.A. Harrison, J. Simpson and  
M. Weil

## *N*-Phenyl-*tert*-butanesulfinamide

Mritunjoy Datta, Alan J. Buglass and Mark R. J. Elsegood

*Acta Cryst.* (2009). E65, o2034

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

## N-Phenyl-tert-butanesulfinamide

Mritunjoy Datta,<sup>a</sup> Alan J. Buglass<sup>a\*</sup> and Mark R. J. Elsegood<sup>b</sup>

<sup>a</sup>Department of Chemistry, Korea Advanced Institute of Science and Technology, Daejeon 305-701, Republic of Korea, and <sup>b</sup>Chemistry Department, Loughborough University, Loughborough LE11 3TU, England  
Correspondence e-mail: ajbuglass@kaist.ac.kr


Received 11 July 2009; accepted 20 July 2009

Key indicators: single-crystal X-ray study;  $T = 150$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.034;  $wR$  factor = 0.092; data-to-parameter ratio = 25.2.

In the racemic title compound,  $\text{C}_{10}\text{H}_{15}\text{NOS}$ , the packing exhibits centrosymmetric pairs of molecules linked by  $\text{N}-\text{H}\cdots\text{O}=\text{S}$  hydrogen bonds in a head-to-tail fashion. The  $\text{N}-\text{C}_{\text{aryl}}$  bond [1.4083 (12) Å] is considerably shorter than the  $\text{N}-\text{C}_{\text{alkyl}}$  bonds typically found in  $N$ -alkylalkanesulfinamides (1.470–1.530 Å).

### Related literature

For  $N$ -arylalkanesulfinamides, see: Datta *et al.* (2008) and for cyclic  $N$ -arylalkanesulfinamides (sultims), see: Schulze *et al.* (2005). For  $N$ -alkylalkanesulfinamides, see: Sato *et al.* (1975); Schuckmann *et al.* (1978); Ferreira *et al.* (2005). For the synthesis, see: Stretter *et al.* (1969).



### Experimental

#### Crystal data

$\text{C}_{10}\text{H}_{15}\text{NOS}$   
 $M_r = 197.29$   
Monoclinic,  $P2_1/c$

$a = 7.4822$  (3) Å  
 $b = 15.7881$  (6) Å  
 $c = 8.8333$  (4) Å

$\beta = 99.3865$  (6)<sup>°</sup>  
 $V = 1029.50$  (7) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 0.28$  mm<sup>-1</sup>  
 $T = 150$  K  
 $0.54 \times 0.49 \times 0.39$  mm

#### Data collection

Bruker APEXII CCD diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2007)  
 $T_{\text{min}} = 0.866$ ,  $T_{\text{max}} = 0.900$

12022 measured reflections  
3150 independent reflections  
2861 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.020$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.092$   
 $S = 1.05$   
3150 reflections  
125 parameters  
1 restraint

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.38$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.36$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^i$	0.808 (12)	2.085 (12)	2.8882 (11)	173.1 (14)

Symmetry code: (i)  $-x, -y + 1, -z + 1$ .

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* and local programs.

MD and AJB thank KAIST for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2002).

### References

- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Datta, M., Buglass, A. J., Hong, C. S. & Lim, J. H. (2008). *Acta Cryst.* **E64**, o1393.  
Ferreira, F., Audoin, M. & Chemla, F. (2005). *Chem. Eur. J.* **11**, 5269–5278.  
Sato, S., Yoshioka, T. & Tamura, C. (1975). *Acta Cryst.* **B31**, 1385–1392.  
Schuckmann, W., Fuess, H., Möisinger, O. & Ried, W. (1978). *Acta Cryst.* **B34**, 1516–1520.  
Schulze, B., Taubert, K., Siegemund, A., Freysoldt, T. H. E. & Sieler, J. (2005). *Z. Naturforsch. Teil B*, **60**, 41–47.  
Sheldrick, G. M. (2007). *SADABS*. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Stretter, H., Krause, M. & Last, W.-D. (1969). *Chem. Ber.* **102**, 3357–3363.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o2034 [ doi:10.1107/S1600536809028633 ]

## *N*-Phenyl-*tert*-butanesulfinamide

M. Datta, A. J. Buglass and M. R. J. Elsegood

### Comment

The molecular structure of (I) (Fig. 1) exhibits a short *N*—(aryl)C Bond (1.4083 (12) Å), like that in *N*-phenyladamantane-1-sulfinamide (1.409 (2) Å) (Datta *et al.*, 2008), and in contrast with *N*—(alkyl)C bonds in *N*-alkylalkanesulfinamides (1.470–1.530 Å) (Sato *et al.*, 1975; Schuckmann *et al.*, 1978; Ferreira *et al.*, 2005). Otherwise, molecular geometry is similar to other sulfinamides. The molecules of (I) in the crystal lattice are linked by pairs of N—H···O=S hydrogen bonds (Fig. 2 and Table 1). There is no evidence of weak intermolecular C—H···O=S hydrogen bonds, as in the packing of *N*-phenyladamantane-1-sulfinamide (Datta *et al.*, 2008). The crystal system and space group for (I) and *N*-phenyladamantane-1-sulfinamide are the same (namely monoclinic and *P*2<sub>1</sub>/*c*, respectively).

### Experimental

Compound (I) was prepared by the method of Stretter *et al.* (1969), using *tert*-butanesulfinyl chloride (702.5 mg, 5 mmol) and aniline (930 mg, 10 mmol) in dry chloroform (30 ml). After 6 h (with TLC monitoring) the white solid amine salt was filtered off and the solvent was removed under reduced pressure. Column chromatography (silica gel, dichloromethane) provided (I) as white crystals (950 mg, 96%), m.p. 376–377 K. Single crystals suitable for X-ray analysis were obtained by evaporation of a solution of (I) in dichloromethane at room temperature. Spectroscopic analysis: FTIR (KBr) (cm<sup>-1</sup>) 3015, 2599, 2330, 1496, 1469, 1420, 1370, 1274, 1068, 1026, 888, 859. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub> p.p.m. with respect to TMS) δ 7.26–7.22 (m, 2H), 7.00–6.98 (m, 3H), 5.48 (bs, 1H), 1.30 (s, 9H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub> p.p.m. with respect to TMS) δ 142.0, 129.4, 123.0, 118.4, 56.5, 22.4. EIMS *m/z* (%) 197 (*M*<sup>+</sup>, 18) 141 (100), 140 (*M*<sup>+</sup>-<sup>t</sup>Bu, 28), 105 (*M*<sup>+</sup>-PhNH, 86), 92 (*M*<sup>+</sup>-<sup>t</sup>BuSO, 72), 78 (77), 57 (93).

### Refinement

H atoms were located in a difference Fourier map and refined geometrically using a riding model except for NH for which the coordinates were freely refined. Bond lengths and displacement parameters were constrained as follows: C—H = 0.95–0.98 Å and *U*<sub>iso</sub>(H) = 1.2 (1.5 for CH<sub>3</sub>) times *U*<sub>eq</sub>(C, N).

### Figures




Fig. 1. Molecular structure of (I), with atom labels and 50% probability displacement ellipsoids for non-H atoms.




Fig. 2. Intermolecular hydrogen bonding between a pair of opposite enantiomers of (I) in the crystal packing. Symmetry code  $i = -x, -y + 1, -z + 1$ .

## *N*-Phenyl-*tert*-butanesulfinamide

### Crystal data

$C_{10}H_{15}NOS$	$F_{000} = 424$
$M_r = 197.29$	$D_x = 1.273 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Melting point = 376–377 K
Hall symbol: $-P\ 2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.4822 (3) \text{ \AA}$	Cell parameters from 6597 reflections
$b = 15.7881 (6) \text{ \AA}$	$\theta = 2.3\text{--}30.5^\circ$
$c = 8.8333 (4) \text{ \AA}$	$\mu = 0.28 \text{ mm}^{-1}$
$\beta = 99.3865 (6)^\circ$	$T = 150 \text{ K}$
$V = 1029.50 (7) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.54 \times 0.49 \times 0.39 \text{ mm}$

### Data collection

Bruker APEXII CCD diffractometer	3150 independent reflections
Radiation source: fine-focus sealed tube	2861 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.020$
$T = 150 \text{ K}$	$\theta_{\text{max}} = 30.6^\circ$
$\omega$ rotation with narrow frames scans	$\theta_{\text{min}} = 2.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2007)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.866, T_{\text{max}} = 0.900$	$k = -22 \rightarrow 22$
12022 measured reflections	$l = -12 \rightarrow 12$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.092$	$w = 1/[\sigma^2(F_o^2) + (0.0516P)^2 + 0.2529P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
3150 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
125 parameters	$\Delta\rho_{\text{max}} = 0.38 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$

l restraint

Extinction correction: SHELXL97 (Sheldrick, 2008),  
 $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Primary atom site location: structure-invariant direct methods

Extinction coefficient: 0.020 (3)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{iso}^*/U_{eq}$
O1	-0.06049 (10)	0.60738 (4)	0.41648 (9)	0.02295 (16)
S1	0.10835 (3)	0.601595 (13)	0.34614 (3)	0.01896 (9)
N1	0.17155 (13)	0.50050 (5)	0.35165 (10)	0.02446 (18)
H1	0.1463 (19)	0.4727 (9)	0.4219 (15)	0.029*
C1	0.19492 (12)	0.45439 (6)	0.21979 (11)	0.01998 (18)
C2	0.24483 (16)	0.49296 (7)	0.09080 (12)	0.0281 (2)
H2	0.2589	0.5527	0.0877	0.034*
C3	0.27381 (17)	0.44344 (7)	-0.03327 (13)	0.0318 (2)
H3	0.3053	0.4700	-0.1218	0.038*
C4	0.25757 (15)	0.35617 (7)	-0.02991 (13)	0.0295 (2)
H4	0.2790	0.3230	-0.1148	0.035*
C5	0.20955 (14)	0.31769 (6)	0.09922 (12)	0.0257 (2)
H5	0.1990	0.2578	0.1029	0.031*
C6	0.17690 (13)	0.36629 (6)	0.22303 (11)	0.02134 (18)
H6	0.1422	0.3396	0.3101	0.026*
C7	0.29072 (13)	0.64579 (6)	0.48984 (11)	0.02172 (18)
C8	0.24368 (16)	0.73950 (7)	0.50283 (14)	0.0316 (2)
H8A	0.3444	0.7687	0.5668	0.047*
H8B	0.1342	0.7448	0.5497	0.047*
H8C	0.2223	0.7650	0.4003	0.047*
C9	0.46486 (15)	0.63537 (8)	0.42278 (14)	0.0312 (2)
H9A	0.5658	0.6613	0.4919	0.047*
H9B	0.4512	0.6632	0.3224	0.047*
H9C	0.4894	0.5750	0.4110	0.047*
C10	0.30068 (15)	0.60073 (6)	0.64357 (12)	0.0265 (2)
H10A	0.3290	0.5408	0.6314	0.040*
H10B	0.1839	0.6057	0.6794	0.040*
H10C	0.3956	0.6267	0.7188	0.040*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0218 (3)	0.0215 (3)	0.0264 (4)	0.0011 (2)	0.0067 (3)	0.0021 (2)
S1	0.02263 (13)	0.01561 (12)	0.01895 (13)	0.00052 (7)	0.00429 (8)	0.00155 (7)
N1	0.0397 (5)	0.0153 (3)	0.0206 (4)	0.0013 (3)	0.0114 (3)	0.0012 (3)
C1	0.0214 (4)	0.0187 (4)	0.0202 (4)	0.0003 (3)	0.0045 (3)	-0.0011 (3)
C2	0.0397 (6)	0.0210 (4)	0.0254 (5)	-0.0014 (4)	0.0112 (4)	0.0010 (4)
C3	0.0422 (6)	0.0327 (6)	0.0224 (5)	0.0011 (4)	0.0111 (4)	0.0006 (4)
C4	0.0324 (5)	0.0316 (5)	0.0251 (5)	0.0014 (4)	0.0063 (4)	-0.0080 (4)
C5	0.0247 (5)	0.0218 (4)	0.0304 (5)	-0.0010 (3)	0.0043 (4)	-0.0060 (4)
C6	0.0207 (4)	0.0186 (4)	0.0252 (4)	-0.0012 (3)	0.0053 (3)	-0.0006 (3)
C7	0.0218 (4)	0.0186 (4)	0.0242 (4)	0.0006 (3)	0.0022 (3)	-0.0001 (3)
C8	0.0352 (5)	0.0183 (4)	0.0387 (6)	-0.0002 (4)	-0.0013 (4)	-0.0037 (4)
C9	0.0230 (5)	0.0366 (6)	0.0346 (6)	0.0004 (4)	0.0065 (4)	0.0042 (4)
C10	0.0290 (5)	0.0278 (5)	0.0220 (5)	0.0026 (4)	0.0020 (4)	0.0006 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O1—S1	1.4988 (7)	C5—H5	0.9500
S1—N1	1.6632 (9)	C6—H6	0.9500
S1—C7	1.8426 (10)	C7—C10	1.5241 (14)
N1—C1	1.4083 (12)	C7—C9	1.5255 (14)
N1—H1	0.808 (12)	C7—C8	1.5294 (14)
C1—C2	1.3954 (13)	C8—H8A	0.9800
C1—C6	1.3982 (12)	C8—H8B	0.9800
C2—C3	1.3918 (15)	C8—H8C	0.9800
C2—H2	0.9500	C9—H9A	0.9800
C3—C4	1.3838 (17)	C9—H9B	0.9800
C3—H3	0.9500	C9—H9C	0.9800
C4—C5	1.3903 (15)	C10—H10A	0.9800
C4—H4	0.9500	C10—H10B	0.9800
C5—C6	1.3898 (13)	C10—H10C	0.9800
O1—S1—N1	107.53 (4)	C10—C7—C9	112.02 (8)
O1—S1—C7	105.72 (4)	C10—C7—C8	111.29 (9)
N1—S1—C7	99.69 (5)	C9—C7—C8	110.80 (9)
C1—N1—S1	123.02 (7)	C10—C7—S1	111.11 (7)
C1—N1—H1	115.5 (10)	C9—C7—S1	105.97 (7)
S1—N1—H1	116.3 (10)	C8—C7—S1	105.33 (7)
C2—C1—C6	119.36 (9)	C7—C8—H8A	109.5
C2—C1—N1	122.39 (8)	C7—C8—H8B	109.5
C6—C1—N1	118.16 (8)	H8A—C8—H8B	109.5
C3—C2—C1	119.65 (10)	C7—C8—H8C	109.5
C3—C2—H2	120.2	H8A—C8—H8C	109.5
C1—C2—H2	120.2	H8B—C8—H8C	109.5
C4—C3—C2	121.12 (10)	C7—C9—H9A	109.5
C4—C3—H3	119.4	C7—C9—H9B	109.5



C2—C3—H3	119.4	H9A—C9—H9B	109.5
C3—C4—C5	119.21 (9)	C7—C9—H9C	109.5
C3—C4—H4	120.4	H9A—C9—H9C	109.5
C5—C4—H4	120.4	H9B—C9—H9C	109.5
C6—C5—C4	120.43 (9)	C7—C10—H10A	109.5
C6—C5—H5	119.8	C7—C10—H10B	109.5
C4—C5—H5	119.8	H10A—C10—H10B	109.5
C5—C6—C1	120.21 (9)	C7—C10—H10C	109.5
C5—C6—H6	119.9	H10A—C10—H10C	109.5
C1—C6—H6	119.9	H10B—C10—H10C	109.5
O1—S1—N1—C1	123.13 (8)	C4—C5—C6—C1	-1.02 (15)
C7—S1—N1—C1	-126.86 (8)	C2—C1—C6—C5	0.40 (14)
S1—N1—C1—C2	28.58 (14)	N1—C1—C6—C5	-176.09 (9)
S1—N1—C1—C6	-155.05 (8)	O1—S1—C7—C10	54.89 (8)
C6—C1—C2—C3	0.76 (16)	N1—S1—C7—C10	-56.55 (8)
N1—C1—C2—C3	177.09 (10)	O1—S1—C7—C9	176.79 (7)
C1—C2—C3—C4	-1.32 (18)	N1—S1—C7—C9	65.35 (7)
C2—C3—C4—C5	0.70 (17)	O1—S1—C7—C8	-65.73 (8)
C3—C4—C5—C6	0.48 (16)	N1—S1—C7—C8	-177.17 (7)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N1—H1 $\cdots$ O1 <sup>i</sup>	0.808 (12)	2.085 (12)	2.8882 (11)	173.1 (14)

Symmetry codes: (i)  $-x, -y+1, -z+1$ .

Fig. 1




Fig. 2

