



This item was submitted to Loughborough's Institutional Repository by the author and is made available under the following Creative Commons Licence conditions.



CC creative commons
COMMONS DEED

Attribution-NonCommercial-NoDerivs 2.5

You are free:

- to copy, distribute, display, and perform the work

Under the following conditions:

BY: **Attribution.** You must attribute the work in the manner specified by the author or licensor.

Noncommercial. You may not use this work for commercial purposes.

No Derivative Works. You may not alter, transform, or build upon this work.

- For any reuse or distribution, you must make clear to others the license terms of this work.
- Any of these conditions can be waived if you get permission from the copyright holder.

Your fair use and other rights are in no way affected by the above.

This is a human-readable summary of the [Legal Code \(the full license\)](#).

[Disclaimer](#) 

For the full text of this licence, please go to:
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368


Editors: W. Clegg and D. G. Watson

N'-(2,4-Dinitrophenyl)acetohydrazide

Muhammad Zia-ur-Rehman, Mark R. J. Elsegood, Shahid Mahmud and Hamid Latif Siddiqui

Acta Cryst. (2008). E64, o1441

This article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***N'*-(2,4-Dinitrophenyl)acetohydrazide**Muhammad Zia-ur-Rehman,^{a*} Mark R. J. Elsegood,^b
Shahid Mahmud^a and Hamid Latif Siddiqui^c^aApplied Chemistry Research Centre, PCSIR Laboratories Complex, Lahore 54600, Pakistan, ^bChemistry Department, Loughborough University, Loughborough LE11 3TU, England, and ^cInstitute of Chemistry, University of The Punjab, Lahore 54590, Pakistan

Correspondence e-mail: drhamidlatif@yahoo.com


Received 23 June 2008; accepted 27 June 2008

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.031; wR factor = 0.079; data-to-parameter ratio = 11.1.

In the title compound, $C_8H_8N_4O_5$, the nitro groups *ortho* and *para* to the hydrazone group are twisted by 10.0 (2) and 3.6 (2)°, respectively, relative to the aromatic ring. The structure exhibits an intramolecular N—H···O hydrogen bond between the hydrazide and *ortho*-nitro groups. There is a strong intermolecular C=O···H—N hydrogen bond, giving rise to chains, and weaker ONO···NO₂ [2.944 (2) Å] and C—H···O—N interactions linking the molecules into a three-dimensional network.

Related literature

For related literature, see: Domiano *et al.* (1984); Guo (2007); Li *et al.* (1988); Rudnicka & Osmialowska (1979); Sakamoto *et al.* (1993); Siddiqui *et al.* (2007); Zia-ur-Rehman *et al.* (2005, 2006).



Experimental

Crystal data

 $C_8H_8N_4O_5$ $M_r = 240.18$ Orthorhombic, $P2_12_12_1$ $a = 4.8585$ (4) Å $b = 10.7703$ (8) Å $c = 19.1059$ (14) Å $V = 999.76$ (13) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.14$ mm⁻¹ $T = 150$ (2) K $0.57 \times 0.09 \times 0.06$ mm

Data collection

Bruker APEXII CCD

diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 2007)

 $T_{min} = 0.927$, $T_{max} = 0.992$

11843 measured reflections

1794 independent reflections

1616 reflections with $I > 2\sigma(I)$ $R_{int} = 0.031$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.030$ $wR(F^2) = 0.078$ $S = 1.03$

1794 reflections

161 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{max} = 0.27$ e Å⁻³ $\Delta\rho_{min} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-------------------------|----------|-------------|-------------|---------------|
| $N3-H3\cdots O2$ | 0.83 (2) | 2.001 (18) | 2.5942 (16) | 127.9 (16) |
| $N4-H4\cdots O5^i$ | 0.85 (2) | 1.95 (2) | 2.7748 (16) | 164.0 (17) |
| $C5-H5\cdots O4^{ii}$ | 0.95 | 2.44 | 3.249 (2) | 143 |
| $C8-H8A\cdots O2^{iii}$ | 0.98 | 2.58 | 3.269 (2) | 128 |
| $C8-H8C\cdots O3^{iv}$ | 0.98 | 2.57 | 3.527 (2) | 165 |

Symmetry code: (i) $x + 1, y, z$; (ii) $-x + 2, y - \frac{1}{2}, -z + \frac{3}{2}$; (iii) $x + \frac{1}{2}, -y + \frac{1}{2}, -z + 2$; (iv) $x - 1, y - 1, z$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* and local programs.

The authors are grateful to the Pakistan Council of Scientific & Industrial Research Laboratories Complex, Lahore, for providing the necessary facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2733).

References

- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Domiano, P., Pelizzi, C. & Predieri, G. (1984). *Polyhedron*, **3**, 281–286.
- Guo, H.-M. (2007). *Acta Cryst.* **E63**, o3123.
- Li, X. R., Sun, Z. M. & Chang, J. C. (1988). *Synth. React. Inorg. Met. Org. Chem.* **18**, 657–665.
- Rudnicka, G. & Osmialowska, Z. (1979). *Acta Pol. Pharm.* **36**, 411–419.
- Sakamoto, H., Goto, H., Yokoshima, M., Dobashi, M., Ishikawa, J., Doi, K. & Otomo, M. (1993). *Bull. Chem. Soc. Jpn.* **66**, 2907–2914.
- Sheldrick, G. M. (2007). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siddiqui, H. L., Zia-ur-Rehman, M., Ahmad, N., Weaver, G. W. & Lucas, P. D. (2007). *Chem. Pharm. Bull.* **55**, 1014–1017.
- Zia-ur-Rehman, M., Choudary, J. A. & Ahmad, S. (2005). *Bull. Korean Chem. Soc.* **26**, 1771–1775.
- Zia-ur-Rehman, M., Choudary, J. A., Ahmad, S. & Siddiqui, H. L. (2006). *Chem. Pharm. Bull.* **54**, 1175–1178.

supplementary materials

Acta Cryst. (2008). E64, o1441 [doi:10.1107/S1600536808019685]

N'-(2,4-Dinitrophenyl)acetohydrazide

M. Zia-ur-Rehman, M. R. J. Elsegood, S. Mahmud and H. L. Siddiqui

Comment

The chemistry of hydrazones has been intensely investigated in recent years due to their excellent coordinating capability (Domiano *et al.*, 1984) and pharmacological activities (Li *et al.*, 1988). These compounds are also being used as precursors for the efficient synthesis of various condensed heterocycles in organic chemistry (Rudnicka & Osmialowska, 1979) and as highly selective metal scavengers (Sakamoto *et al.*, 1993) in analytical chemistry. In continuation of our ongoing work on the synthesis of various heterocyclic compounds (Zia-ur-Rehman *et al.*, 2005, 2006; Siddiqui *et al.*, 2007), the title compound, (I), was synthesized by reacting 2,4-dinitrophenylhydrazine with acetic anhydride.

Most of the bond lengths and angles in (I) are similar to those in related molecules (Guo, 2007). The nitro groups *ortho* and *para* to the hydrazone group are twisted out of this plane by 10.0 (2) and 3.6 (2)°, respectively. The larger twist of the *ortho*-nitro group arises due to the desire to form an intramolecular hydrogen bond which results in a six-membered ring (Fig. 1 and Table 1). Each molecule also forms an intermolecular N—H···O=C hydrogen bond giving rise to stacks of molecules parallel to *a* (Fig. 2). The hydrogen-bonded chains of (I) are further linked together into a three-dimensional network (Fig. 3) *via* weaker C—H···O—N interactions involving the nitro groups and methyl and aryl H atoms (range 2.4–2.6 Å) along with some weak ONO···NO₂ interactions [O1···N1ⁱ = 2.944 (2) Å; symmetry code: (i) -0.5+x, 1.5-y, 2-z].

Experimental

A mixture of 2,4-dinitrophenylhydrazine (1.981 g; 10.0 mmoles) and acetic anhydride (5.0 ml) was stirred for a period of six hours at room temperature. Then, this mixture was poured into ice cooled water and neutralized with 10% sodium bicarbonate solution. The precipitated solids were collected by filtration, washed and dried. Crystals suitable for X-ray crystallography were grown by slow evaporation of solution of the title compound in a mixture of ethanol and water (90:10); m.p. 471 K; yield: 82%.

Refinement

1255 Friedel pairs were merged. H atoms bound to C were placed in geometric positions (C—H distance = 0.95 Å for aryl-H; 0.98 Å for methyl-H) using a riding model. H atoms on N had coordinates freely refined. U_{iso} values were set to 1.2 U_{eq} of the carrier atom (1.5 U_{eq} for methyl-H).

Figures




Fig. 1. The asymmetric unit of the title compound showing the intramolecular hydrogen bond. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Perspective view of molecules linked *via* intermolecular N—H...O=C hydrogen bonds parallel to *a*.




Fig. 3. Perspective view of the three-dimensional crystal packing showing hydrogen-bonds and other intermolecular interactions (dashed lines). H atoms not involved in hydrogen bonding have been omitted for clarity.

N'-(2,4-Dinitrophenyl)acetohydrazide

Crystal data

$C_8H_8N_4O_5$

$M_r = 240.18$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 4.8585$ (4) Å

$b = 10.7703$ (8) Å

$c = 19.1059$ (14) Å

$V = 999.76$ (13) Å³

$Z = 4$

$F_{000} = 496$

$D_x = 1.596$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 3491 reflections

$\theta = 2.9$ – 27.8°

$\mu = 0.14$ mm⁻¹

$T = 150$ (2) K

Lath, orange

$0.57 \times 0.09 \times 0.06$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 150$ (2) K

1794 independent reflections

1616 reflections with $I > 2\sigma(I)$

$R_{int} = 0.031$

$\theta_{max} = 30.6^\circ$

ω rotation with narrow frames scans $\theta_{\min} = 2.1^\circ$
 Absorption correction: multi-scan $h = -6 \rightarrow 6$
 (SADABS; Sheldrick, 2007)
 $T_{\min} = 0.927, T_{\max} = 0.992$ $k = -15 \rightarrow 15$
 11843 measured reflections $l = -27 \rightarrow 27$

Refinement

Refinement on F^2 Secondary atom site location: all non-H atoms found by direct methods
 Least-squares matrix: full Hydrogen site location: geom except NH coords freely refined
 $R[F^2 > 2\sigma(F^2)] = 0.030$ H atoms treated by a mixture of independent and constrained refinement
 $wR(F^2) = 0.078$ $w = 1/[\sigma^2(F_o^2) + (0.0462P)^2 + 0.078P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 1.03$ $(\Delta/\sigma)_{\max} < 0.001$
 1794 reflections $\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$
 161 parameters $\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. 1255 Friedel pairs. Friedels merged.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|------------|--------------|-------------|----------------------------------|
| C1 | 0.3396 (3) | 0.64098 (13) | 0.89834 (7) | 0.0206 (3) |
| N1 | 0.1186 (2) | 0.67553 (11) | 0.94555 (6) | 0.0239 (2) |
| O1 | 0.0352 (2) | 0.78314 (10) | 0.94499 (6) | 0.0298 (2) |
| O2 | 0.0219 (2) | 0.59512 (11) | 0.98457 (6) | 0.0349 (3) |
| C2 | 0.4679 (3) | 0.73783 (13) | 0.86286 (7) | 0.0234 (3) |
| H2 | 0.4153 | 0.8215 | 0.8710 | 0.028* |
| C3 | 0.6722 (3) | 0.70968 (14) | 0.81593 (7) | 0.0245 (3) |
| N2 | 0.8161 (3) | 0.81093 (13) | 0.78132 (7) | 0.0323 (3) |
| O3 | 0.7483 (3) | 0.91873 (12) | 0.79517 (7) | 0.0428 (3) |
| O4 | 1.0012 (3) | 0.78354 (13) | 0.73998 (7) | 0.0462 (3) |
| C4 | 0.7508 (3) | 0.58725 (15) | 0.80240 (7) | 0.0253 (3) |
| H4A | 0.8903 | 0.5700 | 0.7689 | 0.030* |

supplementary materials

| | | | | |
|-----|-------------|--------------|-------------|------------|
| C5 | 0.6255 (3) | 0.49242 (13) | 0.83774 (7) | 0.0231 (3) |
| H5 | 0.6791 | 0.4093 | 0.8283 | 0.028* |
| C6 | 0.4174 (3) | 0.51517 (13) | 0.88817 (7) | 0.0200 (3) |
| N3 | 0.3089 (3) | 0.42103 (11) | 0.92557 (7) | 0.0243 (3) |
| H3 | 0.172 (4) | 0.4329 (16) | 0.9508 (10) | 0.029* |
| N4 | 0.3548 (3) | 0.29805 (11) | 0.90659 (7) | 0.0222 (2) |
| H4 | 0.518 (4) | 0.2710 (16) | 0.9126 (10) | 0.027* |
| C7 | 0.1401 (3) | 0.21961 (13) | 0.91490 (7) | 0.0222 (3) |
| O5 | -0.0894 (2) | 0.25786 (10) | 0.93114 (6) | 0.0291 (2) |
| C8 | 0.2055 (4) | 0.08501 (14) | 0.90370 (9) | 0.0310 (3) |
| H8A | 0.1747 | 0.0392 | 0.9473 | 0.047* |
| H8B | 0.3984 | 0.0764 | 0.8895 | 0.047* |
| H8C | 0.0861 | 0.0515 | 0.8669 | 0.047* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|----|------------|------------|------------|-------------|-------------|-------------|
| C1 | 0.0161 (6) | 0.0247 (6) | 0.0211 (6) | -0.0007 (5) | -0.0003 (5) | -0.0036 (5) |
| N1 | 0.0187 (5) | 0.0254 (5) | 0.0276 (6) | -0.0018 (5) | 0.0007 (5) | -0.0083 (5) |
| O1 | 0.0243 (5) | 0.0272 (5) | 0.0380 (6) | 0.0046 (4) | -0.0021 (5) | -0.0092 (4) |
| O2 | 0.0334 (6) | 0.0304 (6) | 0.0408 (6) | -0.0044 (5) | 0.0178 (5) | -0.0053 (5) |
| C2 | 0.0218 (6) | 0.0241 (6) | 0.0242 (6) | -0.0029 (5) | -0.0050 (5) | 0.0001 (5) |
| C3 | 0.0229 (6) | 0.0303 (7) | 0.0203 (6) | -0.0070 (6) | -0.0027 (5) | 0.0046 (5) |
| N2 | 0.0323 (7) | 0.0384 (7) | 0.0262 (6) | -0.0125 (6) | -0.0069 (6) | 0.0095 (5) |
| O3 | 0.0475 (8) | 0.0320 (6) | 0.0489 (7) | -0.0124 (6) | -0.0052 (6) | 0.0114 (5) |
| O4 | 0.0430 (7) | 0.0581 (8) | 0.0374 (6) | -0.0177 (7) | 0.0111 (6) | 0.0105 (6) |
| C4 | 0.0203 (7) | 0.0352 (7) | 0.0203 (6) | -0.0025 (6) | 0.0020 (5) | 0.0001 (6) |
| C5 | 0.0205 (6) | 0.0267 (6) | 0.0221 (6) | 0.0012 (5) | 0.0029 (5) | -0.0024 (5) |
| C6 | 0.0161 (6) | 0.0238 (6) | 0.0200 (6) | -0.0007 (5) | -0.0006 (5) | -0.0014 (5) |
| N3 | 0.0210 (6) | 0.0217 (5) | 0.0301 (6) | 0.0006 (5) | 0.0084 (5) | -0.0019 (5) |
| N4 | 0.0154 (5) | 0.0200 (5) | 0.0312 (6) | 0.0011 (4) | 0.0006 (5) | -0.0004 (5) |
| C7 | 0.0182 (6) | 0.0256 (6) | 0.0228 (6) | -0.0011 (5) | -0.0022 (5) | 0.0029 (5) |
| O5 | 0.0159 (5) | 0.0339 (6) | 0.0375 (6) | -0.0002 (4) | 0.0012 (4) | 0.0046 (5) |
| C8 | 0.0308 (8) | 0.0235 (6) | 0.0389 (8) | -0.0011 (6) | -0.0008 (7) | 0.0019 (6) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|-------|-------------|--------|-------------|
| C1—C2 | 1.3915 (19) | C5—C6 | 1.4178 (18) |
| C1—C6 | 1.4202 (19) | C5—H5 | 0.9500 |
| C1—N1 | 1.4508 (18) | C6—N3 | 1.3477 (18) |
| N1—O1 | 1.2278 (16) | N3—N4 | 1.3913 (17) |
| N1—O2 | 1.2354 (16) | N3—H3 | 0.83 (2) |
| C2—C3 | 1.371 (2) | N4—C7 | 1.3519 (18) |
| C2—H2 | 0.9500 | N4—H4 | 0.85 (2) |
| C3—C4 | 1.397 (2) | C7—O5 | 1.2282 (17) |
| C3—N2 | 1.4543 (19) | C7—C8 | 1.500 (2) |
| N2—O4 | 1.233 (2) | C8—H8A | 0.9800 |
| N2—O3 | 1.2355 (19) | C8—H8B | 0.9800 |
| C4—C5 | 1.367 (2) | C8—H8C | 0.9800 |

| | | | |
|-------------|--------------|-------------|--------------|
| C4—H4A | 0.9500 | | |
| C2—C1—C6 | 121.96 (13) | C6—C5—H5 | 119.2 |
| C2—C1—N1 | 116.25 (12) | N3—C6—C5 | 120.61 (13) |
| C6—C1—N1 | 121.79 (12) | N3—C6—C1 | 122.76 (12) |
| O1—N1—O2 | 122.79 (12) | C5—C6—C1 | 116.59 (12) |
| O1—N1—C1 | 118.74 (12) | C6—N3—N4 | 121.01 (12) |
| O2—N1—C1 | 118.47 (12) | C6—N3—H3 | 120.3 (12) |
| C3—C2—C1 | 118.50 (13) | N4—N3—H3 | 115.2 (12) |
| C3—C2—H2 | 120.8 | C7—N4—N3 | 116.14 (12) |
| C1—C2—H2 | 120.8 | C7—N4—H4 | 119.2 (12) |
| C2—C3—C4 | 121.83 (13) | N3—N4—H4 | 116.0 (13) |
| C2—C3—N2 | 118.65 (14) | O5—C7—N4 | 121.37 (13) |
| C4—C3—N2 | 119.49 (13) | O5—C7—C8 | 123.55 (14) |
| O4—N2—O3 | 123.81 (14) | N4—C7—C8 | 115.07 (13) |
| O4—N2—C3 | 117.56 (14) | C7—C8—H8A | 109.5 |
| O3—N2—C3 | 118.63 (15) | C7—C8—H8B | 109.5 |
| C5—C4—C3 | 119.47 (13) | H8A—C8—H8B | 109.5 |
| C5—C4—H4A | 120.3 | C7—C8—H8C | 109.5 |
| C3—C4—H4A | 120.3 | H8A—C8—H8C | 109.5 |
| C4—C5—C6 | 121.60 (13) | H8B—C8—H8C | 109.5 |
| C4—C5—H5 | 119.2 | | |
| C2—C1—N1—O1 | 9.02 (18) | N2—C3—C4—C5 | 176.53 (13) |
| C6—C1—N1—O1 | -170.04 (13) | C3—C4—C5—C6 | -0.2 (2) |
| C2—C1—N1—O2 | -171.10 (12) | C4—C5—C6—N3 | -175.76 (14) |
| C6—C1—N1—O2 | 9.84 (19) | C4—C5—C6—C1 | 2.1 (2) |
| C6—C1—C2—C3 | 1.1 (2) | C2—C1—C6—N3 | 175.25 (13) |
| N1—C1—C2—C3 | -177.94 (12) | N1—C1—C6—N3 | -5.7 (2) |
| C1—C2—C3—C4 | 0.9 (2) | C2—C1—C6—C5 | -2.53 (19) |
| C1—C2—C3—N2 | -177.00 (13) | N1—C1—C6—C5 | 176.47 (12) |
| C2—C3—N2—O4 | 179.13 (13) | C5—C6—N3—N4 | -13.1 (2) |
| C4—C3—N2—O4 | 1.2 (2) | C1—C6—N3—N4 | 169.18 (13) |
| C2—C3—N2—O3 | -0.3 (2) | C6—N3—N4—C7 | -142.12 (14) |
| C4—C3—N2—O3 | -178.27 (14) | N3—N4—C7—O5 | 7.5 (2) |
| C2—C3—C4—C5 | -1.3 (2) | N3—N4—C7—C8 | -171.84 (12) |

Hydrogen-bond geometry (\AA , $^\circ$)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|-----------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| N3—H3 \cdots O2 | 0.83 (2) | 2.001 (18) | 2.5942 (16) | 127.9 (16) |
| N4—H4 \cdots O5 ⁱ | 0.85 (2) | 1.95 (2) | 2.7748 (16) | 164.0 (17) |
| C5—H5 \cdots O4 ⁱⁱ | 0.95 | 2.44 | 3.249 (2) | 143 |
| C8—H8A \cdots O2 ⁱⁱⁱ | 0.98 | 2.58 | 3.269 (2) | 128 |
| C8—H8C \cdots O3 ^{iv} | 0.98 | 2.57 | 3.527 (2) | 165 |

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+2, y-1/2, -z+3/2$; (iii) $x+1/2, -y+1/2, -z+2$; (iv) $x-1, y-1, z$.

Fig. 1




Fig. 2




Fig. 3

