

# N-cycling and balancing of the N-deficit generated in the oxygen minimum zone over the Namibian shelf—An isotope-based approach

Birgit Nagel,<sup>1,2</sup> Kay-Christian Emeis,<sup>1,2</sup> Anita Flohr,<sup>3</sup> Tim Rixen,<sup>3</sup> Tim Schlarbaum,<sup>1,2</sup> Volker Mohrholz,<sup>4</sup> and Anja van der Plas<sup>5</sup>

Received 12 April 2012; revised 5 February 2013; accepted 9 February 2013; published 28 March 2013.

[1] The northern Benguela upwelling system is a nutrient-replete region with high plankton biomass production and a seasonally changing oxygen minimum zone. Nitrate:phosphate ratios in fresh upwelling water are low due to denitrification in the near-seafloor oxygen minimum zone and phosphate efflux from sediments. This makes the region a candidate for substantial dinitrogen fixation, for which evidence is scarce. Nutrient and oxygen data, N isotope data of nitrate, nitrogen isotope ratios of particulate matter, particulate organic carbon content, and suspended matter concentrations on a transect across the shelf and upper slope at 23°S illustrate N-cycling processes and are the basis for estimating the contribution of N-sources and N-sinks to the reactive nitrogen pool. It appears that N-removal due to denitrification exceeds N gain by N<sub>2</sub> fixation and physical mixing processes by a factor of >6, although inorganic N:P ratios again increase as surface water is advected offshore. Nitrate and ammonium regeneration, nutrient assimilation with N:P < 16, shelf break mixing, atmospheric input, and N<sub>2</sub> fixation all contribute to the restoration of inorganic N:P ratios back to Redfield conditions, but in seasonally changing proportions. The Benguela upwelling system thus is a nutrient source for the oceanic-mixed layer where N-sources and N-sinks are not in balance and Redfield conditions can only re-adjust by advection and mixing processes integrated over time.

**Citation:** Nagel, B., K.-C. Emeis, A. Flohr, T. Rixen, T. Schlarbaum, V. Mohrholz, and A. van der Plas (2013), N-cycling and balancing of the N-deficit generated in the oxygen minimum zone over the Namibian shelf—An isotope-based approach, *J. Geophys. Res. Biogeosci.*, 118, 361–371, doi:10.1002/jgrg.20040.

## 1. Introduction

[2] Upwelling systems are the largest marine source of CO<sub>2</sub> to the atmosphere and their immense primary production fueled by upwelled nutrients far outweighs their modest regional extent [Watson, 1995; Wetzel *et al.*, 2005]. Productivity eventually re-captures the outgassed CO<sub>2</sub> by phytoplankton assimilation. But it is questionable if CO<sub>2</sub> outgassing and uptake are in direct balance in individual

upwelling systems. Because they often tap oxygen-deficient intermediate water masses, upwelling shelves are prone to oxygen deficiency aggravated by very high oxygen demand generated by remineralization of sinking organic matter [Monteiro *et al.*, 2006]. Oxygen-deficient coastal upwelling often does not supply nutrients into the euphotic zone in Redfield ratios, but with surplus of phosphate over nitrate (termed P\* by Deutsch *et al.* [2007]), because either is nitrate used to oxidize organic matter in the absence of dissolved O<sub>2</sub> [Lipschultz *et al.*, 1990], or phosphate is added from anoxic sediments high in organic carbon and phosphorus overlying the shelf [Bailey, 1991]. In the case that inorganic N:P ratios of upwelled waters are smaller than 16, current understanding is that there will be a zone at the fringes of the high productivity belt where nitrate is depleted by assimilation, and phosphate remains in aged upwelling waters, triggering the “Redfield homeostat” of diazotrophic N<sub>2</sub> fixation until the nitrate deficit is filled [Rixen *et al.*, 2005; Rixen *et al.*, 2009; Tyrrell, 1999]. If this basic concept is true, the CO<sub>2</sub> budget of an upwelling system remains constant on time scales of years to decades and on spatial scales far beyond the immediate coastal upwelling area, since the N-deficits produced in hypoxic water masses are balanced by N<sub>2</sub> fixation at the fringes of the upwelling region and the hemipelagic ocean.

Additional supporting information may be found in the online version of this article.

<sup>1</sup>Helmholtz-Zentrum Geesthacht, Centre for Material and Coastal Research, Geesthacht, Germany.

<sup>2</sup>Institute for Biogeochemistry and Marine Chemistry, University of Hamburg, Hamburg, Germany.

<sup>3</sup>Leibniz Center for Tropical Marine Ecology, Bremen, Germany.

<sup>4</sup>Leibniz Institute for Baltic Sea Research Warnemünde, Rostock, Germany.

<sup>5</sup>NatMIRC, Ministry of Fisheries and Marine Resources, Swakopmund, Namibia.


Corresponding author: B. Nagel, Helmholtz-Zentrum Geesthacht, Centre for Material and Coastal Research, Max-Planck-Str. 1, D-21502 Geesthacht, Germany. (birgit.nagel@hzg.de)

©2013 American Geophysical Union. All Rights Reserved.  
2169-8953/13/10.1002/jgrg.20040

[3] One example of such a system is the Benguela upwelling system (BUS) where several upwelling cells exist along the Namibian and South African coast due to the interaction of trade winds with coastal topography [Shannon and Nelson, 1996]. The northward component of the Southeast Trades drives upwelling and causes offshore Ekman transport in the surface mixed layer. Near-seafloor onshore transport and upwelling from 50 to 200 m water depth compensate for the wind-driven water deficit. The BUS is part of the eastern boundary current system of the south Atlantic. Off central and northern Namibia, the confluence of the northward flowing Benguela Current and the poleward Angola current forms a transition zone between central water of tropical and subtropical origin. Eastern South Atlantic Central Water (ESACW; Figure 1a) is advected within the Benguela current from the Agulhas region equatorward as far as the Angola Benguela Frontal Zone at 17°S [Mohrholz et al., 2001]. The Angola current and its subsurface extension, the poleward undercurrent (PU), transport nutrient-rich and oxygen-depleted South Atlantic Central Water (SACW) from the Angola Gyre along the shelf break (Figure 1a). Its proportion in the upwelling water controls the intensity of the oxygen minimum zone

(OMZ), which is most pronounced on the northern Benguela shelf [Inthorn et al., 2006; Mohrholz et al., 2008; Monteiro et al., 2006]. Upwelling mainly occurs not only at the coast [Mohrholz et al., 2008], but also over the shelf due to a belt of negative wind stress curl [Fennel et al., 2012]. Also shelf break upwelling was observed [Barange and Pillar, 1992].

[4] Water mass composition and thus oxygenation on the shelf vary throughout the year owing to seasonal changes in the wind-driven Ekman regime. High SACW contributions (80–100%) are found along the 23°S transect off Walvis Bay in austral summer (February/March) when the PU is most active and the cross-shelf circulation is weak, so that ventilation of the bottom water is impeded [Mohrholz et al., 2008]. The cross-shelf circulation is strongest in the peak upwelling season in August/September and brings oxygen-rich ESACW from the shelf break onto the shelf [Mohrholz et al., 2008]. Hypoxia and anoxia on the shelf are thus mainly controlled by the water mass composition, but oxygen demand from high organic matter fluxes and degradation reduce original O<sub>2</sub> concentrations [Monteiro et al., 2006] and create conditions conducive for anaerobic pathways of organic matter degradation. Oxygen concentrations below 10 and ~5 μM stimulate anammox and heterotrophic



**Figure 1.** (a) Map of the study area showing nitrate concentrations in surface water (0–5 m) from data available in the Namibian collection of bottle data. The red line marks the 23°S transect where detailed sampling was carried out during the three cruises in 2008, 2010, and 2011. The dashed line indicates the shelf break and arrows mark the Benguela Coastal Current (BCC) and the Poleward Undercurrent (PU) that supply ESACW (Eastern South Atlantic Central Water) and SACW (South Atlantic Central Water), respectively. (b) Water mass distribution (% SACW, colors) and temperature (contours) on the transect at 23°S during expedition M76-2 (May–June 2008).

denitrification, respectively [Cline and Kaplan, 1975; Jensen et al., 2008], and cause nitrate deficits over phosphate in OMZs that are transferred into surface waters by upwelling [Tyrrell and Lucas, 2002].

[5] Nitrate reduction in the water column—the first step of heterotrophic denitrification—and presumably also anammox are associated with kinetic fractionation of stable N isotopes causing an enrichment of the heavier  $^{15}\text{N}$  isotope in the residual nitrate pool [Konovalov et al., 2008; Sigman et al., 2005]. The fractionation factor ( $\epsilon$ ) is the difference between the isotopic composition of the source and the product [Mariotti et al., 1981]. Heterotrophic denitrifying bacteria and nitrate assimilating phytoplankton discriminate against the heavier  $^{15}\text{N}$  isotope with a fractionation factor of 25‰ and 5‰, respectively [Granger et al., 2004; Granger et al., 2008]. Although large reactive nitrogen deficits over phosphate occur in upwelling water, there is no experimental evidence for significant  $\text{N}_2$  fixation in the BUS [Sohm et al., 2011; Wasmund, personal communication, 2011]. This is puzzling, because inorganic N:P ratios are evidently restored to Redfield ratios in modified upwelling water advected offshore [Wasmund et al., 2005] and seem to exemplify the P\*-concept [Deutsch et al., 2007] that links  $\text{N}_2$  fixation in time and space to the denitrification zones of the ocean. In the absence of evidence for  $\text{N}_2$  fixation, several alternative processes have been invoked, such as shelf break upwelling in a second upwelling cell [Barange and Pillar, 1992; Emeis et al., 2009], or non-Redfield assimilation stoichiometry [Weber and Deutsch, 2010].

[6] Here, we use data of oxygen and nutrient concentrations as well as stable isotope compositions of nitrogen in nitrate ( $\delta^{15}\text{N}_{\text{NO}_3}$ ) and  $\delta^{15}\text{N}$  ratios of particulate nitrogen ( $\delta^{15}\text{N}_{\text{PN}}$ ) as well as particulate organic carbon content (POC) to investigate N-cycling processes in the BUS and the isotopic composition of nitrate as a fingerprint of its origin. With the data, we quantify the nitrate deficit generated by denitrification and phosphate release, estimate how much denitrification contributes to the regional nitrate deficit, and identify mechanisms that restore the inorganic N:P ratios in the mixed layer of the adjacent South Atlantic.

## 2. Materials and Methods

### 2.1. Sampling

[7] Water samples were taken with a conductivity-temperature-depth (CTD)/rosette sampler during cruises M76-2 (May/June 2008), D-356 (August/September 2010), and MSM17-3 (February/March 2011) and were filtered over precombusted GF/F Filters onboard. Filtrates were stored frozen in acid-washed polypropylene bottles for nutrient analysis and isotope analysis. Since the main focus of all cruises was on the 23°S transect and sampling was most detailed there, we concentrate on this transect.

[8] Particulate matter sampling was carried out by filtration onboard. Water was filtered over preweighed and precombusted (450°C) GF/F Filters (Whatman). Filters were dried immediately after filtration at 40°C for 48 h.

### 2.2. Analytical Methods

[9] Nutrients of M76-2 samples were measured with a Bran & Luebbe autoanalyzer in the home laboratory in Hamburg by applying standard methods. Nutrient analyses

of D-356 were carried out onboard and samples of MSM17-3 were measured in the home laboratory in Bremen. Standard methods and a Skalar autoanalyzer were used for the samples of D-356 and MSM17-3. Oxygen concentrations in discrete samples were measured by using Winkler methods and were further used to calibrate oxygen data obtained from the CTD.

[10] Particulate organic carbon (POC) was measured using a Thermo Flash EA 1112. Briefly, homogenized samples were weighed into silver capsules and inorganic carbon was removed by adding 1 M HCl suprapur prior to analyses. All POC data are given in weight%.

[11]  $\delta^{15}\text{N}$  of nitrate was determined using the “denitrifier method” [Casciotti et al., 2002; Sigman et al., 2001a] and are reported in ‰ using the delta notation

$$\delta^{15}\text{N}_{\text{sample}} = \left( \left( \frac{^{15}\text{N}/^{14}\text{N}}{\text{sample}} \right) / \left( \frac{^{15}\text{N}/^{14}\text{N}}{\text{reference}} \right) - 1 \right) * 1000 \quad (1)$$

with air  $\text{N}_2$  as reference for  $^{15}\text{N}/^{14}\text{N}$ . The values were calibrated using IAEA-N3 ( $\delta^{15}\text{N}_{\text{NO}_3} = +4.7\text{‰}$ ) and USGS-34 [ $\delta^{15}\text{N}_{\text{NO}_3} = -1.8\text{‰}$ ; Böhlke et al., 2003]. A potassium nitrate standard was measured twice within each run for quality assurance. Isotope values were corrected using the single point correction referring to IAEA-N3 for  $\delta^{15}\text{N}_{\text{NO}_3}$ . The standard deviation for IAEA-N3 was 0.3‰, which is within the same specification for  $\delta^{15}\text{N}_{\text{NO}_3}$  for at least duplicate measurements of the samples. Since nitrite was not removed prior to isotope analyses and since *Pseudomonas aureofaciens* used for the “denitrifier method” consume both nitrate and nitrite, we have to perform a nitrite correction of  $\delta^{15}\text{N}$  values in samples with significant nitrite concentrations (more than 1  $\mu\text{M}$ ). The correction was done using the equation:

$$\delta^{15}\text{N}_{\text{NO}_2} = \left( (\text{NO}_3 + \text{NO}_2) * \delta^{15}\text{N}_{\text{NO}_3 + \text{NO}_2} - (\text{NO}_3 * \delta^{15}\text{N}_{\text{NO}_3}) / \text{NO}_2 \right) \quad (2)$$

for samples from the bottom water of station #206 and #252. Because we lack genuine nitrite  $\delta^{15}\text{N}$  values for the BUS, we have to resort to literature data. In the Arabian Sea OMZ,  $\delta^{15}\text{N}_{\text{NO}_2}$  values are calculated from equation (2) to average  $-18.1\text{‰}$  (B. Gaye et al., Evidence of coupled denitrification and nitrification in the ODZ of the Arabian Sea from paired stable isotopes of nitrate and nitrite, resubmitted to *Global Biogeochemical Cycles*) which is in accordance with nitrite isotope data from the Eastern Tropical North Pacific OMZ [Casciotti and McIlvin, 2007].

[12]  $\delta^{15}\text{N}_{\text{PN}}$  ratios were measured using a Thermo Delta Plus XP mass spectrometer after combustion in a Thermo Flash EA 1112. A single-point calibration was performed with IAEA-N1 standard [Bahlmann et al., 2010]. Within each run, IAEA-N2 and IVA sediment standards have been measured to assure measurement quality and to assess the laboratories long-term measurement performance. The standard deviation for IAEA-N1 was less than 0.2‰ and duplicate measurements of suspended matter samples differ less than 0.3‰ due to inhomogeneous particle coverage of filters.

### 3. Results

[13] We concentrate on the data obtained for the sampling period in May/June 2008, when the shelf was partially anoxic, and use nutrient data from D-356 and MSM17-3 to demonstrate the high seasonal variability on the Namibian shelf. In May/June 2008, moderate coastal upwelling of water from 100 to 150 m water depth occurred at 23°S. On the shelf, the subsurface SACW contribution (Figure 1b) was highest (<85%, 50 m) on the inner shelf in the coastal upwelling region and decreased toward the outer shelf (<65%, 200 m). A patch of low SACW fraction (<40%) was found above the bottom at the shelf break (station #202). A distinct thermocline was present within the upper 50 m above the shelf break and further offshore. We found subsoxic conditions (<20  $\mu\text{M}$ ) in the bottom water of the inner shelf at 23°S and even surface waters near the coast were oxygen depleted ( $\text{O}_2 < 80 \mu\text{M}$ ; Figure 2a). The subsurface OMZ spread into the hemipelagic ocean where minimum  $\text{O}_2$  concentrations ranged between 40 and 60  $\mu\text{M}$ .


[14] Nitrate concentrations (Figure 2b) in the mixed layer were lowest near the coast and at the offshore station #201 (<2  $\mu\text{M}$ ). A “nitrate plume” above the shelf break with significantly elevated concentrations by >5  $\mu\text{M}$  compared to the adjacent stations lifted the nitracline up into the thermocline. A similar pattern was observed for phosphate (Figure 2c): elevated concentrations occurred above the shelf break, but differing from nitrate, highest concentrations (>1.5  $\mu\text{M}$ ) were found near the coast. Nitrite concentrations (Figure 2d) above 1  $\mu\text{M}$  were restricted to subsoxic bottom water on the inner shelf with a maximum of 4.4  $\mu\text{M}$  at 60 cm above the seafloor at station #252. Ammonium concentrations (Figure 2e) were above 1  $\mu\text{M}$  throughout the inner shelf, with maximum values in coastal surface water between 6 and 7  $\mu\text{M}$ , and above 8  $\mu\text{M}$  at the seafloor (#206).

[15] Suspended matter concentrations (SPM, see Figure S1 in the supporting information) and particulate organic carbon content (POC, see Figure S2) correlated near the seafloor. Both increased in bottom water toward the inner shelf from 4.9 to ~30 mg/L in the bottom nepheloid layer and from 2.8% to 6.0%, respectively. SPM concentrations were high in surface water near the coast (~19 mg/L) at co-occurring low POC concentrations of 2.7–2.9%. SPM concentrations decreased in surface water in offshore direction to 4.2 mg/L at the most offshore station. POC content in surface water was high at stations #206 and #198. Further offshore, high POC contents in surface waters were observed at co-occurring low SPM concentrations and vice versa. At offshore station #201, a pronounced deep surface SPM maximum (18 mg/L) occurred at 17 m water depth at a remarkably low POC content of 1.2%. Bottom waters at the shelf break were rich in SPM (12.5 mg/L at 5 m above seafloor) and POC (4%) compared to the surrounding waters.


[16] Bottom waters at the shelf break had excess reactive nitrogen with respect to phosphate. DIN/P ratios (see Figure S3) were calculated as  $(\text{NO}_3^- + \text{NO}_2^- + \text{NH}_4^+)/\text{PO}_4^{3-}$  and ranged between 21 and 16 at the outer shelf stations. DIN/P ratios decrease on the inner shelf down to 3.4 in coastal surface water.

[17] The  $\delta^{15}\text{N}_{\text{NO}_3}$  of water at the shelf edge—the source for upwelling waters over the shelf—ratios ranged from 5.7‰ to 6.7‰ (Figure 3a). These values are slightly higher

than deep ocean nitrate originating from subduction around the polar front (5.5‰) [Sigman *et al.*, 2001b]. By contrast, bottom water nitrate on the inner shelf OMZ was most



**Figure 2.** Interpolated concentrations during M76-2 (May–June 2008) on the 23°S transect normal to the coast. (a) Oxygen, (b) nitrate, (c) phosphate, (d) nitrite, and (e) ammonium. Numbers mark individual concentrations that deviate from the color code due to steep concentration gradients.



**Figure 3.** Interpolated isotopic compositions in ‰ during M 76-2. (a)  $\delta^{15}\text{N}_{\text{NO}_3}$  and (b)  $\delta^{15}\text{N}_{\text{PN}}$ .

enriched in  $^{15}\text{N}$  ( $\delta^{15}\text{N}_{\text{NO}_3}$  values between 28‰ and 29.6‰) at station #252. The nitrite correction (equation (2)) was performed for these samples because nitrite content was 32% of the combined  $\text{NO}_2^- + \text{NO}_3^-$  pool. High  $\delta^{15}\text{N}_{\text{NO}_3}$  values (<13.9‰) were also found within the mixed layer at the most distal station #201 on the 23°S transect, where nitrate concentrations dropped below  $2 \mu\text{M}$ . Stable nitrogen isotope ratios of particulate nitrogen ( $\delta^{15}\text{N}_{\text{PN}}$ ) ratios ranged between 3.0‰ in the surface water at the most distal station (#201) and 11.0‰ in intermediate water (50 m) at near-shore station #252 (Figure 3b).  $\delta^{15}\text{N}_{\text{PN}}$  was low (<5.5‰) in surface waters at the outer shelf and upper slope. High  $\delta^{15}\text{N}_{\text{PN}}$  (>7‰) characterized all suspensions in intermediate and bottom water with the exceptions of shelf break bottom water (6.2‰) and bottom water at station #206 (6.7‰).

#### 4. Discussion

[18] N-loss processes such as heterotrophic denitrification and anammox are known to occur in the Benguela OMZ [Kuypers *et al.*, 2005; Tyrrell and Lucas, 2002], but there is only little experimental evidence for  $\text{N}_2$  fixation balancing the N-loss, and the rates observed were too low to compensate the N-loss [Sohm *et al.*, 2011]. Despite a pronounced N-deficit on the inner Namibian shelf, however, we find a balanced inorganic N:P ratio ( $(\text{NO}_3^- + \text{NO}_2^- + \text{NH}_4^+)/\text{PO}_4^{3-}$ ) of  $\sim 16$  in surface water offshore the shelf break in May/June 2008. This suggests a source of inorganic N in surface water advected from the shelf to the open ocean. Below we address the question if the BUS is a closed system in which N-losses are compensated by N-supply restoring Redfield conditions. If this is not the case and N-losses exceed N-supply, the BUS would be a sink of fixed N, thus weakening the  $\text{CO}_2$  uptake capacity of the regional biological pump. We then investigate seasonal variability linked to water mass and oxygen dynamics, and attempt to quantify the deficit of inorganic N transferred into the adjacent hemipelagic ocean.

#### 4.1. Processes of N-Cycling on the Namibian Shelf

[19] Our discussion of data and transformations of reactive N follows mainly a 2D cross shelf circulation model view. This is applicable to the circulation on the shelf, but has some weaknesses for the shelf break and adjacent oceanic stations. Mohrholz *et al.* [2008] have shown that the cross-shelf circulation on the shelf is controlled by upwelling dynamics with offshore Ekman transport in the mixed layer and an onshore compensation flow below the thermocline. Offshore of the shelf edge and below the surface layer, the circulation is dominated by meridional currents, namely the Benguela current and the poleward undercurrent. Although the meridional flow on the shelf is prominent [Monteiro and van der Plas, 2006; Monteiro *et al.*, 2008], one can interpret the conditions observed on the transect off Walvis Bay as a projection of the 3D picture onto a 2D transect. This is because the meridional extent of the upwelling area is large compared to the shelf width. Further assuming that processes are uniform along the coast, a 2D transect depiction is mainly controlled by cross-shelf dynamics. This assumption is also applicable for the surface mixed layer beyond the shelf edge, because the Ekman offshore transport dominates the flow on longer time scales.

[20] Phosphate and ammonium concentrations (Figures 2c and 2e) increase in bottom water from the outer shelf toward the coast. The increase is caused by intense remineralisation of organic matter in sediments of the inner shelf. Ammonium and phosphate diffuse out of the sediments under low oxygen conditions [Neumann, 2012]. The benthic ammonium supply is entrained into the cross shelf flow and wells up near the coast.

[21] Significant amounts of nitrate in sub-thermocline water on the shelf are consumed within the OMZ on the inner shelf (Figure 2b) as soon as decreasing oxygen concentrations reach the threshold of  $\sim 5\text{--}10 \mu\text{M}$  for heterotrophic denitrification and anammox [Cline and Kaplan, 1975; Jensen *et al.*, 2008]. In sub-thermocline water on the outer shelf and at station #206,  $\delta^{15}\text{N}_{\text{NO}_3}$  and  $\delta^{18}\text{O}_{\text{NO}_3}$  (data not shown) change in parallel. This pattern is consistent with nitrate reduction (the first step of heterotrophic denitrification) [Granger *et al.*, 2008], which was detected in the bottom nepheloid layer throughout the OMZ in low rates [Füssel *et al.*, 2011]. The  $\delta^{15}\text{N}_{\text{NO}_3}$  and  $\delta^{18}\text{O}_{\text{NO}_3}$  values continue to rise toward the inner shelf while the nitrate deficit ( $\text{N}_{\text{def}}$ ; see Figure S4) increases. This indicates progressive nitrate consumption by nitrate reduction. Other nitrate and nitrite consuming processes which could account for the evolving nitrate deficit in the deep water over the shelf at 23°S are anammox and dissimilative nitrate reduction to ammonium (DNRA). Anammox is possibly the dominant reason for the N-deficit on the Namibian shelf [Kuypers *et al.*, 2005] and anammox rates exceed nitrate reduction rates at station #252 [Füssel *et al.*, 2011]. Nitrate ammonifiers were not evident at station #252 [Füssel *et al.*, 2011] and not responsible for decreasing nitrate concentrations in waters of the OMZ. From our data, we cannot definitely state which process is the most effective in eliminating reactive N before the water upwells into the euphotic mixed layer.

[22] As the upwelled water moves across the shelf, the  $\delta^{15}\text{N}_{\text{NO}_3}$  values in surface waters decrease from the stations closest the coast to roughly the shelf break. Near the coast,

particulate matter that is apparently produced in the surface layer by phytoplankton has higher  $\delta^{15}\text{N}_{\text{PN}}$  values than  $\delta^{15}\text{N}$  of the ambient nitrate. This suggests that a relatively  $^{15}\text{N}$ -enriched N-source is assimilated by phytoplankton, whereas a relatively  $^{15}\text{N}$ -depleted source adds to the dissolved nitrate pool. For both, the branching of high ambient ammonium concentrations into either biomass (ammonium assimilation) or nitrate (nitrification) is a good candidate [Wankel *et al.*, 2007].

[23] High ammonium concentrations (Figure 2e) in coastal surface water most likely originate from two sources. The first is benthic remineralization of organic matter, because both ammonium and phosphate concentrations are elevated in freshly upwelled waters near the coast. In addition to the benthic source, an ammonium source from ammonification of particulate nitrogen in the water column is indicated by a vertical decrease of SPM concentrations from the mixed layer (~10 mg/L) to 50 m water depth (~5 mg/L; see Figure S1). Additionally, POC content decreases in the upper 50 m of station #252 (see Figure S2). At the same time, the average  $\delta^{15}\text{N}_{\text{PN}}$  ratio of the sinking material increases from 7.8‰ in the mixed layer to 9.5‰ at 50 m water depth (Figure 3b), indicating a preferential release of  $^{15}\text{N}$ -depleted ammonium during ammonification. The ammonium mixture produced is partly assimilated and causes the mismatch between  $\delta^{15}\text{N}_{\text{NO}_3}$  and  $\delta^{15}\text{N}_{\text{PN}}$  seen at station #252. Here, particles in the surface water have higher  $\delta^{15}\text{N}$  ratios than ambient nitrate ( $\delta^{15}\text{N}_{\text{PN}} = 8.3\text{‰}$  versus  $\delta^{15}\text{N}_{\text{NO}_3} = 7.5\text{‰}$  and 7.8‰), whereas biomass from assimilation of that nitrate should have a  $\delta^{15}\text{N}_{\text{PN}}$  that is 5‰ more negative than ambient  $\delta^{15}\text{N}_{\text{NO}_3}$  [Granger *et al.*, 2004]. This implies that assimilation of isotopically enriched ammonium—associated with a fractionation factor of 6 to 27‰ [Hoch *et al.*, 1992; Wankel *et al.*, 2007; Waser *et al.*, 1998]—raises the  $\delta^{15}\text{N}_{\text{PN}}$ .


[24] Additions from nitrification of ammonium on the other hand decrease the  $\delta^{15}\text{N}$  of the nitrate pool, because marine nitrification discriminates strongly against  $^{15}\text{N}$  ( $^{15}\epsilon$  between 14‰ and 19‰) [Casciotti *et al.*, 2003]. The branching of ammonium fluxes to either nitrification or ammonium assimilation, associated with different fractionation factors, thus leads to nitrate that is more  $^{15}\text{N}$ -depleted than the suspended matter produced by ammonium assimilation of the residual [Wankel *et al.*, 2007].

[25]  $\delta^{15}\text{N}_{\text{PN}}$  ratios (Figure 3b) decrease in offshore direction from 7.2‰ at station #206 to values between 5.5‰ and 3.8‰ at the shelf break (station #198) coincident with decreasing ammonium concentrations (Figure 2e). Nitrate rather than ammonium is the predominant N-source for primary production offshore of station #206 and low  $\delta^{15}\text{N}_{\text{PN}}$  ratios mimic low  $\delta^{15}\text{N}_{\text{NO}_3}$  ratios of newly produced nitrate [see also Granger *et al.*, 2011]. The production of nitrate in the oxygen gradient of OMZs is a typical feature also known from the Eastern Tropical North Pacific [Ward and Zafriou, 1988], the Eastern Tropical South Pacific [Lipschultz *et al.*, 1990; Ward *et al.*, 1989], and from the Arabian Sea (B. Gaye *et al.*, resubmitted manuscript). As surface water moves offshore with the Ekman transport, decreasing primary production at decreasing nitrate concentrations causes decreasing SPM concentrations (see Figure S1). A contribution of  $\text{N}_2$  fixation is seen in the oceanic end member (station #201), where nitrate is almost completely consumed (Figure 2b), but SPM concentration

is high and has a low  $\delta^{15}\text{N}_{\text{PN}}$  value of 3.0‰ (Figure 3b). Nitrate in surface water of station #201 is almost consumed (0.14  $\mu\text{M}$  at 11 m water depth) and  $\delta^{15}\text{N}_{\text{NO}_3}$  and  $\delta^{15}\text{N}_{\text{PN}}$  at 17 m water depth are 13.9‰ and 3.0‰, respectively. Beyond the shelf break, *Trichodesmium* occasionally occurs and 1–10 *Trichodesmium* colonies per cubic meter were encountered in plankton nets in February 2011 in the northern BUS (H. Auel, personal communication, 2011).

#### 4.2. Seasonal Variability of Water Masses and Nutrient Composition on the Namibian Shelf

[26] Nutrient concentrations and nutrient ratios in the northern BUS are highly variable due to the seesaw of two distinct water masses caused by physical processes. Nutrient-depleted and oxygenated ESACW mixes on the shelf with nutrient-rich and oxygen-depleted SACW between the Lüderitz upwelling cell (26°S) and the Angola-Benguela-Frontal-Zone at 17°S [Mohrholz *et al.*, 2008]. Since ESACW and SACW have completely different oxygen and nutrient concentrations (as well as differing nitrate/phosphate ratios), prevailing upwelling conditions control oxygenation and nutrient conditions over the shelf. This is illustrated in Figure 4 that depicts two different situations on the 23° transect. In August/September 2010 (cruise D356), DIN/P ratios on the shelf ranged between 13.6 and 22.8 (Figure 4b) when intense upwelling occurred in the northern BUS. In this season, a weak poleward undercurrent and strong cross-shelf compensation currents transported waters with high ESACW proportions onto the shelf [Mohrholz *et al.*, 2008]. This mixture had high inorganic N/P ratios. Poole and Tomczak [1999] propose low N:P ratios of ~9 for the ESACW in their water mass definitions of the Atlantic Ocean. But they define the core depth of ESACW between 350 and 600 m, which is below our sampling depths. Because oxygen concentrations on the shelf are  $>30 \mu\text{M}$  (Figure 4a), denitrification and release of ammonium and phosphate from sediments were inhibited so that the high inorganic N:P ratios of upwelling source waters reached the surface. Bottom water DIN/P ratios increased from the outer to the inner shelf, where sulfur bacteria abundantly inhabiting the benthic boundary layer take up phosphate during oxic periods to store it within the cells as polyphosphate [Brock and Schulz-Vogt, 2011]. Consequently, arriving high DIN:P ratios further increase and are transported into surface water by upwelling and Ekman transport. By contrast, a weak cross-shelf compensation current and strong poleward undercurrent activity controlled the water mass composition in the northern BUS during cruise MSM17-3 in February/March 2011. ESACW had retracted to the south and the dominance of oxygen-poor SACW (with DIN:P ratios around 16:1) promoted the development of suboxic conditions on the shelf (Figure 4c). This stimulated heterotrophic denitrification and anammox and created an N-deficit by either denitrification or phosphate release from sediments leading to lowered DIN/P ratios (Figure 4d). Phosphate can be released from the seafloor under anoxic conditions from internal sources from the inner-shelf organic-rich diatomaceous mud [Bailey, 1991]. Under changing oxic/anoxic conditions, phosphate-sequestering bacteria play a role in generating recent concretary phosphorites on the landward flank of the diatomaceous mud belt and in releasing mediate



**Figure 4.** (a)  $O_2$  concentrations on the  $23^\circ S$  transect show enhanced oxygenation during August/September 2010 that was caused by strong upwelling and cross-shelf currents transporting ESACW with (b) elevated molar DIN:P ratios onto the shelf. In February/March 2011, the (c) OMZ covered almost the entire shelf and (d) molar DIN:P ratios decreased from 17.5 above the upper slope to 5 in coastal surface water.

phosphorous from sediments [Nathan *et al.*, 1993]. This source likely contains a contribution from large sulfur bacteria inhabiting the diatomaceous mud on the inner shelf: Under anoxic conditions and sulfide exposure, intracellular polyphosphate stored under oxic conditions is decomposed and released to pore water [Brock and Schulz-Vogt, 2011]. Because sediment pore water and ambient bottom water are in isotopic disequilibrium regarding oxygen isotopes of phosphate ( $\delta^{18}O_P$ ), the imprint of regenerated phosphate is evident under anoxic conditions [Goldhammer *et al.*, 2011] and further decreases the DIN:P ratio in the water column on the inner shelf. Phosphate concentrations in pore water of the diatomaceous mud reach up to  $<500 \mu M$

in 10–20 cm below seafloor [Goldhammer *et al.*, 2011] and increase to  $>1000 \mu M$  during prolonged suboxic periods over the central Namibian shelf [Bartholomae and van der Plas, 2007; van der Plas *et al.*, 2007].

### 4.3. Quantification of the Nitrate Deficit and Denitrification

[27] N-loss and P-gain both affect the local deficit of inorganic N ( $N_{def}$ ) in the system. Devol *et al.* [2006] defined  $N_{def}$  as the difference between the preformed N:P conditions ( $N^{pref}$ ) and the measured inorganic N-pool ( $NO_3^-$ ,  $NO_2^-$  and  $NH_4^+$ ):


$$N_{def} = N^{pref} - N^{meas} \quad (3)$$

[28] To estimate the local  $N^{pref}$ , we plotted DIN ( $[NO_3^-] + [NO_2^-] + [NH_4^+]$ ) versus  $[PO_4^{3-}]$  from water samples that were not affected by denitrification and phosphate release ( $O_2 > 65 \mu M$ ) nor by assimilation processes (50–500 m water depth). The slope of the regression in Figure 5 is 18.22 and the intercept on the  $PO_4^{3-}$  axis is 0.01, so that

$$N^{pref} = 18.22 \times (PO_4^{3-} - 0.01) \quad (4)$$

[29] Offshore of the shelf break,  $N_{def}$  (see Figure S4) is negative between 100 and 650 m water depth and mirrors a slight N excess over phosphate.  $N_{def}$  increases toward the mixed layer and laterally toward the shelf, to maximum values in coastal surface water at station #192 ( $27.2 \mu M$ ) and within the bottom nepheloid layer at #252 ( $28.6 \mu M$ ).

[30] During the expedition M76-2 in May/June 2008, upwelling source water at the shelf break had DIN:P ratios  $\sim 20:1$  (see Figure S3) that correspond to a  $N_{def}$  of approximately  $-2 \mu M$ . This is in accordance with nutrient data for SACW of van der Plas *et al.* [2007]. During onshore advection with the cross-shelf compensation current,  $O_2$  concentrations decrease from 70 to  $0 \mu M$  in inner shelf bottom water. Assuming a threshold for heterotrophic denitrification of  $\sim 5$  and of  $\sim 10 \mu M$  for anammox [Cline and Kaplan, 1975; Jensen *et al.*, 2008], both processes are



**Figure 5.** Linear correlation between dissolved inorganic nitrogen ( $NO_3^- + NO_2^- + NH_4^+$ ) and phosphate of sampling sites from M76-2 on the  $23^\circ S$  transect between 50 and 500 m water depth and  $O_2 > 65 \mu M$ .

feasible on the inner shelf below 70 m water depth. There, heterotrophic denitrification, anammox, and phosphate release decrease the inorganic N:P ratio to 7.1 (see Figure S3) and cause a  $N_{\text{def}}$  of 28.6  $\mu\text{M}$  in the bottom water of station #252 (see Figure S4). Integrated over the water column at 23°S and at  $\text{O}_2$  concentrations below 10  $\mu\text{M}$ , a cumulative  $N_{\text{def}}$  of 3 M N per  $\text{m}^2$  is generated by all three processes. Taking into account that the OMZ is roughly situated above the diatomaceous mud belt, which has an extension of 18,000  $\text{km}^2$  [Emeis *et al.*, 2004], we calculate an  $N_{\text{def}}$  amounting 0.75 Tg N for that area in May/June 2008. Water on the shelf moves eastward at a rate of 550 km per annum [Mohrholz *et al.*, 2008], so that water in the OMZ is completely exchanged 4.5 times a year. Assuming that the lower 30 m of the water column are suboxic for 9 months per year [Bartholomae and van der Plas, 2007; Joubert, 2006], our estimate for the inorganic N-deficit relative to phosphate amounts 2.5 Tg N per annum for the water column above the diatomaceous mud belt. This is in the same range of calculations by Kuypers *et al.* [2005] who attributed a loss of fixed N of  $1.4 \pm 1$  Tg on an area of 100,000  $\text{km}^2$  to anammox.

[31] To calculate the contribution of heterotrophic denitrification to  $N_{\text{def}}$ , we use the stable nitrate isotope data set. Heterotrophic denitrification is associated with a fractionation factor of 25‰ for stable N and O isotopes of nitrate [Granger *et al.*, 2004; Sigman *et al.*, 2009a; Voss *et al.*, 2001], leaving the residual nitrate isotopically enriched. In an upwelling system, nitrate is supplied constantly to the OMZ so that heterotrophic denitrification follows the “open system model” rather than the “Rayleigh closed system” model [Sigman *et al.*, 2009b]. Using the equation

$$\delta^{15}\text{N}_{\text{reactant}} = \delta^{15}\text{N}_{\text{initial}} + \varepsilon(1 - f) \quad (5)$$

for nitrate N isotopes, we calculated the contribution of denitrification to the  $N_{\text{def}}$ . Source nitrate at the shelf break had a  $\delta^{15}\text{N}_{\text{NO}_3}$  of 6.1‰, and  $f$  was calculated according to [Mariotti *et al.*, 1981]:

$$f = (\text{NO}_3^-) / (\text{NO}_3^- + N_{\text{def}}). \quad (6)$$

[32] The observed depth-integrated and nitrite-corrected average  $\delta^{15}\text{N}_{\text{NO}_3}$  of 18.1‰ within the OMZ requires that ~48% of the nitrate arriving at the core of the OMZ have to be eliminated by heterotrophic denitrification. Looking at the nitrate concentrations (Figure 2b), we find that this isotope-based calculation underestimates the nitrate elimination due to heterotrophic denitrification: water enters the OMZ containing ~18  $\mu\text{M}$  nitrate and lowest nitrate concentrations in the core of the OMZ are 7.0  $\mu\text{M}$ . Annual heterotrophic water column denitrification above the diatomaceous mud belt amounts to 0.25 Tg N assuming an average suboxic depth interval of 30 m ( $\text{O}_2 < 10 \mu\text{M}$ ) for 9 months per year and a mean residence time of water of 80 days. According to this calculation only 10% of the total  $N_{\text{def}}$  was due to heterotrophic denitrification during M76-2 in May/June 2008. Other candidates causing the  $N_{\text{def}}$  are phosphate release from sediments and anammox that is thought to eliminate the majority of reactive N within the northern BUS [Füssel *et al.*, 2011; Kuypers *et al.*, 2005].

#### 4.4. Restoration of Surface Water N:P Ratio

[33] The N-deficit in surface water is advected offshore from the coast by the Ekman transport and decreases successively from 27.2 to 0.2  $\mu\text{M}$  at station #201 (see Figure S4). Possible N-sources for filling the N-deficit in the Benguela region are (1) atmospheric or land input, (2)  $\text{N}_2$  fixation, (3) assimilation of nutrients with N:P ratio  $< 16$ , (4) nitrification, or (5) shelf break upwelling or mixing.

[34] Atmospheric dry deposition in the Atlantic Ocean between 30°S and 10°S has N:P ratios much above 16 and therefore is a potential N-source. The atmospheric nitrate input in the open Atlantic Ocean between 25° and 17°S is relatively low [ $< 20 \mu\text{M m}^{-2} \text{d}^{-1}$ ; Baker *et al.*, 2003]. Many arid environments have highly alkaline soils where ammonia is emitted to the atmosphere [McCalley and Sparks, 2008]. Dry deposition of ammonium is unlikely since  $\text{NH}_4^+$  and organic N are associated with fine mode aerosols ( $< 1 \mu\text{m}$ ) [Baker *et al.*, 2003] but fine sands dominate the sediments immediately along the Namibian coast where elevated ammonium concentrations occur. Ammonium concentrations in coastal surface water range between  $< 7 \mu\text{M}$  in May/June 2008 (Figure 2e) and 0.7  $\mu\text{M}$  in February/March 2011 (data not shown) during times of weak trade wind and upwelling activity. We attribute changing ammonium concentrations in surface water to remineralization processes at the sediment-water interface and the upwelling intensity rather than to atmospheric inputs.

[35]  $\text{N}_2$  fixation rates in the BUS or the adjacent South Atlantic Gyre are generally low. Highest  $\text{N}_2$  fixation rates ( $< 8 \text{ nmol N L}^{-1} \text{d}^{-1}$ ) were found within or near the BUS where high nitrate concentrations did not exclude  $\text{N}_2$  fixation [Sohm *et al.*, 2011]. This agrees with our data from May/June 2008 where patches of negative  $\delta^{15}\text{N}_{\text{PN}}$  (−1.8‰) occurred on the Namibian shelf at 20°S (data not shown). We find no isotopic indication of  $\text{N}_2$  fixation in surface water above the shelf at 23°S in either  $\delta^{15}\text{N}$  of nitrate or in  $\delta^{15}\text{N}_{\text{PN}}$ . Over the continental slope at the most offshore station #201, however, we find balanced N:P ratios where nitrate is almost completely consumed in the mixed layer (Figure 2b) and organic matter is isotopically enriched in  $^{15}\text{N}$  (13.9‰; Figure 3a). Suspended matter concentrations in the deep mixed layer (17 m water depth) are high and comparable to those in surface water near the coast (18 mg/L; see Figure S1). This indicates stimulated phytoplankton activity in surface water. Intense remineralization reduces POC content (see Figure S2) from surface water (4.3%) to the deep mixed layer (1.2%). Summarizing, we find nutrient-depleted surface water at station #201 (Figure 2), a deep-surface SPM maximum, and low  $\delta^{15}\text{N}_{\text{PN}}$  values of 3.0‰ at co-occurring high  $\delta^{15}\text{N}_{\text{NO}_3}$  values of 13.9‰. This pattern is consistent with  $\text{N}_2$  fixation. If it is indeed present, this is likely a small contribution, because the  $N_{\text{def}}$  in surface water (0–20 m) drops by only 0.05 M/ $\text{m}^2$  between stations #202 and #201. The corresponding N-supply by fixation to the BUS at 23°S is 6 times less than the N-loss estimated from nitrate isotopes.

[36] Another mechanism for N:P ratio restoration is the consumption of nutrients with N:P  $< 16$  under nutrient-replete conditions [Geider and La Roche, 2002; Weber and Deutsch, 2010]. In our study area, diatoms dominate phytoplankton on the inner shelf and nutrient-replete conditions favor nutrient assimilation with N:P  $< 16$ .



[37] Even if organic matter N:P ratios are below Redfield, remineralization releases nutrients with proportional N:P of  $\sim 16$  and thus helps to increase the initially low inorganic N:P ratio at the coast. Nitrification of ammonium from benthic and water-column sources plays an important role in the surface water of the inner shelf (see discussion above). Although the supply of preformed nitrate in upwelling systems is large, evidence for quantitatively significant nitrification within OMZs is accumulating [Füssel *et al.*, 2011; Lam *et al.*, 2011; B. Gaye *et al.*, resubmitted manuscript], and recycling of ammonium from particles and sediments contributes to the increase of N:P ratios offshore.

[38] Finally, shelf break upwelling and turbulent mixing across the thermocline are episodic processes that dilute and mask the N-deficit in modified upwelling water because inorganic N:P ratios in subsurface waters at the shelf break are above 16. Shelf break upwelling is reported to be active during high wind speed periods [Barange and Pillar, 1992; Pichevin *et al.*, 2005] and is caused by negative wind stress curl that roughly coincides with the shelf break. Using an analytical model, Fennel and Lass [2007] and Fennel *et al.* [2012] have shown that the structure of the wind field off Namibia forces curl-driven upwelling on the entire shelf. This process supplies nutrients from the subsurface layers into the euphotic zone. Additionally, the breaking of internal waves and tides at the shelf edge increases the turbulent diapycnal mixing and also contributes to the entrainment of nutrients into the upper layer [Mohrholz and Heene, 2011]. Elevated SPM concentrations (Figure S1) and high  $\delta^{15}\text{N}_{\text{PN}}$  ratios (Figure 3c) in warm stratified surface water at the 23°S transect image a recent shelf break upwelling phase and a late stage plankton bloom due to an uplift of the nitracline (Figure 2b) and  $\delta^{15}\text{N}_{\text{NO}_3}$  isolines (Figure 3a). Shelf break mixing thus is an episodic phenomenon, but elevated biomass concentrations in surface water above the shelf break have been encountered on other cruises (N. Wasmund, personal communication, 2011). The water mass analysis (Figure 1b) illustrates a dominating ESACW portion at station #201. As shown above, ESACW imports nutrients with elevated inorganic N/P ratios and thus contributes to balance the  $N_{\text{def}}$ .

[39] The N-deficit generated in the OMZ cannot always be balanced as suggested by inorganic N/P ratios in surface water encountered during expeditions D356 and MSM17-3 (Figure 4). During late austral summer (MSM17-3; February/March 2011), when the OMZ spreads over the entire shelf (Figure 4c), DIN/P ratios in surface water were constantly  $< 7.5$  at nitrate concentrations offshore the inner shelf break  $< 1.5 \mu\text{M}$ . Even surface water temperatures around 20.5°C apparently did not stimulate  $\text{N}_2$  fixation anywhere on the transect. DIN/P ratios in surface water from September 2010 (expedition D356; Figure 4b) drop from  $\sim 22$  to  $\sim 6$  in offshore direction at surface water temperatures of 16.6°C at the most offshore station. This implies that the modified upwelling water still hosted the N-deficit generated during times when the water parcel was situated in the OMZ weeks to months before.

## 5. Summary and Conclusions

[40] The northern BUS is a highly variable system, because it depends on the seasonally changing water-mass

mixture over the shelf. This creates conditions leading to rapidly changing nutrient supply to surface waters over the shelf and in the adjacent hemipelagic ocean, to highly variable nutrient ratios in the modified upwelling waters, and to seasonally changing oxygenation on the shelf. Owing to the seesaw of two distinct water masses, N-cycling processes change or shift spatially. In May/June 2008 denitrification occurred in the moderately extended OMZ. Ammonium produced during benthic remineralization and during ammonification of organic matter in the water column accumulated in the OMZ and welled up near the coast. The branching of ammonium to either ammonium assimilation or nitrification leads to  $^{15}\text{N}$ -enriched particulate matter and replenishes the nitrate pool. Nitrate assimilation dominated primary production above the outer shelf.

[41] In the BUS, we calculate that the total N-deficit of 2.5 Tg N per year is created in the oxygen deficient water column overlying the shelf. Using the isotope-based “open system model” from Sigman *et al.* [2009b], we calculate that only 0.05 Tg N of this deficit is generated by denitrification of preformed nitrate. The global annual N-loss due to water-column denitrification matches 150 Tg [Codispoti *et al.*, 2001]. About one half of reactive N is removed in the three major OMZs located in the Arabian Sea (about 50 Tg  $\text{N a}^{-1}$ ) [Devol *et al.*, 2006], in the Eastern Tropical North Pacific and in the Eastern Tropical South Pacific (25–50 Tg  $\text{a}^{-1}$  in each region) [Codispoti *et al.*, 2001; Devol, 2008]. Compared to these large-volume OMZs, the Benguela OMZ is only a minor sink of oceanic reactive N.

[42] The P\*-concept of Deutsch *et al.* [2007] links oceanic  $\text{N}_2$  fixation with zones of reactive N removal in time and space. Possibly due to the intense seasonal variability, the BUS  $\text{N}_2$  fixation rate is low and differs from other upwelling areas [e.g., Fernandez *et al.*, 2011; Deutsch *et al.*, 2007], although conditions of modified upwelling water advected into the adjacent hemipelagic ocean are conducive for  $\text{N}_2$  fixation. Our isotope-based estimate and mass-balanced calculation illustrate that denitrification in the OMZ exceeds  $\text{N}_2$  fixation offshore by a factor of 6. In spite of this, the N-deficit generated in the OMZ was filled in offshore surface waters in May/June 2008 most probably due to physical mixing of sub-thermocline water with surface water at the shelf-break that diluted the N-deficit. The BUS is a dynamic system where N-loss and N-supply are temporally and spatially decoupled. The associated variability in nutrient ratios exported to the hemipelagic ocean may be the reason why diazotrophic  $\text{N}_2$  fixation is a sporadic and subordinate N-source in the system.

[43] **Acknowledgments.** We thank the captains and crews of R/V *Meteor*, R/V *Discovery*, and R/V *Maria S Merian* for their excellent technical support as well as T. Heene from IOW and Steffen Oesterle and Paloma Ellitson from NatMIRC Institute, Swakopmund, for operating the CTD and water samplers. Ilse Büns is gratefully acknowledged for nutrient analyses of water samples from cruise M76-2. Furthermore, we thank Birgit Gaye and two anonymous reviewers for their constructive feedbacks on this manuscript. Financial support by DFG (SPP511), BMBF (project GENUS), and the Helmholtz-Zentrum Geesthacht are gratefully acknowledged.

## References

Bahlmann, E., *et al.* (2010), Performance evaluation of nitrogen isotope ratio determination in marine and lacustrine sediments: An inter-laboratory comparison, *Org. Geochem.*, 41(1), 3–12, doi:10.1016/j.orggeochem.2009.05.008.

- Bailey, G. W. (1991), Organic carbon flux and development of oxygen deficiency on the modern Benguela continental shelf south of 22°S: Spatial and temporal variability, in *Modern and Ancient Continental Shelf Anoxia*, edited by E. V. Tyson, and E. A. Pearson, pp. 171–183, The Geological Society, London.
- Baker, A. R., S. D. Kelly, K. F. Biswas, M. Witt, and T. D. Jickells (2003), Atmospheric deposition of nutrients to the Atlantic Ocean, *Geophys. Res. Lett.*, *30*(24), doi:10.1029/2003GL018518
- Barange, M., and S. C. Pillar (1992), Cross-shelf circulation, zonation and maintenance mechanisms of *Nyctiphanes capensis* and *Euphausenia hanseni* (Euphausiacea) in the northern Benguela upwelling system, *Cont. Shelf Res.*, *12*(9), 1027–1042.
- Bartholomae, C. H., and A. K. van der Plas (2007), Towards the development of environmental indices for the Namibian shelf with particular reference to fisheries management, *Afr. J. Mar. Sci.*, *29*(1), 25–35.
- Böhlke, J. K., S. J. Mroczkowski, and T. B. Coplen (2003), Oxygen isotopes in nitrate: New reference materials for <sup>18</sup>O:<sup>17</sup>O:<sup>16</sup>O measurements and observations on nitrate-water equilibration, *Rapid Commun. Mass Spectrom.*, *17*, 1835–1846.
- Brock, J., and H. N. Schulz-Vogt (2011), Sulfide induces phosphate release from polyphosphate in cultures of a marine *Beggiatoa* strain, *ISME J.*, *5*, 497–506, doi:10.1038/ismej.2010.135.
- Casciotti, K. L., and M. R. McIlvin (2007), Isotopic analyses of nitrate and nitrite from reference mixtures and application to Eastern Tropical North Pacific waters, *Mar. Chem.*, *107*(2), 184–201, doi:10.1016/j.marchem.2007.06.021.
- Casciotti, K. L., D. M. Sigman, and B. B. Ward (2003), Linking diversity and stable isotope fractionation in ammonia-oxidizing bacteria, *Geomicrobiol. J.*, *20*, 335–353.
- Casciotti, K. L., D. M. Sigman, M. Galanter Hastings, J. K. Böhlke, and A. Hilker (2002), Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method, *Anal. Chem.*, *74*(19), 4905–4912.
- Cline, J. D., and I. R. Kaplan (1975), Isotopic fractionation of dissolved nitrate during denitrification in the eastern tropical North Pacific, *Mar. Chem.*, *3*, 271–299.
- Codispoti, L. A., J. A. Brandes, J. P. Christensen, A. H. Devol, S. W. A. Naqvi, H. W. Paerl, and T. Yoshinari (2001), The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the Anthropocene?, *Scientia Marina*, *65*, 85–105.
- Deutsch, C., J. L. Sarmiento, D. M. Sigman, N. Gruber, and J. P. Dunne (2007), Spatial coupling of nitrogen inputs and losses in the ocean, *Nature*, *445*, 163–167, doi:10.1038/nature05392.
- Devol, A. H., A. G. Uhlenhopp, S. W. A. Naqvi, J. A. Brandes, D. A. Jayakumar, H. Naik, S. Gaurin, L. A. Codispoti, and T. Yoshinari (2006), Denitrification rates and excess nitrogen gas concentrations in the Arabian Sea oxygen deficient zone, *Deep-Sea Res. I*, *53*, 1533–1547, doi:10.1016/j.dsr.2006.07.005
- Devol, A. H. (2008), Denitrification including anammox, in *Nitrogen in the Marine Environment*, 2nd ed., edited by D. G. Capone, D. A. Bronck, M. R. Mulholland, and E. J. Carpenter, pp. 263–302, Elsevier, Amsterdam, doi:10.1016/B978-0-12-372522-6.00006-2.
- Emeis, K.-C., U. Struck, T. Leipe, and T. G. Ferdelman (2009), Variability in upwelling intensity and nutrient regime in the coastal upwelling system offshore Namibia: Results from sediment archives, *Int. J. Earth Sci.*, *98*, 309–326.
- Emeis, K. C., V. Brüchert, B. Currie, R. Endler, T. Ferdelman, A. Kiessling, T. Leipe, K. Noli-Pearl, U. Struck, and T. Vogt (2004), Shallow gas in shelf sediments of the Namibian coastal upwelling ecosystem, *Cont. Shelf Res.*, *24*, 627–642.
- Fennel, W., and H. U. Lass (2007), On the impact of wind curls on coastal currents, *J. Mar. Syst.*, *68*, 128–142.
- Fennel, W., T. Junker, M. Schmidt, and V. Mohrholz (2012), Response of the Benguela Upwelling Systems to spatial variations in the wind stress, *Cont. Shelf Res.*, (<http://dx.doi.org/10.1016/j.csr.2012.06.004>).
- Fernandez C., L. Farias, and O. Ulloa (2011), Nitrogen fixation in denitrified marine waters, *PLoS One*, *6*(6), e20539, doi:10.1371/journal.pone.0020539
- Füssel, J., P. Lam, G. Lavik, M. M. Jensen, M. Holtappels, M. Günter, and M. M. M. Kuypers (2011), Nitrite oxidation in the Namibian oxygen minimum zone, *ISME J.*, doi:10.1038/ismej.2011.178.
- Geider, R. J., and J. La Roche (2002), Redfield revisited: Variability of C:N:P in marine microalgae and its biochemical basis, *Eur. J. Phycol.*, *37*, 1–17.
- Goldammer, T., B. Brunner, S. M. Bernasconi, T. G. Ferdelman, and M. Zabel (2011), Phosphate oxygen isotopes: Insights into sedimentary phosphorus cycling from the Benguela upwelling system, *Geochim. Cosmochim. Acta*, *75*, 3741–3756, doi:10.1016/j.gca.2011.04.006.
- Granger, J., D. M. Sigman, J. A. Needoba, and P. J. Harrison (2004), Coupled nitrogen and oxygen isotope fractionation of nitrate during assimilation by cultures of marine phytoplankton, *Limnol. Oceanogr.*, *49*(5), 1763–1773, doi:10.4319/lo.2004.49.5.1763.
- Granger, J., D. M. Sigman, M. F. Lehmann, and P. D. Tortell (2008), Nitrogen and oxygen isotope fractionation during nitrate reduction by denitrifying bacteria, *Limnol. Oceanogr.*, *53*(6), 2533–2545, doi:10.4319/lo.2008.53.6.2533.
- Granger, J., M. G. Prokopenko, D. M. Sigman, C. W. Mordy, Z. M. Morse, L. V. Morales, R. N. Sambrotto, and B. Plessen (2011), Coupled nitrification-denitrification in sediment of the eastern Bering Sea shelf leads to <sup>15</sup>N enrichment of fixed N in shelf waters, *J. Geophys. Res.*, *116*, C11006, doi:10.1029/2010JC006751.
- Hoch, M. P., M. L. Fogel, and D. L. Kirchman (1992), Isotope fractionation associated with ammonium uptake by a marine bacterium, *Limnol. Oceanogr.*, *37*, 1447–1459.
- Inthorn, M., V. Mohrholz, and M. Zabel (2006), Nepheloid layer distribution in the Benguela upwelling area offshore Namibia, *Deep Sea Res. I*, *53*, 1423–1438.
- Jensen, M. M., M. M. M. Kuypers, G. Lavik, and B. Thamdrup (2008), Rates and regulations of anaerobic ammonium oxidation and denitrification in the Black Sea, *Limnol. Oceanogr.*, *53*(1), 23–36.
- Joubert, W. (2006), Seasonal variability of sediment oxygen demand and biogeochemistry on the Namibian Inner Shelf, Masters thesis, 97 pp, University of Cape Town, Cape Town.
- Konovalov, S. K., C. A. Fuchsman, V. Belokopitov, and J. W. Murray (2008), Modeling the distribution of nitrogen species and isotopes in the water column of the Black Sea, *Mar. Chem.*, *111*, 106–124.
- Kuypers, M. M. M., G. Lavik, D. Wobken, M. Schmid, B. M. Fuchs, R. Amann, B. B. Jörgensen, and M. S. M. Jetten (2005), Massive nitrogen loss from the Benguela upwelling system through anaerobic ammonium oxidation, *Proc. Natl. Acad. Sci. U.S.A.*, *102*(18), 6478–6483, doi:10.1073/pnas.0502088102.
- Lam, P., M. M. Jensen, A. Kock, K. A. Lettmann, Y. Plancherel, G. Lavik, H. W. Bange, and M. M. M. Kuypers (2011), Origin and fate of the secondary nitrite maximum in the Arabian Sea, *Biogeosci.*, *8*, 1565–1577, doi:10.5194/bg-8-1565-2011.
- Lipschultz, F., S. C. Wofsy, B. B. Ward, L. A. Codispoti, G. Friedrich, and J. W. Elkins (1990), Bacterial transformations of inorganic nitrogen in the oxygen-deficient waters of the eastern tropical south Pacific Ocean, *Deep Sea Res.*, *37*, 1513–1541.
- Mariotti, A., J. C. Germon, P. Hubert, P. Kaiser, R. Letolle, A. Tardieux, and P. Tardieux (1981), Experimental determination of nitrogen kinetic isotope fractionation: Some principles; illustration for the denitrification and nitrification processes, *Plant and Soil*, *62*, 413–430.
- McCalley, C., and J. Sparks (2008), Controls over nitric oxide and ammonia emissions from Mojave Desert soils, *Ecosyst. Ecol.*, *156*, 871–881.
- Mohrholz, V., M. Schmidt, and J. R. E. Lutjeharms (2001), The hydrography and dynamics of the Angola-Benguela frontal zone and environment in April 1999, *S. Afr. J. Mar. Sci.*, *97*, 199–208.
- Mohrholz, V., C. H. Bartholomae, A. K. van der Plas, and H. U. Lass (2008), The seasonal variability of the northern Benguela undercurrent and its relation to the oxygen budget on the shelf, *Cont. Shelf Res.*, *28*, 424–441, doi:10.1016/j.csr.2007.10.001.
- Mohrholz, V., and T. Heene (2011), Observations of nonlinear internal waves at the Namibian shelf. 3rd Annual Science Forum of Benguela Current Commission, Swakopmund, Namibia 17–19 Oct. 2011.
- Monteiro, P. M. S., A. van der Plas, V. Mohrholz, E. Mabilhe, A. Pascall, and W. Joubert (2006), Variability of natural hypoxia and methane in a coastal upwelling system: Oceanic physics or shelf biology?, *Geophys. Res. Lett.*, *33*, L16614, doi:10.1029/2006GL026234.
- Monteiro, P. M. S., and A. K. van der Plas (2006), Low oxygen water (LOW) variability in the Benguela System: Key processes and forcing scales relevant to forecasting, in *Large Marine Ecosystems*, vol. 14, edited by V. Shannon, G. Hempel, P. Malanotte-Rizzoli, C. Moloney, and J. Woods, pp. 71–90, Elsevier, Amsterdam.
- Monteiro, P. M. S., A. K. van der Plas, J.-L. Mélice, and P. Florenchie (2008), Interannual hypoxia variability in a coastal upwelling system: Ocean-shelf exchange, climate and ecosystem-state implications, *Deep Sea Res. I*, *55*, 435–450, doi:10.1016/j.dsr.2007.12.010.
- Nathan, Y., J. M. Bremner, R. E. Loewenthal, and P. M. S. Monteiro (1993), Role of bacteria in phosphorite genesis, *Geomicrobiol. J.*, *11*, 69–76.
- Neumann, A. (2012), Elimination of reactive nitrogen in continental shelf sediments measured by membrane inlet mass spectrometry. PhD thesis, 162 pp, University of Hamburg.
- Pichevin, L., P. Martinez, B. Philippe, R. Schneider, and J. Giraudeau (2005), Nitrogen cycling on the Namibian shelf and slope over the last two climatic cycles: Local and global forcings, *Paleoceanography*, *20*, PA2006, doi:10.1029/2004PA001001.
- Poole, R., and M. Tomczak (1999), Optimum multiparameter analysis of the water mass structure in the Atlantic Ocean thermocline, *Deep Sea Res. I*, *46*, 1895–1921.

- Rixen, T., M. V. S. Guptha, and V. Ittekkot (2005), Deep ocean fluxes and their link to surface ocean processes and the biological pump, *Prog. Oceanogr.*, *65*, 240–259.
- Rixen, T., V. Ramaswamy, B. Gaye, B. Herunadi, E. Maier-Reimer, H. W. Bange, and V. Ittekkot (2009), Monsoonal and ENSO impacts on particle fluxes and the biological pump in the Indian Ocean, in *Indian Ocean Biogeochemical Processes and Ecology Variability*, edited by J.D. Wiggert, pp. 429, American Geophysical Union, Washington D.C.
- Shannon, L. V., and G. Nelson (1996), The Benguela: Large-scale features and processes and system variability, in *The South Atlantic: Present and Past Circulation*, edited by G. Wefer, et al., pp.163–210, Springer, Berlin.
- Sigman, D., J. Granger, P. J. DiFiore, M. F. Lehmann, R. Ho, G. Cane, and A. van Green (2005), Coupled nitrogen and oxygen isotope measurements of nitrate along the eastern North Pacific margin, *Global Biogeochem. Cycles*, *19*, GB4022, doi:10.1029/2005GB002458.
- Sigman, D. M., K. L. Casciotti, M. Andreani, C. Barford, M. Galanter, and J. K. Böhlke (2001a), A Bacterial Method for the Nitrogen Isotopic Analysis of Nitrate in Seawater and Freshwater, *Anal. Chem.*, *73*(17), 4145–4153.
- Sigman, D. M., K. L. Casciotti, and H. S. John (2001b), Nitrogen Isotopes in the Ocean, in *Encyclopedia of Ocean Sciences*, edited by J. H. Steele, pp. 1884–1894, Academic Press, Oxford.
- Sigman, D. M., K. L. Karsh, and K. L. Casciotti (2009a), Ocean process tracers: Nitrogen isotopes in the ocean, in *Encyclopedia of Ocean Sciences* (update from 2001), edited by J. H. Steele, et al., pp. 4138–4153, Academic Press, London.
- Sigman, D. M., P. J. DiFiore, M. P. Hain, C. Deutsch, Y. Wang, D. M. Karl, A. N. Knapp, M. F. Lehmann, and S. Pantoja (2009b), The dual isotopes of deep nitrate as a constraint on the cycle and budget of oceanic fixed nitrogen, *Deep Sea Res. Part I*, *56*, 1419–1439, doi:10.1016/j.dsr.2009.04.007.
- Sohm, J. A., J. A. Hilton, A. E. Noble, J. P. Zehr, M. A. Saito, and E. A. Webb (2011), Nitrogen fixation in the South Atlantic Gyre and the Benguela Upwelling System, *Geophys. Res. Lett.*, *38*, doi:10.1029/2011GL048315
- Tyrrell, T. (1999), The relative influences of nitrogen and phosphorus on oceanic primary production, *Nature*, *400*, 525–531.
- Tyrrell, T., and M. I. Lucas (2002), Geochemical evidence of denitrification in the Benguela upwelling system, *Cont. Shelf Res.*, *22*, 2497–2511.
- van der Plas, A. K., P. M. S. Monteiro, and A. Pascall (2007), Cross-shelf biogeochemical characteristics of sediments in the central Benguela and their relationship to overlying water column hypoxia, *Afr. J. Mar. Sci.*, *29*(1), 37–47.
- Voss, R., J. W. Dippner, and J. P. Montoya (2001), Nitrogen isotope pattern in the oxygen-deficient waters of the Eastern Tropical North Pacific Ocean, *Deep-Sea Res. I*, *48*, 1905–1921.
- Wankel, S. D., C. Kendall, J. T. Pennington, F. P. Chavez, and A. Paytan (2007), Nitrification in the euphotic zone as evidenced by nitrate dual isotopic composition: Observations from Monterey Bay, California, *Global Biogeochem. Cycles*, *21*, GB2009, doi:10.1029/2006GB002723.
- Ward, B. B., and O. C. Zafriou (1988), Nitrification and nitric oxide in the oxygen minimum zone of the eastern tropical north Pacific, *Deep Sea Res.*, *35*, 1127–1142.
- Ward, B. B., H. E. Glover, and F. Lipschultz (1989), Chemoautotrophic activity and nitrification in the oxygen minimum zone off Peru, *Deep Sea Res.*, *36*, 1031–1051.
- Waser, N. A., K. D. Yin, Z. M. Yu, K. Tada, P. J. Harrison, D. H. Turpin, and S. E. Calvert (1998), Nitrogen isotope fractionation during nitrate, ammonium and urea uptake by marine diatoms and coccolithophores under various conditions of N availability, *Mar. Ecol. Prog. Ser.*, *169*, 29–41.
- Wasmund, N., H. U. Lass, and G. Nausch (2005), Distribution of nutrients, chlorophyll and phytoplankton primary production in relation to hydrographic structures bordering the Benguela-Angolan front region, *Afr. J. Mar. Sci.*, *27*(1), 177–190.
- Watson, A. J. (1995), Are upwelling areas sources or sinks of CO<sub>2</sub>?, in *Upwelling in the Ocean: Modern Processes and Ancient Records*, edited by C.P. Summerhayes, et al., pp. 321–336, John Wiley and Sons, Chichester.
- Weber, T. S., and C. Deutsch (2010), Ocean nutrient ratios governed by plankton biogeography, *Nature*, *467*, 550–554, doi:10.1038/nature09403.
- Wetzel, P., A. Winguth, and E. Maier-Reimer (2005), Sea-to-air CO<sub>2</sub> flux from 1948 to 2003: A model study, *Global Biogeochem. Cycles*, *19*, GB2005, doi:10.1029/2004gb002339.