

1 Selenium biofortification of crops on a Malawi Alfisol under conservation agriculture

2

3 I.S. Ligowe^{1,2}, S.D. Young^{3*}, E.L. Ander⁴, V. Kabambe¹, A.D.C. Chilimba², E.H. Bailey³, R.M. Lark³ and P. C. Nalivata¹

4 ¹*Lilongwe University of Agriculture and Natural Resources, Bunda Campus, P.O. Box 219, Lilongwe, Malawi*

5 ²*Department of Agricultural Research Services, P.O. Box 30779, Lilongwe 3, Malawi*

6 ³*University of Nottingham, School of Biosciences, Sutton Bonington Campus, Loughborough, LE12 5RD, UK*

7 ⁴*Centre for Environmental Geochemistry, British Geological Survey, Nottingham NG12 5GG, UK*

8 *Corresponding author: - E-mail: scott.young@nottingham.ac.uk

9

10

11 **Abstract**

12 Biofortification with selenium (Se) may rely on rapid uptake by crops, following application, to offset progressive fixation into
13 unavailable organic forms of Se in soil. A biofortification study was conducted on an Alfisol within a long-term conservation agriculture
14 (CA) field trial at Chitedze Research Station, Malawi. The aim was to assess the dynamics of selenium bioavailability to a staple cereal
15 (*Zea mays*) and a range of legumes (cowpeas, groundnuts, pigeon peas and velvet beans) under CA management, as well as residual Se
16 effects in the year following biofortification. Isotopically labelled selenate (>99% enriched ⁷⁷Se^{VI}) was applied to each plot, in solution,
17 at a rate of 20 g ha⁻¹, at maize flowering (75 days after planting), in February, 2017. Samples of grain and stover from maize and legumes,
18 and topsoil, were collected at harvest in May, 2017 and May, 2018. Plant and soil samples were analyzed by ICP-MS for selenium
19 isotopes (⁷⁷Se and ⁷⁸Se). The concentration of ⁷⁷Se in the grain of maize and single-cropped legumes exceeded 200 µg kg⁻¹ in all the
20 treatments. This would contribute approximately 56 – 64 µg day⁻¹ to the Malawi diet, as refined maize flour. The fertilizer derived Se

21 concentration ratio of maize grain-to-stover Se were >1 in 2017 but <1 in 2018; which followed the same trend as the native soil-derived
22 Se in the residual year. In legumes the grain-to-stover concentration ratio was consistently < 1 in both years, except for the velvet beans.
23 Differences in CA management had minimal influence on ⁷⁷Se concentration in plant grain but the low yield in the single conventional
24 treatment reduced ⁷⁷Se uptake. Residual ⁷⁷Se in the soil (35 % of the applied) measured at harvest in 2017 was still present at harvest
25 in the residual year (2018) but was completely unavailable to any of the crops. Almost none of the remaining ⁷⁷Se was present in soluble
26 or phosphate-extractable forms and virtually all was present in the ‘organic’ (TMAH-extractable) fraction. Thus, annual Se applications
27 to maize would be necessary to maintain concentrations which could improve dietary supply and reduce current Se deficiency in Malawi.

28

29

30 Highlights.

31 Selenium at 20 g ha⁻¹ raises dietary levels above the recommended dietary allowance

32 Selenium fixation into humus-bound forms eliminates residual availability

33 Conservation agriculture treatments have little effect on selenium uptake by maize

34 Legumes (e.g. groundnuts) accumulate greater concentrations of selenium than maize

35

36

37

38 Keywords.

39 Maize, Malawi, conservation agriculture, selenium biofortification, micronutrient deficiency.

40

41

42

43

44

45

46

47

48

49

50

51 **1. Introduction**

52

53 Selenium (Se) is a trace element with an essential nutritional role in human and animal health. Selenium deficiency in humans has been
54 linked to thyroid gland dysfunction, irreversible brain damage, peripheral vascular diseases, chronic and degenerative osteoarthropathy
55 (Kashin–Beck disease), impaired immune response to viral infections, male infertility, pre-eclampsia in women, heart diseases and
56 higher risks for several types of cancers (Cardoso et al., 2015; Fairweather-Tait et al., 2011; Riaz and Mehmood, 2012). According to
57 the Institute of Medicine of the USA National Academy, the Se recommended dietary allowance (RDA) for Se is 55 $\mu\text{g day}^{-1}$ for adults
58 while the tolerable upper intake for adults is 400 $\mu\text{g day}^{-1}$ (Bendich, 2001).

59

60 Dietary Se intake can be strongly related to the availability of Se in soil (Fairweather-Tait et al., 2011; Mehdi et al., 2013; Rayman,
61 2008), especially where populations depend on local food production in Se-deficient regions. The bioavailability of selenium in soil
62 depends on supply factors, such as parent material and atmospheric inputs, and on soil factors that affect the strength of Se sorption such
63 as pH and the concentration of soil organic matter and hydrous oxides of Fe, Al and Mn (Fordyce et al., 2000; Lopez et al., 2017;
64 Rayman, 2008). It is recognized that selenite ions (HSeO_3^- , SeO_3^{2-}) are adsorbed strongly on hydrous oxides at low pH (3.5 – 6.5).
65 However, most Se in soils is usually organically bound in humus and there is evidence that different forms of humus can give rise to
66 differences in Se bioavailability (Qin et al., 2012). It has also been shown that soil microbial processes have some involvement in the
67 control of inorganic Se availability (Tolu et al., 2014). In highly weathered, acidic, oxide-rich tropical soils, as found in Malawi, low
68 bioavailable Se presents a serious restriction to dietary Se supply with estimates of 70 % of the population eating insufficient Se with
69 an average daily intake range of 27 – 45 $\mu\text{g capita}^{-1} \text{ day}^{-1}$ (Chilimba et al., 2011; Hurst et al., 2013; Joy et al., 2015) which is in close
70 agreement with national plasma Se data for adult women (Phiri et al., 2019).

71

72 Dietary intake of Se can be increased through agronomic ‘biofortification’ of crops (Broadley and White, 2006; Chilimba et al., 2012a;
73 Mathers et al., 2017) with application as Se-containing fertilizers applied to soil or as foliar sprays (Lopez et al., 2017). In particular,
74 the success of agronomic biofortification of staple cereal crops is well recognized (Broadley et al., 2010; Chilimba et al., 2012a; Chilimba
75 et al., 2014). The use of selenium-enriched fertilizers was the public-health solution successfully adopted in 1984 by Finland, that
76 resulted in an increase in Se intake from 38 $\mu\text{g d}^{-1}$, before fortification, to 80 $\mu\text{g d}^{-1}$ in 2000, assuming a daily energy intake of 10 MJ
77 (Broadley et al., 2006; Euroola and Hietaniemi, 2000; Hartikainen, 2005).

78

79 The use of enriched Se isotopes as tracers in field experimentation permits discrimination between soil-derived and fertilizer-derived Se
80 in crops and, potentially, allows the examination of residual effects of Se application in soil and crops. The approach has been made
81 possible by advances in ICP-MS technology, the commercial availability of several enriched stable Se isotopes and the fact that
82 comparatively small Se additions are required on a field plot scale (c. 1 mg m^{-2}) for realistic biofortification results. To date, there have
83 been relatively few studies that have utilized this approach in field experiments. Chilimba et al. (2012b) studied uptake of ^{74}Se in maize
84 in Malawi; Mathers et al. (2017) used ^{77}Se to audit the fate of Se applied to wheat in U.K. soils. More recently, Ligowe et al. (2020)
85 used enriched ^{77}Se to characterize uptake and residual availability to green vegetables grown in an Oxisol, Alfisol and Vertisol from
86 Malawi (Ligowe et al., 2020).

87

88 Conservation agriculture (CA) that focuses on minimum soil disturbance, the retention of crop residues and crop diversification, is one
89 of the cropping systems which has been heavily promoted in recent years in southern Africa (Thierfelder and Wall, 2010; Thierfelder et
90 al., 2017; Kassam et al., 2009). In addition to providing a coping strategy for climate change (Branca et al., 2011; Lipper et al., 2014;
91 Steward et al., 2019), conservation agriculture is widely reported to improve soil health and crop yield through partly increased soil
92 organic matter (Ligowe et al., 2017; Ngwira et al., 2012; Powlson et al., 2016).

93

94 The aim of this study was to examine the viability of Se biofortification in an Alfisol, managed under CA, and representing typical
95 agronomic circumstances in Malawi and, more broadly, in sub-Saharan countries. We used application of ^{77}Se -enriched selenate
96 ($^{77}\text{Se}_{\text{Fert}}$) to sub-plots of maize and selected legumes (cowpea, ground nuts, pigeon peas and velvet beans) within an established CA
97 rotational and intercropped trial. The objectives were (i) to assess Se availability, uptake and translocation within crops and (ii) to
98 quantify residual effects of Se application in the following season.

99

100 **2. Materials and Methods**

101

102 2.1. Study location

103 The study was carried out within a long-term CA trial situated at Chitedze Research Station (CRS), Malawi. The CRS is located on the
104 Lilongwe-Kasungu plains (13.973 S, 33.654 E) at 1145 m above sea level. The soils are Ferruginous Latosols, classed as Alfisols under
105 the USDA Soil Taxonomy (USDA, 1975) which are deep and free-draining with a well-developed structure. In the first year of the trial
106 establishment (2007), baseline topsoil (0-10 cm) analysis showed the following average values for the site: pH (H_2O) = 5.2 ± 0.02 , soil
107 organic matter content = 3.3 ± 0.13 %, total C content = 2.1 ± 0.07 %, total N content = 0.17 ± 0.01 %, available P = 14.2 ± 0.6 $\mu\text{g g}^{-1}$
108 and available K = 0.44 ± 0.03 $\text{cmol}_c \text{ kg}^{-1}$ (Ligowe et al., 2017).

109

110 2.2. Conservation agriculture (CA) trial design and treatments

111 The CA trial was designed with ten management treatments including the conventional approach to maize cultivation ('control') in a
112 randomized complete block design, with four replicates per treatment. The trial covers an approximate area of one hectare (Plate 1). The
113 long-term CA experiment treatments are described in Table 1.

114

115 2.3. Fertilizer application

116 A uniform recommended fertilizer rate of 69 kg ha⁻¹ N, 21 kg ha⁻¹ P₂O₅ and 4 kg ha⁻¹ S is applied, yearly, to all the treatments. The
117 application comprised 23 kg ha⁻¹ N, 21 kg ha⁻¹ P₂O₅ and 4 kg ha⁻¹ S as a basal fertilizer dressing at planting and 46 kg ha⁻¹ N, in the form
118 of urea, 21 days after planting.

119

120 2.4. Sub-plots for isotopically labelled selenate (⁷⁷Se_{Fert}) biofortification treatment

121 Within the middle of each plot of the CA trial, sub-plots (2 m x 2 m), were marked. Each sub-plot included 3 maize rows, with 24 maize
122 plants, in the case of plots in which maize was the sole crop (T1, T2, T3 and T5C and T5G). For the intercropped plots (T6, T7, and T8
123 in Table 1) the main plot was the middle 1 m² of the plot, consisting of one maize row and 2 side-by-side rows of legumes. In rotational
124 legume plots (T4C and T4G), having 27 legume plants per sub plot.

125

126 2.5. Application of ⁷⁷Se_{Fert}

127 Isotopically enriched selenate (⁷⁷Se^{VI}) was prepared from an elemental stock of enriched ⁷⁷Se⁰ (50 mg; 99.2% atom %) from Isoflex,
128 USA, following the method described by Mathers et al. (2017). Isotopically enriched selenate (⁷⁷Se_{Fert}) was then applied to the sub-plots
129 at a single rate of 20 g ha⁻¹ of Se, (2 mg m⁻²), as a potassium selenate solution, 75 days after planting, at the maize crop-tasseling stage.
130 Aliquots of the ⁷⁷Se stock solution containing 8 mg ⁷⁷Se were diluted to 500 mL with irrigation water then applied evenly to the sub-
131 plot (4 m²) using a watering can. After Se application, 500 mL of water were applied to each sub-plot to aid Se distribution and wash
132 Se solution off the leaves.

133

134 2.6. Soil sampling and preparation

135 Soil samples were taken from all the sub-plots soon after harvest (May 2017), approximately 80 days after Se application, and again a
136 year later (May 2018). Topsoil samples (0 – 20 cm), using an Edelman soil auger (Dow AgroSciences Limited, U.K.) were taken within
137 the middle row of the sub-plots, leaving a discard of 0.25 m from each end of the row. Eight sampling points per plot were used, four

138 from each side of the maize rows. Similarly, in legume plots, soil samples were taken within the two central rows, at eight sampling
139 points within the plot. The soil samples from the eight sampling points were mixed by hand in a bucket in the field shortly after
140 collection. Using the quartering method, a homogenized composite sample, c. 500 g, was collected from each plot. The soil samples
141 were air-dried, under shade, in the laboratory and sieved to <2 mm.

142

143 2.7. Plant sampling and preparation

144 Samples of maize and legumes were collected within the sub-plot rows. Two maize plants from each side of the middle row were
145 discarded and the four middle maize plants were harvested. Eight legume plants were sampled from the two middle rows, four plants
146 from each row. For the intercropped legumes, four plants were sampled from each side of the middle maize row, next to where the maize
147 plants were sampled. Maize cobs and legume pods were removed from the harvested crop samples; the maize grain was de-husked and
148 the legume grain shelled. A composite sample from each sub-plot was dried to constant weight in an oven (Scientific Laboratory Supply
149 Ltd) at 40 °C. The crop stover was carefully cut into small portions (5 - 10 cm), washed using tap water and distilled water to remove
150 soil dust and dried in perforated bags at 40 °C. Representative sub-samples (c. 200 g) of the biomass and grain were ground to <40 µm
151 using an ultra-centrifugal mill (Model ZM-200, Retsch, Haan, Germany).

152

153 The remaining crop residues were retained on the soil surface in the CA plots (T2 -T8) while in the conventional plot (T1), residues
154 were removed soon after harvest. The ⁷⁷Se demarcated, 2 m x 2 m, sub plots were fenced to prevent residue mixture and blowing off to
155 the treatment area.

156

157 2.8. Soil analysis

158 2.8.1. Total soil elemental analysis

159 Finely ground soil samples (~0.2 g DW) were digested with 2 mL 68 % Primar Plus™ HNO₃, 1 mL 60 % HClO₄ and 2.5 mL 70 % HF,
160 (Fisher Scientific Ltd., U.K.) in PFA digestion vessels using a 48-place Teflon-coated graphite heated block digester (Model A3,
161 Analysco Ltd, Chipping Norton, U.K.), as described in Mathers et al. (2017). Digested samples were initially diluted to 50 mL and then
162 1-in-10, both using Milli-Q water (18.2 MΩ cm), prior to analysis by a single-quadrupole inductively coupled plasma mass spectrometer
163 (ICP-MS) (Thermo-Fisher Scientific icap-Q) operating in He and H₂ cell modes. To measure ‘residual’ ⁷⁷Se_{Fert}, the samples were
164 analysed using a triple quadrupole ICP-MS (Thermo-Fisher Scientific icap-TQ) in O₂-cell mode.

165

166 2.8.2. ‘Soluble’ soil selenium - potassium nitrate (0.01 M KNO₃) extraction

167 Potassium nitrate extraction followed the procedure described by Chilimba et al. (2012a). Air dried < 2 mm sieved soil (4.0 g) was
168 shaken end-over-end in 20 mL of 0.01 M KNO₃ for 2 hr. The suspension was then centrifuged for 30 minutes at 2500 g and the
169 supernatant filtered to < 0.22 μm using a syringe filter (Millipore, Cork, Ireland). Nine mL of the filtered solution was matrix-adjusted
170 with 1 mL of 10 % tetramethyl ammonium hydroxide (TMAH) prior to analysis for ‘Soluble Se’ by ICP-MS operating in H₂-cell mode.

171

172 2.8.3. ‘Adsorbed’ soil selenium - potassium phosphate (0.016 M KH₂PO₄) extraction

173 Centrifuge tubes containing the (wet) soil plug remaining after the KNO₃ extraction were weighed and 20 mL of 0.016 M KH₂PO₄ (pH
174 4.8) was added. The soil suspensions were shaken again for 1 hr, then centrifuged and filtered before retaining an aliquot, preserved in
175 1% TMAH, for Se analysis by ICP-MS in H₂-cell mode.

176

177 2.8.4. ‘Organic’ soil selenium – Tetramethyl ammonium hydroxide (10 % TMAH) extraction

178 After weighing the tubes again, the remaining soil plug was re-suspended in 10 mL of 10 % (v/v) TMAH. The tubes were heated in
179 an oven at 70 °C, for 3 hr, and gently shaken by hand at 1.5 hr. The tubes were then centrifuged at 3500 rpm and the supernatant

180 was filtered to $< 0.22 \mu\text{m}$ using a Millex syringe driven filter unit (Millipore, Cork, Ireland). One mL of the supernatant was diluted
181 with 9 mL Milli-Q water and analyzed by ICP-MS in H_2 cell mode.

182

183 2.9. Plant analysis

184 2.9.1. Crop (grain and stover) elemental analysis

185 The elemental composition of the crops was determined following microwave digestion (Multiwave 3000; Anton Paar, U.K.) of 0.2 g
186 finely ground material in 6.0 mL 68% HNO_3 (Primar Plus™ grade; Fisher Scientific, U.K.). Digested samples were initially diluted to
187 20 mL with Milli-Q water and with a further 1-in-10 dilution step prior to analysis by ICP-MS.

188

189 2.10. Data processing and interference correction

190 Raw intensity data, counts-per-second (CPS), for ^{77}Se and ^{78}Se analyzed in H_2 -cell mode using single quadrupole ICP-MS were
191 processed to correct for drift and for the formation of Se hydrides ($^{76}\text{SeH}^+$, $^{77}\text{SeH}^+$) that interfere with determination of ^{77}Se and ^{78}Se
192 respectively, as described by Mathers et al. (2017). Triple-quadrupole ICP-MS analysis employing O_2 as a cell gas to mass-shift ^{77}Se
193 and ^{80}Se to m/z 93 and 96 respectively was necessary to reduce interference from ^{76}GeH hydride (m/z = 77) from soil-derived
194 germanium. In addition, a small correction for $^{76}\text{GeOH}$ (m/z 93) was also implemented based on measurement of ^{72}Ge and using a Ge
195 standard to determine the m/z ratio (72/93).

196

197 2.11. Estimation of dietary Se intake

198

199 Dietary Se intake was estimated for maize flour and legumes assuming daily consumption of either refined ($276 \text{ g capita}^{-1} \text{ day}^{-1}$) or
200 unrefined ($249 \text{ g capita}^{-1} \text{ day}^{-1}$) maize flour and an average legume consumption of $71 \text{ g capita}^{-1} \text{ day}^{-1}$, derived from a legume

201 consumption range of 55 to 87 g capita⁻¹ day⁻¹, as estimated by Joy et al. (2015b) in Malawi. The product of the crop Se concentration
202 ($\mu\text{g kg}^{-1}$) and the daily estimated flour/grain intake (kg) provides an estimated Se daily intake ($\mu\text{g day}^{-1}$).

203

204 2.12. Statistical data analysis

205 Analytical data for soil and plant analysis were statistically analyzed using linear mixed models fitted with the lme command in the
206 nlme library (Piheiro et al., 2017) for the R platform (R Core Team, 2017). For analyses in which observations from the two seasons
207 were combined, the linear mixed model was used to accommodate, in its random effects structure, correlations between observations on
208 the same plots in successive seasons, and the correlation between subplots in the rotation treatments (where the plots in different phases
209 of the rotation are split plots of the original design, and are not allocated independently to plots within the blocks). In the case of
210 measurements on crop Se (legume and maize, grain and stover) there were marked differences in the variability between the 2017 and
211 2018 harvests, so analyses were done separately for these two seasons. The combined analysis was used for soil measurements

212

213 The effects of treatments and seasons, and their interaction, were analyzed as fixed effects. Specific orthogonal contrasts among
214 treatments were identified a priori. These were as shown in Table 2.

215 Because the main effect of season has just one degree of freedom, the interaction of season and treatment can also be partitioned into
216 single degree of freedom-contrasts, for which the null hypothesis in each case is that any contrast is the same in the two seasons.

217

218 Note that contrasts C3, C5, C6 and C7 could be used to examine differences in measurements on legume crops, corresponding
219 specifically to (C3) a comparison between legume in monocrop (in rotation with maize) and legume in intercrop with maize, (C5) a
220 comparison between cowpea and groundnut monocrop, (C6) a comparison between velvet bean and cowpea and pigeon pea (sown at
221 the same time as the maize) in intercrop and (C7) a comparison between cowpea and pigeon pea both in intercrop.

222

223 3. Results and discussion

224

225 3.1. Concentration of $^{77}\text{Se}_{\text{Fert}}$ and Se_{Soil} in maize and legumes

226 *Maize*

227 In the year of $^{77}\text{Se}_{\text{Fert}}$ biofortification (2017) the concentrations of $^{77}\text{Se}_{\text{Fert}}$ in maize grain had a restricted range across all CA treatments
228 ($217 \pm 27 \mu\text{g kg}^{-1}$), including the conventionally cultivated maize plots, and showed no difference between treatments (Tables 3 and 4).
229 The results confirm the potential of fertilizer to increase Se in staple grain as has been demonstrated in previous studies (Broadley et al.,
230 2010, 2006; Chilimba et al., 2012a; Mathers et al., 2017).

231

232 In the following year, 2018, the concentration of $^{77}\text{Se}_{\text{Fert}}$ in maize grain was much lower, $0.7 - 3 \mu\text{g kg}^{-1}$, with the largest values recorded
233 in the maize grown in rotation with cowpea (Table 3). Thus, the applied $^{77}\text{Se}_{\text{Fert}}$ was mainly unavailable for plant uptake beyond the
234 single year of application (Fig. 2). Very similar results for a second crop were reported by Mathers et al. (2017) for wheat grown with
235 applied ^{77}Se on a site in the U.K. and Chilimba et al., (2012b) for maize with ^{74}Se applied grown at two field sites in Malawi. Non-
236 isotopically labelled studies by Stroud et al. (2010) and Gupta et al. (1993), with wheat and barley respectively, found no evidence of
237 residual Se in crops the year following application, regardless of application rate.

238

239 In the first harvest after biofortification (2017) the concentration of $^{77}\text{Se}_{\text{Fert}}$ was greater in the crop grain than in the stover whereas, in
240 the following year (2018) $^{77}\text{Se}_{\text{Fert}}$ concentration was greater in the crop stover (Table 3). The concentration ratio of $^{77}\text{Se}_{\text{Fert}}$ (grain:stover)
241 was between 2 and 3 in 2017 but was <1 ($0.11 - 0.94$) in 2018 (EA Table 1). Therefore, the residual $^{77}\text{Se}_{\text{Fert}}$ (in 2018) was translocated
242 less efficiently to the grain compared to the year of application (2017). This may reflect a change in speciation in inorganic Se between
243 the two years: selenate was applied in 2017 but selenite is more likely to be the main inorganic form of $^{77}\text{Se}_{\text{Fert}}$ in 2018 given the high
244 solubility of selenate and that the soil pH of the trial treatments was 4.2 to 6.0 (Ligowe et al., 2017). Stroud et al. (2010) noted that

245 selenite is the dominant inorganic form under ‘moderate redox conditions’ and is adsorbed strongly by soil, especially at low pH. They
246 measured the speciation of phosphate-extractable Se one year after applying selenate fertilizer and found only selenite and soluble
247 organic-Se species (Stroud et al., 2010).

248

249 In the 2017 season there was a difference in the mean concentrations of soil-derived selenium (Se_{Soil}) in the maize grain between the
250 mono-cropped and mix-cropped CA maize treatments (contrast C2, Table 4), and in 2018 there was a significant difference in this
251 variable between the rotation and intercropped CA systems (C3, Table 4). However, in 2017 the concentration of (Se_{Soil}) was 39 %
252 greater than the grain Se concentration in 2018. This might be due to the inter-seasons climatic, c. rainfall, variations. In the maize
253 stover the concentration of Se_{Soil} differed between treatments (Table 4, C1 (both years); C3, C4, C7 (2018)) and also between years (EA
254 Table 1). However, the Se_{Soil} grain-to-stover concentration ratios in 2017 and 2018 were approximately equal (both < 1), while the grain-
255 to-stover ratio of $^{77}Se_{Fert}$ in 2017 ranged between 1.9 and 3.2 (AE Table 2). The timing of the $^{77}Se_{Fert}$ application to plots, at maize
256 tasseling, may also have contributed to rapid translocation to the developing seed in 2017. Sankaran and Grusak (2014) and Sperotto
257 (2013) suggested that continued root uptake and translocation of minerals to the flower tissues during grain filling is more important
258 than the remobilization of previously stored minerals within the plant cells.

259

260 *Legumes*

261 The concentration of $^{77}Se_{Fert}$ in the legume grains differed significantly with cropping pattern (Table 5, contrast C3, both seasons); in
262 the year of $^{77}Se_{Fert}$ application, single-cropped legumes had greater $^{77}Se_{Fert}$ concentrations, 400 and 711 $\mu g\ kg^{-1}$ in cowpea and
263 groundnuts, respectively than in legumes intercropped with maize, with an average mean $^{77}Se_{Fert}$ concentration of 78 $\mu g\ kg^{-1}$. The
264 reduced $^{77}Se_{Fert}$ concentration in the intercropped legumes as compared to sole cropped, might be attributed to poor development and
265 growth of intercropped legumes due to the shielding effect (of maize) but also damage to the legumes planted in between the maize
266 rows, caused by manual weeding of the plots. Partitioning of $^{77}Se_{Fert}$ between the 2 crops (maize and legume) grown in close association

267 is likely to have contributed to the low $^{77}\text{Se}_{\text{Fert}}$ concentration in the intercropped legume grain. For example, 10 % and 0.44 % of the
268 applied $^{77}\text{Se}_{\text{Fert}}$ was taken up by maize and cowpeas, respectively, grown in association in 2017 showing the significance of the
269 intercropped legume in recovering the applied Se. In sole cropped legumes, groundnut $^{77}\text{Se}_{\text{Fert}}$ grain concentration was 43 % more
270 compared to the cowpea sole cropped grain in the 2017 season (Table 5, contrast 5, $P=0.004$). The variation of $^{77}\text{Se}_{\text{Fert}}$ concentration
271 between these sole cropped legumes might have been attributed to the variation in maturity and hence variation in the crop development
272 stage at the time of $^{77}\text{Se}_{\text{Fert}}$ application. The legume grain $^{77}\text{Se}_{\text{Fert}}$ in the following year after biofortification was negligible, with an
273 average mean of not more than $1.31\mu\text{g kg}^{-1}$ suggesting the non-accessibility of the residual $^{77}\text{Se}_{\text{Fert}}$ beyond the cropping season in which
274 it is applied.

275

276 The concentrations of Se_{Soil} in the same mono-cropped grain legumes of cowpea and groundnuts were indistinguishable (30 to $33\mu\text{g kg}^{-1}$,
277 respectively), implying that the legume type and differences in maturity, did not affect the accessibility of native soil Se.

278

279 *Dietary consequences*

280 In Malawi, more than 83% of the population live in rural areas and depend on locally produced crops (Erick et al., 2009). Cereals,
281 legumes and vegetables constitute the main dietary components (Ferguson et al., 1989; Donovan et al., 1991) for rural people not living
282 close to Lake Malawi. In the current study, the contribution to Se dietary intake from Se_{Soil} was very small. Using values for the mean
283 per-capita daily consumption of maize flour which is refined (*ufa oyera*; 249 g) and unrefined (*mgaiwa*; 276 g) (Joy et al. 2015a), Se
284 intake was estimated as $< 6\mu\text{g day}^{-1}$ from both sources (EA Table 3). This is far below the minimum recommended Se intake of $40\mu\text{g}$
285 day^{-1} required to prevent Se deficiency (Winkel et al., 2012). These results substantiate similar findings in a national survey of Se maize
286 grain concentration of $22\mu\text{g kg}^{-1}$ Chilimba et al. (2011) and an estimated Se dietary contribution of $< 7\mu\text{g day}^{-1}$ (Hurst et al., 2013)
287 from maize grown on non-calcareous soils that cover the majority (c. 95 %) of the cultivated land in Malawi (Joy et al., 2015b). Joy et
288 al., (2015b) suggested that food crops produced on non-calcareous soils in Malawi contribute approximately $19\mu\text{g day}^{-1}$ to Se intake,

289 presenting a very high risk of Se deficiency. By contrast, maize grain biofortified with Se_{Fert} at an application rate of 20 g ha^{-1} , would
290 contribute 51 to $58 \mu\text{g day}^{-1}$ when using unrefined maize flour and $56\text{-}64 \mu\text{g day}^{-1}$ for refined maize flour in the year of Se fertilizer
291 application. The CA maize grown in rotation with cowpea would have provided the greatest intake in both cases of maize flour
292 processing (EA Table 3). Thus, possible dietary intakes of Se would be within the daily recommended intake range of $50 - 70 \mu\text{g day}^{-1}$
293 (IMFNB, 2000; WHO, 2009; Malagoli et al., 2015) and well below the chronic toxicity level of $400 \mu\text{g day}^{-1}$ (Winkel et al., 2012). The
294 implication is that, even in combination with other dietary components from the region, biofortification of maize with Se could not lead
295 to excessive Se intake and would provide clear nutritional benefits.

296

297 The equivalent contribution of the legumes to Se dietary intake, assuming daily consumption of 71 g of legumes, Joy et al. (2015a), are
298 reported in EA Table 4. Single-cropped cowpea and groundnuts would provide 29 and $50 \mu\text{g day}^{-1}$, respectively, whereas intercropped
299 cowpea would contribute only $13 \mu\text{g day}^{-1}$ and intercropped pigeon peas and velvet beans would provide $< 2 \mu\text{g day}^{-1}$. The restrictions
300 imposed on legumes by the intercropped maize appear to negate the viability of intercropped legumes within a Se biofortification
301 program.

302

303 3.2. Offtake of $^{77}\text{Se}_{\text{Fert}}$ in the grain and stover of maize and legumes

304 *Maize*

305 The uptake of $^{77}\text{Se}_{\text{Fert}}$ was $>95\%$ larger in 2017 than in 2018, the second season after $^{77}\text{Se}_{\text{Fert}}$ application (Fig. 1). As hypothesized,
306 based on the crop Se concentration data, the $^{77}\text{Se}_{\text{Fert}}$ retained in the soil after the harvest of the first crop was largely in a form which was
307 not available for plant uptake in the following year. Uptake of $^{77}\text{Se}_{\text{Fert}}$ differed between the treatment combinations (Table 4, C1 and C3,
308 2017; C3, 2018). The conventionally cultivated maize (T1) produced the smallest $^{77}\text{Se}_{\text{Fert}}$ uptake among all the treatments (Table 4, C1,
309 $P=0.01$ in 2017) but the effect was not significant in 2018 (Table 4, C1, $P=0.536$ and Fig. 1). Similarly, maize grain $^{77}\text{Se}_{\text{Fert}}$ uptake was

310 slightly greater in the CA legume rotation treatments than in the CA legume intercropped plots, contrast C3, (P=0.035 in 2017; P=0.0002
311 in 2018). The trend in $^{77}\text{Se}_{\text{Fert}}$ uptake in the grain was also seen in the maize stover (Table 4).

312

313 The differences between treatments in $^{77}\text{Se}_{\text{Fert}}$ uptake in maize arose mainly because of differences in yield (rather than concentration)
314 in grain and stover (EA Table 5). Yields of maize grain in the conventional (T1) $^{77}\text{Se}_{\text{Fert}}$ sub-plots in 2017 and 2018 (2180 and 3510 kg
315 ha^{-1} respectively) were significantly smaller than from the CA plots (T2-T8) with average yields of 7820 and 7810 kg ha^{-1} in 2017 and
316 2018 respectively, which drives the significantly higher offtake on the CA treatments (Table 4; C1, 2017). This difference reflects the
317 improved soil fertility and physical characteristics in CA plots (Ligowe et al., 2017; Thierfelder and Wall, 2009). Uptake of Se_{Soil} was
318 also less in conventional (T1) with a mean concentration of 0.032 g ha^{-1} while the CA treatments recorded a mean Se_{Soil} uptake range
319 of $0.083 - 1.53 \text{ g ha}^{-1}$, hence the significance differences in C1 (Table 4). As with the $^{77}\text{Se}_{\text{Fert}}$, differences in Se_{soil} uptake were due to the
320 depletion of soil nutrients due to yearly soil tillage and residue removal that lead to the differences in crop yield (EA Table 5).

321

322 In the year of $^{77}\text{Se}_{\text{Fert}}$ application (2017), the grain:stover ratios of $^{77}\text{Se}_{\text{Fert}}$ uptake were 1.5 in the conventional cultivation plot and an
323 average of 3.7 in CA plots. However, in the residual year (2018) the ratios were less < 1.0 in all treatments (EA Table 6). The difference
324 between 2017 and 2018 may reflect differences in (i) timing and concentration of available $^{77}\text{Se}_{\text{Fert}}$ within the growing season or (ii) the
325 speciation of $^{77}\text{Se}_{\text{Fert}}$ in soil. In particular, the (residual) available $^{77}\text{Se}_{\text{Fert}}$ in 2018 was clearly much smaller and it is likely that it was
326 present as inorganic Se^{IV} (selenite) and humus-bound Se rather than the Se^{VI} (selenate) originally applied. An additional factor may have
327 been some unavoidable foliar application of $^{77}\text{Se}_{\text{Fert}}$ when the Se was originally sprayed on the plots – this might have contributed to the
328 greater ratio seen for the CA plots where ground cover by the retained crop residues was greater than in conventionally cultivated maize
329 plots. Whatever the combination of mechanisms, the data suggests that application of a large concentration of selenate at maize flowering
330 promotes transfer into the grain. By contrast, a greater proportion of Se taken up throughout the growing season appears to remain

331 deposited in the crop stover. The 2018 grain:stover ratios followed a trend similar to that of Se_{soil} in that more of the native (soil-derived)
332 Se was in the maize stover than in the maize grain (EA Table 3).

333

334 *Legumes*

335 Uptake of both $^{77}\text{Se}_{\text{Fert}}$ and Se_{soil} , by legume grain and stover, differed markedly between the two years, with negligible $^{77}\text{Se}_{\text{Fert}}$ uptake
336 in 2018 (Fig 1). In both years, $^{77}\text{Se}_{\text{Fert}}$ uptake under single-crop rotations (T4C and T4G), was significantly greater than for intercropped
337 legumes (Table 5; C3). As found for $^{77}\text{Se}_{\text{Fert}}$ concentrations, the uptake of $^{77}\text{Se}_{\text{Fert}}$ in intercropped plots, was hindered by the vigorous
338 growth of maize that suppressed the root and leaf development of the intercropped plants and thereby limited $^{77}\text{Se}_{\text{Fert}}$ uptake. Among
339 the single-crop rotation legumes (contrast C5), groundnuts grain and stover (T4G) recorded the greatest mean (2.76 and 2.34 g ha⁻¹ in
340 grain and stover, respectively) offtake in 2017 – significantly greater than cowpea (0.805 and 0.950 g ha⁻¹ in grain and stover) under
341 single-crop rotation (Fig. 1a). This may partly have arisen due to differences in maturity between the two crops. At the time of $^{77}\text{Se}_{\text{Fert}}$
342 application, 75 days after planting, cowpea was at the post-blossoming stage whereas groundnuts were still at the flowering stage.
343 Cowpea matures in 60-90 days (Abadassi, 2015) and groundnuts take between 110-120 days to mature (Nautiyal et al., 2010). This
344 might have contributed to the lower uptake of $^{77}\text{Se}_{\text{Fert}}$ in cowpea than in groundnuts. Hence the timing of $^{77}\text{Se}_{\text{Fert}}$ application in the maize
345 legume intercropping cropping system is crucial to maximize Se uptake and translocation to edible parts of both crops.

346

347 3.3. Fractionation of $^{77}\text{Se}_{\text{Fert}}$ in soil for the application year (2017) and residual year (2018)

348 A sequential extraction procedure to measure ‘Soluble’, ‘Adsorbed’, ‘Organic’ and ‘Total’ concentrations of $^{77}\text{Se}_{\text{Fert}}$ and Se_{Soil} was
349 undertaken on soil samples collected at the end of the two growing seasons: 2017 and 2018. The main aim was to assess how much of
350 the soil-retained $^{77}\text{Se}_{\text{Fert}}$ was fixed into organic matter (TMAH-extractable) and what proportion of the $^{77}\text{Se}_{\text{Fert}}$ applied was retained in a
351 bioavailable form (Soluble and Adsorbed) after the harvest of the biofortified crop in 2017 and after the harvest of the following crop
352 in 2018. The $^{77}\text{Se}_{\text{Fert}}$ results are presented in Fig. 2.

353

354 3.3.1. Soluble and adsorbed $^{77}\text{Se}_{\text{Fert}}$

355 Soluble $^{77}\text{Se}_{\text{Fert}}$ after crop harvest was 94 %, greater in 2017 as compared to 2018 (Figs 2a and 2b; Table 6; $p < 0.0001$). The
356 concentration of soluble $^{77}\text{Se}_{\text{Fert}}$ in 2017 was similarly greater in conventional (T1), 0.357 unlike in CA treatments with a mean of 0.156
357 ($p = 0.002$). Among the CA treatments the ‘groundnuts in rotation after maize’ (T4 G) plots gave higher values. The greater soluble
358 $^{77}\text{Se}_{\text{Fert}}$ in T1 and T4G suggests greater retention of the soluble $^{77}\text{Se}_{\text{Fert}}$ after the first crop.

359

360 The same trend was observed for adsorbed $^{77}\text{Se}_{\text{Fert}}$ where more was retained in the conventional and the rotation plots planted with
361 legumes, (contrast p values of 0.03 and 0.04, for contrast C1 and contrast C3, respectively; Table 6). Generally, it was observed that
362 both soluble and adsorbed $^{77}\text{Se}_{\text{Fert}}$ were retained to a greater extent in plots with single-cropped maize and legumes compared to rotational
363 and intercropped treatments (Figs 2a - 2d). This might be partly attributed to differences in uptake of $^{77}\text{Se}_{\text{Fert}}$ (Fig. 1) where total uptake
364 of $^{77}\text{Se}_{\text{Fert}}$ was greater in maize under rotation (T5C and T5G) and in intercropped plots (T6, T7 and T8) with a combined uptake by both
365 maize and legumes, as compared to the sole maize treatments (T1, T2 and T3). The results are consistent with Chilimba et al. (2014)
366 who investigated agronomic biofortification in intercropped systems and observed a greater recovery of Se in maize-groundnut
367 intercropped treatments compared to single cropped plots. Additionally, these findings may suggest improved $^{77}\text{Se}_{\text{Fert}}$ efficiency use
368 when two or more crops are grown in association, as reported by Salehi et al. (2018) for other plant nutrients (N). Thus, the reduced
369 soluble and adsorbed $^{77}\text{Se}_{\text{Fert}}$ in the maize grown after cowpea and groundnuts rotation (T5C and T5G) plots is consistent with greater
370 uptake of the applied $^{77}\text{Se}_{\text{Fert}}$ (Fig. 1), facilitated by greater yields of both maize grain and stover as a result of improved soil fertility
371 (EA Table 5).

372

373 Over the course of the residual year (after harvest 2017 until harvest 2018), there was a loss of soluble and adsorbed $^{77}\text{Se}_{\text{Fert}}$, (Figs 2a -
374 2d) representing average percentage reductions of 93 % and 18 % respectively. This was not due to uptake by the crops, because

375 concentrations of $^{77}\text{Se}_{\text{Fert}}$ in the crops from 2018 were very small. It was also not due to leaching or volatilization because there was very
376 little loss of ‘total’ or ‘organic’ $^{77}\text{Se}_{\text{Fert}}$ between harvests in 2017 and 2018 (Figs 2e – 2h). The losses of soluble and adsorbed $^{77}\text{Se}_{\text{Fert}}$
377 therefore, must have been due to a change in Se fractionation in the soil – essentially conversion into organic forms.

378

379

380

381 *Correlation between Adsorbed and Soluble soil $^{77}\text{Se}_{\text{Fert}}$ following harvest.*

382 The relationship (slope) between adsorbed and soluble $^{77}\text{Se}_{\text{Fert}}$ (Fig. 3) is effectively a dimensionless distribution coefficient (k_d) which
383 reflects the strength of adsorption of ‘available’ Se in the soil. At harvest in 2017 the k_d value was 0.89 and showed remarkable
384 consistency across all the treatments despite values of soluble $^{77}\text{Se}_{\text{Fert}}$ varying by an order of magnitude (Fig. 3). However, in 2018 the
385 available Se appeared to be nearly 7 times more strongly adsorbed with a k_d value of 6.8 (Fig. 3). The relationships shown in Fig. 3
386 indicates that in 2018 the soluble $^{77}\text{Se}_{\text{Fert}}$ was very low and restricted to between 0.00 to 0.025 $\mu\text{g kg}^{-1}$, suggesting occupancy of more
387 strongly adsorbing sites. Comparing the two trends in Fig. 3 (data from 2017 and 2018), there does appear to have been a systematic
388 increase in the strength of available $^{77}\text{Se}_{\text{Fert}}$ adsorption during the residual year.

389

390 3.3.2. Total (HF digest) and Organic (TMAH extractable) $^{77}\text{Se}_{\text{Fert}}$

391 The organic and total $^{77}\text{Se}_{\text{Fert}}$ data are shown in Fig. 2e - 2h. There were no statistically significant differences in the two fractions
392 between treatments and across the two years (Table 6). A strong correlation between organic and total $^{77}\text{Se}_{\text{Fert}}$ was seen for both years,
393 with a slope of 0.99 (Fig. 4). The data strongly suggest that the $^{77}\text{Se}_{\text{Fert}}$ that remains in the soil is in an organic (humus-bound) form
394 rather than, for example, occluded within an intractable mineral phase. TMAH (pH c. 12 at 10%) would also solubilize selenite bound
395 to Fe oxide surfaces so such a conclusion relies on complete solubilization of oxide-bound Se by the phosphate extraction (Adsorbed
396 Se) prior to TMAH extraction. Previously, measurement of ^{77}Se in an HF soil digest would be problematic due to (for example)

397 interference from germanium hydride (^{76}GeH) (Mathers et al, 2017). The similarity of the TMAH-extracted Se and HF-digested
398 therefore validates the use of oxygen mass shifting with a triple-quadrupole ICP-MS as a preferred technique.

399

400 Soil-derived $^{77}\text{Se}_{\text{soil}}$ was determined on acid digestions of the soil as a check on possible systematic errors in the assay between the two
401 harvest years; the values in 2017 and 2018 would not be expected to change. However, there was a small significant difference, $< 7\%$,
402 between the two years ($p < 0.001$) (Fig. 5; Table 5). It is likely that this arose due to a small systematic error in the year-on-year sampling
403 or analysis and probably reflects the difficulty in maintaining the quality of analysis rather than a real process.

404

405 Whilst 35 % of the Se fertilizer is stored in a residual fraction in the soil before the following year's crop, the accumulation of fertilizer
406 Se to the existing pool of soil Se is small. At an application rate of 20 g ha^{-1} only $0.0072 \text{ mg kg}^{-1}$ of fertilizer Se would be retained
407 annually, in addition to the natural total soil Se concentration of 0.332 mg kg^{-1} at this site. Taking a soil toxicity level of 20 mg kg^{-1}
408 (Walsh and Fleming, 1952), yearly application of fertilizer Se would take > 3000 years to reach this concentration

409

410 **4. Conclusions**

411 A single application of ^{77}Se to crops grown either under CA or conventional cropping systems, at the grain filling stage, provides a
412 viable approach to Se biofortification. Application of 20 g ha^{-1} Se produced sufficient grain Se enrichment in maize and legumes to
413 provide the recommended dietary Se requirement. The additional organic inputs to the soil through CA cropping systems has no apparent
414 influence on the rapid fixation of applied ^{77}Se . The transformation of applied selenate was almost entirely into humus-bound forms
415 which completely negates any residual agronomic value of the applied Se despite its retention in the soil. Within the CA trial, low yield
416 in the conventional maize cultivation treatment, caused by loss of soil fertility due to crop residue removal, reduced the recovery of
417 applied Se uptake compared to the higher-yielding CA treatments. In particular the greatest overall uptake occurred in a maize-legume
418 rotation cropping system. By contrast, the concentration and uptake of Se by intercropped legumes was affected by poor crop and root
419 development as a consequence of competition for resources by the tall maize crop and possibly disturbances during manual weeding.

420

421 Given that applied Se remaining in the soil is not available for plant uptake beyond the growing season in which it is applied, seasonal
422 application of Se would be required in order to continuously maintain recommended dietary Se intake. This could be achieved through
423 adoption of Se fertilizer amendments to existing fertilizers. Further work should investigate (i) the speciation of residual Se to assess
424 the effects of long-term Se biofortification and accumulation in the soil and (ii) the effects of timing (of application) on plant uptake
425 and, particularly, translocation to the edible crop grain.

426

427

428

429

430

431

432

433

434

435

436

437 Acknowledgements.

438 Funding and support for this study were provided by the Royal Society - Department for International Development (RS-DFID) Africa
439 Capacity Building Initiative Grant AQ140000 “Strengthening African Capacity in Soil Geochemistry to inform Agricultural and
440 Health
441 Policies”, The Malawi Government (Ministry of Agriculture and Food Security – Department of Agricultural Research), Lilongwe
442 University of Agriculture and Natural Resources, the University of Nottingham and the British Geological Survey. ISL, SDY, RML,
443 ELA, EHB, PCN, ADCC and VK conceived and designed the experiments. ISL, SDY, ELA, EHB, PCN, ADCC and VK conducted
444 the experiments. ISL collected and processed the samples. ISL, SDY and EHB analysed the samples. ISL, RML, SDY, ELA and EHB
445 processed and analysed data. All authors were involved in drafting and reviewing the manuscript. We thank Professor Martin
446 Broadley (University of Nottingham) for helpful comments on the manuscript.

447

448 References

449

450 Abadassi, J., 2015. Cowpea (*Vigna unguiculata* (L.) Walp.) agronomic traits needed in tropical zone. International Journal of Pure
451 and Applied Biosciences. 3, 158-165.

452 Bendich, A., 2001. Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. Journal of Nutrition. Philadelphia.
453 17 (4), 364-365.

454 Branca, G., McCarthy, N., Lipper, L., Jolejole, M.C., 2011. Climate-smart agriculture: a synthesis of empirical evidence of food
455 security and mitigation benefits from improved cropland management. Mitigation of climate change in agriculture series 3.
456 FAO Report. Rome, Italy. Available via DIALOG: cgspace.cgiar.org/handle/10568/33460. Accessed 21 Nov, 2019.

457 Broadley M.R., Alcock J., Alford, J., Cartwright, P., Foot, I., Fairweather-Tait, S.J., Hart, D.J., Hurst, R., Knott, P., Mcgrath, S.P.,
458 Meacham, M.C., Norman, K., Mowat, H., Scott, P., Stroud, J.L., Tovey, M., Tucker, M., White, P.J., Young, S.D., Zhao, F.,
459 2010. Selenium biofortification of high-yielding winter wheat (*Triticum aestivum* L.) by liquid or granular Se fertilisation.
460 Plant Soil. 332, 5-18.

461 Broadley, M.R., White, P.J., Bryson, R.J., 2006. Biofortification of UK food crops with selenium. Proceedings of Nutrition Society.
462 65, 169-181.

463 Cardoso, B.R., Roberts, B.R., Bush, A.I., Hare, D.J., 2015. Selenium, selenoproteins and neurodegenerative diseases. Metallomics,
464 7 (8), 1213-1228.

465 Chilimba, A.D.C., Young, S.D., Black, C.R., Rogerson, K.B., Ander, E.L., Watts, M.J., Lammel, J. and Broadley, M.R., 2011. Maize
466 grain and soil surveys reveal suboptimal dietary selenium intake is widespread in Malawi. Scientific Reports. 1, 72.

467 Chilimba A.D.C., Young, S.D., Black, C.R., Meacham, M.C., Lammel, J., Broadley, M.R., 2012a. Agronomic biofortification of
468 maize with selenium (Se) in Malawi. Field Crops Research. 125, 118–128.

469 Chilimba A.D.C., Young, S.D., Black, C.R., Meacham, M.C., Lammel, J., Broadley, M.R., 2012b Assessing residual availability of
470 selenium applied to maize crops in Malawi. *Field Crop Research*. 134, 11–18.

471 Chilimba A.D.C., Young, S.D., Joy, E.J.M., 2014. Agronomic bio-fortification of maize, soyabean and groundnuts with selenium in
472 intercropping and sole cropping systems. *Afric Journal of Agricultural Research*. 9, 3620-3626.

473 Dhillon, K., Dhillon, S., 1999. Adsorption–desorption reactions of selenium in some soils of India. *Geodema*. 93, 19-31.

474 Donovan, U., Gibson, R.S., Ferguson, E.L., Ounpuu, S., Heywood, P., 1991. The selenium content of staple foods from Malawi and
475 Papua New Guinea. *Food Composition and Analysis*. 4, 329–336.

476 Eurola, M., Hietaniemi, V., 2000. Report of the Selenium Monitoring Programme 1997–1999. Publications of Agricultural Research
477 Centre of Finland, Series B 24. Jokioinen: Agricultural Research Centre of Finland. 181.

478 Eick, F., Maleta, K., Govasmark, E., Duttaroy, A.K., Bjune, A.G., 2009. Food intake of selenium and sulphur amino acids in
479 tuberculosis patients and healthy adults in Malawi. *Internationa Jounal of Tuberculosis and Lung Diseases*. 13, 1313-1315.

480 Fairweather-tait, S. J. et al. 2011. Selenium in human health and disease. *Antioxidants & Redox Signaling*. 14, 1337-1383.

481 Ferguson, E.L., Gibson, R.S., Weaver, S.D., Heywood, P., Heywood, A., Yaman, C., 1989. The mineral content of commonly
482 consumed Malawian and Papua New Guinean foods. *Food Composition and Analysis*. 2, 260 - 272.

483 Fordyce, F.M., Guangdi, Z., Green, K., Xinping, L., 2000. Soil, grain and water chemistry in relation to human selenium responsive
484 diseases in Enshi District, China. *Applied Geochemistry*. 15, 117-132.

485 Gupta, U.C., Winter, K.A., Sanderson, J.B., 1993. Selenium content of barley as influenced by selenite-enriched and selenite enriched
486 fertilizers. *Communication in Soil Science and Plant Analysis*. 24, 1165-1170.

487 Hartikainen, H., 2005. Biogeochemistry of selenium and its impact on food chain quality and human health. *Trace Elements in*
488 *Medicine Biology*. 18, 309–318.

489 Hamdy, A.A., Gissel-Nielsen, G., 1977. Fixation of selenium by clay minerals and iron oxides. *Z. Pflanzenernachr.Bodenkd*. 140,
490 63-70.

491 Hurst, R., Siyame, E.W.P., Young, S.D., Chilimba, A.D.C., Joy, E.J.M., Black, C.R., Ander, E.L., Watts, M.J., Chilima, B., Gondwe,
492 J., Kang'ombe, D., Stein, A.J., Fairweather-Tait, S.J., Gibson, R.S., Kalimbara, A.A., Broadley, M.R., 2013, Soil-type
493 influences human selenium status and underlies widespread selenium deficiency risks in Malawi. *Scientific Reports*. 3:1425.
494 Institute of Medicine, Food and Nutrition Board (IMFNB), 2000. *Dietary Reference Intakes: Vitamin C, Vitamin E, Selenium, and*
495 *Carotenoids*. National Academy Press, Washington.

496 Johnson, C.C., Ge, X., Green, K.A., Liu, X., 2000. Selenium distribution in the local environment of selected villages of the Keshan
497 Disease belt, Zhangjiakou District, Hebei Province, People's Republic of China. *Applied Geochemistry*. 15, 385-401.

498 Joy, E.J.M., Kumssa, D.B., Broadley, M.R., Watts, M.J., Young, S.D., Chilimba, A.D.C., Ander, E.L., 2015a. Dietary mineral
499 supplies in Malawi: spatial and socioeconomic assessment. *BMC Nutrition*. 1, 1-25.

500 Joy, E.J.M., Broadley, M.R., Young, S.D., Black, C.R., Chilimba, A.D.C., Ander, E.L., Barlow, T.S. Watts, M.J., 2015b. Soil type
501 influences crop mineral composition in Malawi. *Science of the Total Environment*. 505, 587 – 595.

502 Kassam, A., Friedrich, T., Shaxson, F., Pretty, J., 2009. The spread of conservation agriculture: justification, sustainability and uptake.
503 *International Journal of Agricultural Sustainability*. 7, 292-320.

504 Ligowe, I.S., Nalivata, P.C., Njoloma, J., Makumba, W., Thierfelder, C., 2017. Medium-term effects of conservation agriculture on
505 soil quality. *African Journal of Agricultural Research*. 12, 2412-2420.

506 Ligowe I.S., Young S.D., Ander E.L., Kabambe V., Chilimba A.D., Bailey E.H., Lark R.M., Nalivata P.C. (2020). Agronomic
507 biofortification of leafy vegetables, grown in an Oxisol, Alfisol and Vertisol with isotopically labelled selenium
508 (⁷⁷Se). *Geoderma* 361, 114106.

509 Lipper, L., Thornton, P., Campbell, B.M., Baedeker, T., Braimoh, A., Bwalya, M., Caron, P., Cattaneo, A., Garrity, D., Henry, K.,
510 2014. Climate-smart agriculture for food security. *Nature Climate Change*. 4, 1068-1072.

511 Lopes, G., Ávila F.W., Guilherme, L.R.G., 2017. Selenium behavior in the soil environment and its implication for human health.
512 *Ciência e Agrotecnologia* 41 (6), 605-15.

513 Malagoli M., Schiavon M., dall'Acqua S., Pilon-Smits E.A., 2015. Effects of selenium biofortification on crop nutritional quality.
514 *Frontiers in Plant Science*. 6, 280.

515 Mathers, A.W., Young, S.D., McGrath, S.P., Zhao, F.J., Crout, N.M.J., Bailey, E.H., 2017. Determining the fate of selenium in wheat
516 bio-fortification: an isotopically labelled field trial study. *Plant Soil*. 420, 61-77.

517 McCarthy, N., Lipper, L., Branca, G., 2011. Climate-smart agriculture: smallholder adoption and implications for climate change
518 adaptation and mitigation. *Mitigation of Climate Change in Agriculture Working Paper*. 3, 1-37.

519 Nautiyal, P.C., Misra, J.B., Zala, P. V., 2010. Influence of seed maturity stages on germinability and seedling vigor in groundnut.
520 *Journal of ICRISAT Agricultural Research* 8.

521 Ngwira, A.R., Sleutel, S., De Neve, S., 2012. Soil carbon dynamics as influenced by tillage and crop residue management in loamy
522 sand and sandy loam soils under smallholder farmers' conditions in Malawi. *Nutrient Cycling in Agroecosystems*. 92, 315-
523 328.

524 Phiri, F.P., Ander, E.L. Bailey, E.H., Chilima, B., Chilimba, A.D.C., Gondwe, J., Joy E.J.M., Kalimpira, A.A., Kumssa D.B., Lark,
525 R.M., Phuka J.C., Salter, A., Suchdev, P.S., Watts M.J., Young, S.D., Broadley M.R., 2019. The risk of selenium deficiency
526 in Malawi is large and varies over multiple spatial scales. *Scientific Reports*. 9 (1), 6566.

527 Pinheiro J, Bates D, DebRoy S, Sarkar D and R Core Team. 2017. nlme: Linear and Nonlinear Mixed Effects Models_. R package
528 version 3.1-131, <URL: Available via DIALOG: [https://CRAN.R project.org/package=nlme](https://CRAN.R-project.org/package=nlme)>. Accessed 12 Sept, 2019.

529 Powlson, D.S., Stirling, C.M., Thierfelder, C., White, R.P., Jat, M.L., 2016. Does conservation agriculture deliver climate change
530 mitigation through soil carbon sequestration in tropical agro-ecosystems? *Agriculture Ecosystems and Environment*. 220,
531 164-174.

532 Qin, H.B.; Zhu, J.M. Su, H., 2012. Selenium fractions in organic matter from Se-rich soils and weathered stone coal in selenosis
533 areas of China. *Chemosphere*. 86 (6), 626-633.

534 Rayman, M.P., 2008. Food-chain selenium and human health: emphasis on intake. *British Journal of Nutrition*. 100, 254-68.

535 R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna,
536 Austria. Available via DIALOG:URL <https://www.R-project.org/>. Accessed 12 Sept, 2019.

537 Riaz, M., Mehmood, K.T., 2012. Selenium in human health and disease: A review. Journal of Postgrad. Medicine Institute. 26 (2),
538 120-133.

539 Salehi, A., Mehdi, B., Fallah, S., Kaul, H., Neugschwandtne, R.W., 2018. Productivity and nutrient use efficiency with integrated
540 fertilization of buckwheat–fenugreek intercrops. Nutrient Cycling in Agroecosystem. 110, (3), 407.

541 Sankaran, R. P., Grusak, M. A., 2014. Whole shoot mineral partitioning and accumulation in pea (*Pisum sativum*). Frontiers in Plant
542 Science. 5, 149.

543 Sperotto, R.A., 2013. Zn/Fe remobilization from vegetative tissues to rice seeds: should I stay or should I go? Ask Zn/Fe supply!
544 Frontiers in Plant Science. 4, 464.

545 Steward, P.R., Thierfelder, C., Dougill, A.J., Ligowe, I., 2019. Conservation agriculture enhances resistance of maize to climate stress
546 in a Malawian medium-term trial. Agriculture Ecosystems and Environment. 277, 95-104.

547 Stroud J. L., Broadley, M.R., Foot, I., Fairweather-Tait, S.J., Hart, D.J., Hurst, R., Knott, P., Mowat, H., Norman, K., Scott, P.,
548 Tucker, M., White, P.J., Mcgrath, S.P., Zhao, F.J., 2010. Soil factors affecting selenium concentration in wheat grain and the
549 fate and speciation of Se fertilisers applied to soil. Plant Soil. 332, 19-30.

550 Tolu, J., Thiry, Y., Bueno, M., Jolivet, C., Potin-Gautier, M., Le Hécho, I., 2014. Distribution and speciation of ambient selenium in
551 contrasted soils, from mineral to organic rich. Science of the Total Environment. 479–480:93–101.

552 Thierfelder, C., Chivenge, P., Mupangwa, W., Rosenstock, T.S., Lamanna, C., Eyre, J.X., 2017. How climate-smart is conservation
553 agriculture (CA)? - its potential to deliver on adaptation, mitigation and productivity on smallholder farms in southern Africa.
554 Food Security. 9, 537-560.

555 Thierfelder, C., Wall, P.C., 2010. Investigating Conservation Agriculture (CA) Systems in Zambia and Zimbabwe to Mitigate Future
556 Effects of Climate Change. Crop Improvement. 24, 113-121.

557 Thierfelder, C., Wall, P.C., 2009. Effects of conservation agriculture techniques on infiltration and soil water content in Zambia and
558 Zimbabwe. *Soil and Tillage. Research.* 105, 217-227.

559 Walsh, T., Fleming, G.A., 1952. Selenium levels in rocks, soils and herbage from a high selenium locality in Ireland *Trans.*
560 *International Society of Soil Science.* 2,178-185.

561 WHO, 2009. *Global Health Risks: Mortality and Burden of Disease Attributable to Selected Major Risks.* Available via DIALOG:
562 http://www.who.int/healthinfo/global_burden_disease/GlobalHealth2009:Risks_report_annex.pdf. Accessed 21 Nov, 2019.

563 Winkel L. H., Johnson C. A., Lenz M., Grundl T., Leupin O. X., Amini M., Charlet, L., 2012. Environmental selenium research:
564 From microscopic processes to global understanding. *Environmental Science and Technology.* 46, 571–579.

565

566 List of Tables
567
568

569 **Table 1.** Treatment descriptions of the Chitedze long-term Conservation Agriculture trial.

Treatment	Description
T1	Check plot: using traditional farmers' practice using a hand hoe (ridge and furrow system); maize is the sole crop and no crop residues are returned to the soil
T2	Basin: maize planted as the sole crop in basins with dimensions of 0.15 x 0.15 x 0.15m, that are re-constructed yearly; crop residues are retained.
T3	Direct Seeding: maize as a sole crop, planted with a dibble stick; crop residues are retained.
T4C	Crop rotation: direct seeding, with a dibble stick, of the rotation sequence 'cowpea–maize–cowpea'; crop residues are retained; split plot with T4G.
T4G	Crop rotation: direct seeding, with a dibble stick, of the rotation sequence 'groundnut–maize–groundnut'; crop residues are retained; split plot with T4C.
T5C	Crop rotation: direct seeding, with a dibble stick, of the rotation sequence 'maize-cowpea-maize' (in contrast to T4C); crop residues are retained; split plot with T5G.
T5G	Crop rotation: direct seeding, with a dibble stick, of the rotation sequence maize-groundnut-maize (in contrast to T4G); crop residues are retained; split plot with T5C.
T6	Intercropping: direct seeding, with a dibble stick, of maize intercropped with pigeon pea; crop residues are retained.
T7	Intercropping: direct seeding, with a dibble stick, of maize intercropped with cowpea; crop residues are retained
T8	Intercropping: direct seeding, with a dibble stick, of maize intercropped with velvet beans at 8 weeks; crop residues are retained

570

571

572 **Table 2:** List of treatment contrasts

Contrast number	Contrast code	Contrast Description
1	C1	Check plots - conventional cultivation (T1) compared with the mean for all CA treatments (T2-T8) - an assessment of the general impact of CA practices.
2	C2	Mono-cropped maize compared (T2 and T3) with mixed-cropped maize (T4C, T4G, T5C, T5G, T6, T7 and T8) - an assessment of the impacts of diversification by rotation or intercropping.
3	C3	Maize under legume rotation (T5C and T5G) compared with maize intercropped with legumes (T6, T7, and T8)
4	C4	Maize planted in basins (T2) compared with monocrop maize planted directly into residues with a dibble stick (T3).
5	C5	A comparison between maize rotation with cowpea (T5C) and with groundnut (T5G).
6	C6	A comparison between intercropping with a late-sown legume -velvet bean (T8) and intercropping with cowpea or pigeon pea-sown with the maize (T6 and T7)
7	C7	A comparison between the cowpea (T7) and pigeon pea intercrop (T8)

573
574

575

576

Table 3: Concentration of $^{77}\text{Se}_{\text{Fert}}$ ($\mu\text{g kg}^{-1}$) in the crops (maize and legume, stover and grain) in 2017 and 2018. Treatments are fully described in Table 1 and

577

include: T1 = conventional cultivation; T2 = CA basins; T3 = CA sole maize; T4C = CA rotation cowpea after maize; T4G = CA rotation groundnuts after maize;

578

T5C = CA rotation maize after cowpea; T5G = CA rotation maize after groundnuts; T6 = CA intercropped: maize + Pigeon peas; T7 CA intercropped: maize +

579

cowpeas T8 = CA intercropped: maize + velvet beans.

Crop part	Maize Grain $^{77}\text{Se}_{\text{Fert}}$ ($\mu\text{g kg}^{-1}$)		Maize Stover $^{77}\text{Se}_{\text{Fert}}$ ($\mu\text{g kg}^{-1}$)		Legume Grain $^{77}\text{Se}_{\text{Fert}}$ ($\mu\text{g kg}^{-1}$)		Legume Stover $^{77}\text{Se}_{\text{Fert}}$ ($\mu\text{g kg}^{-1}$)	
	Treatment	2017	2018	2017	2018	2017	2018	2017
T1	209	1.89	111	5.17				
T2	210	1.24	65.5	1.32				
T3	203	0.77	108	2.77				
4C		2.16		10.5	404		305	
4G		2.78		6.83	711		427	
5C	232		119			2.49		4.08
5G	228		102			1.32		7.93
T6	226	1.11	96.4	4.36	16.5	1.37	11.6	0.46
T7	223	0.72	72.9	0.85	188	0.79	199	2.92
T8	206	1.35	67.3	2.20	28.4	0.54	52.0	0.27
SE\pm	33.6	0.45	18.8	2.75	60.6	0.48	45.5	0.52

580

581

582 **Table 4:** Statistical contrasts of maize grain and stover $^{77}\text{Se}_{\text{Fert}}$ and Se_{soil} concentrations and plant uptake. Contrasts (C) numbers are described in Table 2.

Contrast number	Denominator DF	P-values for Maize Grain				P-values for Maize Stover			
		$^{77}\text{Se}_{\text{Fert}}$ Concentration	$^{77}\text{Se}_{\text{Fert}}$ Uptake	Se_{Soil} Concentration	Se_{Soil} Uptake	$^{77}\text{Se}_{\text{Fert}}$ Concentration	$^{77}\text{Se}_{\text{Fert}}$ Uptake	Se_{Soil} Concentration	Se_{Soil} Uptake
2017									
C1	21	0.790	<0.0001	0.584	0.0002	0.308	0.010	0.006	0.043
C2	21	0.572	0.057	0.013	0.412	0.772	0.196	0.101	0.555
C3	21	0.718	0.035	0.201	0.009	0.078	0.005	0.448	0.058
C4	21	0.882	0.589	0.206	0.489	0.121	0.214	0.985	0.937
C5	21	0.942	0.918	0.972	0.957	0.533	0.398	0.941	0.964
C6	21	0.665	0.787	0.741	0.834	0.459	0.311	0.717	0.392
C7	21	0.944	0.316	0.775	0.368	0.384	0.197	0.272	0.181
2018									
C1	21	0.370	0.085	0.741	0.129	0.718	0.536	0.013	0.050
C2	21	0.112	0.032	0.238	0.109	0.226	0.217	0.069	0.408
C3	21	0.002	0.0002	0.010	0.003	0.021	0.019	0.039	0.0007
C4	21	0.459	0.431	0.609	0.642	0.713	0.805	0.021	0.031
C5	21	0.337	0.274	0.347	0.307	0.352	0.340	0.281	0.195
C6	21	0.433	0.360	0.036	0.042	0.929	0.903	0.733	0.970
C7	21	0.540	0.502	0.667	0.698	0.351	0.458	0.037	0.022

583

584

585
586
587
588
589

Table 5: Statistical contrasts for legume grain and stover $^{77}\text{Se}_{\text{Fert}}$ and Se_{soil} concentration and plant uptake. The contrasts are: C3 = legume in monocrop (in rotation with maize) and legume in intercrop with maize, C5 cowpea and groundnut monocrop in rotation with maize, C6 = velvet bean and cowpea and pigeon pea (sown at the same time as the maize) in intercrop and C7 = comparison between cowpea and pigeon pea both in intercrop (see Table 2).

Contrast number	Denominator DF	P-values for Legume Grain				P-values for Legume Stover			
		$^{77}\text{Se}_{\text{Fert}}$ Concentration	$^{77}\text{Se}_{\text{Fert}}$ Uptake	Se_{soil} Concentration	Se_{soil} Uptake	$^{77}\text{Se}_{\text{Fert}}$ Concentration	$^{77}\text{Se}_{\text{Fert}}$ Uptake	Se_{soil} Concentration	Se_{soil} Uptake
2017									
C3	12	<0.0001	<0.0001	0.367	0.157	<0.0001	<0.0001	0.506	<0.0001
C5	12	0.004	0.0001	0.478	0.0006	0.080	<0.0001	0.375	0.067
C6	12	0.338	0.670	0.007	<0.0001	0.359	0.810	0.755	0.025
C7	12	0.068	0.985	0.045	0.0005	0.013	0.952	0.187	0.001
2018									
C3	12	0.042	0.008	0.012	0.042	<0.0001	<0.001	0.039	<0.0001
C5	12	0.113	0.763	0.814	0.142	0.0002	<0.001	0.169	0.001
C6	12	0.380	0.685	0.004	<0.0001	0.045	0.696	0.249	0.060
C7	12	0.415	0.091	0.150	0.010	0.006	0.701	0.227	0.004

590

591

592 **Table 6:** Statistical contrasts for the soil $^{77}\text{Se}_{\text{Fert}}$ fractionation (Soluble, Adsorbed, Total) and Se_{Soil} for both year (2017
 593 and 2018). Contrasts are: C1= Check plot (conventional) vs all CA treatment; C2 = Mono-cropped maize vs mixed-
 594 cropped maize; C3 = Maize under legume rotation vs maize in legume intercrop; C4 = Basin planted maize vs dibble
 595 stick planted maize; C5 = Maize under cowpea rotation vs maize under groundnut rotation; C6 = Maize after
 596 groundnuts in rotation vs groundnuts after maize rotation; C7 = Maize after cowpea in rotation vs cowpea after maize
 597 rotation; C8 = Maize velvet bean intercropped (late sown) vs maize cowpea or pigeon peas intercropped; C9 = Maize
 598 cowpea intercropped vs maize pigeon peas intercropped (see Table 2).
 599

Contrast number	Denominator DF	P-values for Soluble $^{77}\text{Se}_{\text{Fert}}$	P-values for Adsorbed $^{77}\text{Se}_{\text{Fert}}$	P-values for Total TMAH $^{77}\text{Se}_{\text{Fert}}$	P-values for Total HF $^{77}\text{Se}_{\text{Fert}}$	P-values for Se_{Soil}
C1	22	0.016	0.027	0.070	0.065	0.318
C2	22	0.4081	0.631	0.707	0.532	0.942
C3	22	0.034	0.041	0.831	0.219	0.001
C4	22	0.894	0.074	0.143	0.749	0.913
C5	7	0.411	0.082	0.662	0.864	0.856
C6	28	0.001	0.003	0.513	0.351	0.256
C7	28	0.08	0.086	0.511	0.24	0.388
C8	22	0.618	0.32	0.885	0.202	0.077
C9	22	0.986	0.712	0.311	0.329	0.557
Year (Yr)	28	<0.0001	0.05	0.115	0.856	<0.0001
C1 x Yr	28	0.004	0.013	0.068	0.177	0.387
C2 x Yr	28	0.268	0.723	0.442	0.639	0.773
C3 x Yr	28	0.006	0.114	0.79	0.745	0.297
C4 x Yr	28	0.9	0.91	0.976	0.992	0.457
C5 x Yr	28	0.469	0.91	0.458	0.417	0.38
C6 x Yr	28	0.041	0.003	0.407	0.238	0.033
C7 x Yr	28	0.147	0.036	0.714	0.621	0.236
C8 x Yr	28	0.548	0.039	0.582	0.059	0.121
C9 x Yr	28	0.828	0.097	0.01	0.146	0.891

600

601

602

603 Plates.
604
605



606
607
608
609
610
611

Plate 1: Aerial photograph of the long-term conservation agriculture trial (c. 1 ha) at Chitedze, Malawi (February 2017).

612 List of Figures.

613 **Figure 1:** Offtake of $^{77}\text{Se}_{\text{Fert}}$ (g ha^{-1}) in maize and legume stover and grain in 2017 and 2018.
614 Treatments included: T1 = conventional cultivation; T2 = CA basins; T3 = CA sole maize; T4C =
615 CA cowpea rotation after maize; T4G = CA groundnuts rotation after maize; T5C = CA maize
616 rotation after cowpea; T5G = CA maize rotation after groundnuts; T6 = CA intercropped maize +
617 pigeon peas; T7 CA intercropped maize + cowpeas; T8 = CA intercropped maize + velvet beans.
618 Note the change in Y-axis scale from 2017 to 2018. Error bars = \pm SEM.

619
620 **Figure 2:** Soil fractions of fertilizer-derived ^{77}Se ($^{77}\text{Se}_{\text{Fert}}$) in 2017 and 2018, including: 'Soluble' (a, b),
621 'Adsorbed' (c, d), 'Organic' (e, f) and 'Total' (g, h). Note the difference in y-axis scales for a-d and e-h.
622 Values presented are the average of four replicate plots; error bars are \pm standard error of means.
623 Treatments include: T1 = conventional; T2 = CA basins; T3 = CA sole maize; T4C = CA rotation cowpea
624 after maize; T4G = CA rotation groundnuts after maize; T5C = CA rotation maize after cowpea; T5G =
625 CA rotation maize after groundnuts; T6 = CA intercrop: maize + Pigeon peas; T7 CA intercrop: maize +
626 cowpeas T8 = CA intercrop: maize + velvet beans. Error bar = \pm SEM.

627

628 **Figure 3:** Correlation between Adsorbed and Soluble soil $^{77}\text{Se}_{\text{Fert}}$ at harvest of the bio-fortified
629 crop in 2017 (open circle; slope = 0.89) and in 2018 (closed circle; slope = 6.8).

630

631 **Figure 4:** Correlation between total Organic and Total soil $^{77}\text{Se}_{\text{Fert}}$ at harvest of the bio-fortified
632 crop in 2017 (open circle; slope = 0.99) and in the following year, 2018 (closed circle; slope =
633 1.15). The line represents a 1:1 relationship.

634

635 **Figure 5:** Soil-derived ^{77}Se in 2017 and 2018. Error bars = \pm SE for 4 replicates

636

637

638
 639
 640
 641
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664




Figure 1: Offtake of $^{77}\text{Se}_{\text{Fert}}$ (g ha⁻¹) in maize and legume stover and grain in 2017 and 2018. Treatments included: T1 = conventional cultivation; T2 = CA basins; T3 = CA sole maize; T4C = CA cowpea rotation after maize; T4G = CA groundnuts rotation after maize; T5C = CA maize rotation after cowpea; T5G = CA maize rotation after groundnuts; T6 = CA intercropped maize + pigeon peas; T7 CA intercropped maize + cowpeas; T8 = CA intercropped maize + velvet beans. Note the change in Y-axis scale from 2017 to 2018. Error bars = ±SEM.

665
 666
 667
 668
 669
 670
 671
 672
 673
 674
 675
 676
 677
 678
 679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693
 694
 695
 696
 697





Figure 2: Soil fractions of fertilizer-derived ^{77}Se ($^{77}\text{Se}_{\text{Fert}}$) in 2017 and 2018, including: ‘Soluble’ (a, b), ‘Adsorbed’ (c, d), ‘Organic’ (e, f) and ‘Total’ (g, h). Note the difference in y-axis scales for a-d and e-h. Values presented are the average of four replicate plots; error bars are \pm standard error of means.

698 Treatments include: T1 = conventional; T2 = CA basins; T3 = CA sole maize; T4C = CA rotation cowpea
699 after maize; T4G = CA rotation groundnuts after maize; T5C = CA rotation maize after cowpea; T5G =
700 CA rotation maize after groundnuts; T6 = CA intercrop: maize + Pigeon peas; T7 CA intercrop: maize +
701 cowpeas T8 = CA intercrop: maize + velvet beans. Error bar = \pm SEM.
702

703
704



705
706


707

708 **Figure 3:** Correlation between Adsorbed and Soluble soil $^{77}\text{Se}_{\text{Fert}}$ at harvest of the bio-fortified
709 crop in 2017 (open circle; slope = 0.89) and in 2018 (closed circle; slope = 6.8).

710

711


712
713
714



715
716
717
718
719
720

Figure 4: Correlation between total Organic and Total soil ⁷⁷Se_{Fert} at harvest of the bio-fortified crop in 2017 (open circle; slope = 0.99) and in the following year, 2018 (closed circle; slope = 1.15). The line represents a 1:1 relationship.

721
722



723
724
725
726
727
728
729

Figure 5: Soil-derived ⁷⁷Se in 2017 and 2018. Error bars = ±SE for 4 replicates