



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Current Opinion in
Microbiology

CrossMark

Nucleotide second messengers in bacterial decision making

Catriona MA Thompson^{1,2} and Jacob G Malone^{1,2}

Since the initial discovery of bacterial nucleotide second messengers (NSMs), we have made huge progress towards understanding these complex signalling networks. Many NSM networks contain dozens of metabolic enzymes and binding targets, whose activity is tightly controlled at every regulatory level. They function as global regulators and in specific signalling circuits, controlling multiple aspects of bacterial behaviour and development. Despite these advances there is much still to discover, with current research focussing on the molecular mechanisms of signalling circuits, the role of the environment in controlling NSM pathways and attempts to understand signalling at the whole cell/community level. Here we examine recent developments in the NSM signalling field and discuss their implications for understanding this important driver of microbial behaviour.

Addresses

¹ Molecular Microbiology Department, John Innes Centre, Norwich, UK² School of Biological Sciences, University of East Anglia, Norwich, UKCorresponding author: Malone, Jacob G (jacob.malone@jic.ac.uk)**Current Opinion in Microbiology** 2020, **55**:34–39This review comes from a themed issue on **Cell regulation**Edited by **Charles J Dorman** and **Joan Geoghehan**<https://doi.org/10.1016/j.mib.2020.02.006>1369-5274/© 2020 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Models for bacterial nucleotide second messenger (NSM) signalling have evolved substantially since the discovery of cAMP catabolite repression in the late 1960s [1]. The first pathways discovered, for example, cAMP signalling [1] and the stringent response [2], typically contained few discrete NSM metabolic proteins. However, the advent of whole genome sequencing revealed that bacterial NSM signalling is substantially more complex than previously thought. This has led to the discovery of more NSMs, such as cyclic-di-GMP (cdG, [3]), cyclic-di-AMP (cdA, [4]) and recently cyclic-GAMP [5], which have been shown to control a wide variety of bacterial behaviours. Furthermore, many NSM pathways have been shown to contain far more metabolic enzymes than were initially suspected, with dozens of cdG cyclases and phosphodiesterases identified

in many bacterial species [6]. Finally, the complexity and variability of NSM binding domains have become more apparent, with NSMs shown to bind specifically to diverse protein and RNA folds, defying straightforward bioinformatic prediction [7].

To explain this remarkable complexity, sophisticated models for NSM networks have been developed that incorporate several key findings. NSM signalling proteins function under tight temporal and spatial control, with NSM synthases and degradative enzymes controlled at multiple regulatory levels in response to the cell cycle and the surrounding environment [7,8]. Discrete signalling circuits, where specific NSM enzymes control a small number of downstream proteins have been identified, often accompanied by direct protein–protein interaction between signalling enzymes and NSM effectors [7,8]. Related to this, unconventional enzyme functions have been discovered that explain certain specific NSM responses, for example, trigger enzymes, whose enzyme activity alters their interaction with an effector protein [9]. Supported by bioinformatic and computational analyses, comprehensive models have been built for several NSM networks, particularly cdG signalling in *Caulobacter crescentus* [10] and *Escherichia coli* [11,12].

NSM-binding effector proteins

Building on their initial characterisation as controllers of the stringent response ((p)ppGpp, [2]), the motile-sessile transition (cdG, [13]) and potassium/glutamate homeostasis (cdA, [14]) NSM signalling molecules have been implicated in the control of a diverse range of bacterial processes. In many cases these are quite different to the initial NSM regulon, for example, decreased cdA levels are implicated in eDNA release during *Staphylococcus aureus* biofilm formation [15], while cdG controls *Sinorhizobium meliloti* peptidoglycan biosynthesis [16]. The *Streptomyces* life cycle is also controlled by σ -factor binding to cdG [17]. Given the inherently leaky nature of second messenger signal transduction, a major factor in shaping this expanded NSM regulon has been the development of reliable biochemical methods to identify and characterise NSM binding proteins [18,19]. For example, Perez-Mendoza *et al.* used surface plasmon resonance to show that *S. meliloti* β -glucan synthesis is controlled by cdG binding to the BgsA synthase subunit [20]. The absence of an obvious NSM binding fold suggests that BgsA contains a novel binding motif.

The cdG-binding transcriptional regulator BrlR mediates the correlation between biofilm formation and antibiotic tolerance in *Pseudomonas aeruginosa*. Intriguingly, BrlR was also recently shown to bind to a second intracellular regulatory molecule: pyocyanin, a phenazine [21[•]]. This finding, alongside other recent work [22] identify mechanisms for regulatory crosstalk between cdG and phenazine signalling, more closely integrating processes such as redox adaptation and antibiotic tolerance with biofilm formation and motility control. This dual regulatory mechanism is reminiscent of the master regulator FleQ, which responds to both cdG and ATP levels and integrates biofilm formation with flagella-driven motility [23]. This raises the prospect of finding other multiple-input regulators that integrate input signals from discrete signalling pathways.

Alongside the identification of new NSM binding proteins, recent biochemical and structural biology research has led to significant advances in our mechanistic understanding of NSM regulatory pathways. For example, Anderson *et al.* showed how (p)ppGpp inhibits the purine salvage enzyme HPRT by binding a conserved motif with its substrate phosphoribosyl pyrophosphate (PRPP), and used their findings to propose a more general mechanism for the evolution of ligand binding sites [24]. Wood *et al.* [25] conducted an analysis of Era, a GTPase involved in *S. aureus* ribosomal assembly and showed it interacted with the DEAD-box helicase CshA and the (p)ppGpp synthase Rel_{Sau}. Interaction with the latter was shown to boost Era GTPase activity, impacting its function as a hub protein for enzymes involved in ribosomal assembly and rRNA processing/degradation [25].

Two new studies from the Sondermann lab examined previously identified NSM regulatory systems; the LapD cdG-binding adhesin regulator and the Orn di-ribonucleotidase, which degrades linearised cyclic di-nucleotides [26,27]. In both cases, these studies refined our understanding of these proteins and uncovered new insights into the molecular mechanisms of NSM signalling. Orn was shown to possess a unique specificity for di-ribonucleotides in contrast to other cellular ribonucleotidases [26]. Meanwhile, LapD binding was shown to strongly enhance the DGC activity of its cognate cyclase, GcbC [27], suggesting a mechanism for the establishment of signal specificity, a long-standing problem in the field of NSM regulation.

Advances in our understanding of NSM input signals

While our understanding of the diversity and mechanisms underlying NSM-effector binding has rapidly advanced, comparatively few of the direct sensory inputs controlling NSM metabolism have been identified. The low number of unambiguously identified NSM input proteins stems from the pleiotropic, leaky nature of NSM signalling, making it difficult to interpret overexpression or deletion

studies with confidence. Those direct sensors that have been identified often have predictable input domains, for example, for phosphorylation [28] or light-sensing [29], or have been identified following detailed structural analysis, for example, the DgcZ zinc-binding diguanylate cyclase [30], or the redox/L-tryptophan/vitamin B6-regulated YfiBNR system [31,32]. Nonetheless, in recent years several new input signals have been attributed to specific NSM proteins, including a novel class of phosphodiesterases (PDEs) identified in *E. coli* [33]. These PDEs contain an N-terminal periplasmic CSS domain with two highly conserved cysteine residues, and are controlled by a cysteine disulphide REDOX switch. The presence of a cysteine cross-bridge inhibits the cytoplasmic EAL (PDE) domain, while disulphide bond reduction is associated with increased PDE activity [33].

The transmembrane GGDEF-EAL hybrid protein RmcA from *P. aeruginosa* is another important NSM signalling protein whose inputs have recently been discovered [22,34]. RmcA controls colony morphology in response to changing redox conditions, which are transmitted through phenazine binding to one of four cytosolic PAS domains leading to PDE stimulation under oxidising conditions. Intriguingly, RmcA has also been implicated in direct sensing of the amino-acid L-arginine [34] that binds to its periplasmic Venus Flytrap domain and again stimulates PDE activity. This dual-input mechanism is reminiscent of the BrlR pyocyanin/cdG binding regulator [21[•]], which also contributes to phenazine-linked regulation. Going forward, it will be interesting to determine how many more NSM signalling pathways are controlled by proteins with multiple signal inputs, both at the NSM-binding and metabolic levels.

Molecular characterisation of NSM signalling pathways

The identification and characterisation of individual NSM proteins has been accompanied by detailed analysis of several well-known NSM-controlled phenotypes. Recent studies have progressed beyond simple community-level phenotyping to define the underlying molecular mechanisms of NSM control. An excellent example of this is initial surface attachment in *P. aeruginosa* and *C. crescentus*. Recent work implicates the flagellar motor proteins MotA and MotB (MotAB) in initial surface detection by both species [35,36]. In *C. crescentus*, the initial chemotaxis input comes from the Cle family of chemotaxis proteins, where cdG-binding promotes interactions with the MotAB complex [35]. This interaction in *C. crescentus* re-orientates the cell [13] and increases the tactile response initiated by surface proximity [13,36]. A similar feedback system that triggers the initial cascade of surface attachment is also likely to exist in other bacteria, with recent work showing that MotAB are essential for the switch from flagellar tethering to surface attachment in *P. aeruginosa* [37].

Once activated, the flagellar motor complex triggers an increase in cellular cdG levels through the upregulation of DGCs [13,36,38,39^{***}]. *C. crescentus* has a specialised flagellar motor-coupled DGC, DgcB, which is specifically upregulated during surface sensing. Meanwhile, several DGCs are implicated in the case of *P. aeruginosa* surface attachment [13,36,39^{***}]. The increase in cellular cdG leads to an upregulation of type IV pili (T4P) and adherence of the cell to the surface (illustrated in Figure 1) [13,39^{***}]. In *P. aeruginosa*, the synthesised cdG specifically binds to FimW, causing it to localise to the cell poles and induce multiple T4P to adhere to the surface [39^{***}]. However, in *C. crescentus* the high levels of cdG bind to the protein HfsJ, which is essential for the biogenesis of the holdfast, the retraction of T4P and consequently to irreversible surface adherence [36]. The commitment to surface attachment in *P. aeruginosa* has been shown to lead to asymmetric division driven by different cdG levels in mother and daughter cells, ultimately leading to a mixed population of surface attached and motile cells reminiscent of the lifecycle of *C. crescentus* [39^{***}].

Complex microbial communities and NSM signalling networks

Much of our knowledge of NSM signalling has been developed for pathways isolated from the wider bacterial signalling network, and for model organisms growing in defined laboratory conditions. Refining our current models to incorporate crosstalk with other NSM networks, and using these models to explain bacterial responses to complex, dynamic environments remain major challenges in molecular microbiology. It is easy (and often practical) to assume that bacterial gene expression is reasonably

uniform throughout a clonal bacterial colony. However, work over the last decade has shown that this is generally not the case. Spatial and temporal diversity within colonies and biofilms has been shown for species including *P. aeruginosa* and *E. coli* [12,40], suggesting in turn that NSM signalling is a major driver of stratification and diversification within bacterial communities.

Biofilm matrix heterogeneity in *E. coli* is generated by a cdG-dependent network characterised by nested positive and negative feedback loops [12]. The transcription factor CsgD is a key regulator in this complex signalling cascade, controlling localised expression of key cdG enzymes [41]. The cascade begins when two DGC/PDE pairs; PdeH/DgcE and DgcM/PdeR activate the transcription factor MlrA, which in turn drives expression of CsgD. Another input to biofilm architecture comes from the PDE PdeC [33], whose activity is linked to oxygen levels that decrease along a steep vertical gradient within the biofilm. Micro-aerobic conditions deep within biofilms promote the free-thiol form of the PdeC periplasmic domain, activating the enzyme and decreasing intracellular cdG levels, and suppressing EPS production [33].

Pseudomonas spp. have particularly complex cdG-signalling networks, with dozens of metabolic enzymes and binding targets. Recently, several groups have used high-throughput analysis methods to define the functions and environmental responses of different *Pseudomonas* spp. cdG networks [42^{*},43,44]. Dahlstrom *et al.* [42^{*}] investigated the role of a large number of *Pseudomonas fluorescens* cdG-signalling proteins under multiple nutrient conditions. This work uncovered a large scale multimodal network

Figure 1




Illustration of flagellar-based surface sensing in *P. aeruginosa* (adapted from Ref. [39^{***}]) and *C. crescentus* (adapted from Ref. [13]) showing the role of cdG signalling in surface attachment. The presence of a surface is detected through the flagellar motor complex (green), which upregulates DGCs and in turn increases the intracellular level of cdG (blue) leading to type IV pili assembly (red) and subsequent surface attachment. For *C. crescentus* the increased cdG level leads to the deployment of the holdfast (purple). Asymmetrical intracellular levels of cdG during cell division give rise to one bacterium favouring a motile lifestyle and the other committed to surface attachment [39^{***}].

of PDEs and DGCs that were able to respond to a variety of signals, giving rise to a complex, context-dependent regulatory network based on a combination of ligand-binding, protein–protein interaction, and transcriptional regulation to fine-tune cdG outputs and tightly control biofilm formation [42^{*}]. Nie *et al.* recently conducted a similar study in *Pseudomonas putida*. By screening the phenotypes associated with deletion/overexpression mutants in 42 predicted cdG-metabolic genes, the authors assigned preliminary roles of each gene in colony morphology, biofilm formation and swimming motility [43]. These papers show that the roles of individual PDEs and DGCs play in bacterial behaviour are dynamic, with individual proteins responding to a diverse set of external stimuli and producing specific phenotypic responses. These studies go some way towards answering the question: ‘why do bacteria encode so many NSM metabolic enzymes?’.


Finally, Little *et al.* investigated the functions of a large group of *P. fluorescens* PDEs and DGCs during plant colonisation. They showed that cdG-gene upregulation

was not uniform during root colonisation, nor did patterns of gene regulation group exclusively by enzyme family [44]. These data suggest that different groups of NSM enzymes are required at different stages of the root colonisation process, likely due to differing environmental stimuli (illustrated in Figure 2). This study was one of a series of recently published papers examining different aspects of cdG-signalling during bacterial plant colonisation [44–48]. While there is still a great deal of work needed to fully understand how NSM signalling functions during complex host–microbe interactions, the past few years have seen encouraging steps in this direction.

Novel NSM molecules: more than microbial signals?

One of the most exciting recent discoveries in NSM signalling has uncovered the true diversity of NSM signalling found in nature [49^{••}]. Whiteley *et al.* combined bioinformatic analyses with a systematic biochemical screen to discover a large family of cGAS/DncV-like nucleotidyl-transferases (CD-NTases) that synthesise a diverse range

Figure 2



A predicted model for signalling during rhizosphere colonisation, (adapted from Ref. [44]). In the early stages of root colonisation, the balance of DGC (pentagons) and PDE (circle) transcription suggests an increase in cdG favouring root attachment and siderophore production (red). As colonisation progresses, transcriptional levels of the tested genes suggest a balance between motile, surface-attached and siderophore producing bacteria. During the later stages of colonisation, upregulation of the PDE PFLU6074 leads to a shift towards cdG degradation and a more motile lifestyle. The size of each shape denotes the level of transcription with larger shapes corresponding to higher transcript abundance.

of cyclic-NSMs, including asymmetric molecules and ones based on pyrimidine nucleotides. CD-NTase enzymes were also discovered that produced cyclic trinucleotides, further expanding the NSM signalling space [49^{••}]. While the biological function and importance of individual novel NSM molecules is still unclear, the implications of this finding for our understanding of the diversity and complexity of NSM signalling cannot be understated.

Since their discovery, research into NSM molecules has focussed almost exclusively on their roles as microbial signalling molecules. However, a recent study by Ahmad *et al.* [50^{••}] has uncovered a new role for the NSM molecule (p)ppApp as an interbacterial toxin. *P. aeruginosa* (p)ppApp synthesis is catalysed by a (p)ppGpp synthase homolog; Tas1, which is injected into neighbouring cells through the type VI secretion system effector [50^{••}]. This novel mechanism of bacterial warfare acts by depleting the target cells pool of ATP and inhibiting ATP synthesis [50^{••}]. Furthermore, Tas1 mediated (p)ppApp synthesis was shown to decrease essential pathway metabolites, dysregulate the proton motive force and interact with known (p)ppGpp binding proteins to disrupt their function.

Conclusions

The last few years have seen remarkable advances in our understanding of NSM signalling in bacteria. New binding proteins and signal inputs have been characterised. Molecular and structural analyses have established new principles of signalling, crosstalk and network formation. Research into whole-cell, multicellular and host-microbe signalling networks have begun to uncover how NSM signalling operates in increasingly complex environments. Finally, our understanding of the nature of NSM function in bacteria has expanded, both in terms of the known NSM repertoire, and the discovery of new, non-signalling functions for these molecules. On this basis, the future for NSM signalling research promises to be equally exciting.

Conflict of interest statement

Nothing declared.

Acknowledgements

JGM was supported in this work by the University of East Anglia and the Biotechnology and Biological Sciences Research Council Institute Strategic Program Grant; Plant Health [BBS/E/J/000PR9797], to the John Innes Centre. CMAT was supported by Biotechnology and Biological Sciences Research Council Responsive Mode Grant [BB/R018154]. The authors would like to thank Dr Richard Little for critical comments on the manuscript.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Perlman RL, De Crombrughe B, Pastan I: **Cyclic AMP regulates catabolite and transient repression in *E. coli***. *Nature* 1969, **223**:810-812.

2. Cashel M, Kalbacher B: **The control of ribonucleic acid synthesis in *Escherichia coli*. V. Characterization of a nucleotide associated with the stringent response**. *J Biol Chem* 1970, **245**:2309-2318.
3. Ross P, Weinhouse H, Aloni Y, Michaeli D, Weinberger-Ohana P, Mayer R, Braun S, de Wroom E, van der Marel GA, van Boom JH *et al.*: **Regulation of cellulose synthesis in *Acetobacter xylinum* by cyclic diguanylic acid**. *Nature* 1987, **325**:279-281.
4. Corrigan RM, Abbott JC, Burhenne H, Kaever V, Grundling A: **c-di-AMP is a new second messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope stress**. *PLoS Pathog* 2011, **7**:e1002217.
5. Ming ZH, Wang W, Xie YC, Ding PF, Chen YC, Jin DZ, Sun Y, Xia B, Yan LM, Lou ZY: **Crystal structure of the novel di-nucleotide cyclase from *Vibrio cholerae* (DncV) responsible for synthesizing a hybrid cyclic GMP-AMP**. *Cell Res* 2014, **24**:1270-1273.
6. Galperin MY, Nikolskaya AN, Koonin EV: **Novel domains of the prokaryotic two-component signal transduction systems**. *FEMS Microbiol Lett* 2001, **203**:11-21.
7. Jenal U, Reinders A, Lori C: **Cyclic di-GMP: second messenger extraordinaire**. *Nat Rev Microbiol* 2017, **15**:271-284.
8. Hengge R, Haussler S, Pruteanu M, Stulke J, Tschowri N, Turgay K: **Recent advances and current trends in nucleotide second messenger signaling in bacteria**. *J Mol Biol* 2019, **431**:908-927.
9. Hengge R: **Trigger phosphodiesterases as a novel class of c-di-GMP effector proteins**. *Philos Trans R Soc Lond B Biol Sci* 2016, **371**.
10. Abel S, Bucher T, Nicollier M, Hug I, Kaever V, Abel Zur Wiesch P, Jenal U: **Bi-modal distribution of the second messenger c-di-GMP controls cell fate and asymmetry during the *Caulobacter* cell cycle**. *PLoS Genet* 2013, **9**:e1003744.
11. Sarenko O, Klauck G, Wilke FM, Pfiffer V, Richter AM, Herbst S, Kaever V, Hengge R: **More than enzymes that make or break cyclic Di-GMP-local signaling in the interactome of GGDEF/EAL domain proteins of *Escherichia coli***. *mBio* 2017, **8**.
12. Serra DO, Hengge R: **A c-di-GMP-based switch controls local heterogeneity of extracellular matrix synthesis which is crucial for integrity and morphogenesis of *Escherichia coli* macrocolony biofilms**. *J Mol Biol* 2019, **431**:4775-4793.
13. Sangermani M, Hug I, Sauter N, Pfohl T, Jenal U: **Tad pili play a dynamic role in *Caulobacter crescentus* surface colonization**. *mBio* 2019, **10**.
14. Woodward JJ, Iavarone AT, Portnoy DA: **c-di-AMP secreted by intracellular *Listeria monocytogenes* activates a host type I interferon response**. *Science* 2010, **328**:1703-1705.
15. DeFrancesco AS, Masloboeva N, Syed AK, DeLoughery A, Bradshaw N, Li GW, Gilmore MS, Walker S, Losick R: **Genome-wide screen for genes involved in eDNA release during biofilm formation by *Staphylococcus aureus***. *Proc Natl Acad Sci U S A* 2017, **114**:E5969-E5978.
16. Schaper S, Yau HCL, Krol E, Skotnicka D, Heimerl T, Gray J, Kaever V, Sogaard-Andersen L, Vollmer W, Becker A: **Seven-transmembrane receptor protein RgsP and cell wall-binding protein RgsM promote unipolar growth in rhizobiales**. *PLoS Genet* 2018, **14**:e1007594.
17. Gallagher KA, Schumacher MA, Bush MJ, Bibb MJ, Chandra G, Holmes NA, Zeng W, Henderson M, Zhang H, Findlay KC *et al.*: **c-di-GMP arms an anti-sigma to control progression of multicellular differentiation in streptomyces**. *Mol Cell* 2019, **77**:586-599.e6.
18. Laventie BJ, Nesper J, Ahrne E, Glatter T, Schmidt A, Jenal U: **Capture compound mass spectrometry—a powerful tool to identify novel c-di-GMP effector proteins**. *J Vis Exp* 2015:51404.
19. Trampari E, Stevenson CE, Little RH, Wilhelm T, Lawson DM, Malone JG: **Bacterial rotary export ATPases are allosterically**

- regulated by the nucleotide second messenger cyclic-di-GMP. *J Biol Chem* 2015, **290**:24470-24483.
20. Perez-Mendoza D, Bertinetti D, Lorenz R, Gallegos MT, Herberg FW, Sanjuan J: **A novel c-di-GMP binding domain in glycosyltransferase BgsA is responsible for the synthesis of a mixed-linkage beta-glucan.** *Sci Rep* 2017, **7**:8997.
 21. Wang F, He Q, Yin J, Xu S, Hu W, Gu L: **BrIR from *Pseudomonas aeruginosa* is a receptor for both cyclic di-GMP and pyocyanin.** *Nat Commun* 2018, **9**:2563.
- The authors show the co-binding of both an NSM and a virulence factor to a transcriptional regulator, with co-binding enhancing transcriptional regulation. This furthers our understanding of the complex interplay between signalling molecules, virulence factors and gene expression.
22. Okegbe C, Fields BL, Cole SJ, Beierschmitt C, Morgan CJ, Price-Whelan A, Stewart RC, Lee VT, Dietrich LEP: **Electron-shuttling antibiotics structure bacterial communities by modulating cellular levels of c-di-GMP.** *Proc Natl Acad Sci U S A* 2017, **114**: E5236-E5245.
 23. Matsuyama BY, Krasteva PV, Baraquet C, Harwood CS, Sondermann H, Navarro MV: **Mechanistic insights into c-di-GMP-dependent control of the biofilm regulator FleQ from *Pseudomonas aeruginosa*.** *Proc Natl Acad Sci U S A* 2016, **113**: E209-218.
 24. Anderson BW, Liu K, Wolak C, Dubiel K, She F, Satyshur KA, Keck JL, Wang JD: **Evolution of (p)ppGpp-HPRT regulation through diversification of an allosteric oligomeric interaction.** *eLife* 2019, **8**.
 25. Wood A, Irving SE, Bennison DJ, Corrigan RM: **The (p)ppGpp-binding GTPase Era promotes rRNA processing and cold adaptation in *Staphylococcus aureus*.** *PLoS Genet* 2019, **15**:e1008346.
 26. Kim SK, Lormand JD, Weiss CA, Eger KA, Turdiev H, Turdiev A, Winkler WC, Sondermann H, Lee VT: **A dedicated diribonucleotidase resolves a key bottleneck for the terminal step of RNA degradation.** *eLife* 2019, **8**.
 27. Giacalone D, Smith TJ, Collins AJ, Sondermann H, Koziol LJ, O'Toole GA: **Ligand-mediated biofilm formation via enhanced physical interaction between a diguanylate cyclase and its receptor.** *mBio* 2018, **9**.
 28. De N, Pirruccello M, Krasteva PV, Bae N, Raghavan RV, Sondermann H: **Phosphorylation-independent regulation of the diguanylate cyclase WsP_R.** *PLoS Biol* 2008, **6**:e67.
 29. Tschowri N, Busse S, Hengge R: **The BLUF-EAL protein YcgF acts as a direct anti-repressor in a blue-light response of *Escherichia coli*.** *Genes Dev* 2009, **23**:522-534.
 30. Zahringer F, Lacanna E, Jenal U, Schirmer T, Boehm A: **Structure and signaling mechanism of a zinc-sensory diguanylate cyclase.** *Structure* 2013, **21**:1149-1157.
 31. Xu M, Yang X, Yang XA, Zhou L, Liu TZ, Fan ZS, Jiang T: **Structural insights into the regulatory mechanism of the *Pseudomonas aeruginosa* YfiBNR system.** *Protein Cell* 2016, **7**:403-416.
 32. Yang X, Yang XA, Xu M, Zhou L, Fan ZS, Jiang T: **Crystal structures of YfiR from *Pseudomonas aeruginosa* in two redox states.** *Biochem Biophys Res Commun* 2015, **461**:14-20.
 33. Herbst S, Lorkowski M, Sarenko O, Nguyen TKL, Jaenicke T, Hengge R: **Transmembrane redox control and proteolysis of PdeC, a novel type of c-di-GMP phosphodiesterase.** *EMBO J* 2018, **37**.
 34. Paiardini A, Mantoni F, Giardina G, Paone A, Janson G, Leoni L, Rampioni G, Cutruzzola F, Rinaldo S: **A novel bacterial l-arginine sensor controlling c-di-GMP levels in *Pseudomonas aeruginosa*.** *Proteins* 2018, **86**:1088-1096.
 35. Nesper J, Hug I, Kato S, Hee CS, Habazettl JM, Manfredi P, Grzesiek S, Schirmer T, Emonet T, Jenal U: **Cyclic di-GMP differentially tunes a bacterial flagellar motor through a novel class of CheY-like regulators.** *eLife* 2017, **6**.
 36. Hug I, Deshpande S, Sprecher KS, Pfohl T, Jenal U: **Second messenger-mediated tactile response by a bacterial rotary motor.** *Science* 2017, **358**:531-534.
 37. Schniederberend M, Williams JF, Shine E, Shen C, Jain R, Emonet T, Kazmierczak BI: **Modulation of flagellar rotation in surface-attached bacteria: a pathway for rapid surface-sensing after flagellar attachment.** *PLoS Pathog* 2019, **15**: e1008149.
 38. Baker AE, Webster SS, Diepold A, Kuchma SL, Bordeleau E, Armitage JP, O'Toole GA: **Flagellar staters stimulate c-di-GMP production by *Pseudomonas aeruginosa*.** *J Bacteriol* 2019, **201**.
 39. Laventie BJ, Sangermani M, Estermann F, Manfredi P, Planes R, Hug I, Jaeger T, Meunier E, Broz P, Jenal U: **A surface-induced asymmetric program promotes tissue colonization by *Pseudomonas aeruginosa*.** *Cell Host Microbe* 2019, **25**:140-152 e146.
- A detailed analysis of the signalling cascade involved in *P. aeruginosa* surface attachment, as well as an exploration of asymmetrical cell division due to uneven NSM distributions between mother and daughter cells.
40. Werner E, Roe F, Bugnicourt A, Franklin MJ, Heydorn A, Molin S, Pitts B, Stewart PS: **Stratified growth in *Pseudomonas aeruginosa* biofilms.** *Appl Environ Microbiol* 2004, **70**:6188-6196.
 41. Klauck G, Serra DO, Possling A, Hengge R: **Spatial organization of different sigma factor activities and c-di-GMP signalling within the three-dimensional landscape of a bacterial biofilm.** *Open Biol* 2018, **8**.
 42. Dahlstrom KM, Collins AJ, Doing G, Taroni JN, Gauvin TJ, Greene CS, Hogan DA, O'Toole GA: **A multimodal strategy used by a large c-di-GMP network.** *J Bacteriol* 2018, **200**.
- An extensive and methodical large-scale study that begins to dissect the role of individual DGC and PDE networks in *P. fluorescens*, giving some insight into the discrete roles of individual proteins in the overall network.
43. Nie H, Xiao Y, He J, Liu H, Nie L, Chen W, Huang Q: **Phenotypic-genotypic analysis of GGDEF/EAL/HD-GYP domain-encoding genes in *Pseudomonas putida*.** *Environ Microbiol Rep* 2019.
 44. Little RH, Woodcock SD, Campilongo R, Fung RKY, Heal R, Humphries L, Pacheco-Moreno A, Paulusch S, Stigliano E, Vikeli E et al.: **Differential regulation of genes for cyclic-di-GMP metabolism orchestrates adaptive changes during rhizosphere colonization by *Pseudomonas fluorescens*.** *Front Microbiol* 2019, **10**.
 45. Perez-Mendoza D, Felipe A, Ferreira MD, Sanjuan J, Gallegos MT: **AmrZ and FleQ Co-regulate cellulose production in *Pseudomonas syringae* pv. tomato DC3000.** *Front Microbiol* 2019, **10**.
 46. O'Neal L, Akhter S, Alexandre G: **A PilZ-containing chemotaxis receptor mediates oxygen and wheat root sensing in *Azospirillum brasilense*.** *Front Microbiol* 2019, **10**.
 47. Lopez-Farfan D, Reyes-Darias JA, Matilla MA, Krell T: **Concentration dependent effect of plant root exudates on the chemosensory systems of *Pseudomonas putida* KT2440.** *Front Microbiol* 2019, **10**.
 48. Lopez-Baena FJ, Vinardell JM, Medina C: **Regulation of protein secretion systems mediated by cyclic diguanylate in plant-interacting bacteria.** *Front Microbiol* 2019, **10**.
 49. Whiteley AT, Eaglesham JB, de Oliveira Mann CC, Morehouse BR, Lowey B, Nieminen EA, Danilchanka O, King DS, Lee ASY, Mekalanos JJ et al.: **Bacterial cGAS-like enzymes synthesize diverse nucleotide signals.** *Nature* 2019, **567**:194-199.
- An interesting development showing the potential and diversity of NSMs that can be created and used by bacteria. This paper challenges current models of bacterial signalling, suggesting the existence of a far more diverse NSM signalling landscape than previously suspected.
50. Ahmad S, Wang B, Walker MD, Tran HR, Stogios PJ, Savchenko A, Grant RA, McArthur AG, Laub MT, Whitney JC: **An interbacterial toxin inhibits target cell growth by synthesizing (p)ppApp.** *Nature* 2019, **575**:674-678.
- The authors describe both the discovery and novel function of (p)ppApp in interbacterial warfare. This insight is an exciting development in the field of NSMs, moving away from classical signalling pathways into bacterial competition.