

Oxygen stability in the new [FeFe]-hydrogenase from *Clostridium beijerinckii* SM10 (CbA5H)

Simone Morra[‡], Mariaconcetta Arizzi[‡], Francesca Valetti*, Gianfranco Gilardi*

Department of Life Sciences and Systems Biology, University of Torino, Via Accademia Albertina 13, Torino 10123, Italy.

Supporting Information Placeholder

ABSTRACT: The newly isolated [FeFe]-hydrogenase CbA5H was characterized by FTIR spectroscopy coupled to enzymatic activity assays. This showed for the first time that in this enzyme the oxygen-sensitive active state H_{ox} can be simply and reversibly converted to the oxygen-stable inactive H_{inact} state. This suggests that oxygen sensitivity is not an intrinsic feature of the catalytic center of [FeFe]-hydrogenases (H-cluster), opening new challenging perspectives on the oxygen sensitivity mechanism as well as new possibilities for the exploitation in industrial applications.

[FeFe]-hydrogenases are a vast class of redox enzymes that catalyze reversibly the reaction of H_2 evolution by the catalytic center H-cluster (Fig. 1A).¹ Several hundreds of sequenced genes can potentially express enzymes that are members of this class,² but so far only few of them have been characterized.³⁻⁵ Thus, the isolation and characterization of novel [FeFe]-hydrogenases is expected to provide novel important information.^{6,7,8}

A common feature to all [FeFe]-hydrogenases is oxygen sensitivity: these enzymes are found in strict anaerobic or facultative anaerobic organisms and they are inactivated by molecular oxygen. Oxygen inactivation is a complex mechanism, still under active debate. Variability in the inactivation rate has been described between enzymes from different organisms.⁹⁻¹⁵

Oxygen sensitivity is the main factor that hinders the exploitation of [FeFe]-hydrogenases for applicative purposes^{3,16,17} as generally exposure to O_2 leads to irreversible inactivation on a time scale of seconds. This is believed to be due to a partial degradation of the H-cluster, via a multistep process.^{14,15,18} Partial reversible reactivation in some [FeFe]-hydrogenases was previously reported by electrochemical studies, but resulting from a very complex interplay of electrode potential and anaerobic/aerobic inactivation processes.^{12,13,19} Protein engineering was used to generate [FeFe]-hydrogenase mutants with slightly decreased oxygen sensitivity.²⁰

So far, the clearest exceptions to irreversible inactivation are the [FeFe]-hydrogenases from *Desulfovibrio desulfuricans* (DdH),^{21,22} *D. vulgaris* (DvH)^{23,24,25} and *D. gigas*.²⁶ DdH is known to produce the inactive and oxygen-stable redox state called H_{inact} (Fig. 1B).^{21,22} This state was originally identified in DdH preparations that have been purified aerobically.²⁷




Figure 1. A) Scheme of the H-cluster structure. B) Scheme of the H-cluster redox states and their interconversion, as known in DdH (black arrows).^{4,21,22} The red arrow marks the reversible transition described in this paper in CbA5H.

Spectroscopic experiments showed that this form of the DdH enzyme can be activated by a reductive treatment but, subsequently, the H_{inact} state cannot be formed again.^{13,21,22} Also, when the activated protein is exposed to oxygen, the H-cluster is degraded.^{13,28} In particular, spectro-electrochemical experiments on DdH have shown that H_{inact} can be reversibly converted into the intermediate state H_{trans} by a single electron reduction and that, subsequently, H_{trans} can be converted irreversibly into the active state H_{ox} by a two electrons reduction. Attempts to prepare intentionally *in vitro* the H_{inact} state were usually unsuccessful^{21,22}, with the exception of a single report of a complex procedure²⁸ that could not be successfully reproduced^{21,22}. Cyclic voltammetry experiments showed the reversible formation of an inactive species under anaerobic conditions, but no spectroscopic evidence has determined its identity.^{9,16,29}

In this work, we report the first characterization of a novel monomeric [FeFe]-hydrogenase from the strain *Clostridium beijerinckii* SM10 that was isolated from an efficient bio-hydrogen pilot plant fed with vegetable wastes.⁶ The gene sequence was deposited in the NCBI database (accession KX147468) and the enzyme was recombinantly produced in *E. coli* under strict anaerobic conditions.^{30,31}

Phylogenetically, the enzyme belongs to cluster A5 and has a modular structure M2c.^{5,32} For this reason, the enzyme has been named as CbA5H. Starting from the N-terminus, it is composed by 1) a domain hosting the poorly characterized SLBB motif (soluble-ligand-binding β -grasp fold);³³ 2) a domain hosting two [4Fe4S] centers that is homologous to bacterial ferredoxins and that is widely distributed in hydrogenases; 3) the H-domain hosting the catalytic center H-cluster, which shows high sequence similarity to other [FeFe]-hydrogenases (See Fig. S1).

When the enzyme is purified anaerobically in the presence of 2 mM sodium dithionite as oxygen scavenger, it is active both in the H₂ uptake assay and the H₂ evolution assay, with rates that are 158±34 and 751±91 $\mu\text{mol H}_2 \text{ min}^{-1} \text{ mg protein}^{-1}$, respectively.

A detailed FTIR analysis was performed to characterize the specific influence of the protein environment on the H-cluster properties in this hydrogenase. The assignment of the various signals to known redox states as discussed below, was obtained by comparison to data previously reported for other [FeFe]-hydrogenases.^{21,22,33}

The anaerobically purified enzyme mainly equilibrated in the H_{ox} state, with minor contributions from the H_{red} state (fig. 2A). The FTIR spectral signature of the H_{ox} is composed by five peaks at 2091, 2080, 1964, 1940 and 1800 cm⁻¹. The FTIR spectral signature of H_{red} was more difficult to identify, due to low intensity signals that could be identified at 2040, 1915 and 1893 cm⁻¹. Treatment of this sample with carbon monoxide caused a complete shift to the H_{ox}-CO state (fig. 2B), whose spectral signature is composed by six peaks at 2094, 2090, 2016, 1971, 1963 and 1807 cm⁻¹.

These data show that the structure of CbA5H H-cluster is highly consistent with that of other [FeFe]-hydrogenases:^{21,22,34,35} two terminal CO, two terminal CN and a bridging CO are coordinated to the [2Fe] sub-cluster in the H_{ox} state; upon reduction, the bridging CO is shifted to a semi-bridging position (the signal at 1800 cm⁻¹ in H_{ox} shifts to 1893 cm⁻¹ in H_{red}); upon CO exposure, the new ligand binds to the vacant coordination of Fe_d, causing a vast rearrangement in the vibrational modes of all the ligands.

Much more interestingly, treatment of the anaerobically purified protein with different organic oxidants, such as thionine ($E_{m7} = +60 \text{ mV}$), 2,6-dichlorophenolindophenol (DCIP, $E_{m7} = +220 \text{ mV}$) or air (*i.e.* oxygen) produce samples displaying an homo-

geneous spectrum with the H_{inact} signature with signals at 2107, 2080, 2011, 1992 and 1840 cm⁻¹ (fig. 2C, 2D, 2E). The H_{inact} could be reconverted to a mixture of H_{ox} and H_{red} by reducing treatment with sodium dithionite or H₂ (fig. 2F and 2G).




Figure 2. FTIR spectra of *Clostridium beijerinckii* [FeFe]-hydrogenase CbA5H. A) After anaerobic purification, without further treatment. B) After CO treatment. C) After thionine oxidation (8-fold molar ratio). D) After DCIP oxidation (24-fold molar ratio). E) After oxidation with air (10 min). F) After oxidation with air and reduction with sodium dithionite (10-fold molar ratio). G) After oxidation with air and reduction with H₂ (50 min). The protein concentration was 0.3-1 mM and opportune scaling factors were applied to the spectra for simpler comparison, as follows: B) x2. C) x3.5. D) x1.8. E) x1.6. F) x1.2. G) x2.4.

These first evidences show that in *Clostridium beijerinckii* [FeFe]-hydrogenase it is possible to convert the H_{ox} state into the H_{inact} state and that the conversion is spontaneous upon oxidation by thionine or DCIP or by the oxygen present in air, while this has never been observed in DdH, the H_{inact} to H_{ox} reaction being irreversible, according to FTIR evidences.^{21,22} More detailed experiments were carried out to correlate the spectral change to the catalytic rate of CbA5H. To confirm the functional properties of the different redox states of the H-cluster, the same sample was split: a part was used to acquire FTIR spectra (fig. 3) and the other part tested for H₂ uptake activity (fig. 4).

As already presented above, the anaerobically purified enzyme was a catalytically active mixture of H_{ox} and

H_{red} (Fig. 3A and 4) and it could be pushed towards the inactive H_{inact} by air treatment (Fig. 3B and 4). Subsequently, the enzyme could be completely reactivated by H_2 treatment; after the reactivation, CbA5H equilibrated again in a mixture of H_{ox} and H_{red} states, where the reduced state was more abundant (Fig. 3C and 4).




Figure 3. FTIR spectra of CbA5H after anaerobic or aerobic purification and after various cycles of activation/inactivation. A) After anaerobic purification, without further treatment. B) The anaerobically purified sample after exposure to air (10 min). C) The anaerobically purified sample after exposure to air (10 min) followed by exposure to H_2 (50 min). D) The anaerobically purified sample after exposure to air (10 min), followed by exposure to H_2 (50 min) and then air (10 min). E) The anaerobically purified sample after two cycles of exposure to air, followed by exposure to H_2 . F) After aerobic purification, without further treatment. G) The aerobically purified sample after exposure to H_2 (50 min). H) The aerobically purified sample after exposure to H_2 (50 min) followed by exposure to air (10 min). The protein concentration was 0.8-1 mM and opportune scaling factors were applied to the spectra for simpler comparison, as follows: C) x1.5. D) x1.5. E) x1.5.

Low intensity peaks at 2075 and 1980 cm^{-1} might suggest the presence of a minor proportion of H_{trans} in this sample. Furthermore, after the reactivation, the enzyme could be converted again completely into H_{inact} by air treatment (Fig. 3D and 4). Once more, the resulting sample could be converted into an active mixture of H_{ox} and H_{red} (Fig. 3E and 4), with low intensity peaks at

2075 and 1987 cm^{-1} that might suggest the presence of a minor proportion of H_{trans} .

These results show that, in CbA5H, the transition H_{ox}/H_{inact} is completely reversible in both directions and that several cycles can be repeated without damaging the structural and functional properties of the enzyme. Control experiments on the O_2 sensitive CaHydA confirmed instead all the signatures¹⁵ of O_2 damage (see Fig. S4).




Figure 4. Enzyme activity of samples presented in figure 3.

Since CbA5H produced very intense and homogeneous H_{inact} spectra when exposed to air and the enzyme was stable, an aerobic purification was carried out. The enzyme was expressed under anaerobic conditions, but the purification was entirely carried out aerobically in the absence of dithionite.

The aerobically purified sample displayed a homogeneous H_{inact} spectrum and it was catalytically inactive (Fig. 3F and 4). Also in this case, it was possible to reactivate completely the enzyme with a H_2 treatment: this sample showed a mixture of H_{ox} and H_{red} signals and full H_2 evolution activity (Fig. 3G and 4). Furthermore, this sample could be re-inactivated by air, displaying a homogeneous H_{inact} spectrum and strong activity loss (Fig. 3H and 4). The low intensity signal at 2014-2015 cm^{-1} present in most of the active CbA5H samples might be an unassigned vibrational mode that was previously observed in reduced samples of DdH;^{21,22} alternatively, it might be due to a very low amount of H_{ox} -CO, which is often found in [FeFe]-hydrogenases preparations. The low intensity signals at 2098-2099 and 2003 cm^{-1} present in some of the inactive CbA5H samples could not be assigned to any previously described H-cluster signal or redox state.

In conclusion, the data presented here show that in *Clostridium beijerinckii* [FeFe]-hydrogenase CbA5H the transition between the oxygen-sensitive catalytically active H_{ox} state and the oxygen-stable inactive H_{inact} state is completely reversible. The transition can be driven by mild organic oxidants, such as thionine or DCIP, or simply by exposure to the oxygen present in air. Moreover, the transition H_{ox}/H_{inact} can be repeated several times (at least two), while in DdH after the first transition from H_{inact} to H_{ox} the reverse reaction was impossible and oxygen damage was observed.

These results give an original contribution to the discussion on oxygen sensitivity in [FeFe]-hydrogenases, because they show that irreversible oxygen sensitivity is not an intrinsic property of the H-cluster, but it is very strongly influenced by the protein environment. It seems clear that, upon oxygen exposure, the protein environment of different enzymes can tune the fate of the H-cluster towards an irreversible O₂ damage or a completely reversible transition to the oxygen-stable H_{inact}.

The exact mechanism in CbA5H is currently under investigation: more detailed experiments will be performed to assess if there are specific residues in the H-domain that can influence the H_{ox}/H_{inact} transition or if this novel effect is more specifically due to the accessory domains present in this new enzyme (SLBB, 2[4Fe4S] ferredoxin-like).

Another important conclusion is that air does not cause any irreversible damage to the enzyme, since the transition H_{ox}/H_{inact} can be performed several times without any activity loss or spectral influence. The treatment did not result in significant appearance of H_{ox}-CO state, typical signature of H-cluster decomposition¹⁵.

Consequently, these data open completely new perspectives for the exploitation of [FeFe]-hydrogenases in real applications. The use of CbA5H (or engineered enzymes with similar properties) would make the enzyme preparation, manipulation and storage much simpler than with any other [FeFe]-hydrogenase and pave the way for applicative exploitation of these highly efficient biocatalysts in hydrogen producing devices.

ASSOCIATED CONTENT

Supporting Information.

The Supporting Information is available free of charge on the ACS Publications website (<http://pubs.acs.org>). Experimental details, Sequence alignment, FTIR spectra with all signals, control experiments on CaHydA (PDF).

AUTHOR INFORMATION

Corresponding Authors

*francesca.valetti@unito.it, *gianfranco.gilardi@unito.it

Author Contributions

‡These authors contributed equally.

Notes

The authors declare no competing financial interests.

List of abbreviations

CbA5H: *Clostridium beijerinckii* [FeFe]-hydrogenase cluster A5. CaHydA: *C. acetobutylicum* [FeFe]-hydrogenase. DCIP: 2,6-dichlorophenolindophenol. DdH: *Desulfovibrio desulfuricans* [FeFe]-hydrogenase, DvH: *D. vulgaris* [FeFe]-hydrogenase. SLBB:soluble-ligand-binding β-grasp.

Funding

This work was supported by project HyStrEM (E.U. Structural Funds N.1083/2006 F.E.S.R. 2007-2013).

REFERENCES

- (1) Vignais, P.M., and Billoud, B. (2007) *Chem. Rev.* 107, 4206-4272.
- (2) Meyer, J. (2007) *Cell. Mol. Life Sci.* 64, 1063-1084.
- (3) Winkler, M., Esselborn, J., and Happe, T. (2013) *Biochim Biophys Acta* 1827, 974-985.
- (4) Lubitz, W., Ogata, H., Rüdiger, O., and Reijerse, E. (2014) *Chem. Rev.* 114, 4081-4148.
- (5) Peters, J.W., Schut, G.J., Boyd, E., Mulder, D.W., Shepard, E.M., Broderick, J.B., King, P.W., and Adams, M.W.W. (2015) *Biochim. Biophys. Acta* 1853, 1350-1369.
- (6) Morra, S., Arizzi, M., Allegra, P., La Licata, B., Sagnelli, F., Zitella, P., Gilardi, G., and Valetti, F. (2014) *Int. J. Hydrogen Energy* 39, 9018-9027.
- (7) Morra, S., Valetti, F., Sarasso, V., Castrignanò, S., Sadeghi, S.J., and Gilardi, G. (2015) *Bioelectrochemistry* 106(Pt B), 258-262.
- (8) Morra, S., Mongili, B., Maurelli, S., Gilardi, G., and Valetti, F. (2016) *Biotechnol. Appl. Biochem.* 63, 305-311.
- (9) Vincent, K.A., Parkin, A., Lenz, O., Albracht, S.P.J., Fontecilla-Camps, J.C., Cammack, R., Friedrich, B., and Armstrong, F.A. (2005) *J. Am. Chem. Soc.* 127, 18179-18189.
- (10) Stripp, S.T., Goldet, G., Brandmayr, C., Sanganas, O., Vincent, K.A., Haumann, M., Armstrong, F.A., and Happe, T. (2009) *Proc. Natl. Acad. Sci.* 106, 17331-17336.
- (11) Lautier, T., Ezanno, P., Baffert, C., Fourmond, V., Cournac, L., Fontecilla-Camps, J.C., Soucaille, P., Bertrand, P., Meynial-Salles, I., and Léger, C. (2011) *Faraday Discuss.* 148, 385-407.
- (12) Orain, C., Saujet, L., Gauquelin, C., Soucaille, P., Meynial-Salles, I., Baffert, C., Fourmond, V., Bottin, H., and Léger, C. (2015) *J. Am. Chem. Soc.* 137, 12580-12587.
- (13) Goldet, G., Brandmayr, C., Stripp, S.T., Happe, T., Cavazza, C., Fontecilla-Camps, J.C., and Armstrong, F.A. (2009) *J. Am. Chem. Soc.*, 131, 14979-14989.
- (14) Lambertz, C., Neidel, N., Havelius, K.G.V., Noth, J., Chernev, P., Winkler, M., Happe, T., and Haumann, M. (2011) *J. Biol. Chem.* 286, 40614-40623.
- (15) Swanson, K.D., Ratzloff, M.W., Mulder, D.M., Artz, J.H., Ghose, S., Hoffman, A., White, S., Zadovnyy, O.A., Broderick, J.B., Bothner, B., King, P.W., and Peters, J.W. (2015) *J. Am. Chem. Soc.* 137, 1809-1816.
- (16) Vincent, K.A., Parkin, A., and Armstrong, F.A. (2007) *Chem. Rev.* 107, 4366-4413.
- (17) King, P.W. (2013) *Biochim. Biophys. Acta* 1827, 949-657.
- (18) Adams, M.W.W. (1990) *Biochim. Biophys. Acta* 1020, 115-145.
- (19) Fourmond, V., Greco, C., Sybirna, K., Baffert, C., Wang, P.H., Ezanno, P., Montefiori, M., Bruschi, M., Meynial-Salles, I., Soucaille, P., Blumberger, J., Bottin, H., De Gioia, L., Léger, C. (2014) *Nature Chem.* 6, 336-342.
- (20) Bingham, A.S., Smith, P.R., Swartz, J.R. (2012) *Int. J. Hydrogen Energy* 37, 2965-2976.
- (21) Roseboom, W., De Lacey, A.L., Fernandez, V.M., Hatchikian, E.C., and Albracht, S.P.J. (2006) *J. Biol. Inorg. Chem.* 11, 102-118.
- (22) De Lacey, A.L., and Fernandez, V.M. (2007) *Chem. Rev.* 107, 4304-4330.
- (23) van der Westen, H.M., Mayhew, S.G., and Veeger, C. (1978) *FEBS Lett.* 86, 122-126.
- (24) Pierik, A.J., Hagen, W.R., Redeker, J.S., Wolbert, R.B., Boldersma, M., Verhagen, M.F., Grande, H.J., Veeger, C., Mutsaers, P.H., Sands, R.H., and Dunham, W.R. (1992) *Eur. J. Biochem.* 209, 63-72.
- (25) Pierik, A.J., Hulstein, M., Hagen, W.R., and Albracht, S.P.J. (1998) *Eur. J. Biochem.* 258, 572-578.
- (26) Fernandez, V.M., Hatchikian, E.C., and Cammack, R. (1985) *Biochim. Biophys. Acta* 832, 69-79.
- (27) Hatchikian, E.C., Forget, N., Fernandez, V.M., Williams, R., and Cammack, R. (1992) *Eur. J. Biochem.* 209, 357-365.
- (28) van Dijk, C., van Berkel-Arts, A., and Veeger, C. (1983) *FEBS Lett.* 156, 340-344.
- (29) Parkin, A., Cavazza, C., Fontecilla-Camps, J.C., and Armstrong, F.A. (2006) *J. Am. Chem. Soc.* 128, 16808-16815.

(30) King, P.W., Posewitz, M.C., Ghirardi, M.L., and Seibert, M. (2006) *J. Bacteriol.* 188, 2163-2172.

(31) Morra, S., Cordara, A., Gilardi, G., and Valetti, F. (2015) *Protein Sci.* 24, 2090-2094.

(32) Calusinska, M., Happe, T., Joris, B., and Wilmotte, A. (2010) *Microbiology* 156, 1575-1588.

(33) Burroughs, A.M., Balaji, S., Iyer, L.M., and Aravind, L. (2007) *Biol. Direct.* 2, 18-45.

(34) Silakov, A., Kamp, C., Reijerse, E., Happe, T., and Lubitz, W. (2009) *Biochemistry* 48, 7780-7786.

(35) Nicolet, Y., De Lacey, A.L., Vernède, X., Fernandez, V.M., Hatchikian, E.C., and Fontecilla-Camps, J.C. (2001) *J. Am. Chem. Soc.* 123, 1596-1601.

Authors are required to submit a graphic entry for the Table of Contents (TOC) that, in conjunction with the manuscript title, should give the reader a representative idea of one of the following: A key structure, reaction, equation, concept, or theorem, etc., that is discussed in the manuscript. Consult the journal's Instructions for Authors for TOC graphic specifications.

Insert Table of Contents artwork here

