

Northumbria Research Link

Citation: Wilhelm, Eurico N., Mourot, Laurent and Rakobowchuk, Mark (2018) Exercise-Derived Microvesicles: A Review of the Literature. *Sports Medicine*, 48 (9). pp. 2025-2039. ISSN 0112-1642

Published by: Adis International

URL: <https://doi.org/10.1007/s40279-018-0943-z> <<https://doi.org/10.1007/s40279-018-0943-z>>

This version was downloaded from Northumbria Research Link: <http://nrl.northumbria.ac.uk/42470/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University**
NEWCASTLE



UniversityLibrary

1 Exercise-derived microvesicles: a review of literature

2 Eurico N. Wilhelm^{1*}, Laurent Mourot², Mark Rakobowchuk³

3

4 ¹ School of Physical Education, UFPel, Pelotas – RS, Brazil.

5 ² EA3920 Prognostic Factors and Regulatory Factors of Cardiac and Vascular Pathologies, (Exercise
6 Performance Health Innovation - EPHI), Univ. Bourgogne Franche-Comté, F-25000 Besançon, France
7 and Tomsk Polytechnic University, Tomsk, Russia.

8 ³ Department of Biological Sciences, Faculty of Science, Thompson Rivers University, Kamloops – BC,
9 Canada.

10 * Corresponding author

11

12 * Corresponding author contact details:
13 School of Physical Education, UFPel
14 Rua Luís de Camões, 625 - Três Vendas
15 Pelotas – RS, Brazil
16 Postal Code: 96055-630
17 Telephone: +55 53 32732752
18 Email: euricowilhelm@gmail.com

19

20 Short Title: Microvesicles and Exercise

1 Abstract

2 Initially suggested as simple cell debris, cell-derived microvesicles (MVs) have now gained acceptance
3 as recognized players in cellular communication and physiology. Shed by most, and perhaps all, human
4 cells, these tiny lipid-membrane vesicles carry bioactive agents, such as proteins, lipids, and microRNA
5 from their cell source, and are produced under orchestrated events in response to a myriad of stimuli.
6 Physical exercise introduces systemic physiological challenges capable of acutely disrupting cell
7 homeostasis and stimulating the release of MVs into the circulation. The novel and promising field of
8 exercise-derived MVs is expanding quickly, and the following work provides a review of the influence
9 of exercise on circulating MVs, considering both acute and chronic aspects of exercise and training.
10 Potential effects of the MV response to exercise are highlighted and future directions suggested as
11 exercise and sports sciences extend the realm of extracellular vesicles.

12

13 Key points

- 14
- 15 • Cells naturally release microvesicles (MVs) known to regulate a variety of physiological
16 processes including haemostasis and vascular adaptations.
 - 17 • Acute exercise can influence the production and clearance of certain circulating MVs, with
18 remarkable effect in stimulating a transient increase in the concentration of MVs derived from
19 platelets.
 - 20 • In selected populations, chronic exercise improves vascular function while reducing the blood
21 concentration of endothelial MVs linked to vascular damage at rest.
 - 22 • Training variables, such as exercise intensity, can be modified to stimulate an acute increase
23 in the concentration of exercise-derived MVs with pro-angiogenic potential.

1. Introduction

When stimulated, different cell types release plasma membrane-derived vesicles into the extracellular compartment, which can enter the circulation and interact with remote tissues. Among these, microvesicles (MVs), also known as microparticles, transport lipids, proteins, and transcripts from their parental cells. Despite the long debate about whether platelets should be classified as cells [1], thrombocytes actively release MVs and comprise a large (if not the largest) fraction of circulating MVs [2–4].

Previously viewed as biomarkers of the parental cell state, we now understand that MVs are not only passive by-products of cell membrane cytoskeleton reorganisation, but active agents released under specific environmental stresses and capable of triggering functional and structural alterations in recipient cells [5–7]. The profile [8–10] and content [9,11] of circulating MVs differ between healthy and clinical conditions, with the concentration of certain MV populations related to impaired vascular health [10,12].

Exercise training is one of the best interventions to maintain cardiovascular function, and recent evidence demonstrates that acute exercise alters circulating MVs concentrations [13–17], especially platelet-derived MVs (PMVs). Although exercise-mediated vascular adaptations like improved endothelial function are mediated in large part by haemodynamic forces (i.e. increased antero-grade shear stress) [18–20], systemic circulating factors have also been suggested as involved [21], and circulating MVs have arisen as putative mediators of local and systemic adaptations to exercise. The precise role played by these tiny vesicles is still being unravelled, and the study of extracellular vesicles has received intensive attention due to its potential in physiology and medicine. As such, this review focuses on the promising field of cell-derived MVs considering their application in exercise physiology and medicine and aims to provide future directions for this novel research area.

1.1. Literature search criteria

Electronic databases were searched (Pubmed/Medline, Scopus and Google Scholar) between 2017 and 2018, with no restriction regarding publication date. English language scientific articles were selected based on content concerning (but not restricted to) the exercise and MV literature. Although the present manuscript is not a systematic review, the literature search criteria included the terms “exercise”, “training”, “microvesicles”, “microparticles”, “cell-derived microparticles”. The reference list of selected manuscripts, as well as from PhD theses and MSc dissertations on the topic, were also used searching sources. For the review of specific literature relating to the influence of exercise upon circulating MVs, articles were selected only if they included at least one acute or chronic exercise intervention group and MV outcome.

2. Background – From extracellular vesicles to cell-derived microvesicles

Cells have been known to release biologically active small vesicles with many functions but not all vesicles are created the same, so the term extracellular vesicles has been introduced as an all-encompassing descriptor of vesicles released by cells into the extracellular compartment. Such vesicles can be sub-classified as exosomes, MVs, and apoptotic bodies, depending on their mechanism of formation, size, and the presence of specific markers. Importantly, nomenclature in this field varies with MVs and microparticles referred to as similar constructs, while others defining MVs as an extracellular vesicle category which encompasses both microparticles and exosomes. Although we appreciate that future efforts are necessary for nomenclature standardization, for clarity in this review MVs and microparticles will be used interchangeably, whereas exosomes are considered distinct.

1 By standard definition, MVs are anucleate vesicular populations derived from plasma membranes,
2 ranging from approximately 0.1 to 1 μm in diameter, and with no synthetic capabilities. MVs differ
3 from exosomes and apoptotic bodies not only in size, but also in how they are formed, their content,
4 and since MVs originate from the plasma membrane, they also carry cell membrane-specific antigens.
5 In contrast, exosomes are produced through a constitutive intracellular process and released by
6 exocytosis upon fusion of multivesicular bodies with the plasma membrane, whereas apoptotic bodies
7 are released as blebs from apoptotic cells and can also carry cell-membrane specific antigens.
8 Although some overlap exists and size definitions vary slightly, exosomes are generally described as
9 ranging from ~ 40 to 100 nm in diameter, and apoptotic bodies are normally characterised as vesicles
10 larger than 1.5 μm that often carrying nuclear content [22,23].

11 Differentiating extracellular vesicle populations is important as each differs in function and may
12 exhibit broad physiological and pathophysiological effects. This review focuses on MVs, but the reader
13 is referred to previous publications for specific reviews on exosomes and exercise [24,25].

14 2.1. Introduction to circulating microvesicles

15 In 1967 Peter Wolf published a detailed manuscript describing that clot-formation occurred even in
16 platelet free-plasma as long as platelet-derived elements were present, which he named “platelet
17 dust” [26]. These pro-coagulant fragments were later found to have originated from the plasma
18 membrane and several years later, this was confirmed by electron microscopy after platelet
19 stimulation with thrombin [27], providing the first evidence of what we now call MVs.

20 Over decades, the field advanced with MVs characterized and, to some extent, content was
21 determined, and evidence emerged that platelets were not the sole source of circulating MVs [22].
22 Presently, diverse MV populations have been identified from cell culture media, and biological fluids
23 including plasma [3,28], urine [29], saliva [28], and synovial fluid [3]. The current view is that upon
24 agonist stimulation most, if not all, cells release MVs carrying specific markers that enable cell origin
25 to be determined like endothelial (EMVs), PMVs, and red blood cell MVs (RBCMVs).

26 Currently, altered circulating MV concentrations have been associated with subclinical and clinical
27 conditions. For instance, increased circulating concentrations of EMVs occur with obesity [9,12], the
28 metabolic syndrome [30], those with coronary artery disease (CAD) [8], and type 2 diabetes mellitus
29 [10], to list a few. They have even provided prognostic information about cardiovascular mortality in
30 renal failure patients [31]. These conditions all exhibit vascular dysfunction, with increased EMVs
31 concentrations likely reflecting chronic vascular damage, as observed in patients with known poor
32 vascular outcomes such as CAD [8], and in acute stroke [32].

33 Accordingly, MVs are naturally produced and found in the circulation fluctuating within a physiological
34 range, with chronic alterations in MV concentrations identified as potential biomarkers of pathology.
35 Of particular relevance and based on *in vitro* and *in vivo* evidence, EMVs have received great attention
36 as a surrogate circulating marker of endothelial cell health [33–36].

37 2.1.1. Microvesicle formation and phenotype

38 Shedding of MVs from the cell membrane is initiated by complex events that lead to cytoskeleton
39 proteolysis, cell shrinkage, and eventually MV sprouting [37,38]. Our understanding of the overall
40 mechanisms governing MV formation and release derives mainly from platelets, since they were
41 among the first to be identified, and constitute the predominant MV phenotype in human blood.

42 Briefly, in the basal state the content of phospholipid cell membranes are asymmetric with negatively
43 charged phospholipids, such as phosphatidylserine (PS), mainly a part of the inner leaflet [37]. When

1 activated or undergoing apoptosis, a randomisation in the phospholipid plasma membrane content
2 occurs, increasing the appearance of PS on the outer membrane, which, coupled with cytoskeleton
3 reorganisation and membrane remodelling, culminates in the shedding of newly formed MVs that may
4 express PS on their surface [38]. Increases in intracellular Ca^{2+} concentration [37,39] with activation of
5 calcium-dependent proteins [39,40] is accepted as a general mechanism leading to these processes,
6 but calcium-independent pathways have also been identified in platelets [41]. Several stimuli that
7 bring about PMV release have been identified *in vitro* and include high shear stress, catecholamines,
8 adenosine diphosphate (ADP), and thrombin [42–44], but some agonists for platelet vesiculation can
9 downregulate the release of MVs from other sources such as endothelial cells [45]. An excellent
10 discussion of biological mechanisms related to MV formation has been published elsewhere [38].

11 Methods to quantify MVs in body fluids are still limited, but recent developments have facilitated
12 quick progression from time-consuming and semi-quantitative microscopy, to high throughput
13 quantitative MV analyses using enzyme-linked immunosorbent assays (ELISA), traditional flow
14 cytometry, imaging flow cytometry, resistive pulse sensing, and nanoparticle tracking analysis
15 [29,46,47]. Knowing that MVs express PS in the outer leaflet enabled PS-binding agents (e.g. annexin-
16 V staining) as general MV markers; however, PS exposure alone does not necessarily induce
17 vesiculation in platelets [39], and only a fraction of MVs released by unstimulated platelets express
18 sufficient PS to facilitate annexin-V binding [48]. Hence, more recently quantifying specific MV
19 populations has been based on the expression of cell-specific antigens associated with the MV
20 membrane, independent of annexin-V binding [16,34,48]. For example, PMVs can be identified by flow
21 cytometry as glycoprotein IIb (CD41) or glycoprotein Iba (CD42b) positive events, whereas E-selectin
22 (CD62E) can be used for EMV quantification, and CD45 is employed as a common maker for leukocyte-
23 derived MVs (Table 1).

24

25 [TABLE1]

26

27 Previous studies indicate that PMVs are the most common MV phenotype in plasma [2–4], although
28 reports of greater abundance of RBCMVs also exists [47]. Such discrepancies in the proportion of
29 RBCMVs may relate to differences in blood sampling and handling procedures, as phlebotomy
30 conditions (e.g. needle Gauge, tourniquet use, and anticoagulant choice) and sample processing
31 protocols (e.g. centrifugation steps, time, and speed) can ultimately affect final MV concentrations
32 [49]. Although a complete list of all antigens expressed by circulating MVs would be useful, it is beyond
33 our scope. However, the most frequently observed MVs antigens in the circulation are also expressed
34 by platelets [12,48], erythrocytes [35,50], monocytes [4,47], neutrophils [47,50], lymphocytes [47] and
35 endothelial cells [11,33,51] suggesting these are the origins of most MVs.

36 2.1.2. Overall microvesicle function

37 Circulating MVs likely regulate physiological processes, and the interaction of MVs with target cells
38 occurs through at least three mechanisms. First, MVs may bind to membrane receptors and mediate
39 cell modifications through signalling pathway activation. Next, cell-MV interactions at the plasma
40 membrane may deliver vesicular content upon fusion with the cell membrane [6,22]. Finally, MVs may
41 be internalized by certain cells, altering the function and structure or their recipient targets [52,53].
42 The latter may also serve as a MV clearance pathway if they are directed to lysosomes. As a result,
43 complex and sometimes contrasting responses arise from studies investigating isolated MV
44 populations.

1 Platelet-derived MVs are linked to the haemostatic system [22,23] with initial evidence supporting
2 their role in coagulation, even in the absence of intact platelets [26]. Although not all MVs are pro-
3 coagulant [48], a number of MV populations present in a variety of body fluids, including blood and
4 saliva, display thrombogenic functions [3,28,54] similar to PS-rich PMVs [48]. The pro-thrombotic
5 potential of many blood MVs is highly related to their surface PS and tissue factor (TF) content [28,48],
6 and the natural shedding of MVs seems like a necessary physiological process. For instance, Scott
7 syndrome, a rare haemorrhagic disorder, is linked to impaired translocation of PS to the outer leaflet
8 cell membranes and reduced MV shedding [38], which results in defective blood coagulation.

9 Even though production of MVs is continuous, only a fraction are PS and TF-rich [4,33,48], suggesting
10 roles beyond coagulation. As will be explained, blood MV functions vary widely. For example, specific
11 EMV populations can trigger inflammation [5] and oxidative stress [55] in recipient endothelial cells,
12 and lymphocyte-derived MVs can suppress angiogenesis through vascular endothelial growth factor
13 (VEGF) downregulation [56]. Furthermore circulating MVs isolated from myocardial infarction patients
14 [57] and pre-eclamptic women [58] lead to endothelial dysfunction *in situ*, suggesting that the link
15 between augmented MV concentrations in pathological conditions is beyond correlational.

16 Conversely, other MVs display opposite functions than those described above. Grasser and Schifferli
17 [59] have shown that granulocyte-derived MVs induce anti-inflammatory properties by
18 downregulating macrophage activity, and using an ischaemic-limb model Leroyer *et al.* [60]
19 demonstrated a 3.5-fold increase in pro-angiogenic MV concentration in mouse muscle homogenates.
20 These ischaemia-related MVs increased progenitor cell differentiation into an endothelial phenotype
21 *in vitro* and *in vivo*, and other studies have shown that endothelial cells undergoing migration release
22 pro-angiogenic EMVs [61]. PMVs may also promote activation of pro-angiogenic pathways, resulting
23 in endothelial cell proliferation, chemotaxis, and protection from apoptosis [7,62].

24 The identified function of MVs is constantly growing as it depends on a variety of factors, including
25 the MV population, content, and the recipient cell with which they are interacting, which illustrates
26 the complexity of these physiological interactions. More complex than local ischemia, physical
27 exertion imposes unique challenges upon many organ systems often introducing MVs into the human
28 circulation. As such the dynamics of MVs with exercise, and their potential relevance in exercise-
29 induced adaptation is an intriguingly understudied area.

30 3. Exercise and microvesicles

31 In the following sections a review of the current state of knowledge on microvesicles and exercise will
32 be presented, but as the reader will see the current understanding about the interaction between
33 MVs and exercise is still incomplete. As such, the subsequent sections aim to provide a starting point
34 and guidance for future research. Certainly, further experiments are warranted to answer specific
35 questions related to exercise and training variables (e.g. exercise volume, intensity, etc.), before more
36 definitive conclusions can be drawn.

37 3.1. Acute exercise

38 The phenomenon of microvesicle release with exercise has been appreciated only recently, with most
39 experiments in the field focused on acute aerobic exercise and published in the current decade (Table
40 2). The dynamics of this response to exercise, however, is MV population specific. For example,
41 circulating RBCMVs concentrations did not change in the single study that examined this MV phenotype
42 with exercise, but increases in the concentration of PMVs and MVs with polymorphonuclear
43 neutrophil antigens was observed following a series of maximal cycling protocols [50]. Since MVs are
44 physiologically active, their timed release may be involved in exercise responses.

1 Most studies have investigated the effect of performing an aerobic cycling session on subsequent
2 blood EMV concentrations. Although some authors observed an increase in certain EMV populations
3 after exercise in healthy individuals [4,16,51,63,64], most experiments report no change [4,13–
4 17,50,65–67], or even a decrease in EMV concentration [63,68], with one study finding an increase in
5 blood EMV in men, but not women [16]. The discrepancies in the EMV response between studies may
6 arise from a number of confounding factors. For example, exercise brings about many acute
7 physiological adjustments that may stimulate or blunt EMV shedding. Cytokines and vascular cyclic
8 strain stimulate the formation of MVs from cultured endothelial cells [33,54,69], whereas high
9 vascular shear stress reduces EMV formation *in vitro* [45]. Since each of these factors increases during
10 exercise, an antagonistic environment may be created where little additional endothelial vesiculation
11 will occur and this may account for inconsistent findings. In addition, plasma volume changes may also
12 confound finding as observed by Wilhelm and colleagues [17] who noted increased EMV
13 concentration with heat stress and strenuous exercise, but this response was abolished when plasma
14 volume shifts were taken into account. This suggests changes in circulating EMVs were mediated by
15 haemoconcentration rather than altered MV dynamics. Most studies, however, do not report blood
16 volume corrected EMV concentrations, making it difficult to establish whether EMV shedding actually
17 occurs.

18 In contrast to other MV phenotypes, PMVs are the most responsive to an acute exercise bout. Most
19 studies report an increase in these MVs after exercise [4,13–15,50,64,70], with increases up to 2 to 3-
20 fold from baseline, and several agonists known to stimulate PMV formation increase during exertion.
21 Plasma concentrations of ADP originating from exercising limbs in humans increase [71,72], and may
22 lead to platelet activation [72] since it is a PMV production agonist [44]. Similarly, sympathetic
23 activation during exercise can lead to blood noradrenaline spill-over that could stimulate platelet MV
24 formation, similar to *in vitro* effects [43]. Exercise also increases mean and anterograde vascular wall
25 shear rate (surrogate markers of shear stress) in conduit arteries of exercising [19,20], and even non-
26 exercising limbs with prolonged exercise [18,73]. Increased shear stimulates PMV release from *ex vivo*
27 platelets [7,42] and recently a positive correlation between vascular shear rate and plasma PMV
28 concentration was noted in exercising men [15], although a follow-up set of experiments revealed this
29 correlation to not necessarily represents causation [17].

30 When comparing these results, however, the reader must be aware of intrinsic methodological
31 limitations of the field. Distinct quantification techniques result in different absolute MV
32 concentrations. For instance, in the study by Maruyama et al. [70] PMVs were assessed by ELISA and
33 concentration differed by orders of magnitude from those using traditional flow cytometry, which in
34 turn are not necessarily comparable to studies employing imaging flow cytometry. Nevertheless, the
35 trend of response was the same between techniques (i.e. an increase in PMV with exercise),
36 suggesting that the same physiological phenomenon was being assessed. As shown in Table 2, the vast
37 majority of experiments employed flow cytometry as the main MV quantification technique, but it is
38 recognized that both pre-analytical and analytical procedures influence the MV content in a sample
39 [49]. Although efforts to standardize traditional flow cytometry protocols have been made [74,75]
40 there still lacks methodological agreement between experiments. Furthermore and according to the
41 iceberg concept [29,46], flow cytometers can not identify all MVs due to light scattering limitations of
42 small particles, and as a result swarm detection of small MVs confounds absolute concentration
43 measurements. These factors can complicate comparisons made between studies and should be taken
44 into consideration when interpreting individual results. Future experiments combining quantification
45 techniques such as flow cytometry and nanoparticle-tracking analysis will certainly be useful to
46 robustly quantify absolute MV concentrations after acute exercise bouts.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

[TABLE2]

3.1.1. Time-course of circulating microvesicle appearance with acute exercise

The dynamics of blood MV appearance depend on the vesicle source, and until the middle of the current decade our understanding of their time-course of release with exercise was limited to one study exploring EMVs [65], with the remaining experiments exploring post-exercise responses. More recently, plasma PMVs, but not EMV, were shown to increase and stabilise within 30 minutes of continuous cycling, with peak post-exercise values similar to those observed during exercise [15]. Blood PMVs are augmented from a few minutes [4,14,50,70] up to 1 [4,15,70] and 2 h [4,13,50] after exercise, and return towards baseline values thereafter. As such, the current literature indicates that the rise in blood PMVs is a short-lived phenomenon, with these vesicles reaching peak concentrations as early as 30 minutes after the onset of a training session, with values continuing to be elevated for a few minutes to hours into the post-exercise recovery period (Figure 1).

The time-course dynamics of other MV populations are less clear. A single experiment has reported an increase in the concentration of neutrophil-derived MV by the end of a ramp cycling protocol, with a tendency to return to baseline after 2 h of recovery [50], and a delayed rise in circulating monocyte MVs was observed after exercise in well-trained men [4], while it remained unchanged or undetectable in less fit individuals [4,50]. Studies that observed a rise in EMVs expressing markers of endothelial activation were limited to the post-exercise period [4,16,51,63]. From time-course experiments, a late EMV increase may be expected, with peak values between 45-90 min after exercise that returns towards baseline thereafter [4,51], but it is important to recall that several investigations failed to report increases in circulating EMV concentrations (Table 2).

[FIGURE1]

Of relevance is an EMV subpopulation carrying apoptotic antigens (e.g. CD31⁺ and negative for platelet markers). These MVs have been thought to reflect endothelial apoptosis and vascular damage by several authors [33–35]. One should note that a number of studies in the field of exercise have considered CD31⁺ EMVs as an apoptotic subpopulation irrespective of annexin-V staining, and this review will follow the same definition when referring to apoptotic EMVs.

The current literature regarding apoptotic EMVs is ambiguous and may depend on the assessment time-point and participant sex/population. For example, Durrer et al. [63] reported reduced concentrations of circulating apoptotic EMVs in the morning following continuous and interval exercise bouts in overweight/obese males, but not females, which was also evident 1 to 3 hours after exercise bouts in healthy men [68]. Conversely, plasma CD31⁺ EMV remained unchanged in healthy individuals when assessed 5 min after moderate intensity cycling [16]. On the other hand, Schwarz *et al.* [64] observed a 30% increase in plasma CD31⁺ MVs in men and women within 30 min of completing a marathon, but it is unclear whether the authors' flow cytometry gating strategy excluded platelet markers to ensure appropriate apoptotic EMV quantification. Hence, the limited evidence available suggests that short to moderate duration exercise sessions (i.e. from a few minutes up to 2 h) do not

1 lead to endothelial injury, as reflected by stable circulating apoptotic EMVs and, if anything, this MV
2 subpopulation may even decrease below baseline values after exercise.

3 The exact fate of apoptotic EMVs remains unknown but it may include uptake by endothelial cells in
4 the post-exercise period [68]. Internalization of MVs by native endothelial cells have been reported in
5 several tissues [53,76,77] and may serve as a clearance mechanism since haemodynamic forces push
6 circulating MVs towards the vascular endothelium. Previous experiments have identified that
7 endocytosis of MVs by such vascular cells occurs through the anchoring of MV surface PS to
8 endothelial integrins, in a process mediated by endothelial locus-1 glycoproteins [53]. Macrophages
9 have also been reported to remove MVs through phagocytic pathways regulated by the presence of
10 PS, CD31 and Immunoglobulin M on the MV surface [78], and clearance likely involves MV
11 opsonisation by complement components and subsequent uptake by phagocytes [79]. Furthermore,
12 systemically infused MVs localise in the spleen, lungs, and liver, which are all thought to be important
13 organs involved in MV clearance [78]. It is unknown, however, whether these mechanisms apply to
14 the exercise context. Nonetheless, a decrease in apoptotic EMVs compared to resting values has also
15 been reported 1 h into recovery in patients who undergo dobutamine-induced cardiac stress [80],
16 indicating rapid clearance of EMVs which further supports findings from physical stress (exercise)
17 challenges. An extensive review of MV release and clearance has been published by Ayers et al. [78].

18 3.1.2. Exercise type

19 Little information exists regarding exercise modalities and MV responses. Most studies have employed
20 continuous or interval cycling and only a single study involved (continuous, incremental) treadmill
21 exercise, where a rise in circulating PMVs occurred [70]. Cycling and running differ in terms of
22 contraction type and muscle mass engaged, which could be factors influencing the MV response.
23 Furthermore, footstrike during running has been proposed to induce mechanical damage of
24 erythrocytes and contribute to exercise-related haemolysis [81,82], which may influence RBCMVVs.
25 Although no direct comparison between exercise modalities has not been performed, some
26 conclusions regarding those variables can be drawn from published data. Using an adapted cycling
27 model, Rakobowchuk et al. [14] compared aerobic power matched concentric vs. eccentric exercise
28 and noted similar patterns of increased post-exercise plasma PMV and unaltered EMV concentrations,
29 indicating no influence of contraction type upon MV dynamics. Furthermore, MV release appears to
30 require the activation of only small quantities of skeletal muscle, as circulating PMV increase even
31 with small muscle mass exercise, like with incremental knee extensor exercise [17].

32 Resistance exercise, on the other hand, does not seem to affect blood EMV levels [67], the only MV
33 population studied to date. One could speculate that PMVs would be released under such conditions
34 since resistance exercise can acutely increase platelet activation as assessed by increased plasma β -
35 thromboglobulin concentrations [83,84] which highly correlates with plasma PMV ($r=0.95$) [85].

36 Hence, cycling and running may be expected to induce MV or at least PMV formation, indicating that
37 rhythmic endurance-like exercise can stimulate cell vesiculation, whereas data pertaining to resistance
38 exercise is scarce.

39 3.1.3. Exercise intensity

40 The relative intensity of an exercise session often governs acute physiological adjustments and
41 potentially impacts MV responses to physical exertion. If the exercise intensity domains
42 recommended by the American College of Sports Medicine are taken into consideration [86], most
43 studies of MVs and acute exercise have employed moderate to vigorous exercise stimuli, with
44 intensities generally $\geq 50\%$ of peak oxygen uptake. Post-exercise increase in blood PMV concentration

1 is a consistent response to exercise, even reported with cycling performed below the first ventilatory
2 threshold [14]. A between-study analysis, however, may lead to uncertainty about the influence of
3 exercise intensity on MV dynamics, since augmented appearance of certain blood MVs is reported
4 with near maximal incremental exercise [50,70], but variable responses have been noted after high
5 intensity interval exercise [51,63,66,68], or light to moderate intensity protocols [14,65].

6 Recent evidence comparing continuous cycling within different intensity domains has helped to clarify
7 this topic. Based on 60 min of moderate or vigorous continuous cycling performed by healthy men, it
8 has become apparent that relative intensity plays an important role in MVs dynamics. Specifically,
9 circulating PMVs increased from baseline during and after vigorous exercise (i.e. $\geq 64\%$ of maximal
10 oxygen uptake), whereas moderate intensity cycling (i.e. $\geq 46\%$ of maximal oxygen uptake) resulted
11 in a very modest non-significant rise [15]. The plasma EMVs concentrations were unaltered, no matter
12 the exercise intensity.

13 3.1.4. Exercise volume

14 The influence of exercise volume on acute MV responses lacks systematic evaluation at this point in
15 time. Studies designed to directly isolate exercise volume are needed, and comparison of exercise
16 protocols that induce changes in blood MV concentrations have been limited in terms of exercise
17 duration. Nevertheless, pronounced increases in circulating PMVs occurs even with small volume
18 exercise (e.g. a few minutes of incremental whole-body or isolated-limb exercise) [17,50,70], whereas
19 the longest duration studies (i.e. 4 h cycling performed below the anaerobic threshold or marathon
20 running) showed smaller increases or unchanged concentrations of circulating MVs [64,65]. One could
21 speculate that MVs may have increased and subsequently returned towards baseline concentrations
22 during the latter experiments, but the small number of sequential blood sampling time points limits
23 conclusions.

24 At this stage, data from Wilhelm et al. [15] helps shed further light on this topic as blood samples were
25 taken throughout 1 h of moderate and vigorous intensity cycling. Plasma PMVs increased from
26 baseline by 30 min of vigorous exercise and remained stable until the end of the 1 h protocol,
27 suggesting little influence of exercise volume upon PMV, since doubling the exercise duration did not
28 further increase plasma MV concentrations. Moreover, if simplistic energy expenditure estimations
29 are made considering a general O_2 caloric equivalent of 5 kcal/l O_2 (i.e. disregarding protocol-specific
30 respiratory exchange rate) it becomes apparent that exercise volume has little influence on circulating
31 MV appearance, since the concentration of plasma PMV was unchanged throughout the moderate
32 cycling session, even though the energy expenditure at the end of 1 h of this protocol (~ 450 kcal) was
33 greater than at 30 min of vigorous exercise (~ 330 kcal), when a rise in PMV was already evident.

34 Together, although still limited, the current body of evidence points toward a greater influence of
35 exercise intensity rather than volume upon MV release.

36 3.1.5. Physiological significance of exercise-derived microvesicles

37 The introduction of MVs into the circulation of exercising humans plays a regulatory role in
38 haemostatic control. Sossdorf et al. [4] isolated MVs exposing PS and reported an increase in MV-
39 related prothrombinase activity from post-exercise samples. Fibrin formation was greater in samples
40 from well-trained participants, reinforcing the procoagulant potential of MVs. As one exercises, both
41 the pro and anticoagulant systems may be activated [70,72,87], so the increased MV pro-coagulant
42 potential with acute exercise may play a natural role in fine-tuning the fibrinolytic and thrombotic
43 balance.

1 Exercise training traditionally improves endothelial function mainly through a vascular shear stress-
2 mediated mechanism [18–20], but circulatory factors are also involved [21]. Since circulating MVs,
3 particularly PMVs, can be acutely increased after an exercise session, speculation that these vesicles
4 are involved in vascular adaptation to training has emerged. Early experimental evidence that exercise
5 increases the interaction between endothelial cells and MVs was provided by Wahl *et al.* [68]. They
6 fluorescently labelled-MVs (PKH26 staining) *in vitro* and loaded these into serum samples obtained
7 from male athletes prior to and after exercise. Human coronary artery endothelial cells were then
8 incubated with the MV-rich sera which increased EMV (but not monocyte MV) uptake by cultured
9 endothelial cells exposed to post-exercise serum, providing some of the first evidence that blood
10 milieu alterations after exercise stimulate the uptake of selected MV populations from the circulation.
11 Moreover, a concomitant decrease in endothelial cell apoptosis was observed as assessed by, caspase-
12 3 activity but unfortunately the lack of a MV-poor serum control condition precludes that the
13 antiapoptotic effect was mediated by MVs specifically.

14 In subsequent experiments, MVs isolated from the plasma of exercising humans stimulated
15 angiogenesis of cultured endothelial cells when compared to MVs obtained from baseline resting
16 condition [15]. The angiogenic potential of exercise-derived MVs may stem from enhanced endothelial
17 proliferation and migratory capacity induced by these MVs compared to those obtained at baseline.
18 The mechanisms through which exercise-derived MVs induced alterations in endothelial phenotype
19 were not investigated but may relate to delivery of angiogenic growth factors. For example, VEGF is
20 considered a potent regulator of skeletal muscle capillary formation, and MVs shed by platelets *ex*
21 *vivo* increase endothelial wound-healing and angiogenesis partially through the delivery of VEGF to
22 endothelial cells [62]. Part of the pro-angiogenic effect of VEGF depends on activating the NO pathway
23 [88,89], and MVs from healthy individuals are a source of bloodborne eNOS [9,11]. Moreover, in
24 rodent limb ischaemia models MVs likely facilitate compensatory angiogenesis through the
25 stimulation of progenitor cell differentiation into an endothelial phenotype [60]. Together, these are
26 potential, yet speculative, mechanistic alternatives through which exercise-derived MVs may bring
27 about endothelial adaptation. Although exercise-derived PMVs remain logical candidates for their
28 physiological effects, the specific MV population responsible for the pro-angiogenic and proliferative
29 endothelial stimulation, as well as the exact biochemical pathways through which exercise-derived
30 MVs stimulate endothelial cells still needs to be determined.

31 As postulated with different extracellular vesicle populations [24], it is possible exercise-derived MVs
32 transport “exerkines” and promote systemic adaptations. An important recent study by Whitham *et*
33 *al.* [90] suggests that acute exercise might modify the cargo of sampled extracellular vesicles, which
34 contained a fraction of small MVs, as hundreds of proteins transported in extracellular vesicles were
35 altered in the circulation after a cycling bout. Beyond the vascular system, they further demonstrated
36 that systemically infused exercise-derived extracellular vesicles accumulated in the liver of recipient
37 mice, indicating a pathway for communication with non-exercising tissues during exercise, which
38 could include the lungs, kidneys, liver, and brain [53,76,77]. Hence, all these tissues could be prone to
39 phenotypical modifications mediated by the exercise-derived MVs. As such, given the systemic nature
40 of circulating MVs, one can speculate that their timed release with exercise can positively influence
41 several tissues and organs by altering remote cell function and morphology (Figure 2). Future studies
42 ought to further unravel the relevance of these MVs in vascular and systemic adaptations to exercise
43 *in vivo*.

44

45 [FIGURE2]

1

2 3.2. Physical activity and exercise training

3 Observational studies reporting altered circulating MVs concentrations in patients with vascular and
4 metabolic dysfunctions, as well as established diseases suggest that MVs respond to chronic
5 physiological challenges. Moreover, the ability of MVs to carry bioactive makers and participate in
6 horizontal gene transfer at remote sites suggests involvement in the pathophysiology. Accordingly,
7 environmental factors and chronic lifestyle modifications, such as exercise training, may influence the
8 basal concentration of circulating MVs.

9 Early evidence of the influence of physical activity and exercise upon the blood MV profile comes from
10 small, yet well conducted, bedrest and restricted physical activity experiments. Navasiolava et al. [35]
11 limited physical activity of 8 healthy men to a minimum for 7 days, and observed an early elevation in
12 apoptotic EMVs by the third day of inactivity, with reduced microvascular vasodilatory function that
13 was likely prostaglandin-related. In agreement with this previous study, reducing daily physical activity
14 levels of recreationally active men by ~50% (from > 10,000 to < 5,000 steps/day) for 5 consecutive
15 days impaired popliteal artery endothelial function augmented circulating concentrations of apoptotic
16 EMVs [36]. Since inactivity-mediated vascular dysfunction seems to reflect increases in EMVs, a
17 decrease in basal EMVs could be expected with increased physical activity through, for example,
18 exercise training.

19 Only a few studies have investigated changes in blood MVs after a training period. Babbitt et al. [91]
20 were among the first and reported a near 50% decreased in plasma CD62⁺ EMVs (a MV subset
21 theoretically derived from activated endothelial cells) of middle-aged and older African-Americans
22 after 6 months of light to moderate intensity endurance exercise, which was accompanied by an
23 improvement in blood inflammatory markers and brachial artery flow-mediated dilation. A subgroup
24 analysis also revealed a decrease in apoptotic EMVs [92], suggesting reduced vascular damage and
25 vesiculation at rest after the training period. Unfortunately, the lack of a control group hinders
26 categorical conclusions that the reported effects were exclusive to the training program itself.
27 However, these results importantly indicate that chronic improvements in endothelial function are
28 reflected by a reduction in blood EMV concentrations in a population at elevated risk for
29 cardiovascular events [93–95].

30 In light of current evidence, we can only hypothesize about the chronic effect of exercise training on
31 resting PMV adaptations. As platelet hyperreactivity to ADP has been associated with long-term
32 cardiovascular complications in some studies [96,97], and considering that: (1) baseline PMV
33 concentration is increased in patients with established or at risk for cardiovascular diseases [8,10,12];
34 and (2) exercise training can reduce markers of platelet activation at rest and diminish their sensitivity
35 to activation agonists [87]; it appears reasonable that circulating PMV content would drop after a
36 training programme in selected patient populations. However, it is important to stress that the impact
37 of exercise training on resting platelet activation is not unequivocal [87]. Alternatively, the
38 concentration of PMVs might remain unchanged while their content could be modified towards a less
39 thrombogenic and more vasoprotective phenotype, but again this idea remains merely hypothetical.

40 4. Further directions in physiology and medicine

41 Even though alterations in certain circulating MVs concentrations with exercise seems like a normal
42 response in healthy adults, little is known about the influence of acute and chronic exercise upon the
43 cargo of MVs and their dynamics in specific patient populations. For example, Guiraud *et al.* [66]
44 observed no change in circulating PMV and EMV amongst stable CAD patients after an exercise

1 challenge, whereas subsequent work by Augustine *et al.* [80] revealed that circulating MVs
2 concentrations increased in patients with normal coronary arteries in response to a
3 pharmacologically-mediated cardiac stress, but MVs concentrations were unaltered in patients likely
4 to have further cardiovascular complications. These findings suggest that an increase in selected MV
5 populations with acute stress is an important physiological process that appears blunted in patients
6 at risk. These MV dynamics to stress tests could even be useful in predicting future cardiovascular
7 events in certain populations. Monitoring changes in circulating MVs concentrations prior and after
8 standard tests (such as exercise) in healthy controls and populations with cardiovascular risk factors
9 could provide further insight into the MV dynamics and could progress to the use of exercise-derived
10 MVs as biomarkers in patient stratification.

11 As discussed, our understanding of the MVs responses to exercise and training is far from complete,
12 and this field will benefit from acute experiments exploring the influence of exercise variables (e.g.
13 volume, intensity, and type of exercise) and their chronic influence to optimise training adaptations.
14 Greater standardization on MV quantification protocols within the field of exercise is also warranted.
15 Moreover, beyond changes in the MV profile, the content of MVs is affected by one's physiological
16 state [9,11], and it is likely that exercise-released MV differ in composition compared to those
17 produced under basal or pathological conditions. It is also tempting to speculate that chronic exercise
18 may impact not just the circulating MV profile, but its cargo as well. Future studies that examine
19 proteomic, lipidomic, transcriptomic, and metabolomic profiles with more appropriate species
20 isolation and quantification techniques will certainly shed light upon the composition of MVs.

21 The exciting findings that extracellular vesicles can mediate cell-to-cell signalling through horizontal
22 transfer of mRNA and miRNA [98,99] is of particular interest, as acute and chronic exercise can alter
23 the concentration of circulating miRNA [100], and MVs might act as a transport vehicle of
24 transcription-controlling factors and alter the function of remote cells. Interestingly, there is evidence
25 linking PMVs to adhesion and differentiation of early outgrowth endothelial cells *in vitro* and
26 amplification of early outgrowth endothelial cell reendothelization in murine vascular injury models
27 [6]. This suggests the exercise-induced PMV release phenomenon may improve vascular repair and
28 function by enhancing differentiation of circulating endothelial progenitor cells. Integration of *in vitro*
29 and *in vivo* studies exploring these ideas may help unravel the mechanisms of exercise-mediated
30 vascular adaptations.

31 5. Conclusion

32 Cell-derived MVs have received increased attention in the scientific community due to their potential
33 to serve as biomarkers of intracellular events and their innate biological activities. These extracellular
34 vesicles had been initially thought of as simple by-products of pathological disorders, and
35 subsequently believed to play a role in maladaptation, but more recent evidence has also shown that
36 MVs are not necessarily harmful, and actually necessary for proper physiological function. Exercise is
37 a powerful factor affecting circulating MV dynamics. Available evidence indicates that the blood MV
38 response to acute and chronic training may resemble the cytokine adjustments to exercise: that is,
39 transient physical exertion may lead to a timed release of MVs, in particular PMVs, which are likely to
40 be involved in acute and subacute exercise adjustments, as is the case with the cytokine response to
41 a single bout of exercise. However, in the long term, and in analogy to pro-inflammatory cytokines,
42 exercise training may decrease resting levels EMVs, which may be involved and reflect reduced
43 vascular injury. Last but not least, a body of evidence indicates that these tiny vesicles play a role in
44 the complex haemostatic control to exercise and are potential novel mediators of endothelial
45 adaptations to training. The upcoming decades will certainly benefit from research investigating the

1 precise dynamics of MVs in response to specific exercise variables, and will unravel their relevance in
2 human physiology.

3 Compliance with Ethical Standards

4 Conflict of Interest

5 Eurico N. Wilhelm, Laurent Mourot and Mark Rakobowchuk declare that they have no conflict of
6 interest.

7 Funding

8 The research was funded by grants from the French Ministry of National Education, of Research and
9 of Technology (EA3920), from Tomsk Polytechnic University Competitiveness Enhancement Program
10 grant, Project № ВИУ-ИСГТ-108/2017 - TPU CEP-HSTI-108/2017 and a Natural Science and Engineering
11 Research Council of Canada Discovery Grant to M.R.

12 E.N.W. is supported by the Brazilian Education Ministry Foundation CAPES (Postdoctoral Fellowship).

13 References

14 1. Garraud O, Cognasse F. Are platelets cells? and if yes, are they immune cells? *Front. Immunol.* 2015;6:1–8.

15 2. Berckmans RJ, Nieuwland R, Böing AN, Romijn FP, Hack CE, Sturk A. Cell-derived microparticles circulate in
16 healthy humans and support low grade thrombin generation. *Thromb. Haemost.* 2001;85:639–46.

17 3. Berckmans RJ, Nieuwland R, Tak PP, Böing AN, Romijn FP, Kraan MC, et al. Cell-derived microparticles in
18 synovial fluid from inflamed arthritic joints support coagulation exclusively via a factor VII-dependent
19 mechanism. *Arthritis Rheum.* 2002;46:2857–66.

20 4. Sossdorf M, Otto GP, Claus RA, Gabriel HHW, Lösche W. Cell-derived microparticles promote coagulation after
21 moderate exercise. *Med. Sci. Sport. Exerc.* 2011;43:1169–76.

22 5. Curtis AM, Wilkinson PF, Gui M, Gales TL, Hu E, Edelberg JM. p38 mitogen-activated protein kinase targets the
23 production of proinflammatory endothelial microparticles. *J. Thromb. Haemost.* 2009;7:701–9.

24 6. Mause SF, Ritzel E, Liehn EA, Hristov M, Bidzhekov K, Muller-Newen G, et al. Platelet microparticles enhance
25 the vasoregenerative potential of angiogenic early outgrowth cells after vascular injury. *Circulation.*
26 2010;122:495–506.

27 7. Kim HK, Song KS, Chung J, Lee KR, Lee S. Platelet microparticles induce angiogenesis in vitro. *Br. J. Haematol.*
28 2004;124:376–84.

29 8. Bernal-Mizrachi L, Jy W, Jimenez JJ, Pastor J, Mauro LM, Horstman LL, et al. High levels of circulating
30 endothelial microparticles in patients with acute coronary syndromes. *Am. Heart J.* 2003;145:962–70.

31 9. Dimassi S, Chahed K, Boumiza S, Canault M, Tabka Z, Laurant P, et al. Role of eNOS- and NOX-containing
32 microparticles in endothelial dysfunction in patients with obesity. *Obesity.* 2016;24:1305–12.

33 10. Feng B, Chen Y, Luo Y, Chen M, Li X, Ni Y. Circulating level of microparticles and their correlation with arterial
34 elasticity and endothelium-dependent dilation in patients with type 2 diabetes mellitus. *Atherosclerosis.*
35 2010;208:264–9.

36 11. Horn P, Cortese-krott MM, Amabile N, Hundsöfer C, Kröncke K-D, Kelm M, et al. Circulating microparticles
37 carry a functional endothelial nitric oxide that is decreased in patients with endothelial dysfunction. *J. Am. Heart*
38 *Assoc.* 2012;2:1–12.

39 12. Esposito K, Ciotola M, Schisano B, Gualdiero R, Sardelli L, Misso L, et al. Endothelial microparticles correlate
40 with endothelial dysfunction in obese women. *J. Clin. Endocrinol. Metab.* 2006;91:3676–9.

41 13. Sossdorf M, Otto GP, Claus R a, Gabriel HH, Lösche W. Release of pro-coagulant microparticles after

- 1 moderate endurance exercise. *Platelets*. 2010;21:389–91.
- 2 14. Rakobowchuk M, Ritter O, Wilhelm EN, Isacco L, Bouhaddi M, Degano B, et al. Divergent endothelial function
3 but similar platelet microvesicle responses following eccentric and concentric cycling at a similar aerobic power
4 output. *J. Appl. Physiol.* 2017;122:1031–9.
- 5 15. Wilhelm EN, González-Alonso J, Parris C, Rakobowchuk M. Exercise intensity modulates the appearance of
6 circulating microvesicles with proangiogenic potential upon endothelial cells. *Am. J. Physiol. - Hear. Circ. Physiol.*
7 2016;311:H1297–310.
- 8 16. Lansford KA, Shill DD, Dicks AB, Marshburn MP, Southern WM, Jenkins NT. Effect of acute exercise on
9 circulating angiogenic cell and microparticle populations. *Exp. Physiol.* 2016;101:155–67.
- 10 17. Wilhelm EN, González-Alonso J, Chiesa ST, Trangmar SJ, Kalsi KK, Rakobowchuk M. Whole-body heat stress
11 and exercise stimulate the appearance of platelet microvesicles in plasma with limited influence of vascular
12 shear stress. *Physiol. Rep.* 2017;5:e13496.
- 13 18. Birk GK, Dawson EA, Atkinson C, Haynes A, Cable NT, Thijssen DHJ, et al. Brachial artery adaptation to lower
14 limb exercise training: role of shear stress. *J. Appl. Physiol.* 2012;112:1653–8.
- 15 19. Tinken TM, Thijssen DHJ, Hopkins N, Dawson EA, Cable NT, Green DJ. Shear Stress Mediates Endothelial
16 Adaptations to Exercise Training in Humans. *Hypertension*. 2010;55:312–8.
- 17 20. Credeur DP, Hollis BC, Welsch MA. Effects of handgrip training with venous restriction on brachial artery
18 vasodilation. *Med. Sci. Sports Exerc.* 2010;42:1296–302.
- 19 21. Padilla J, Simmons GH, Bender SB, Arce-Esquivel AA, Whyte JJ, Laughlin MH. Vascular effects of exercise:
20 endothelial adaptations beyond active muscle beds. *Physiology*. 2011;26:132–45.
- 21 22. Hargett LA, Bauer NN. On the origin of microparticles: From “platelet dust” to mediators of intercellular
22 communication. *Pulm. Circ.* 2013;3:329–40.
- 23 23. Shantsila E, Kamphuisen PW, Lip GYH. Circulating microparticles in cardiovascular disease: implications for
24 atherogenesis and atherothrombosis. *J. Thromb. Haemost.* 2010;8:2358–68.
- 25 24. Safdar A, Saleem A, Tarnopolsky MA. The potential of endurance exercise-derived exosomes to treat
26 metabolic diseases. *Nat. Rev. Endocrinol.* 2016;12:504–17.
- 27 25. Théry C, Zitvogel L, Amigorena S. Exosomes: composition, biogenesis and function. *Nat. Rev. Immunol.*
28 2002;2:569–79.
- 29 26. Wolf P. The nature and significance of platelet products in human plasma. *Br. J. Haematol.* 1967;13:269–88.
- 30 27. Webber AJ, Johnson SA. Platelet participation in blood coagulation aspects of hemostasis. *Am. J. Pathol.*
31 1970;60:19–42.
- 32 28. Berckmans RJ, Sturk A, van Tienen LM, Schaap MCL, Nieuwland R. Cell-derived vesicles exposing coagulant
33 tissue factor in saliva. *Blood*. 2011;117:3172–80.
- 34 29. van der Pol E, van Gemert MJC, Sturk A, Nieuwland R, van Leeuwen TG. Single vs. swarm detection of
35 microparticles and exosomes by flow cytometry. *J. Thromb. Haemost.* 2012;10:919–30.
- 36 30. Amabile N, Cheng S, Renard JM, Larson MG, Ghorbani A, McCabe E, et al. Association of circulating
37 endothelial microparticles with cardiometabolic risk factors in the Framingham Heart Study. *Eur. Heart J.*
38 2014;35:2972–9.
- 39 31. Amabile N, Guerin AP, Tedgui A, Boulanger CM, London GM. Predictive value of circulating endothelial
40 microparticles for cardiovascular mortality in end-stage renal failure: a pilot study. *Nephrol. Dial. Transplant.*
41 2012;27:1873–80.
- 42 32. Simak J, Gelderman MP, Yu H, Wright V, Baird AE. Circulating endothelial microparticles in acute ischemic
43 stroke: A link to severity, lesion volume and outcome. *J. Thromb. Haemost.* 2006;4:1296–302.
- 44 33. Jimenez JJ, Jy W, Mauro LM, Soderland C, Horstman LL, Ahn YS. Endothelial cells release phenotypically and

- 1 quantitatively distinct microparticles in activation and apoptosis. *Thromb. Res.* 2003;109:175–80.
- 2 34. Jenkins NT, Padilla J, Boyle LJ, Credeur DP, Laughlin MH, Fadel PJ. Disturbed blood flow acutely induces
3 activation and apoptosis of the human vascular endothelium. *Hypertens. (Dallas, Tex. 1979).* 2013;61:615–21.
- 4 35. Navasolava NM, Dignat-George F, Sabatier F, Larina IM, Demiot C, Fortrat J-O, et al. Enforced physical
5 inactivity increases endothelial microparticle levels in healthy volunteers. *Am. J. Physiol. - Hear. Circ. Physiol.*
6 2010;299:H248–56.
- 7 36. Boyle LJ, Credeur DP, Jenkins NT, Padilla J, Leidy HJ, Thyfault JP, et al. Impact of reduced daily physical activity
8 on conduit artery flow-mediated dilation and circulating endothelial microparticles. *J. Appl. Physiol.*
9 2013;115:1519–25.
- 10 37. Chang CP, Zhao J, Wiedmer T, Sims PJ. Contribution of platelet microparticle formation and granule secretion
11 to the transmembrane migration of phosphatidylserine. *J. Biol. Chem.* 1993;268:7171–8.
- 12 38. Morel O, Jesel L, Freyssinet JM, Toti F. Cellular mechanisms underlying the formation of circulating
13 microparticles. *Arterioscler. Thromb. Vasc. Biol.* 2011;31:15–26.
- 14 39. Dachary-Prigent J, Pasquet JM, Freyssinet JM, Nurden AT. Calcium involvement in aminophospholipid
15 exposure and microparticle formation during platelet activation: a study using Ca²⁺-ATPase inhibitors.
16 *Biochemistry.* 1995;34:11625–34.
- 17 40. Fox JE, Austin CD, Reynolds CC, Steffen PK. Evidence that agonist-induced activation of calpain causes the
18 shedding of procoagulant-containing microvesicles from the membrane of aggregating platelets. *J. Biol. Chem.*
19 1991;266:13289–95.
- 20 41. Cauwenberghs S, Feijge MAH, Harper AGS, Sage SO, Curvers J, Heemskerk JWM. Shedding of procoagulant
21 microparticles from unstimulated platelets by integrin-mediated destabilization of actin cytoskeleton. *FEBS Lett.*
22 2006;580:5313–20.
- 23 42. Miyazaki Y, Nomura S, Miyake T, Kagawa H, Kitada C, Taniguchi H, et al. High shear stress can initiate both
24 platelet aggregation and shedding of procoagulant containing microparticles. *Blood.* 1996;88:3456–64.
- 25 43. Tschuor C, Asmis LM, Lenzlinger PM, Tanner M, Härter L, Keel M, et al. In vitro norepinephrine significantly
26 activates isolated platelets from healthy volunteers and critically ill patients following severe traumatic brain
27 injury. *Crit. Care.* 2008;12:R80.
- 28 44. Horstman LL, Ahn YS. Platelet microparticles: a wide-angle perspective. *Crit. Rev. Oncol. Hematol.*
29 1999;30:111–42.
- 30 45. Vion A-C, Ramkhalawon B, Loyer X, Chironi G, Devue C, Loirand G, et al. Shear stress regulates endothelial
31 microparticle release. *Circ. Res.* 2013;112:1323–33.
- 32 46. Harrison P, Gardiner C. Invisible vesicles swarm within the iceberg. *J. Thromb. Haemost.* 2012;10:916–8.
- 33 47. Headland SE, Jones HR, D'Sa AS V., Perretti M, Norling L V. Cutting-edge analysis of extracellular
34 microparticles using ImageStream(X) imaging flow cytometry. *Sci. Rep.* 2014;4:5237.
- 35 48. Connor DE, Exner T, Ma DDF, Joseph JE. The majority of circulating platelet-derived microparticles fail to bind
36 annexin V , lack phospholipid-dependent procoagulant activity and demonstrate greater expression of
37 glycoprotein Ib. *Thromb. Haemost.* 2010;103:1044–52.
- 38 49. Yuana Y, Bertina R, Osanto S. Pre-analytical and analytical issues in the analysis of blood microparticles.
39 *Thromb. Haemost.* 2011;105:396–408.
- 40 50. Chaar V, Romana M, Tripette J, Broquere C, Huisse MG, Hue O, et al. Effect of strenuous physical exercise on
41 circulating cell-derived microparticles. *Clin. Hemorheol. Microcirc.* 2011;47:15–25.
- 42 51. Kirk RJ, Peart DJ, Madden LA, Vince R V. Repeated supra-maximal sprint cycling with and without sodium
43 bicarbonate supplementation induces endothelial microparticle release. *Eur. J. Sport Sci.* 2014;14:345–52.
- 44 52. Terrisse AD, Puech N, Allart S, Gourdy P, Xuereb JM, Payrastre B, et al. Internalization of microparticles by

- 1 endothelial cells promotes platelet/endothelial cell interaction under flow. *J. Thromb. Haemost.* 2010;8:2810–
2 9.
- 3 53. Dasgupta SK, Le A, Chavakis T, Rumbaut RE, Thiagarajan P. Developmental endothelial locus-1 (Del-1)
4 mediates clearance of Platelet microparticles by the endothelium. *Circulation.* 2012;125:1664–72.
- 5 54. Abid Hussein MN, Böing AN, Biró É, Hoek FJ, Vogel GMT, Meuleman DG, et al. Phospholipid composition of
6 in vitro endothelial microparticles and their in vivo thrombogenic properties. *Thromb. Res.* 2008;121:865–71.
- 7 55. Brodsky S V, Zhang F, Nasjletti A, Goligorsky MS. Endothelium-derived microparticles impair endothelial
8 function in vitro. *Am. J. Physiol. - Hear. Circ. Physiol.* 2004;286:H1910-5.
- 9 56. Yang C, Mwaikambo BR, Zhu T, Gagnon C, Lafleur J, Seshadri S, et al. Lymphocytic microparticles inhibit
10 angiogenesis by stimulating oxidative stress and negatively regulating VEGF-induced pathways. *Am. J. Physiol. -
11 Regul. Integr. Comp. Physiol.* 2008;5:467–76.
- 12 57. Boulanger CM, Scoazec A, Ebrahimian T, Henry P, Mathieu E, Tedgui A, et al. Circulating microparticles from
13 patients with myocardial infarction cause endothelial dysfunction. *Circulation.* 2001;104:2649–52.
- 14 58. VanWijk MJ, Svedas E, Boer K, Nieuwland R, VanBavel E, Kublickiene KR. Isolated microparticles, but not
15 whole plasma, from women with preeclampsia impair endothelium-dependent relaxation in isolated myometrial
16 arteries from healthy pregnant women. *Am. J. Obstet. Gynecol.* 2002;187:1686–93.
- 17 59. Gasser O, Schifferli JA. Activated polymorphonuclear neutrophils disseminate anti-inflammatory
18 microparticles by ectocytosis. *Blood.* 2004;104:2543–8.
- 19 60. Leroyer AS, Ebrahimian TG, Cochain C, Récalde A, Blanc-Brude O, Mees B, et al. Microparticles from ischemic
20 muscle promotes postnatal vasculogenesis. *Circulation.* 2009;119:2808–17.
- 21 61. Arderiu G, Peña E, Badimon L. Angiogenic Microvascular Endothelial Cells Release Microparticles Rich in
22 Tissue Factor That Promotes Postischemic Collateral Vessel Formation. *Arterioscler. Thromb. Vasc. Biol.*
23 2015;35:348–57.
- 24 62. Brill A, Dashevsky O, Rivo J, Gozal Y, Varon D. Platelet-derived microparticles induce angiogenesis and
25 stimulate post-ischemic revascularization. *Cardiovasc. Res.* 2005;67:30–8.
- 26 63. Durrer C, Robinson E, Wan Z, Martinez N, Hummel ML, Jenkins NT, et al. Differential impact of acute high-
27 intensity exercise on circulating endothelial microparticles and insulin resistance between overweight/obese
28 males and females. *PLoS One.* 2015;10:e0115860.
- 29 64. Schwarz V, Düsing P, Liman T, Werner C, Herm J, Bachelier K, et al. Marathon running increases circulating
30 endothelial- and thrombocyte-derived microparticles. *Eur. J. Prev. Cardiol.* 2017;1–8.
- 31 65. Mobius-Winkler S, Hilberg T, Menzel K, Golla E, Burman A, Schuler G, et al. Time-dependent mobilization of
32 circulating progenitor cells during strenuous exercise in healthy individuals. *J. Appl. Physiol.* 2009;107:1943–50.
- 33 66. Guiraud T, Gayda M, Juneau M, Bosquet L, Meyer P, Théberge-Julien G, et al. A single bout of high-intensity
34 interval exercise does not increase endothelial or platelet microparticles in stable, physically fit men with
35 coronary heart disease. *Can. J. Cardiol.* 2013;29:1285–91.
- 36 67. Ross MD, Wekesa AL, Phelan JP, Harrison M. Resistance exercise increases endothelial progenitor cells and
37 angiogenic factors. *Med. Sci. Sport. Exerc.* 2014;46:16–23.
- 38 68. Wahl P, Jansen F, Achtzehn S, Schmitz T, Bloch W, Mester J, et al. Effects of high intensity training and high
39 volume training on endothelial microparticles and angiogenic growth factors. Madeddu P, editor. *PLoS One.*
40 2014;9:e96024.
- 41 69. Vion AC, Birukova A a, Boulanger CM, Birukov KG. Mechanical forces stimulate endothelial microparticle
42 generation via caspase-dependent apoptosis-independent mechanism. *Pulm. Circ.* 2013;3:95–9.
- 43 70. Maruyama K, Kadono T, Morishita E. Plasma levels of platelet-derived microparticles are increased after
44 anaerobic exercise in healthy subjects. *J. Atheroscler. Thromb.* 2012;19:585–7.

- 1 71. Gonzalez-Alonso J, Olsen DB, Saltin B. Erythrocyte and the regulation of human skeletal muscle blood flow
2 and oxygen delivery: role of circulating ATP. *Circ. Res.* 2002;91:1046–55.
- 3 72. Yegutkin GG, Samburski SS, Mortensen SP, Jalkanen S, González-Alonso J. Intravascular ADP and soluble
4 nucleotidases contribute to acute prothrombotic state during vigorous exercise in humans. *J. Physiol.*
5 2007;2:553–64.
- 6 73. Padilla J, Simmons GH, Vianna LC, Davis MJ, Laughlin MH, Fadel PJ. Brachial artery vasodilatation during
7 prolonged lower limb exercise: role of shear rate. *Exp. Physiol.* 2011;96:1019–27.
- 8 74. Lacroix R, Judicone C, Mooberry M, Boucekine M, Dignat-George F. Standardization of pre-analytical
9 variables in plasma microparticle determination: results of the International Society on Thrombosis and
10 Haemostasis SSC Collaborative workshop. *J. Thromb. Haemost.* 2013;11:1190–1193.
- 11 75. Cointe S, Judicone C, Robert S, Mooberry MJ, Poncelet P, Wauben M, et al. Standardization of microparticle
12 enumeration across different flow cytometry platforms: results of a multicenter collaborative workshop. *J.*
13 *Thromb. Haemost.* 2017;15:187–93.
- 14 76. Cantaluppi V, Gatti S, Medica D, Figliolini F, Bruno S, Deregibus MC, et al. Microvesicles derived from
15 endothelial progenitor cells protect the kidney from ischemia – reperfusion injury by microRNA-dependent
16 reprogramming of resident renal cells. *Kidney Int. Elsevier Masson SAS;* 2012;82:412–27.
- 17 77. Faille D, El-assaad F, Mitchell AJ, Alessi M, Chimini G, Fusai T, et al. Endocytosis and intracellular processing
18 of platelet microparticles by brain endothelial cells. *J. Cell. Mol. Med.* 2012;16:1731–8.
- 19 78. Ayers L, Nieuwland R, Kohler M, Kraenkel N, Ferry B, Leeson P. Dynamic microvesicle release and clearance
20 within the cardiovascular system: triggers and mechanisms. *Clin. Sci.* 2015;129:915–31.
- 21 79. Flaumenhaft R. Formation and fate of platelet microparticles. *Blood Cells, Mol. Dis.* 2006;36:182–7.
- 22 80. Augustine D, Ayers L V., Lima E, Newton L, Lewandowski AJ, Davis EF, et al. Dynamic release and clearance
23 of circulating microparticles during cardiac stress. *Circ. Res.* 2014;114:109–13.
- 24 81. Miller BJ, Pate RR, Burgess W. Foot impact force and intravascular hemolysis during distance running. *Int. J.*
25 *Sports Med.* 1988;9:56–60.
- 26 82. Telford RD, Sly GJ, Hahn AG, Cunningham RB, Bryant C, Smith JA. Footstrike is the major cause of hemolysis
27 during running. *J. Appl. Physiol.* 2003;94:38–42.
- 28 83. Ahmadizad S, El-sayed MS, MacLaren DP. Effects of time of day and acute resistance exercise on platelet
29 activation and function. *Clin. Hemorheol. Microcirc.* 2010;45:391–9.
- 30 84. Ahmadizad S, El-sayed MS. The Effects of graded resistance exercise on platelet aggregation and activation.
31 *Med. Sci. Sport. Exerc.* 2003;35:1026–32.
- 32 85. Kim HK, Song KS, Lee ES, Lee YJ, Park YS, Lee KR, et al. Optimized flow cytometric assay for the measurement
33 of platelet microparticles in plasma : pre-analytic and analytic considerations. *Blood Coagul. Fibrinolysis.*
34 2002;13:393–7.
- 35 86. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee I-M, et al. American College of Sports
36 Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory,
37 musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med.*
38 *Sci. Sports Exerc.* 2011;43:1334–59.
- 39 87. El-Sayed MS, Sale C, Jones PG, Chester M. Blood hemostasis in exercise and training. *Med. Sci. Sports Exerc.*
40 2000;32:918–25.
- 41 88. Ziche M, Morbidelli L, Choudhuri R, Zhang HT, Donnini S, Granger HJ, et al. Nitric oxide synthase lies
42 downstream from vascular endothelial growth factor-induced but not basic fibroblast growth factor-induced
43 angiogenesis. *J Clin Invest.* 1997;99:2625–34.
- 44 89. Fukumura D, Gohongi T, Kadambi A, Izumi Y, Ang J, Yun C-O, et al. Predominant role of endothelial nitric
45 oxide synthase in vascular endothelial growth factor-induced angiogenesis and vascular permeability. *Proc. Natl.*

- 1 Acad. Sci. 2001;98:2604–9.
- 2 90. Whitham M, Parker BL, Friedrichsen M, Hingst JR, Hjorth M, Hughes WE, et al. Extracellular vesicles provide
3 a means for tissue crosstalk during exercise. *Cell Metab. Elsevier*; 2018;27:237–251.e4.
- 4 91. Babbitt DM, Diaz KM, Fearheller DL, Sturgeon KM, Perkins AM, Veerabhadrapa P, et al. Endothelial
5 activation microparticles and inflammation status improve with exercise training in african Americans. *Int. J.*
6 *Hypertens.* 2013;2013:1–8.
- 7 92. Fearheller DL, Diaz KM, Kashem MA, Thakkar SR, Veerabhadrapa P, Sturgeon KM, et al. Effects of moderate
8 aerobic exercise training on vascular health and blood pressure in African Americans. *J. Clin. Hypertens.*
9 2014;16:504–10.
- 10 93. Duck MM, Hoffman RP. Impaired endothelial function in healthy African-American adolescents compared
11 with Caucasians. *J. Pediatr.* 2007;150:400–6.
- 12 94. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart disease and stroke
13 statistics—2016 Update. *Circulation.* 2016;133:e38–360.
- 14 95. Heffernan KS, Jae SY, Wilund KR, Woods JA, Fernhall B. Racial differences in central blood pressure and
15 vascular function in young men. *Am. J. Physiol. - Hear. Circ. Physiol.* 2008;61820:2380–7.
- 16 96. Puurunen MK, Hwang S, Larson MG, Vasan RS, O’Donnell CJ, Tofler G, et al. ADP platelet hyperreactivity
17 predicts cardiovascular disease in the FHS (Framingham Heart Study). *J. Am. Heart Assoc.* 2018;7:e008522.
- 18 97. Thaulow E, Erikssen J, Sandvik L, Stormorken H, Cohn PF. Blood platelet count and function are related to
19 total and cardiovascular death in apparently healthy men. *Circulation.* 1991;84:613–7.
- 20 98. Deregibus MC, Cantaluppi V, Calogero R, Lo Iacono M, Tetta C, Biancone L, et al. Endothelial progenitor cell
21 derived microvesicles activate an angiogenic program in endothelial cells by a horizontal transfer of mRNA.
22 *Blood.* 2007;110:2440–8.
- 23 99. Diehl P, Fricke A, Sander L, Stamm J, Bassler N, Htun N, et al. Microparticles: major transport vehicles for
24 distinct microRNAs in circulation. *Cardiovasc. Res.* 2012;93:633–44.
- 25 100. Baggish AL, Hale A, Weiner RB, Lewis GD, Systrom D, Wang F, et al. Dynamic regulation of circulating
26 microRNA during acute exhaustive exercise and sustained aerobic exercise training. *J. Physiol.* 2011;589:3983–
27 94.
- 28 101. Pauw K De, Roelands B, Cheung SS, de Geus B, Rietjens G, Meeusen R. Guidelines to classify subject groups
29 in sport-science research. *Int. J. Sports Physiol. Perform.* 2013;8:111–22.
- 30 102. Decroix L, De Pauw K, Foster C, Meeusen R. Guidelines to classify female subject groups in sport-science
31 research. *Int. J. Sports Physiol. Perform.* 2016;11:204–13.
- 32

1 FIGURE LEGENDS

2

3 Figure 1. Projected time-course of circulating microvesicle (MV) appearance during and after an endurance
4 exercise session based on the current literature. A 2 to 3-fold increase in plasma platelet MV (PMV)
5 concentrations during and few hours after exercise have been consistently reported, but data supporting an
6 increase in activation-derived endothelial MV (EMV) concentrations are inconsistent. The concentration of EMVs
7 carrying apoptotic markers may be decreased in the circulation hours to days after exercise.

8

9 Figure 2. Exercise-derived microvesicle formation and release in the circulation, and their putative role as
10 ubiquitous mediators of adjustments and adaptations to exercise. Representative figure not to scale. Schematic
11 figure developed using images from the Servier Medical Art image bank.