



This is a repository copy of *Dietary behaviours in the context of nutrition transition : a systematic review and meta-analyses in two African countries.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/158521/>

Version: Published Version

---

**Article:**

Rousham, E.K., Pradeilles, R., Akparibo, R. et al. (8 more authors) (2020) Dietary behaviours in the context of nutrition transition : a systematic review and meta-analyses in two African countries. Public Health Nutrition. ISSN 1368-9800

<https://doi.org/10.1017/s1368980019004014>

---

This article has been published in Public Health Nutrition [<https://doi.org/10.1017/S1368980019004014>]. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © 2020 The Authors.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**



If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



# Dietary behaviours in the context of nutrition transition: a systematic review and meta-analyses in two African countries

Emily K Rousham<sup>1,\*</sup> , Rebecca Pradeilles<sup>1,2</sup>, Robert Akparibo<sup>2</sup>, Richmond Aryeetey<sup>3</sup> , Kristin Bash<sup>2</sup>, Andrew Booth<sup>2</sup>, Stella K Muthuri<sup>4</sup>, Hibbah Osei-Kwasi<sup>2,5</sup>, Colette M Marr<sup>2</sup>, Tom Norris<sup>1</sup> and Michelle Holdsworth<sup>2,6</sup>

<sup>1</sup>School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, United Kingdom: <sup>2</sup>School of Health and Related Research (ScHARR), University of Sheffield, Sheffield, United Kingdom: <sup>3</sup>Department of Population, Family and Reproductive Health, School of Public Health, University of Ghana, Accra, Ghana: <sup>4</sup>UK Department of International Development (DFID), Nairobi, Kenya: <sup>5</sup>Department of Geography, University of Sheffield, Sheffield, United Kingdom: <sup>6</sup>NUTRIPASS Unit, University of Montpellier-IRD-Montpellier SupAgro, Montpellier, France

Submitted 3 March 2019: Final revision received 2 September 2019: Accepted 10 September 2019

## Abstract

**Objective:** To synthesise evidence of urban dietary behaviours (macronutrients, types of foods, dietary diversity and dietary practices) in two African countries in relation to postulated changes in the context of nutrition transition.

**Design:** Systematic review and meta-analyses, including six online databases and grey literature, 1971–2018 (Protocol CRD42017067718).

**Setting:** Urban Ghana and Kenya.

**Participants:** Population-based studies of healthy adolescents and adults.

**Results:** The forty-seven included studies encompassed 20 726 individuals plus 6526 households. Macronutrients were within WHO-recommended ranges: mean energy intake was 1867 kcal/d (95 % CI 1764, 1969) and the proportions of macronutrients were carbohydrate 61.2 % (58.4, 64.0), fat 25.3 % (22.8, 28.0) and protein 13.7 % (12.3, 15.1). The proportion of population consuming fruit and vegetables was 51.6%; unhealthy foods, 29.4%; and sugar-sweetened beverages (SSBs), 39.9%. Two-thirds (68.8%) consumed animal-source proteins. Dietary diversity scores were within the mid-range. Meal patterns were structured (typically three meals per day), with evidence lacking on snacking or eating out.

**Conclusions:** Population-level diets fell within WHO macronutrient recommendations, were relatively diverse with structured meal patterns, but some indications of nutrition transition were apparent. The proportion of population consuming fruit and vegetables was low compared to healthy-eating recommendations, and consumption of SSBs was widespread. A paucity of evidence from 1971 to 2010 precluded a longitudinal analysis of nutrition transition. Evidence from these two countries indicates which aspects of dietary behaviours may be contributing to increasing overweight/obesity, namely a low proportion of population consuming fruit and vegetables and widespread consumption of SSBs. These are potential targets for promoting healthier diets.

**Keywords**  
Dietary behaviours  
Macronutrients  
Food consumption  
Dietary diversity  
Dietary practices  
Ghana  
Kenya

## Introduction

Nutrition transition includes changes in dietary patterns and nutrient intakes observed as populations go through economic and social development<sup>(1)</sup>. It has been associated with concomitant increases in obesity and nutrition-related non-communicable diseases (NR-NCDs), which are now among the leading causes of death globally<sup>(2)</sup>. In African

countries, the risk of NR-NCDs is increasing at a faster rate and at a lower economic threshold than in high-income countries. Changes in dietary behaviours could account, in part, for this increased disease risk<sup>(3)</sup>. The generalised patterns of dietary change associated with nutrition transition include increased consumption of fat, particularly vegetable and edible oils, increased added sugar, increased animal-source foods and decreases in cereals and fibre<sup>(1,4)</sup>,

\*Corresponding author: Email e.k.rousham@lboro.ac.uk

specifically coarse grains, staple cereals and pulses<sup>(5)</sup>. However, the exact nature of changes in dietary behaviours and the foods that drive nutrition transition vary according to the region.

Evidence of how dietary behaviours are changing in urban populations in African countries is lacking, although food balance sheets show that most countries have experienced an increase in the national availability per capita energy, fat, protein and sugar<sup>(6)</sup>. Ghana and Kenya are two middle-income countries experiencing rapid urbanisation and increasing levels of overweight/obesity and NR-NCDs<sup>(7)</sup>. We selected these two countries as representing different cultural contexts of nutrition transition (East and West Africa), and because both countries are undergoing rapid economic development, urbanisation and increases in obesity prevalence. In Ghana, overweight and obesity in women aged 15–49 years tripled from 13.2% to 40.1% between 1993 and 2014<sup>(8)</sup>. In Kenya, one-third of women are overweight or obese<sup>(9)</sup>. In Ghana and Kenya, NCDs account for >40% of total morbidity and are now recognised as pressing public health concerns evidenced by interventions to promote healthy diets<sup>(10–12)</sup>. However, dietary behaviours in urban African populations have not been systematically reviewed in relation to how diets are postulated to change in the context of nutrition transition.

We aimed to assess the strength of evidence of nutrition transition in urban Africa based on the reported changes in dietary behaviours stated in the literature in low- and middle-income countries (LMICs) undergoing transition<sup>(1,4,5)</sup>. In line with the reported changes with nutrition transition in other studies, we aimed to examine dietary behaviours related to: macronutrient profiles indicating a higher proportion of energy intake from fats and a lower proportion from carbohydrate<sup>(1,4)</sup>; a high consumption of animal-source foods, edible oils, energy-dense snack foods, sweetened beverages, processed foods and sugary snacks in the food items consumed<sup>(3)</sup>; enhanced dietary diversity<sup>(13)</sup>; and less structured dietary practices reflected in more food or meals eaten away from home and an increase in processed, convenience and ‘fast’ foods. The aim of the study was to synthesise evidence of urban dietary behaviours (macronutrients, types of foods, dietary diversity and dietary practices) in two African countries in relation to postulated changes in the context of nutrition transition.

## Methods

### Review typology

A systematic review of dietary behaviours in populations of urban adolescents and adults living in Ghana and Kenya was undertaken to systematically search for, appraise and synthesise research evidence<sup>(14)</sup>. The review was conducted in line with the PRISMA statement<sup>(15)</sup>. A meta-analysis was conducted on data from quantitative studies

where feasible. The review protocol was registered with Prospero (CRD42017067718).

### Inclusion and exclusion criteria

Inclusion criteria for the review were studies of males and females aged 13–18 years (adolescents) or >18 years of age (adults) living in urban areas of Ghana or Kenya with a quantitative primary outcome of diet or dietary behaviours. Diet was defined as food consumption patterns. Dietary behaviours were defined as a combination of eating habits, preferences, choices and feeding-related mannerisms<sup>(16)</sup>. Eligibility criteria for the searches included evidence dating from 1971 to present in line with the concept of epidemiological transition theory<sup>(17)</sup>. Published articles and academic theses available through online repositories (Masters and PhD) in the English language were included.


Exclusion criteria included studies based on a purposively selected specialist sample that was likely to influence dietary behaviours (e.g., long-distance runners, vegetarians, patients with HIV or AIDS) and clinical studies of inpatients. Where studies included a healthy control or comparison group, data from the comparison group were extracted.

### Search strategy

We developed a search strategy and conducted searches in six electronic databases: MEDLINE, EMBASE, PsycINFO, CINAHL, ASSIA and Africa Index Medicus (see online supplementary material 1 for example of MEDLINE search strategy). Relevant institutional repositories were searched for grey literature followed by searches in Google Scholar to obtain abstracts and full texts. Institutional online repository searches included nine Kenyan and four Ghanaian universities with nutrition departments offering postgraduate courses (see online supplementary material 2); the UK White Rose collection (Universities of Sheffield, York and Leeds); and those of the top ten UK universities for research on food and nutrition in Africa according to the Web of Science. Hand-searches of reference lists of included literature were carried out alongside data extraction. Citation follow-up was used to identify studies missed by electronic and grey literature searches. Electronic database searches were carried out from 25 to 28 July 2017; grey literature searches were completed by 11 October 2017; and hand-searches and requests to authors for data continued up to 31 January 2018.

### Screening

Duplicates were removed before screening. Title and abstract and subsequently full-text items were screened against the inclusion criteria listed above by a team (KB, RP, JT, HO-K). An independent reviewer (ER) checked a 10% random sample of excluded records at the title-and-abstract and full-text screening stages. Any disagreements



**Fig. 1** Framework of dietary behaviours of relevance to nutrition transition

were resolved via discussion. If one study led to more than one publication presenting the same data, only one publication record was included for data synthesis and meta-analyses. If publication records presented different data from a single study, all records were included in data synthesis.

#### Data extraction

Data were extracted using a standardised data extraction spreadsheet in Excel by eight reviewers (ER, RP, KB, HO-K, RAK, RAR, SM, JT). A draft data extraction form was piloted by all extractors, modified following discussion, then finalised. Two reviewers (RP, ER) checked and gave feedback on a sample of data extraction by each assessor to ensure consistency of extraction and recording. Data were extracted on title, author(s), year, study design, study aim, publication type, setting, country (Kenya/Ghana), city, sample characteristics (age, sex, socioeconomic indicators), methods (sample size, sampling strategy, design, method of measurement, data management), outcome measures (type of dietary behaviour studied), analysis and results and study limitations. Where data could not be extracted from the publication, we contacted the authors to request relevant data (e.g., disaggregated data from pooled urban and rural samples, or pooled for more than one country).

Figure 1 presents the framework of dietary behaviours of relevance to nutrition transition. Reported dietary

behaviours were divided into nutrient-based and food-based behaviours, then categorised as: (i) energy and macronutrient intake, (ii) food items consumed, (iii) dietary patterns using data reduction techniques (e.g., principal component analysis), (iv) dietary diversity and food variety and (v) dietary practices. Dietary patterns were defined as ‘the quantities, proportions, variety, or combination of different foods, drinks and nutrients in diets, and frequency of habitual consumption’<sup>(18)</sup>. Dietary practices were defined as the foods consumed (e.g., type of snacks, street foods), where, how and with whom foods were eaten, and where ingredients were purchased<sup>(19)</sup>. Studies often included more than one type of dietary behaviour.

#### Quality assessment

Each document included in the review was rated using a 14-criterion quality assessment tool<sup>(20)</sup>. This was modified from the original tool by replacing the score for each criterion (0, 1, 2) with a qualitative assessment of high quality/green (low risk of bias), medium quality/yellow or low quality/red (high risk of bias) as Cochrane guidance advises against the use of scores<sup>(21)</sup>. Quality assessment was conducted by eight assessors working in four teams. The first and second reviewer independently completed a quality assessment rating for each record. Following this, they compared ratings and came to an agreed rating for each criterion. Two assessors (ER and RP) undertook checks of every agreed rating by the review teams. Online supplementary material 3 provides

a summary of the quality assessment for each of the fourteen criteria as well as the overall quality assessment for included studies.

The overall quality assessment of each publication (high, medium or low quality) was calculated based on the quality rating for four out of fourteen individual criteria that were of most relevance to the types of studies included (observational, cross-sectional studies) and those that reflected the quality and interpretation of dietary data. These included: criteria 3 'Is the method of subject/comparison group selection or source of information/input variables described and appropriate?'; criteria 8 'Is the outcome and (if applicable) exposure measure(s) well defined and robust to measurement bias?'; criteria 10 'Are the analytic methods described/justified and appropriate?'; and criteria 13 'Are results reported in sufficient detail?' Further details of methods to assign overall quality are provided in online supplementary material 3.

### **Data synthesis and analysis**

Data were grouped and analysed according to the relevant dietary behaviour(s): (i) energy and macronutrient intakes (fat, carbohydrates and protein), (ii) food items consumed, (iii) dietary patterns, (iv) dietary diversity and food variety and (v) dietary practices.

#### *Energy and macronutrient intakes*

Data on energy, protein, fat and carbohydrate intakes were extracted for meta-analyses. Mean energy intakes were recorded as kcal/d. The percentage of energy from protein, fat and carbohydrate were either extracted directly or calculated from intakes reported as g/d. Data for male and female samples were retained separately in the meta-analyses where available. Where two records came from the same study, we selected the record that provided the most detailed or more reliable macronutrient intake data<sup>(22,23)</sup>. Intakes were compared to the WHO population nutrient intake goals of carbohydrate 55–75 %, fat 15–30 % and protein 10–15 %<sup>(24)</sup>.

Data synthesis was performed using random-effects meta-analyses to account for the expected heterogeneity between studies, with heterogeneity expressed using the  $I^2$  statistic<sup>(25)</sup>. Total energy intake (kcal/d), stratified by country, was synthesised using the 'metan' command in Stata. The percentage of energy consumed was synthesised using the 'metaprop' command, incorporating the Freeman-Tukey double arcsine transformation of proportions to stabilise variances<sup>(26)</sup>. The pooled result and 95 % CIs were back-transformed for interpretation, and forest plots were generated. Analyses were performed in Stata MP, version 15.1. A narrative synthesis was conducted for studies reporting macronutrient intakes that could not be included in the meta-analyses.

#### *Food items consumed*

A meta-analysis was carried out at the food group (rather than food item) level because heterogeneity between studies was too high at the food item level. To achieve data consistency, a list of 182 food items categorised into eighteen food groups was generated based on the markers of nutrition transition from the literature<sup>(1,3,4,27–29)</sup>. Additional details on the method of grouping food items for the meta-analyses can be found in online supplementary material 4. The same statistical methods were employed as for macronutrient intakes. Results were not stratified by country due to the low number of studies. Studies excluded from the meta-analyses were reviewed by narrative synthesis.

#### *Dietary patterns, dietary diversity and dietary practices*


Data on dietary patterns, dietary diversity and dietary practices were synthesised narratively due to high heterogeneity. Dietary practices were synthesised as: (i) meal patterns and snacking behaviour, (ii) eating out of home and (iii) food provisioning. Eating out of home included: purchase of fast foods, frequency of street food consumption and eating out. Meal patterns were defined as consuming food at the expected three meals a day. Snacks were defined either as food consumed outside of typical meal times or by the authors' own descriptions. This allowed us to investigate whether there was de-structuring of eating towards a less traditional, unstructured meal pattern.

## **Results**

The search yielded 2973 records. After removal of duplicates, 2715 titles and abstracts were screened (see Fig. 2). Of these, 116 records qualified for full-text screening, and fifty-nine records were excluded at the full-text stage. The remaining records, plus an additional five articles from searches of reference lists, met the inclusion criteria and were included in the review.

### **Description of studies**

Fifty-one records were included from forty-seven studies dating from 1988 to 2017: twenty-seven records (twenty-six studies) from Ghana and twenty-four records (twenty-one studies) from Kenya. The records included nineteen theses (eighteen Masters', one PhD). Most studies included adults only (thirty-five studies); four studies included adolescents only; and eight studies included adults and adolescents<sup>(27–34)</sup>. Thirty-two studies comprised mixed samples of males and females, while fifteen studies included only females<sup>(22,27,35–42)</sup>. A total of 12 065 participants and a further 1452 households were included from Ghana; a further 8661 individuals as well as 5074 households were included from Kenya (total 20 726 individuals



**Fig. 2** (colour online) Flow diagram of study selection process

plus 6526 households). The sample size of participants in individual studies ranged from 19 to 4037. Details of all included studies and participant characteristics are provided in online supplementary material 5.

The earliest included study dated from 1988. From the first three decades of the search period (1971–2000), there were only nine studies from both countries. From 2000 to 2010, there were a further fourteen studies, and post-2010, there were twenty-four studies. The measurement tools used to assess dietary behaviours were FFQs (thirty-four studies)<sup>(22,28–33,36–38,41,43–65)</sup>, food practices questionnaires (seven studies)<sup>(34,41,50,64,66–68)</sup> and 24-h recall (twenty-eight studies)<sup>(22,23,27,31,32,35,36,39,40,42,44,46,48,52,55,57,59,61,63,64,67–72)</sup>. One study employed a 7-d food diary<sup>(73)</sup>. There were twenty-three studies on macronutrient intake, twenty-eight studies on food items consumed, nine studies on dietary diversity and food variety, two on dietary patterns and thirteen on dietary practices (see online supplementary material 5).


### Quality assessment summary

Online supplementary material 3 shows a summary of quality assessment for each included record. We assessed thirteen records (25.5%) as high quality<sup>(22,30,40,49,53,58,61–63,70,72,74)</sup>, twenty-five records (49.0%) were of medium quality<sup>(23,27,29,31,32,36,38,41,42,47,48,50–52,56,57,59,60,65–67,69,71,73,75)</sup> and thirteen (25.5%) were of low quality<sup>(28,34,35,37,39,43–46,54,55,64,68)</sup>. Overall, 74.5% were of high or medium quality, indicative of reliable evidence and low risk of bias.

### Summary of dietary behaviours

#### Macronutrient intakes

**Overall summary.** Twenty-three studies reported energy and macronutrient intakes: eleven from Ghana and twelve from Kenya. Most studies (nineteen out of twenty-three) employed a 24-h recall method to quantify intakes, and



**Fig. 3a** (colour online) Total daily energy intake. m = males, f = females, mix = both sexes. \* Dagoretti, \*\* Korogochi. 1 kcal = 4.184 kJ

four studies used an FFQ (see online supplementary material 5). There were no studies from 1970s or 1980s, five studies from 1990s, and the remainder were from 2000 onwards (eighteen) (Online supplementary material 6).

**Meta-analyses.** Figures 3(a)–3(d) presents energy and macronutrient intakes by year of study. No trend in intake over time was apparent, and longitudinal analysis was not conducted due to high heterogeneity. Mean energy intake was 1849 kcal/d (95 % CI 1700, 1998) from nine studies in Ghana *v* 1889 kcal/d (95 % CI 1762, 2017) from six studies in Kenya (Fig. 3(a)). The contribution of carbohydrates to total energy intake was 62.9 % (95 % CI 59.2, 66.6) in Ghana and 59.6 % (95 % CI 55.5, 63.6) in Kenya (Fig. 3(b)). Energy intake from fat was 24.7 % (95 % CI 21.2, 28.3) in Ghana and 25.8 % (95 % CI 22.1, 29.6) in Kenya (Fig. 3(c)). Protein comprised just below 14 % of total energy intake in both countries (Ghana: 13.6 %, 95 % CI 11.8, 15.6; Kenya: 13.8 %, 95 % CI 11.6, 16.0) (Fig. 3(d)). Heterogeneity between studies, as evidenced by the  $I^2$  statistic, was substantial ( $I^2 > 90\%$ ).


**Narrative synthesis.** Among the studies excluded from the meta-analyses, mean energy intake was 1855 kcal/d

(no SD) among working women<sup>(36)</sup>. Two studies reported assessing dietary intakes but did not present data in a usable format<sup>(22,62)</sup>. Mbotela<sup>(60)</sup> reported that 7.8 % of the sample ( $n$  102) consumed 'adequate' carbohydrate (author's definition), 15.6 % consumed 'adequate' protein and 5.9 % 'adequate' fat. Wathome<sup>(65)</sup> reported energy consumptions of 2700 kcal/d (high income), 2500 kcal/d (middle income) and 2400 kcal/d (low income).

#### Food items consumed

**Overall summary.** Twenty-eight studies reported food consumption data: seventeen in Ghana<sup>(27–29,37,43–53,56,57)</sup> and eleven in Kenya<sup>(22,31,32,34,41,58,59,61,64,65,67)</sup>. Most studies (twenty-six out of twenty-eight) used an FFQ to assess food consumption patterns; one study used a 24-h recall; and another one employed a food practices questionnaire. The reporting of food items and methods of data synthesis and coding prior to the meta-analyses is provided in Online supplementary material 5.

**Meta-analyses.** Data from fifteen studies were included in the meta-analyses: nine from Ghana<sup>(27,28,37,43–45,47,52,57)</sup> and six from Kenya<sup>(22,31,58,59,64,67)</sup>. An estimated 68.8 % of the



**Fig. 3b** (colour online) Percentage of total energy from carbohydrates. m = males, f = females, mix = both sexes. \* Dagoretti, \*\* Korogochi

sample consumed animal-source foods (95 % CI 57.9, 78.8) (Fig. 4). Dairy products were consumed most commonly (87.1 %, 95 % CI 68.1, 98.3), followed by red meat (85.9 %, 95 % CI 57.0, 99.8), poultry (69.6 %, 95 % CI 50.0, 86.2), eggs (68.5 %, 95 % CI 36.0, 93.3), fish (46.6 %, 95 % CI 25.2, 68.6) and processed meat (28.0 %, 95 % CI 25.3, 30.9).

Fruit, vegetables or legumes were consumed by half of the sample (51.6 %, 95 % CI 33.9, 69.0) (Fig. 5). Vegetables, excluding potatoes and starchy roots, were the most commonly consumed (63.7 %, 95 % CI 0.0, 100.0), followed by fruit (55.2 %, 95 % CI 9.8, 95.5), vitamin A-rich dark green leafy vegetables (54.5 %, 95 % CI 31.4, 76.6), fruit and vegetables (53.3 %, 95 % CI 2.8, 99.1), legumes (29.7 %, 95 % CI 28.1, 31.3) and vitamin A-rich fruit and other vegetables (27.2 %, 95 % CI 25.4, 29.1).


An estimated 29.4 % (95 % CI 20.7, 39.0) of the sample consumed unhealthy food and drink (i.e., vegetable fats (including palm oil), sugar-sweetened beverages (SSBs), cakes and biscuits, chocolate and sweets, alcohol, animal fats) (Fig. 6). These foods and drinks were classed as

unhealthy based on their properties of being either energy-dense (fats, cakes, biscuits, chocolates and sweets) or nutrient-poor (alcohol and SSBs). We did not distinguish between vegetable fat and animal fat since this information was not always available and because the percentage of energy from total fat is associated with unhealthy weight gain in LMICs. SSBs were consumed by the highest proportion of the population (39.9 %, 95 % CI 20.6, 61.0), followed by cakes and biscuits (32.6 %, 95 % CI 29.2, 36.0), vegetable fats (23.2 %, 95 % CI 21.0, 25.5) and chocolates and sweets (31.3 %, 95 % CI 27.7, 34.9). No studies reported on the consumption of savoury snacks. Only 14.0 % of the sample consumed alcohol (95 % CI 11.4, 16.8) and 15.0 % animal fats (95 % CI 11.7, 19.0). Heterogeneity between studies, as evidenced by the  $I^2$  statistic, was substantial ( $I^2 > 90\%$ ).

#### Narrative synthesis

Data from thirteen studies could not be pooled for meta-analyses: eight from Ghana<sup>(29,46,48–51,53,56)</sup> and five from





**Fig. 3c** (colour online) Percentage of total energy from fat. m = males, f = females, mix = both sexes. \* Dagoretti, \*\* Korogochi

Kenya<sup>(32,34,41,61,65)</sup>. Of these, two studies examined a single item unrelated to nutrition transition (fermented foods<sup>(34)</sup> and groundnut consumption<sup>(51)</sup>). One study examined foods consumed in a specific market area<sup>(50)</sup>. Data from one study were inaccessible<sup>(32)</sup>. The remaining studies are synthesised narratively below.

#### Animal-source foods


**Milk and dairy products.** Seven papers reported milk/dairy consumption. In Kenya, milk consumption/purchase per household ranged from 2923 g/week ( $\pm 2685$  g)<sup>(41)</sup> to 7416 g/week (SD not provided)<sup>(65)</sup>. Per person, 683 g/week ( $\pm 795$  g) of milk was purchased<sup>(41)</sup>. Frequency of milk consumption ranged from 1.8<sup>(61)</sup> to 14.5 times per week<sup>(65)</sup>.

In Ghana, one study reported dairy product consumption of 50.1 g/d (95 % CI 47.4, 52.9, equivalent to 350 g/week)<sup>(49)</sup>. Another study observed almost twice as much milk consumed at 644 g/week per person ( $\pm 581$  g)<sup>(46)</sup>, similar to that reported in Kenya. The frequency of consumption in Ghana ranged from zero servings of milk per week<sup>(48)</sup> to

0.7 serving of milk/dairy products per week<sup>(29)</sup>; the same study reported milk/dairy products as least frequently consumed food group<sup>(29)</sup>.

**Red meat.** Two studies reported on beef consumption in Kenya<sup>(41,65)</sup>, and two studies reported on red meat consumption in Ghana<sup>(48,49)</sup>. In Kenya, the average frequency of beef consumption per week per household was five times per week (SD not provided)<sup>(65)</sup>, and intake was, on average, 66.4 g/week ( $\pm 125.0$ ) per individual<sup>(41)</sup>. Conversely, in Ghana, the frequency of servings of red meat was lower, 1.5 servings per week (IQR 0.5, 3.5)<sup>(48)</sup>, but reported intake was higher than in Kenya at 45 g/d ( $\pm 2.2$  g)<sup>(49)</sup>.

**Fruit and vegetables.** Fruit and vegetable intakes in Ghana were reported by five studies<sup>(29,48,49,53,56)</sup> that were not included in the meta-analyses. One study reported a small proportion of individuals eating five portions of fruit per day (1.3 %) and five portions of vegetables per day (1.1 %)<sup>(56)</sup>. Another study estimated that over two-thirds (67.1 %, 95 % CI 63.1, 70.9) of the sample did not consume the recommended five servings of fruit and



**Fig. 3d** (colour online) Percentage of total energy from protein. m = males, f = females, mix = both sexes. \* Dagoretti, \*\* Korogochi

vegetables on average per day<sup>(53)</sup>. A third study reported 3.5 servings of fruit per week but did not specify serving size<sup>(48)</sup>. One study found higher intake of vegetables ( $267.1 \pm 4.1$  g) than fruits ( $152.8 \pm 7$  g)<sup>(49)</sup>. Likewise, one study found vegetables were consumed more frequently than all other food groups (3.8 times per week; SD not provided)<sup>(29)</sup>.

**Sugar-sweetened beverages.** Three studies found that SSBs were consumed more frequently, on average, in Kenya at 2.6 times per week (males 2.66, SD 2.59; females 2.60, SD 2.42)<sup>(61)</sup> than in Ghana, where consumption ranged from 1.2<sup>(29)</sup> to 1.5 times per week (IQR 0.5–1.5)<sup>(48)</sup>.


#### Dietary patterns

Dietary patterns were reported in two studies in urban Ghana<sup>(48,49)</sup>. Galbete *et al.*<sup>(49)</sup> reported that urban Ghanaians had highest median scores on 'rice, pasta, meat and fish' patterns compared to rural and European

Ghanaians. Diet was characterised by high intakes of dairy, red meat, processed meat, eggs, legumes, rice and pasta, fish, mixed meaty dishes, cakes and sweets and condiments. A second study identified two dietary patterns in urban Ghana<sup>(48)</sup>: a 'purchase' pattern (high intake of sweets, juice, rice, soft drinks, 'Milo', eggs, red meat, margarine, cow's milk, poultry, cucumber, lettuce, carrot and fruits) and a 'traditional pattern' (high intake of fruits, plantain, cassava, smoked fish, palm oil, eggplant, green leaves, *banku* and beans). The purchase pattern was more typical of younger participants of a higher socioeconomic status (SES), whereas the traditional pattern was more typical of older and more deprived participants.

#### Dietary diversity and food variety

Dietary diversity was reported in eight studies, four in each country. The earliest available information on dietary diversity dated from 2012 (Table 1). Four studies focused




**Fig. 4** (colour online) Proportion of sample consuming animal-source foods. m = males; f = females; mix = both sexes; G = Ghana.  $I^2$  not calculated for items with less than four studies

on adult women<sup>(27,40,67,74)</sup>, one on adult women and adolescent girls<sup>(38)</sup> and one on adolescent girls<sup>(54)</sup>. Data pooling was precluded by different methods of assessment and reporting. The dietary diversity of women based on the nine-point FAO method was marginally higher in Ghana ( $4.7 \pm 1.5$ )<sup>(27)</sup> compared to Kenya ( $4.4 \pm 1.1$  in Dagoretti, and  $4.1 \pm 1.4$  in Korogocho, both in Nairobi)<sup>(40)</sup>. Steyn *et al.*<sup>(74)</sup> reported a score of 4.8 (95% CI 4.6, 4.9) out of nine food groups among women in Kenya using a different assessment tool<sup>(78)</sup>. In Ghana, household dietary diversity, using the nine-point FAO food groups, was estimated at 6.8 (SD not reported)<sup>(29)</sup>. Dietary diversity in this sample varied by age, sex, education, occupation of household head, household size and residence of household<sup>(29)</sup>.

Other studies reported dietary diversity using different scoring systems. In Kenya, a mean score of  $5.2 \pm 1.1$  was reported based on an eight-point food group system<sup>(64)</sup>, and a score of  $7.5 \pm 1.4$  based on a fourteen-point food group score for Kenyan women<sup>(67)</sup>. In Ghana, a mean dietary diversity score of  $9.2 \pm 2.9$  out of a sixteen-point food group score was reported<sup>(38)</sup>. In this study, dietary diversity was lower among women ( $8.31 \pm 2.83$ ) than adolescents ( $10.11 \pm 2.76$ ), although the adolescent sample was small ( $n = 13$ ). Also in Ghana, 56% of adolescents consumed diets with  $< 3$  of six food groups<sup>(54)</sup>.

Two studies assessed food variety. In Kenya, the number of foods consumed in the past 24 h from a list of 163 items produced a food variety estimate of 7.8 (95% CI 7.5, 8.2)<sup>(74)</sup>. In Kenya, more than two-thirds of women



**Fig. 5** (colour online) Proportion of sample consuming fruit, vegetables and legumes. m = males; f = females; mix = both sexes; G = Ghana.  $I^2$  not calculated for items with less than four studies

in Viwandani and Korogocho consumed diets with >4 food groups<sup>(30)</sup>.


*Dietary practices*

Overall thirteen studies reported on dietary practices: eight from Ghana and five from Kenya (Online supplementary material 7). Practices were grouped as meal patterns and snacking, eating outside the home and food provisioning. **Meal patterns and snacking.** Seven Ghanaian studies reported on meal patterns<sup>(37,45,46,48,54,55,66)</sup>. Of these, five reported structured meals, that is, breakfast, lunch and dinner<sup>(37,45,48,54,66)</sup>. In the remaining studies, meal patterns were unclear. One study reported almost all participants ate 2–3 meals per day at weekends (91.5%) and on weekdays (90.7%)<sup>(45)</sup>. In Barichella *et al.*<sup>(46)</sup>, participants consumed on average 2.6 meals per day, but did not specify the types of meals consumed. Data on meal patterns were not available for Kenya.

Three of the seven Ghanaian studies reported on snacking behaviours<sup>(37,45,66)</sup>. Snacking at least once per day was reported by 83.7% of respondents<sup>(45)</sup>; of these, 53.2% snacked on pastries and beverages with added sugar, 19% snacked on nuts and roasted corn/plantain, and 11.5% snacked on fruits. Snacks eaten by university students included fruits, soft drinks, or baked, fried or roasted foods, but provided no further breakdown of food items<sup>(66)</sup>. Among female homemakers in Accra, 28% of women reported buying snacks from vendors<sup>(37)</sup>.

**Eating outside the home.** Behaviours around eating outside the home were reported in seven studies (Online supplementary material 7): four in Ghana<sup>(37,45,50,66)</sup>, three in Kenya<sup>(30,62,75)</sup>. Only one study, from Ghana, reported on fast-food purchasing<sup>(45)</sup>, suggesting that 55% of respondents had purchased or eaten fast food in the last 7 d.

No studies from Kenya reported on fast-food purchasing but referred to local ‘street food’. Street food consumption



**Fig. 6** (colour online) Proportion of sample consuming 'unhealthy' food items. m = males; f = females; mix = both sexes; G = Ghana.  $I^2$  not calculated for items with less than four studies

was reported in three studies, but they were not always explicit about what was consumed (snacks or meals). In two Kenyan studies<sup>(30,62)</sup>, street foods were consumed by less than a quarter of respondents. In urban slums, 13% of households purchased street-cooked foods<sup>(30)</sup>, while a non-slum survey reported 22% of household members consuming street foods daily<sup>(62)</sup>. In Ghana, the frequency of eating street foods ranged from one to twenty-one times in a week<sup>(50)</sup>, with men eating street foods more often than women (Online supplementary material 7). Among university students in Ghana, 86% purchased meals from street vendors; and of these, 47% did so on a daily basis<sup>(66)</sup>.

For meals purchased and/or eaten outside the home, 65% of respondents in Kenya ate at least once per day outside the home<sup>(75)</sup>, while women in the two lower SES groups ate significantly fewer meals away from home compared to the two upper SES groups<sup>(75)</sup>. Among female homemakers, 20% always purchased breakfast from vendors, 34% purchased lunch and 20% purchased supper from vendors<sup>(37)</sup>.

**Food provisioning.** Three studies focused on food provisioning practices<sup>(30,68,73)</sup>. In urban slums of Nairobi, 87% of households purchased raw food from the market<sup>(30)</sup>. A study of low SES households in Nairobi reported that urban farmers purchased 67% of their food from the market and produced 25% of their own food, compared to non-farmers who purchased 82% of all household foods<sup>(73)</sup>.

## Discussion

We synthesised evidence of urban dietary behaviours (macronutrients, types of foods, dietary diversity and dietary practices) in two African countries in relation to postulated changes in the context of nutrition transition. This comprehensive search of published studies and grey literature over a period of nearly 50 years reveals how little is known on urban dietary behaviours prior to 2010. No studies reported macronutrient intakes from the 1970s or 1980s. Five studies were available in the 1990s, and the

**Table 1** Summary of studies reporting dietary diversity and food variety

Study identifier	Outcome measure	Number of food groups	Outcome	
			Mean or <i>n</i>	SD or 95% CI
<b>Ghana</b>				
Amugsi <i>et al.</i> <sup>(27)</sup>	Women's diet diversity	Nine food groups (Kennedy <i>et al.</i> ) <sup>(84)</sup>	4.7	1.5
Nti <i>et al.</i> <sup>(54)</sup>	Adolescent diet diversity	Six food groups (no indicated reference)	56.0% had diets with <3 food groups; 20.4% had diets with 3–4 food groups; 23.6% had diets with 5–6 food groups	
Okutu <sup>(29)</sup>	Individual diet diversity (of household head)	Nine food groups – adapted version of Kennedy <i>et al.</i> <sup>(84)</sup>	4.7	1.5
Quarshie <sup>(38)</sup>	Diet diversity of adult women, adolescents	Sixteen food groups – adapted from FAO (Kennedy <i>et al.</i> ) <sup>(84)</sup>	9.2	2.9
<b>Kenya</b>				
Dominguez-Salas <i>et al.</i> <sup>(40)</sup>	Women's diet diversity	Nine food groups (Kennedy <i>et al.</i> ) <sup>(84)</sup>	4.4 (Dagoretti) 4.1 (Korogocho)	1.1 1.4
Kemunto <sup>(67)</sup>	Women's diet diversity	Fourteen food groups (FAO) <sup>(85)</sup>	7.5	1.4
Kimani-Murage <i>et al.</i> <sup>(30)</sup>	Food variety	Not indicated	Two-thirds of women consume diets with >4 food groups	
Steyn <i>et al.</i> <sup>(74)</sup>	Women's diet diversity	Nine food groups (Hatloy <i>et al.</i> ) <sup>(78)</sup>	4.8	4.6, 4.9
Steyn <i>et al.</i> <sup>(74)</sup>	Food variety score	Number of food items consumed from 163 items (Hatloy <i>et al.</i> ) <sup>(78)</sup>	7.8	7.5, 8.2
Thuita <sup>(64)</sup>	Household diet diversity	Eight food groups – adapted from FAO (no reference given)	5.2	1.1

remaining studies were published after 2000. Dietary diversity data were available after 2012. Studies were cross-sectional and included varying population samples with high heterogeneity; hence a meta-analysis of changes in dietary behaviours over time was not conducted. From the forty-seven included studies, there was some evidence of diverse diets, with relatively widespread consumption of unhealthy foods, suggesting that diets may be undergoing nutrition transition at the population level. However, the findings were mixed. This study, therefore, adds evidence of the difficulty of measuring nutrition transition in emerging economies as recently highlighted<sup>(79)</sup>. In contrast to the strong evidence of increasing overweight/obesity and NR-NCDs in Ghana and Kenya, the less conclusive evidence of nutrition transition may be partly due to a lack of good-quality, longitudinal, national-level surveys utilising validated tools and well-developed food composition tables in urbanising middle-income countries<sup>(79)</sup>. The collation of data from existing longitudinal dietary surveys in LMICs could provide a valuable resource for comparing the timing and rate of nutrition transition of individual countries to a 'reference' source.

Meta-analyses of energy and macronutrient intakes showed that mean intakes were within the normal range of WHO-recommended nutrient intake goals for preventing NR-NCDs<sup>(80)</sup>. Protein contributed 13.7% of energy (recommended range 10–15%) and carbohydrate contributed 62.9% of energy in Ghana and 59.6% in Kenya (recommended range 55–75%). Fat intake comprised 25.8% of

energy in Kenya and 24.7% in Ghana, also within the recommended range (15–30%). Diets were relatively diverse, with mean dietary diversity scores within the mid-range of values, using standardised measurement tools. The consumption of animal-source foods (including dairy, red meat, processed meat, eggs and fish) was widespread in both countries.

Indicators of nutrition transition were apparent in the consumption of unhealthy foods and drinks, particularly SSBs, consumed by around 40% of the sample. The proportion of population consuming fruit and vegetables was low, consumed by only half of the included population. Dietary patterns in Ghana showed that traditional and modern patterns co-existed, characterised by widespread intakes of processed meat, cakes, sweets, soft drinks and juices.

In countries undergoing nutrition transition, it is assumed that dietary practices of urban dwellers shift towards greater consumption of sugary snacks, processed food and food eaten away from home<sup>(5)</sup>. There is insufficient evidence from this review to draw conclusions on whether people eat a lot of fast foods, street foods or are eating out more often. Few included studies on eating out of home, and only one study reported specifically on fast-food purchasing<sup>(45)</sup>. Evidence suggests that men<sup>(50)</sup> and those from higher SES groups<sup>(75)</sup> may be more likely to consume food out of home. There was some conflation in the evidence between fast food (as in 'international' fast food) and locally produced street food. In the African context, the latter can also involve selling similar types of foods (from a nutritional and/or culinary perspective) as those



sold in international fast-food chains. The diversity of food landscape in African cities makes the categorisation of different food outlets a challenge.

Evidence suggests that meal patterns remain structured around the conventional pattern of three meals a day<sup>(37,45,48,54,66)</sup>, with foods more likely to be eaten or purchased out of home at breakfast and lunch. There was insufficient evidence to conclude about the extent of snacking and consequences of snacking on eating frequency and impact on the nutritional quality of diets.

Longitudinal evidence of changes in dietary behaviours in the African region is available from the PURE cohort in South Africa. From 2005 to 2010, rapid increases in sugars and SSB consumption have been reported in urban and rural areas, which is associated with increased NCD risk factors<sup>(81)</sup>. The same study, however, observed the beneficial consequences of nutrition transition, with increased vegetables, fruit and milk consumption in most groups<sup>(82)</sup>. This finding underlines the dual transition processes of reduced undernutrition that accompanies greater food availability and economic development, alongside a rapid increase in unhealthy foods and NR-NCDs. Furthermore, nutrition transition can be experienced differently within a population; fruit and vegetable consumption in urban areas may increase among high SES groups but decrease among low SES groups<sup>(5)</sup>. More sensitive indicators of nutrition transition, such as added sugar or saturated fat consumption, were seldom reported in studies in this review, possibly due to a lack of food composition data on these nutrients; hence these could not be assessed. Similarly, the data precluded any breakdown by type of carbohydrate.

### **Strengths and limitations**

The strengths of this review are the inclusion of studies from both grey and published literature, the synthesis of evidence from two middle-income African countries, and the meta-analyses of energy, macronutrient intakes and food items consumed. We also synthesised dietary behaviours in relation to the key indicators of nutrition transition.

Limitations of the review arise from the limited extent of evidence from early periods of urbanisation, with only nine studies dating from 1971 to 2000, and from the high heterogeneity of evidence. This precluded an assessment of changes in dietary behaviours over time. The impact of urbanisation on diet in Africa is thought to have emerged from the mid-1980s onwards<sup>(6)</sup>, but dietary data from this time are lacking. The reviewed studies focussed on general dietary intakes rather than nutrition transition per se and, therefore, did not necessarily report on the key indicators of transition such as added sugar or saturated fat. One-quarter of included studies were assessed as having a high risk of bias. This was usually due to the absence of detailed methods of dietary assessment and heterogeneity of methods employed in assessment tools

and units of measurement. Dietary studies are liable to a range of misreporting biases, and not all data collection methods were standardised or validated; these sources of bias can affect estimates. However, these factors are incorporated into the risk of bias in the quality assessment procedure and the reliability of estimates from meta-analyses, as indicated by high  $I^2$  values. Furthermore, the focus of this review was not on absolute individual dietary intakes but on population-level dietary behaviours and relative intakes. A further limitation is that only narrative syntheses were possible for dietary patterns, dietary diversity and dietary practices. Even within the same dietary practice, different outcome measures were reported, for example, frequency *v.* types of snacks consumed. The use of STROBE-nut reporting guidelines would improve consistency across studies<sup>(83)</sup>.

### **Conclusion**

This systematic review on the availability, nature and quality of data from two middle-income African countries highlights the paucity of data on dietary behaviours between 1971 and 2000 when rapid urbanisation and early stages of nutrition transition began. Evidence from more recent studies, however, provides an indication of nutrition transition in urban dietary behaviours in these two middle-income African countries. Diets were relatively diverse compared with standardised measures, including a variety of animal-source foods consumed by a high proportion of the population. Meal patterns remained structured around the conventional pattern of three meals per day. There was insufficient evidence to conclude whether patterns of eating out or snacking are shifting. Based on the WHO recommendations on healthy diets containing <10 % of total energy intake from free sugars, unhealthy food and drink consumption was relatively widespread, particularly SSBs that were consumed by 39.9 % of the population. Similarly, based on the WHO recommendation for a healthy diet containing a minimum daily consumption of 400 g or five portions per day of fruit and vegetables (i.e., 100 % of the population consuming fruit and vegetables every day), the proportion of population consuming fruit and vegetables was 51.6 %, which is substantially below recommended intakes. Energy and macronutrient intakes were within the normal range of WHO recommendation. The assessment of dietary changes was hindered by limited evidence from early decades and high heterogeneity between studies. There has been a call for greater information on urban diets, disaggregated by SES<sup>(5)</sup>, and the drivers of unhealthy diets<sup>(5)</sup>. In the global context, the inadequacy of standard dietary assessment instruments and comprehensive food composition tables to assess diets undergoing transition has been highlighted<sup>(79)</sup>. In view of a rapid urbanisation underway in many African



countries, and the varying stages of economic development of different countries, significant changes in dietary behaviours will be taking place in future. Longitudinal, nationally representative dietary surveys, designed to track changes in dietary behaviours in relation to non-communicable disease risk in urbanising African countries such as Ghana and Kenya, would enable an assessment of the contribution of dietary behaviours to the risk of overweight and obesity and non-communicable diseases. Based on evidence from these two countries, we highlight areas where dietary behaviours may be contributing to increasing overweight and obesity, namely a low proportion of population consuming fruit and vegetables and a widespread consumption of SSBs.

### Acknowledgements

**Acknowledgements:** We gratefully acknowledge Fiona Graham for assistance with formatting documents, and Jedidah Tetteh for assistance with screening and data extraction. **Financial support:** Medical Research Council, UK (MR/P025153/1). **Conflict of interest:** None. **Authorship:** E.K.R., R.P., A.B., R.Ar., R.Ak., S.K.M. and M.H. designed the research; M.H., R.P., E.K.R., A.B., R.Ar., R.Ak., S.K.M., H.O.-K., K.B., C.M. and T.N. extracted and analysed data; E.K.R., R.P., A.B., R.Ar., R.Ak., H.O.-K., C.M., T.N. and M.H. wrote the paper; E.K.R. had primary responsibility for the final content. All authors read and approved the final manuscript. **Ethics of human subject participation:** This systematic review and synthesis was based on published works and was, therefore, exempt from ethical approval.

### Supplementary material

For supplementary material accompanying this paper visit <https://doi.org/10.1017/S1368980019004014>.

### References

1. Popkin BM (2001) The nutrition transition and obesity in the developing world. *J Nutr* **131**, 871–873.
2. Naghavi M, Abajobir AA, Abbafati C *et al.* (2017) Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease study 2016. *Lancet* **390**, 1151–1210.
3. Vorster HH, Kruger A & Margetts BM (2011) The nutrition transition in Africa: can it be steered into a more positive direction? *Nutrients* **3**, 429–441.
4. Popkin BM & Gordon-Larsen P (2004) The nutrition transition: worldwide obesity dynamics and their determinants. *Int J Obes Relat Metab Disord* **28**, Suppl. 3, S2–S9.
5. Hawkes C, Harris J & Gillespie S (editors) (2017) Chapter 4: Urbanization and the nutrition transition. In *Global Food Policy Report*. Washington, DC: International Food Policy Research Institute (IFPRI).
6. Steyn NP & Mchiza ZJ (2014) Obesity and the nutrition transition in Sub-Saharan Africa. *Ann NY Acad Sci* **1311**, 88–101.
7. Tenkorang EY, Kuuire V, Luginaah I *et al.* (2017) Examining risk factors for hypertension in Ghana: evidence from the study on global ageing and adult health. *Glob Health Promot* **24**, 14–26.
8. Doku DT & Neupane S (2015) Double burden of malnutrition: increasing overweight and obesity and stall underweight trends among Ghanaian women. *BMC Public Health* **15**, 1–9.
9. Kenya National Bureau of Statistics, Ministry of Health/Kenya, National AIDS Control Council/Kenya, Kenya Medical Research Institute, National Council for Population and Development/Kenya, and ICF International (2015) *Kenya Demographic and Health Survey 2014*, pp. 1–601. Rockville, MD, USA: Kenya National Bureau of Statistics, Ministry of Health/Kenya, National AIDS Control Council/Kenya, Kenya Medical Research Institute, National Council for Population and Development/Kenya, and ICF International.
10. de-Graft Aikins A, Addo J, Ofei F *et al.* (2012) Ghana's burden of chronic non-communicable diseases: future directions in research, practice and policy. *Ghana Med J* **46**, 1–3.
11. Ministry of Health Kenya (2015) *Kenya National Strategy for the Prevention and Control of Non-communicable Diseases 2015–2020*, pp. 1–78. <https://www.who.int/nmh/ncd-task-force/kenya-strategy-ncds-2015-2020.pdf> (accessed 01 March 2019).
12. Ministry of Health Ghana (2012) *Strategy for the Management, Prevention and Control of Chronic Non-communicable Diseases in Ghana 2012–2016*, pp. 1–27. <https://extranet.who.int/nutrition/gina/sites/default/files/GHA-2012-NCDs.pdf> (accessed 01 March 2019).
13. Aurino E, Fernandes M & Penny ME (2017) The nutrition transition and adolescents' diets in low- and middle-income countries: a cross-cohort comparison. *Public Health Nutr* **20**, 72–81.
14. Grant MJ & Booth A (2009) A typology of reviews: an analysis of 14 review types and associated methodologies. *Heal Info Libr J* **26**, 91–108.
15. Moher D, Alessandro L, Jennifer T *et al.* (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Br Med J* **339**, b2535.
16. Stok FM, Hoffmann S, Volkert D *et al.* (2017) The DONE framework: creation, evaluation, and updating of an interdisciplinary, dynamic framework 2.0 of determinants of nutrition and eating. *PLoS One* **12**, 1–23.
17. Omran A (1971) The epidemiological transition. *Milbank Mem Fund Q* **49**, 509–538.
18. Reedy J, Subar AF, George SM *et al.* (2018) Extending methods in dietary patterns research. *Nutrients* **10**, 1–8.
19. Tuomainen H (2006) Migration and foodways: continuity and change among Ghanaians in London. PhD thesis, University of Warwick.
20. Kmet LM, Cook LS & Lee RC (2004) *Standard Quality Assessment Criteria for Evaluating Primary Research Papers from a Variety of Fields*. Edmonton: Alberta Heritage Foundation for Medical Research (AHFMR).
21. Higgins J & Green S (2006) *Assessment of Study Quality. Cochrane Handbook for Systematic Reviews of Interventions 4.2.6 Section 6. The Cochrane Library*. Chichester, UK: John Wiley & Sons, Ltd.
22. Mbochi RW, Kuria E, Kimiywe J *et al.* (2012) Predictors of overweight and obesity in adult women in Nairobi Province, Kenya. *BMC Public Health* **12**, 1.
23. Steyn NP, Parker WA, Nel JH *et al.* (2011) Dietary, social, and environmental determinants of obesity in Kenyan women. *Scand J Public Health* **39**, 88–97.





24. WHO – World Health Organization (1990) Diet, nutrition and prevention of chronic disease. Report of a WHO Study Group (WHO Technical Report Series 797). p. 102.
25. Higgins JPT & Thompson SG (2002) Quantifying heterogeneity in a meta-analysis. *Stat Med* **21**, 1539–1558.
26. Freeman MF & Tukey JW (1950) Transformations related to the angular and the square root. *Ann Math Stat* **21**, 607–611.
27. Amugsi DA, Lartey A, Kimani E *et al.* (2016) Women's participation in household decision-making and higher dietary diversity: findings from nationally representative data from Ghana. *J Health Popul Nutr* **35**, 16.
28. Asiedu N (2015) Knowledge of chronic disease and food consumption habits among urban poor communities in Accra. Doctoral Dissertation, University of Ghana.
29. Okutu D (2012) Urban household characteristics and implications for food utilization in Accra. Doctoral Dissertation, University of Ghana.
30. Kimani-Murage EW, Schofield L, Wekesah F *et al.* (2014) Vulnerability to food insecurity in urban slums: experiences from Nairobi, Kenya. *J Urban Heal* **91**, 1098–1113.
31. Nkirigacha EM (2012) Contribution of urban agriculture in food and nutrition security to the low income households of Kasarani District of Kenya. MSc Thesis, University of Nairobi.
32. Onyango AC, Walingo MK & Othunon L (2009) Food consumption patterns, diversity of food nutrients and mean nutrient intake in relation to HIV/AIDS status in Kisumu district Kenya. *Afr J AIDS Res* **8**, 359–366.
33. Shell-Duncan B & Yung SA (2004) The maternal depletion transition in northern Kenya: the effects of settlement, development and disparity. *Soc Sci Med* **58**, 2485–2498.
34. Watson FE, Ngesa A, Onyang'o J *et al.* (1996) Fermentation – a traditional anti-diarhoeal practice lost? The use of fermented foods in urban and rural Kenya. *Int J Food Sci Nutr* **47**, 171–179.
35. Nti CA, Inkumsah D & Fleischer G (1999) Influence of women's workload on their nutritional status in selected communities in Ghana. *J Consum Stud Home Econ* **23**, 165–170.
36. Nti CA, Plahar WA & Larweh PM (2002) Impact of adoption in Ghana of an improved fish processing technology on household income, health and nutrition. *Int J Consum Stud* **26**, 102–108.
37. Opere-Obisaw C, Fianu DAG & Awadzi K (2000) Changes in family food habits: the role of migration. *J Consum Stud Home Econ* **24**, 145–149.
38. Quarshie EM (2014) Comparative nutritional status of adult and adolescent mothers and their infants. Doctoral Dissertation, University of Ghana.
39. Tayie FAK (1995) Nutritional habits of pregnant Ghanaian women and effects on pregnancy outcome. Doctoral Dissertation, University of Ghana.
40. Dominguez-Salas P, Alarcón P, Häsler B *et al.* (2016) Nutritional characterisation of low-income households of Nairobi: socioeconomic, livestock and gender considerations and predictors of malnutrition from a cross-sectional survey. *BMC Nutr* **2**, 47.
41. Cornelsen L, Alarcon P, Häsler B *et al.* (2016) Cross-sectional study of drivers of animal-source food consumption in low-income urban areas of Nairobi, Kenya. *BMC Nutr* **2**, 70.
42. Kamau-Mbuthia E & Elmadfa I (2007) Diet quality of pregnant women attending an antenatal clinic in Nakuru, Kenya. *Ann Nutr Metab* **51**, 324–330.
43. Abbey GA (2004) Prevalence of zinc deficiency among Ghanaian adolescents versus food components of zinc and phytate. Doctoral Dissertation, University of Ghana.
44. Agyeman-Nkansah S (2015) Folate and Vitamin B-12 status of adult vegetarians and non-vegetarians in a Christian community in Mayra, Pokuase. Doctoral Dissertation, University of Ghana.
45. Ansong J (2009) Dietary habits, obesity and elevated blood pressure among workers of the college of health sciences. Doctoral Dissertation, University of Ghana.
46. Barichella M, Akpalu A, Cham M *et al.* (2013) Nutritional status and dietary habits in Parkinson's disease patients in Ghana. *Nutrition* **29**, 470–473.
47. Blay D, Aström AN & Haugejorden O (2000) Oral hygiene and sugar consumption among urban and rural adolescents in Ghana. *Community Dent Oral Epidemiol* **28**, 443–450.
48. Frank LK, Kröger J, Schulze MB *et al.* (2014) Dietary patterns in urban Ghana and risk of type 2 diabetes. *Br J Nutr* **112**, 89–98.
49. Galbete C, Nicolaou M, Meeks KA *et al.* (2017) Food consumption, nutrient intake, and dietary patterns in Ghanaian migrants in Europe and their compatriots in Ghana. *Food Nutr Res* **61**, 1341809.
50. Hiamey SE, Amuquandoh FE & Boison GA (2013) Are we indeed what we eat? Street food consumption in the market circle area of Takoradi, Ghana. *Nutr Health* **22**, 215–235.
51. Jolly CM, Awuah RT, Fialor SC *et al.* (2008) Groundnut consumption frequency in Ghana. *Int J Consum Stud* **32**, 675–686.
52. Kerr-Rabbles SL (2015) Oral health and nutritional status of adults attending dental clinics in Korle-Bu. MSc Thesis, University of Ghana.
53. Wu F, Guo Y, Chatterji S *et al.* (2015) Common risk factors for chronic non-communicable diseases among older adults in China, Ghana, Mexico, India, Russia and South Africa: the study on global AGEing and adult health (SAGE) wave 1. *BMC Public Health* **15**, 1–13.
54. Nti CA, Brown A & Danquah A (2012) Adolescents' knowledge of diet-related chronic diseases and dietary practices in Ghana. *Food Nutr Sci* **3**, 1527–1532.
55. Ofori-Koranteng J (2013) Assessment by dietary intake and haemogram of iron levels of vegetarians in a selected community in Accra, Ghana. Doctoral Dissertation, University of Ghana.
56. Sowah LR (2013) Lifestyle, perception of health and blood pressure among Accra's urban poor. Doctoral Dissertation, University of Ghana.
57. Takyi EEK & Amankwa P (2004) Dietary selenium and copper intake by resident undergraduate students of the University of Ghana. *East Afr Med J* **81**, 34–39.
58. Bloomfield G, Mwangi A, Chege P *et al.* (2013) Multiple cardiovascular risk factors in Kenya: evidence from a health and demographic surveillance system using the WHO STEPwise approach to chronic disease risk factor surveillance. *Heart* **99**, 1323–1329.
59. Ethangatta L (1988) Nutritional status of institutionalized elderly people living in urban Kenya, Nairobi. Doctoral Dissertation, University of Nairobi.
60. Mbotela WNC (1999) Dietary patterns and nutritional status of asymptomatic HIV-1 seropositive and HIV -1 seronegative women in Nairobi-Kenya. Doctoral Dissertation, University of Nairobi.
61. Okoth MA (2013) Overweight and obesity in relation to dietary intake and physical activity among adolescents aged 15–19 years attending day secondary schools in Kisumu District. MSc Thesis, Kenyatta University.
62. Van't Riet H, Den Hartog AP, Mwangi AM *et al.* (2001) The role of street foods in the dietary pattern, of two low-income groups in Nairobi. *Eur J Clin Nutr* **55**, 562–570.
63. Theuri G (2006) Differences in cardiovascular disease, biochemical risk markers, physical activity and nutrition between an urban and pastoral sample in Kenya. Doctoral Dissertation, Kenyatta University.



64. Thuita F, Mwadime K & Wang'ombe J (2013) Influence of access to microfinance credit by women on household food consumption patterns in an urban low income setting in Nairobi, Kenya. *Eur J Sci Technol* **2**, 79–88.
65. Wathome DM (1990) The relationship between income and food consumption patterns in urban Nakuru. Doctoral Dissertation, University of Nairobi.
66. Opere-Obisaw C (1998) Students' patronage and views on the operations and services of food vendors. *J Consum Stud Home Econ* **22**, 139–146.
67. Kemunto ML (2013) Dietary diversity and nutritional status of pregnant women aged 15–49 years attending Kapenguria District Hospital West Pokot County. MSc Thesis, Kenyatta University.
68. Mukasa-Mwanthi F (1990) Assessment of selected maternal attributes and food practices in households with malnourished and households with well nourished children below five years in Peri-Urban Nairobi. Doctoral Dissertation, University of Nairobi.
69. Lokko P, Lartey A, Armar-Klimesu M *et al.* (2007) Regular peanut consumption improves plasma lipid levels in healthy Ghanaians. *Int J Food Sci Nutr* **58**, 190–200.
70. Minicuci N, Biritwum RB, Mensah G *et al.* (2014) Sociodemographic and socioeconomic patterns of chronic non-communicable disease among the older adult population in Ghana. *Glob Health Action* **7**, 1–13.
71. Pereko KKA, Setorglo J, Owusu WB *et al.* (2013) Overnutrition and associated factors among adults aged 20 years and above in fishing communities in the urban Cape Coast Metropolis, Ghana. *Public Health Nutr* **16**, 591–595.
72. Van't Riet H, Den Hartog AP & Van Staveren WA (2002) Non-home prepared foods: contribution to energy and nutrient intake of consumers living in two low-income areas in Nairobi. *Public Health Nutr* **5**, 515–522.
73. Mboganie-Mwangi A & Foeken D (1996) Urban agriculture, food security, and nutrition in low-income areas of the city of Nairobi, Kenya. *Afr Urban Q* **11**, 170–179.
74. Steyn NP, Parker W, Nel JH *et al.* (2012) Urbanisation and the nutrition transition: a comparison of diet and weight status of South African and Kenyan women. *Scand J Public Health* **40**, 229–238
75. Mbochi RW (2010) Overweight and obesity prevalence and associated socio-economic factors, physical activity and dietary intake among women in Kibera division, Nairobi. Doctoral Dissertation, Kenyatta University.
76. Drewnowski A & Popkin BM (1997) The nutrition transition: new trends in the global diet. *Nutr Rev* **55**, 31–43.
77. Popkin BM, Adair LS & Ng SW (2012) Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev* **70**, 3–21.
78. Hatloy A, Torheim LE & Oshaug A (1998) Food variety – a good indicator of nutritional adequacy of the diet? A case study from an urban area in Mali, West Africa. *Eur J Clin Nutr* **12**, 891–898.
79. Walls HL, Johnston D, Mazalale J *et al.* (2018) Why we are still failing to measure the nutrition transition. *BMJ Glob Heal* **3**, e000657.
80. World Health Organization (1990) Diet, nutrition and prevention of chronic disease. Report of a WHO Study Group (WHO Technical Report Series 797). Geneva.
81. Vorster HH, Kruger A, Wentzel-Viljoen E *et al.* (2014) Added sugar intake in South Africa: findings from the adult prospective urban and rural epidemiology cohort study. *Am J Clin Nutr* **99**, 1479–1486.
82. Wentzel-Viljoen E, Lee S, Laubscher R *et al.* (2018) Accelerated nutrition transition in the North West province of South Africa: results from the Prospective Urban and Rural Epidemiology (PURE-NWP-SA) cohort study, 2005 to 2010. *Public Health Nutr* **21**, 1–12.
83. Lachat C, Hawwash D, Ocké MC *et al.* (2016) Strengthening the reporting of observational studies in epidemiology—nutritional epidemiology (STROBE-nut): an extension of the STROBE statement. *PLoS Med* **13**, 1–15.
84. Food and Agriculture Organization (2010) *Guidelines for Measuring Household and Individual Dietary Diversity*, pp. 1–60. Rome: Food and Agriculture Organization.
85. Food and Agriculture Organization (2008) *Guidelines for Measuring Household and Individual Dietary Diversity*, pp. 1–29. Rome: Food and Agriculture Organization.