



This is a repository copy of *Invited commentary - WHO classification of tumors : how should tumours be classified? Expert consensus, systematic reviews or both?*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/158602/>

Version: Accepted Version

Article:

Uttley, L. orcid.org/0000-0003-4603-9069, Indave, B.I., Hyde, C. et al. (3 more authors) (2020) *Invited commentary - WHO classification of tumors : how should tumours be classified? Expert consensus, systematic reviews or both?* *International Journal of Cancer*. ISSN 0020-7136

<https://doi.org/10.1002/ijc.32975>

This is the peer reviewed version of the following article: Uttley, L., Indave, B.I., Hyde, C., White, V., Lokuhetty, D. and Cree, I. (2020), *Invited commentary - WHO classification of tumors: How should tumours be classified? Expert consensus, systematic reviews or both?*. *Int. J. Cancer.*, which has been published in final form at <https://doi.org/10.1002/ijc.32975>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Title: Invited commentary - WHO classification of tumors: How should tumours be classified? Expert consensus, systematic reviews or both?

Authors:

Dr Lesley Uttley¹ l.uttley@sheffield.ac.uk
Dr B. Iciar Indave² Indavel@iarc.fr (corresponding author)
Prof Chris Hyde³ c.i.hyde@exeter.ac.uk
Dr Valerie White² WhiteV@iarc.fr
Dr Dilani Lokuhetty² LokuhettyD@visitors.iarc.fr
Dr Ian Cree² Creel@iarc.fr

1. School of Health and Related Research (SchARR), University of Sheffield, Regent Court, 30 Regent Street, Sheffield, S1 4DA, United Kingdom

2. World Health Organisation, Classification of Tumours Group, International Agency for Research on Cancer, 150 Cours Albert Thomas, 69372 Lyon CEDEX 08, France

3. Exeter Test Group, College of Medicine and Health, University of Exeter, Room 1.22, College House, St Lukes Campus, Exeter EX1 2LU, United Kingdom

List of abbreviations:

- Human Herpesvirus-8 (HHV-8)
- International Agency for Research on Cancer (IARC)
- Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA)
- Randomised controlled trials (RCT)
- Standards for Reporting Diagnostic Accuracy Studies (STARD)
- Transparent Reporting of a Multivariable Prediction Model for Individual Prognosis or Diagnosis (TRIPOD)
- World Health Organisation (WHO)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/ijc.32975

Main text:

The World Health Organisation (WHO) Classification of Tumours series, published as the WHO Blue Books (Figure 1) and the accompanying website, are essential resources for pathologists across the globe, providing the standards against which tumours are classified to aid cancer diagnosis, research, treatment and prognosis. Each book tackles the classifications of up to 300 tumour types, defining for each of these the etiology, pathogenesis, epidemiology, clinical features, macroscopic appearances, histology, cytology, molecular pathology, essential and desirable diagnostic features, staging, prognostic factors, and predictive biomarkers. If each tumour were to require one formal review, this would mean finding or conducting up to 3,600 such reviews for each book which would not be practicable or time efficient. The current approach therefore relies largely on subject experts drawing on published literature searches according to their individually perceived need to inform the content of these books. However, these decisions affect the classification, and hence the diagnosis and management of cancer patients worldwide. Consequently, to minimise the risk of including biased information, the evidence is weighed and decisions made by an editorial board. This editorial board is composed of standing members and content experts (mainly practicing pathologists) who meet, propose, revise and agree on definitions and core criteria for each tumour systematically for every new edition.

Over-reliance on expert consensus at editorial board meetings for this purpose may lead to problems (Table 1). Firstly, evidence needs to be reviewed in order to assess all relevant issues in a WHO Blue Book. But, even if all participating experts are required to perform literature reviews, their different expertise and inevitable time constraints produce variable results with potential for studies that are relevant to decisions to be missed. In the first book

of the new 5th edition, up to 130 tumours or subjects were described as “unknown” and 200 labelled as clinically not relevant, this often being the only description of a whole section(1). Sections such as etiology or pathogenesis are frequently summarized as “unknown”, without providing more details, leaving unclear whether an exhaustive search of the literature has been performed. This does not necessary indicate that systematic reviews should be routinely performed when such questions arise and indeed it would not be possible for standing members and content experts to do so given the number that would be required. Often the statements requiring evidence are more related to background information than issues relevant to classification, and introducing systematic approaches to the literature searches would not improve the evidence

base of the review results. In addition, this would allow authors to focus on the pressing or contentious issues of tumour classification: topics that need to be assessed by systematic reviews, due to their potential controversy. Secondly, there is a potential risk that unintended bias from content experts could influence decisions in the absence of a structured and controlled process of evidence synthesis. This could lead to certain studies being unduly highlighted or overlooked depending on who is reviewing the literature and what their own clinical or research knowledge may be(2). Thirdly, if the convened expert panel is not sufficiently representative(3)· skewed decisions may be made, which may not be fully informed by the best and most relevant evidence(4). Fourthly, due to interpersonal and cultural differences among experts during meetings, it is possible for individuals to dominate discussion(3, 5, 6). Finally, previously included but incorrectly referenced evidence may be carried forward during the updating process .

It is possible that these problems could be mitigated by the addition of evidence-based practices to the editorial process. Systematic reviews are widely regarded as the cornerstone of evidence-based medicine, encouraging comprehensive literature searching, transparency in methods and rigorous study appraisal(7). Performing systematic reviews for some tumour types could certainly improve the reliability of decisions taken by the editorial board, but they may not always be the best solution.

Systematic reviews require training in methods, as well as expertise (8-10). They can be laborious to complete, time-consuming and difficult to interpret, particularly when there is lot of evidence, or when the evidence is low-grade. Systematic review methods have been described as being formulaic(11, 12). Best practice guidelines such as Cochrane(13) and PRISMA(14) are closely aligned with meta-analytic reviews for medical interventions, the methods of which are not necessarily appropriate to pathology. Systematic reviews can be long and unwieldy documents, difficult to interpret and hollow in their conclusions(15, 16). The hierarchy of evidence-based medicine with systematic reviews at the apex has been resisted by some academic groups and clinicians for oversimplifying complex issues relating to clinical care(17). Some argue that clinical judgement based on experience is more exact and applies better to real life clinical situations, because of its emphasis on individual cases rather than evidence derived from randomised controlled trials (RCT) that are produced without social and political confounding effects and of other variables(18). Whilst the scientific rigour of systematic reviews and indeed RCTs are preserved through adhering to reporting guidelines, they have been shown to be less readable than article types which allow some spin or journalistic creativity, such as editorials and narrative reviews(19). They are also subject to influence if not conducted objectively and not properly assessed prior to publication(20).

Managing the influence and expertise of team members in systematic reviews, like non-systematic reviews, remains critical to producing reliable and externally valid conclusions(10).

Although essential for the synthesis of available scientific evidence in controversial questions, it is clear that systematic review methods would need to be adapted to the specific needs of the WHO Classification of Tumors and its editorial process. Timely procurement of a valid evaluation of evidence is one of the major problems for process-driven systems such as the WHO Classification of Tumors, and this is something that traditional, comprehensive systematic reviews struggle to provide. More succinct review methods including, mapping, rapid or scoping reviews could improve the quality of scientific evidence used to make statements in the WHO Blue Books, without the need for full systematic reviews(21-23). While alternative methods that abbreviate systematic reviews may be appealing, they do increase the risk of bias and errors in the review process and may not always provide the expected results(21, 22, 24). Careful assessment of the need for evidence synthesis may show that a structured expert description of the subject is enough. We believe that full systematic reviews of the available evidence should be limited to the assessment of research questions that directly affect the classification of a tumour, or have major consequences in cancer diagnosis.

Nowhere is the need to consider consensus-based versus evidence-based approaches more apparent than in the field of molecular pathology, where research groups may suggest different molecular tumor classifications for the same tumor type. In these instances, evidence synthesis and evaluations to underpin incorporation of molecular advances into the classification will be greatly assisted by the use of the Standards for Reporting Diagnostic Accuracy Studies (STARD)(25), Transparent Reporting of a Multivariable Prediction Model for

Individual Prognosis or Diagnosis (TRIPOD)(26) and other standardised reporting guidance (www.equator.net).

Herein lies the challenge for the WHO Classification of Tumours, currently published as a series of 15 WHO Blue Books, and now as a website. Should it continue to rely primarily on experts without formal systematic review methods, or should it take steps to incorporate systematic reviews and/or adapted methods into the process of using expert consensus? The importance of systematic reviews and expert interpretation do not need to be viewed as mutually exclusive. Mickenautsch argues that to view one as superior to another is erroneous as each represent necessary concepts of analysis and synthesis(27). Indeed, there is an opportunity to ensure that reviews are fit for purpose without enforcing rigid guidelines which can result in review products which are overly detailed and unwieldy to interpret(15). We propose an approach that is adapted to the needs of WHO Blue Books which requires authors to employ a series of 'non-negotiables' when performing the literature reviews that feed into key decisions for the WHO Blue Book series. They are:

1. **Transparency:** clearly stated, pre-specified methods are critical and should be described in detail to assure reproducibility and reliability of the process. A publicly available protocol setting out the intended methods and eligibility criteria would help receiving external input into the review and facilitate updating.
2. **Searching rigor:** at least two relevant bibliographic sources should be searched for relevant literature

3. Double checking: a proportion of randomly selected study selection and data extraction decisions should be double-checked by a second reviewer to verify reliability and consistency.
4. Risk of bias assessment: included individual studies should be assessed using standardised or adapted tools that consider areas of bias to which studies of pathology may be prone. Assessment of the body of the collected evidence may be useful, ranking it using validated criteria.

Detail on the rationale and risk underlying each ‘non-negotiable’ are outlined in Table 2. The issue of using validated methods for WHO classification of tumours is becoming increasingly important because, due to rapid advances in molecular pathology, more information is becoming evident from closer analysis of tumours than was previously available from histopathological analysis alone. Systematic review methods may need to be adapted to assess this information, as some steps of a validated systematic review process may not meet current need, or simply not add sufficient value to the editorial process.

Not every decision made in the WHO Classification of Tumours is contentious; many areas may be regarded as close to fact due to strong logic or pathophysiological basis: they can be considered as equivalent of parachutes – known to be efficacious without the need to conduct a RCT(28). However, when issues are debatable or likely to vary depending on context or interpretation then a systematic review may be deemed necessary. An example of this occurs with Kaposi’s sarcoma where there is some debate over whether the characteristic tumours can reliably be regarded as neoplasms or as a “reactive” proliferations of cells in response to

infections with Human Herpesvirus-8 (HHV-8)(29). Cancerous tumours are commonly regarded to be clonal neoplasms which proliferate uncontrollably, even after the stimulus which evoked the change has been removed(30), Kaposi sarcoma is contentious because there are cases where it does not persist following successful antiretroviral treatment(31). However, reclassification of Kaposi sarcoma could have ramifications for patient diagnosis and care, as well as service delivery and funding for research. In response to this dilemma, a systematic review protocol was designed and registered prospectively by the WHO tumour classification group on the PROSPERO database (CRD42018087595). The review aims to inform and support the decision of the expert panel regarding Kaposi Sarcoma classification using available evidence without restriction by study type.

There are challenges for the classification of tumours in evidence-based pathology. Differing evidence levels and considerations are applied within the field of pathology and consensus is needed to promote a comparable assessment of studies. Review methods should allow reliable assessments of evidence in an appropriate time frame and without being systematic review methodologists. Experts who serve as authors and editors require basic training and methodological support in systematic review and literature searching methods. Furthermore, subjects in need of assessment should be registered and prioritized based on WHO Classification of Tumours requirements and scientific considerations. The WHO Classification of Tumours Group at the International Agency for Research on Cancer (IARC) have set themselves the challenge to revise their procedures to incorporate systematic review methods into the current expert-led approach through the appointment of a systematic reviewer for the Blue Books and by planning an international Joint Action Group. The methods described here aim to start this deliberation and we invite discussion and feedback as to the potential

challenges and solutions to the intended “non-negotiables” proposed to ensure the methods for reviewing literature which feed into tumour classification are robust.

In conclusion, we believe that an evidence-based approach to informing key decisions that feed into tumour classification would allow editors to mitigate any potential risks of bias and also benefit authors by providing structured, transparent and reliable methods for the synthesis of available evidence for each tumour type. Our hope over time is that this approach will increase the rigour of the decisions feeding into WHO tumour classification, by addressing critical questions and identifying research gaps, as well as reaching recommendations for research to inform future editions. Such an approach will help maintain the reliability of tumour classification, and help to provide solutions to challenges like the exponential rise in number of scientific publications and the need to manage new types of information such as evidence from genetics or big data.

Disclaimer

The content of this article represents the personal views of the authors and does not represent the views of the authors' employers and associated institutions. Where authors are identified as personnel of the International Agency for Research on Cancer / World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer / World Health Organization.

Conflict of interest: Dr Lesley Uttley declares that she undertook two paid visiting scientist trips to the International Agency for Research on Cancer to plan and develop this work in July/August 2018. The other authors declare that they have no conflict of interest.

Accepted Article

References

1. WHO Classification of Tumours. Digestive System Tumours. . 5th ed: IARC Publications; 2019.
2. Young SN. Bias in the research literature and conflict of interest: an issue for publishers, editors, reviewers and authors, and it is not just about the money. *Journal of psychiatry & neuroscience* : JPN. 2009;34(6):412-7.
3. McKee M, Priest P, Ginzler M, Black N. How representative are members of expert panels? *Quality assurance in health care : the official journal of the International Society for Quality Assurance in Health Care*. 1991;3(2):89-94.
4. Doust J, Del Mar C. Why do doctors use treatments that do not work? *BMJ (Clinical research ed)*. 2004;328(7438):474-5.
5. Kea B, Sun BC. Consensus development for healthcare professionals. *Internal and emergency medicine*. 2015;10(3):373-83.
6. Tajeddin G, Safayeni F, Connelly CE, Tasa K. The Influence of Emergent Expertise on Group Decision Processes. *Small Group Research*. 2011;43(1):50-74.
7. Mulrow CD. Rationale for systematic reviews. *BMJ (Clinical research ed)*. 1994;309(6954):597-9.
8. Gotsche PC, Ioannidis JP. Content area experts as authors: helpful or harmful for systematic reviews and meta-analyses? *BMJ (Clinical research ed)*. 2012;345:e7031.
9. Piehl JH, Green S, McDonald S. Converting systematic reviews to Cochrane format: a cross-sectional survey of Australian authors of systematic reviews. *BMC health services research*. 2003;3(1):2.
10. Uttley L, Montgomery P. The influence of the team in conducting a systematic review. *Systematic reviews*. 2017;6(1):149.
11. Collins JA, Fauser BC. Balancing the strengths of systematic and narrative reviews. *Human reproduction update*. 2005;11(2):103-4.
12. MacLure K, Paudyal V, Stewart D. Reviewing the literature, how systematic is systematic? *International journal of clinical pharmacy*. 2016;38(3):685-94.
13. Higgins JP GS. *Cochrane handbook for systematic reviews of interventions*. . West Sussex PO19 8SQ, England: The Cochrane Collaboration and John Wiley & Sons Ltd; 2008.
14. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine*. 2009;6(7):e1000097.
15. Greenhalgh T, Thorne S, Malterud K. Time to challenge the spurious hierarchy of systematic over narrative reviews? *European journal of clinical investigation*. 2018;48(6):e12931.
16. Newman TB. The power of stories over statistics. *BMJ (Clinical research ed)*. 2003;327(7429):1424-7.
17. Williams DD, Garner J. The case against "the evidence": a different perspective on evidence-based medicine. *The British journal of psychiatry : the journal of mental science*. 2002;180:8-12.
18. Parker M. False dichotomies: EBM, clinical freedom, and the art of medicine. *Medical humanities*. 2005;31(1):23-30.
19. Loke YK, Derry S. Does anybody read "evidence-based" articles? *BMC medical research methodology*. 2003;3:14.

20. Ioannidis JP. The Mass Production of Redundant, Misleading, and Conflicted Systematic Reviews and Meta-analyses. *The Milbank quarterly*. 2016;94(3):485-514.
21. Dobbins M. Rapid Review Guidebook. Steps for conducting a rapid review. [Guideline]. Canada: The National Collaborating Centre of Methods and Tools (NCCMT) 2017. Available from:
<https://www.nccmt.ca/uploads/media/media/0001/01/a816af720e4d587e13da6bb307df8c907a5dff9a.pdf>.
22. Khangura S, Konnyu K, Cushman R, Grimshaw J, Moher D. Evidence summaries: the evolution of a rapid review approach. *Systematic reviews*. 2012;1:10.
23. Polisena J, Garritty C, Umscheid CA, Kamel C, Samra K, Smith J, et al. Rapid Review Summit: an overview and initiation of a research agenda. *Systematic reviews*. 2015;4:111.
24. Haby MM, Chapman E, Clark R, Barreto J, Reveiz L, Lavis JN. What are the best methodologies for rapid reviews of the research evidence for evidence-informed decision making in health policy and practice: a rapid review. *Health research policy and systems*. 2016;14(1):83.
25. Cohen JF, Korevaar DA, Altman DG, Bruns DE, Gatsonis CA, Hooft L, et al. STARD 2015 guidelines for reporting diagnostic accuracy studies: explanation and elaboration. *BMJ open*. 2016;6(11):e012799.
26. Moons KG, Altman DG, Reitsma JB, Ioannidis JP, Macaskill P, Steyerberg EW, et al. Transparent Reporting of a multivariable prediction model for Individual Prognosis or Diagnosis (TRIPOD): explanation and elaboration. *Annals of internal medicine*. 2015;162(1):W1-73.
27. Mickenautsch S. Systematic reviews, systematic error and the acquisition of clinical knowledge. *BMC medical research methodology*. 2010;10:53.
28. Smith GC, Pell JP. Parachute use to prevent death and major trauma related to gravitational challenge: systematic review of randomised controlled trials. *BMJ (Clinical research ed)*. 2003;327(7429):1459-61.
29. Brooks JJ. Kaposi's sarcoma: a reversible hyperplasia. *Lancet (London, England)*. 1986;2(8519):1309-11.
30. Willis RA. Pathology of Tumours. By R. A. Willis, D.Sc., M.D., F.R.C.P., and Sir William H. Collins Professor of Human and Comparative Pathology, Royal College of Surgeons, London. 95/8 x 65/8 in. Pp. 1044 + xxiii, with 500 illustrations. 1948. London : Butterworth & Co. (Publishers) Ltd. 63s. net. *BJO*. 1948;35(140):446-.
31. Chang Y, Cesarman E, Pessin MS, Lee F, Culpepper J, Knowles DM, et al. Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science (New York, NY)*. 1994;266(5192):1865-9.

Table 1: Problem arising from consensus based approach and proposed solutions by an evidence-based approach

Problem of a consensus based approach	Solution by an evidence-based approach	Potential problems not solved by an evidence-based approach
Risk of missing relevant research	<p>Comprehensive searching which is part of systematic reviewing may improve identification of important literature.</p> <p>A structured, systematic process allows summarizing and evaluating complex information such as big data or basic research information provided for molecular pathology.</p>	<p>Risk of missing research which does not fit into standard study design framework used for systematic reviews.</p> <p>Publication bias may not be addressed if only searching published evidence.</p>
Selection of the literature may be biased	<p>Systematic reviews require clearly stated inclusion criteria, so cherry picking of particular studies to prove a particular point is easier to spot.</p> <p>In addition, the setting of acceptable evidence levels and assessment of risk of bias of studies avoids the use of inappropriate evidence.</p>	Presentation of results may still allow a certain degree of “cherry picking” when presenting only on selected outcomes (Outcome reporting bias).
Interpretation of the literature may be biased	<p>Systematic reviews consider each included study equally, unless there is a specific reason why less emphasis should be placed on it such as small sample or poor study quality.</p> <p>Risk of bias assessment of individual studies, but also of the body of evidence can be undertaken to aid an appropriate interpretation of the retrieved evidence.</p>	The use of several reviewers may not provide the desired control of bias effect and instead interesting information, may not be incorporated due to disagreement.
Panel of experts may be biased in composition or be dominated by particular individuals	A systematic review with clear eligibility criteria made available in a protocol may provide a reference point against which “extreme” views by particular panel members can be mitigated.	Panel may still be biased in developing eligibility criteria, even if an evidence-based approach helps in the discussion.
Difficulties in documentation of included evidence (especially in the updating process)	Systematic review protocol and reports document the biomedical databases searched and over what time period therefore uncertainty about whether a particular study has been included or not is much less likely to occur.	Relies on the well-designed and appropriate literature searches and implementation of reporting standards for systematic reviews.
Credibility of classification may be undermined if not evidence-based	Use of systematic review methods will improve the credibility of the classification, as well as the reliability of tumour classification.	<p>Credibility may also be affected by other factors not addressed by a systematic review process.</p> <p>Experts in the field, important to the credibility of the books, may be put-off by the systematic review process.</p>


Table 2. Methodological non-negotiables for systematic reviews for the purposes of tumour classification

Rationale	Risks (if not considered in the review)
1. Transparency	
<p>Methods should be clearly stated and previously defined. Inclusion and exclusion criteria stated and applied.</p> <p>A review protocol should be written and made publicly available as an explicit statement of intended methods where deviations to these methods can be noted (with justifications). This ensures accountability by authors and facilitates replication of the review.</p> <p>Conflict of interests of the review team, as well as funding information needs to be disclosed.</p>	<ul style="list-style-type: none"> • Methods may not be appropriate to ensure equitable representation of literature globally • Unjustified deviations to planned methods remain unchallenged. • Undeclared conflicts of interest or researcher allegiance from authors may influence conclusions
2. Searching rigor	
<p>Searching two major bibliographic databases, (e.g. PubMed and Web of Science), minimises the chance that a highly relevant study will be missed. Whilst there are overlaps in medical bibliographic databases, indexing varies considerably. Therefore searching only one database means that retrieval of relevant literature is highly dependent on appropriateness/accuracy of the search strategy.</p>	<ul style="list-style-type: none"> • Reliance on one database capturing all relevant studies, reliance on all relevant studies being accurately indexed and reliance on a single search strategy being sufficient to capture all relevant literature • Failure to identify all relevant literature
3. Double checking	
<p>Duplication of the data extraction and a proportion of the total study selection done by the primary author should be completed by a second reviewer for accuracy. Where multiple discrepancies are noted, further checking may be required for consistency.</p>	<ul style="list-style-type: none"> • Reliance on the accuracy and consistency of one author for all study selection and data extraction • Bias in selection of studies • Greater chance of erroneous study selection or data extraction
4. Risk of bias assessment	
<p>A methodological quality assessment tool for pathology reviews should be adapted, based on standardized risk of bias assessment tool. This helps review authors to assign more weight to findings from studies of higher quality or at lower risk of bias in interpretation.</p>	<ul style="list-style-type: none"> • No objective method of appraising studies for higher risk of bias • Biases from primary studies are perpetuated in the review • Bias in interpretation of studies may be applied by review authors

WHO Classification of Tumours • 5th Edition

Breast Tumours

Edited by the WHO Classification of Tumours Editorial Board



The World Health Organisation (WHO) Classification of Tumours series. Breast Tumours. Fifth Edition.